

Clustering of Allergic Outcomes within Families and Households in Areas Endemic for Helminth Infections

Sitti Wahyuni^{a, c} Erliyani Sartono^c Taniawati Supali^b Jaring S. van der Zee^e
Andarias Mangali^a Ronald van Ree^f Jeanine J. Houwing-Duistermaat^d
Maria Yazdanbakhsh^c

^aDepartment of Parasitology, Hasanuddin University, Makassar, and ^bDepartment of Parasitology, University of Indonesia, Jakarta, Indonesia; ^cDepartment of Parasitology and ^dDepartment of Medical Statistics and Bioinformatics, Leiden University Medical Center, Leiden, and ^eDepartment of Pulmonology, Academic Medical Center and ^fDepartment of Immunopathology, Sanquin Research, Amsterdam, The Netherlands

Key Words

Helminth · IgE · Skin prick test · Clustering of allergy

Abstract

Background: Allergy and helminth infections share key immunological features in terms of Th2 responses. Although in industrialized countries clustering of allergic disorders within families has been frequently reported, such information is lacking from areas where helminth infections are endemic. **Methods:** A total of 466 subjects from 29 families and 112 households participated in this study. Filarial infection, skin test reactivity and IgE to mite as well as total IgE were measured in all samples. Clustering of the allergy-related outcomes due to genetic and household factors was tested. **Results:** Genetic factors contributed significantly to the clustering of total IgE and allergen-specific IgE, whereas only household factors contributed to the clustering of SPT positivity. **Conclusion:** Similar to several studies conducted in western populations, total IgE and allergen-specific IgE are influenced by genetic factors in a population resident in a helminth endemic area. However, clustering of SPT positivity due to genetic factors was not significant in the

current study raising the question of whether the presence of helminth infections may override genes that are associated with the expression of tissue reactivity to allergens in the west.

Copyright © 2005 S. Karger AG, Basel

Introduction

Like in industrialized countries, the prevalence of allergy in developing regions of the world, such as in Asia, is also increasing. A study conducted in Thailand reported a high prevalence of asthma and other allergic diseases (wheezing, rhinitis and eczema) among children and indicated that symptoms have been increasing 4-fold over the last decade [1]. In Malaysia, asthma and related symptoms were common among schoolchildren in inner city Kuala Lumpur, where 10% of the children reported to have ever had asthma [2]. However, there is an important difference between urban and rural populations within developing countries with respect to the prevalence of allergic disorders. For example in the West Bank the prevalence of wheeze and asthma was higher in urban than in rural areas [3]; a significantly greater prevalence

of allergic symptoms was found in urban compared with rural children in Saudi Arabia [4]; higher prevalence of exercise-induced bronchospasm was reported in urban compared to rural children in India [5], and several African studies have indicated significant differences in allergic disorders when comparing urban with rural communities [6–9].

Although there is strong evidence that genetic factors influence allergic disorders, this alone cannot explain the increase that is being observed worldwide. However, the gene by environment interactions could play an important role in changing disease patterns in continuously evolving environments [10, 11]. It is therefore important to investigate how genetic factors modify allergic outcomes in varying environments where the expression of allergic disorders is widely different.

In western populations, genetic factors have been shown to influence allergic disorders. Not only asthma [12] but also related phenotypes such as total IgE [13, 14], allergen-specific IgE [15, 16], skin prick test (SPT) positivity [15, 16] and bronchial hyperresponsiveness [17, 18] have been shown to be influenced by several genes. Among environmental factors which may influence allergic disorders, increasing air pollution [19, 20], rise in indoor allergen exposure [21], altered diet [22, 23] and changes in exposure to microbes [24–26] have been studied intensively.

Recently, few studies have indicated that exposure to helminth infections which are often endemic in rural areas of the developing world may influence allergic outcomes, positively and negatively [27, 28]. Helminth infections like allergic diseases are characterized by elevated levels of IgE, tissue eosinophilia, mastocytosis, and CD4⁺ T cells that preferentially secrete the Th2 cytokines [29]. However, chronic helminth infections are also associated with immune suppression, primarily to parasite antigens, but spillover suppression has also been described [30, 31]. This suppression may play an important role for any inverse relationship observed between helminth infections and allergic disorders [26].

Th2 responses have been shown to be under genetic control as reviewed by Wills-Karp and Ewart [12]. Several chromosomal regions, for instance chromosome 4q, 6, 11q and 13q, showed evidence for linkage to asthma, atopy, total IgE, SPT, blood eosinophils and bronchial hyperresponsiveness; chromosomes 2, 6, 9 and 12 showed linkage to asthma, specific IgE, and total IgE, and chromosomes 1p31, 7q, 11p13, 11q13, 12q24, 13q31, 17q12-21 and 19q13 showed linkage to asthma, bronchial hyperresponsiveness, SPT, total IgE, blood eosinophils, and age

at onset of asthma. Thus specific chromosomal regions appear to be important in susceptibility to different phenotypes of allergic responsiveness.

In helminth infections, genetic factors play an important role in controlling the outcome of an infection. A genome scan for human schistosomiasis carried out in Brazil identified chromosome 5q31-q33 as a locus responsible for controlling the intensity of *Schistosoma mansoni* infection [32] and there is also evidence for genetic control of pathology by a region containing the gene for the interferon- γ receptor 1 subunit in this disease [33]. For nematodes, there is also evidence for genetic control of filarial infection [34, 35] and a recent study conducted in Nepal has found a locus controlling *Ascaris lumbricoides* intensity on chromosomes 1 and 13 [36].

To start understanding the relationship between helminths and allergic disorders at the genetic level, it is important to investigate the behavior of allergic disorders in populations residing in areas where helminths are endemic. As a first step towards this, we have characterized the clustering of allergic phenotypes and – using a score statistics model – have dissected the contribution of genetic and nongenetic factors to the clustering.

Material and Methods

Study Design

The study population consisted of life-long residents in Salubarana, Karossa district, South Sulawesi, Indonesia, an area where *Brugia malayi* is endemic. The village is located 3 km from the seaside and most of the houses are situated along the main road and spread over a distance of approximately 1.2 km. The major economic activity of the population is subsistence farming, which mainly involves the growing of cacao and corn. In cooperation with the head of the village and the health workers of the local District Health Center, all inhabitants were informed about the study and invited to participate. Informed consent was obtained from all participants before skin prick testing, clinical and parasitological examination and blood withdrawal in accordance with the guidelines of the Indonesian Department of Health and Human Services.

The households were visited and information was obtained about the members of the household: name, age, sex and how they are related to each other. To maximize information of the link between individuals, multichecks were made before constructing pedigrees. Information on children was obtained from parents or from adults who lived with the children whenever both mother and father were not available.

SPT reactivity for aeroallergens was tested with extract of *Dermatophagoides pteronyssinus*, *Dactylis glomerata* grass pollen, cat dander and dog dander preparation (HAL Allergen Laboratories, The Netherlands). Histamine chloride (10 mg/ml) was used as the positive control and allergen diluent as the negative control (HAL Allergen Laboratories, The Netherlands). SPT was done on the volar side of the lower arm, using separate 25-gauge needles. The re-

sults were measured after 15 min. Skin prick reactivity was determined to be positive if the diameter of erythema was ≥ 10 mm [37–39]. The same investigator performed all the SPTs.

Microfilariae were enumerated by filtration of 1 ml nocturnally collected blood through a Millipore, fixed with methanol, stained with Giemsa's stain and examined microscopically.

Plasma for immunological studies was stored (immediately after separation) at -20°C for several months and subsequently transported to The Netherlands on dry ice, where it was stored at -70°C until use.

Parasite Antigen

Adult *B. malayi* worms were purchased from TRS Laboratories, Athens, Ga., USA. Hundred female worms were freeze-dried, ground to powder on ice, dissolved in phosphate buffer saline (PBS) and slowly stirred at 4°C . The protein concentration was determined by the 2,2'-biquinoline-4,4'-dicarboxylic acid disodium salt hydrate (BCA) method before storage at -20°C [40].

Total IgE Antibodies

Total IgE antibodies were measured by an enzyme-linked immunosorbent assay (ELISA) in serum samples as described previously [41]. Briefly, Maxisorp plates (Nunc, Roskilde, Denmark) were coated at 4°C with 100 μl rabbit anti-human IgE (Dako, Denmark; 1:1,000) in 0.1 M bicarbonate buffer (pH 9.6), and blocked with PBS/2% bovine serum albumin (BSA) at room temperature. After washing with PBS 5% and 0.1% Tween 20, 100 μl of sera diluted in Tris 0.1 M pH 7.5 containing 0.05% Tween 20 (1:200) were added to the wells. As a reference, the WHO standard serum of human IgE (National Institute for Biological Standards and Control, Hertfordshire, UK) was used, applying a 9-fold serial dilution (1:2), starting at a concentration of 50 IU/ml. Plates were incubated for 1 h at room temperature, and after washing goat anti-human IgE-biotinylated antibody (1:1,000; Vector Burlingame, Calif., USA) was added, followed by incubation with streptavidine-alkaline phosphate conjugate (1:3,000; Dako, Glostrup, Denmark) for 1 h at room temperature. Finally 100 μl 1 mg/ml *p*-nitrophenyl phosphatase (Boehringer, Mannheim, Germany) in 0.1 M diethanolamine was added and the reaction was stopped after 20 min by adding 100 μl 3 M NaOH. Absorbance was measured at 405 nm. One international unit is 2.4 ng IgE.

Allergen-Specific IgE

Serum level of mite-, dog-, cat-, grass pollen-specific IgE were determined by radioallergosorbent test as described previously [42]. Briefly, 50 μl serum was incubated overnight with 1.5 mg of Sepharose-coupled allergen in a final volume of 300 μl PBS, 3% BSA, 0.1% Tween 20. After washing away nonbound serum components, radiolabelled sheep antibodies (Sanquin, Amsterdam, The Netherlands) directed to human IgE were added. After overnight incubation and washing, bound radioactivity was measured. The outcomes were expressed as percent binding. To convert these values into IU/ml, the results were plotted to a nonlinear regression curve of chimeric monoclonal IgE antibody dilution series against the major house dust mite allergen Der p 2 and Sepharose-coupled mite extracts [43]. The cutoff of the assay was 0.3 IU/ml and subjects were considered to have high levels of specific IgE when the value was more than 1.0 IU/ml.

ELISA for Detection of Antifilarial Specific IgG4

The levels of antifilarial IgG4 were determined in ELISA as described elsewhere [44]. The optical density of patients' plasma was converted into arbitrary units (AU) by drawing a standard curve of a donor positive for filarial infections from Central Sulawesi who was given a value of 10^6 AU/ml. Antifilarial IgG4 is a valuable diagnostic tool which detects filarial infection not detectable by standard parasitological examination (blood microfilaria). The antifilarial IgG4 drops after treatment and it might detect not only the presence of adult worm and microfilaria but also the earlier stages of infective larvae [45, 46]. The test when using crude filarial antigen extracts has a 100% sensitivity and 95% specificity [47].

Statistical Analysis

Standard statistical analyses were performed in SPSS for windows version 10. More advanced analyses were performed in S-PLUS 2000 for windows. A log transformation was used to obtain normally distributed data; specific IgG4 level, total IgE and mite-specific IgE plus one were transformed to log 10. To derive a threshold between high and low levels of total IgE antibodies, the frequency distribution of this antibody was plotted. Based on this distribution, the cutoff of total IgE was set at 1,400 IU/ml.

Although the presence of microfilariae is a gold standard to assess infection status in individuals, it requires a minimum threshold to visualize in blood and hence fails to establish low parasitemia and cryptic infections. Since measurement of IgG4 is more sensitive and IgG4 levels were shown to be associated with microfilaremia [45, 46], a threshold for the infection status was estimated from the antifilarial IgG4 levels. Briefly, it was assumed that the bimodal distribution of antifilarial IgG4 levels is a mixture of two normal distributions, one representing the uninfected and one representing the infected subjects. The means of these two distributions were estimated using the program k-means in S-PLUS for Windows version 2000. The value of 4.50 AU/ml (log 10) which was just in between the two means was used as a threshold to define infected and uninfected subjects. This threshold was higher than 4.20 AU/ml (log 10) when using 20 Dutch blood donors plus three standard deviations as a cutoff.

For females and males and for various age strata, the prevalence of microfilariae, IgG4 >4.5 AU/ml, mite SPT, mite IgE >0.3 IU/ml and mite IgE >1 IU/ml were estimated and compared using Pearson χ^2 statistic. For the continuous outcomes antifilarial IgG4 level, mite IgE and total IgE level, the mean and standard deviations were computed for the females and males and for the various age strata. The means were compared using ANOVA. Next, a regression model was fitted to assess the relation between age and gender and the outcomes. We verified whether the interaction between age and gender existed.

For filarial-infected and uninfected subjects, the prevalence of mite SPT positivity, of mite IgE >0.3 IU/ml, of mite IgE >1 IU/ml, of total IgE $>1,400$ IU/ml and the mean of mite IgE and total IgE were estimated and compared by using Pearson χ^2 statistic for the binary outcomes and ANOVA for the quantitative outcomes. The relation between total IgE and specific IgE and skin prick reactivity was studied using the logistic regression analysis. Here adjustments for age and gender were made.

Genetic and household clustering of filarial infection were tested by a statistic test of Houwing-Duistermaat et al. [48] using S-PLUS 2000 for windows. This has been applied to test clustering of filarial infection due to genetic and household effects [34, 35,

Table 1. Measurements as a function of gender

	Female (n = 225)	Male (n = 241)	p
Microfilaria prevalence, %	5.8	11.2	0.037
Filarial infection prevalence, %	39.1	67.2	<0.001
Mite SPT prevalence, %	7.1	4.1	0.24
Mite IgE >0.3 IU/ml prevalence, %	56.0	66.4	0.021
Mite IgE >1.0 IU/ml prevalence, %	21.3	33.6	0.003
Total IgE >1,400 IU/ml prevalence, %	28.2	34.8	0.001
Antifilarial IgG4 level, AU/ml	4.24 (3.40–5.16)	4.90 (4.22–5.77)	<0.001
Mite IgE level, IU/ml	0.20 (0.05–0.27)	0.27 (0.09–0.38)	<0.001
Total IgE level, IU/ml	3.30 (2.97–3.72)	3.53 (3.20–3.94)	<0.001

Prevalence of microfilariae, filarial infection (determined by antifilarial IgG4), skin test reactivity to mite (mite SPT), IgE to mite (mite IgE; low level sensitization: >0.3, <1.0 IU/ml, and high sensitization: >1.0 IU/ml), high level of total IgE (>1,400 IU/ml) and the levels of antifilarial IgG4, mite IgE and total IgE are indicated for 466 subjects. p values were measured using Pearson χ^2 for comparing prevalences and one-way ANOVA for comparing antifilarial IgG4, specific and total IgE levels.

48]. Briefly, the genetic and the household correlation structures were based on distances between individuals. The correlation between genetic effects of two individuals equals 2^{-dg} in which dg is the genetic distance in the pedigree. Thus parents and offspring have dg = 1, pairs of siblings have dg = 1, grandparents and grandchildren have dg = 2, aunts/uncles and nephews have dg = 2, cousins have dg = 3, great-grandparents and great-grandchildren have dg = 3, great-uncles/aunts and nephews have dg = 3, and grand-nephews can have dg = 4 or dg = 5 depending on the degree of kinship. Individuals who are not related have infinity as the distance. In the data the distances between relatives varied from 1 to 5. The distances for the household model were defined by dh, with dh = 0 for individuals sharing the same house (correlation 1) and distance infinity for inhabitants of different households (correlation 0). The hypotheses of no correlation due to genetic or household effects were tested versus the alternatives using the test statistic of Houwing-Duistermaat et al. [48]. The allergy-related parameters were adjusted for age and sex.

Results

Study Population: Infection and Allergy

A total of 466 participants from Salubarana aged 5–84 years (mean 28.8 years) participated in this study and the number of females and males in this village was comparable (225 females and 241 males). Filarial infection was determined by conventional blood filtration technique and by specific antifilarial IgG4 as an indicator of active infection. Of the 466 subjects screened 8.6% were microfilaremic whereas 53.15% were found positive for antifilarial IgG4. As already described in Materials and Methods, this test is highly sensitive (100%) with a 95% specificity [47].

In the total population skin prick reactivity was 11.6% and predominantly found against the house dust mite allergen (6.0%). The rest of the reactivity was to grass pollen, cat and dog (4.2, 1.5 and 2.1%, respectively), and all participants showed a positive response to the histamine control.

Effect of Gender and Age on Infection and Allergy

Tables 1 and 2 show the prevalence of microfilaremia, active infection status (detected by antifilarial IgG4), mite SPT and the level of mite IgE (>0.3 and >1 IU/ml) as well as the means of antifilarial IgG4, mite IgE level and total IgE levels as a function of gender and age. In the total population it was shown that microfilaria prevalence and antifilarial IgG4 were higher in males than in females and increased significantly with age. For allergy phenotypes, both the levels and the prevalence of specific mite IgE (using cutoff points 0.3 or 1 IU/ml) as well as total IgE were higher in males compared to females but no difference was found in mite SPT reactivity. When age was divided into four groups, as shown in table 2, the prevalence of mite SPT as well as levels of mite IgE reached a peak in the age group of 15–29 years and then decreased with increasing age. The levels of total IgE also decreased with increasing age. From scatter plots of age versus mite IgE and age versus total IgE (data not shown), it appeared that the relation between age and the levels was best described by using the natural logarithm of age (lnage). The lnage has also been used when analyzing the clustering of filarial infection [35]. Therefore in all further analyses we used lnage as a covariate. Using regression analysis we

Table 2. Measurements as a function of age

	Age ≤ 14 (n = 122)	Age 15–29 (n = 138)	Age 30–44 (n = 111)	Age >44 (n = 95)	< p
Microfilaria prevalence, %	4.1	5.1	11.7	15.8	0.005
Filarial infection prevalence, %	32	55.1	59.5	72.6	<0.001
Mite SPT prevalence, %	6.6	7.2	5.4	3.2	0.59
Mite IgE >0.3 IU/ml prevalence, %	58.2	65.9	62.2	57.9	0.52
Mite IgE >1.0 IU/ml prevalence, %	23.0	35.5	29.7	20	0.035
Total IgE >1,400 IU/ml prevalence, %	30.8	38.3	44.1	48.1	0.32
Antifilarial IgG4 level, AU/ml	4.15 (3.31–5.07)	4.50 (3.82–5.42)	4.80 (4.14–5.75)	5.00 (4.46–5.67)	<0.001
Mite IgE level, IU/ml	0.22 (0.06–0.28)	0.27 (0.06–0.40)	0.24 (0.07–0.37)	0.19 (0.07–0.26)	0.046
Total IgE level, IU/ml	3.54 (3.18–3.94)	3.44 (3.09–3.92)	3.36 (3.05–3.78)	3.29 (2.94–3.77)	0.013

Prevalence of microfilariae, filarial infection (determined by antifilarial IgG4), skin test reactivity to mite (mite SPT), IgE to mite (mite IgE; low level sensitization: >0.3, <1.0 IU/ml and high sensitization: >1.0 IU/ml), high level of total IgE (>1,400 IU/ml) and the levels of antifilarial IgG4, mite IgE and total IgE are indicated for 466 subjects. p values were measured using Pearson χ^2 for comparing prevalences and one-way ANOVA for comparing the levels of antifilarial IgG4, specific and total IgE.

Table 3. Measurements as a function of infection status as determined by antifilarial IgG4

	Noninfected (n = 216)	Infected (n = 250)	< p
Mite SPT prevalence, %	6.0 (13/216)	5.6 (14/250)	0.85
Mite IgE (>0.30 IU/ml) prevalence, %	56.5 (122/216)	65.6 (164/250)	0.04
Mite IgE (>1.0 IU/ml) prevalence, %	21.8 (47/216)	32.8 (82/250)	0.008
Total IgE (>1,400 IU/ml) prevalence, %	62.5 (135/216)	77.6 (194/250)	<0.001
Mite IgE level, IU/ml	0.21 (0.06–0.29)	0.25 (0.07–0.37)	0.021
Total IgE level, IU/ml	3.33 (2.97–3.79)	3.50 (3.22–3.88)	0.001

Skin test reactivity to mite (mite SPT), IgE to mite (mite IgE; low level sensitization: >0.3, <1.0 IU/ml and high sensitization: >1.0 IU/ml), high level of total IgE (>1,400 IU/ml) and the levels of mite IgE and total IgE are indicated for 466 subjects. p values were measured using Pearson χ^2 for comparing prevalence and one-way ANOVA for comparing specific and total IgE levels.

found an interaction between age and gender for infection status and mite IgE, whereas no interaction for mite SPT and total IgE was detected.

Effect of Infection on SPT and IgE Antibodies

When microfilaria presence was used to identify infected subjects, none of the 40 subjects positive for microfilaria were positive for mite SPT, whereas all 27 mite SPT-positive subjects were amicrofilaremic; this difference fell short of statistical significance ($p = 0.10$). When using antifilarial IgG4 to assign study subjects into 'infected' and 'noninfected' categories, prevalence of skin test reactivity to mite was similar in infected and non-

infected groups whereas the infected group showed higher total IgE and sensitization to mite (mite IgE) (table 3).

Determinant of Skin Reactivity

In industrialized countries total IgE and specific IgE are identified as risk factors for skin prick reactivity and allergic diseases. Regression analysis was performed to determine the contributions of total IgE and specific IgE to the outcome of a positive skin test to mite. Association of these independent variables with skin test reactivity was adjusted for age and gender. Mite SPT was strongly associated with mite IgE reactivity and total IgE ($b =$

Table 4. p values for testing the null hypothesis of clustering of mite SPT reactivity, prevalence and the level of mite and total IgE due to genetic and household effects

Outcome	Genetic	Household
Mite SPT prevalence	0.121	0.018
Mite IgE >0.30 IU/ml prevalence	<0.001	<0.001
Mite IgE >1.0 IU/ml prevalence	<0.007	<0.122
Total IgE >1,400 IU/ml prevalence	<0.001	<0.00
Mite IgE level, IU/ml	0.004	0.037
Total IgE level, IU/ml	<0.00	<0.001

Mite SPT and total IgE were adjusted for sex and age, whereas for mite IgE sex, age and interaction of sex and age were used as adjustment.

3.315, CI = 2.63–4.00, $p < 0.001$ and $b = 1.163$, CI = 0.92–1.77, $p = 0.007$, respectively).

Clustering Analysis of Mite SPT, Mite IgE and Total IgE

The study comprised 29 families which ranged in size from 1 to 214 individuals, the mean number of members per family was 16.07. The 466 inhabitants examined were spread over 112 households, the households ranged in size from 1 to 18 individuals and the mean number of members per household was 4.16. Mite SPT reactivity, mite IgE and total IgE were analyzed using the test statistics developed by Houwing-Duistermaat et al. [48]. The result showed that both genetic and household factors could explain the clustering of mite IgE as well as total IgE responses, whereas only household effects significantly explained the clustering of mite SPT reactivity (table 4).

Discussion

Skin Prick Test

The prevalence of skin prick reactivity to aeroallergens in this study population was found to be 11.6%. This number is comparable with the results of studies conducted in developing countries in Africa [41, 49, 50], but much lower than those reported in the developed countries [51]. Moreover, similar to studies carried out in Africa and in South America [41, 49, 50, 52], a positive SPT and IgE sensitization were predominantly against house dust mite in residents of Salubarana.

Differences in Allergy Phenotypes as a Function of Gender and Age

Although we did not find any significant differences in mite SPT as a function of gender, sensitization to mite (mite IgE) was significantly higher in males than in females. A higher prevalence of allergies in males, when described by mite IgE [53, 54] or by mite SPT [55, 56], has been described in several studies in Europe. The absence of a gender-related difference in mite SPT has been observed in other studies conducted in developing countries [8, 30] and the mechanisms behind this discrepancy in the data generated in western populations and those in developing countries have not been investigated.

Mite SPT reactivity as well as mite IgE reached a peak in the age group of 15–29 years and then decreased in older age groups. The decline is similar to studies carried out in the age groups of 20–70 years which found a decline in mite IgE with increasing age [53].

Effect of Infection

The general expansion of IgE-producing B cells during helminth infections may explain the higher levels of IgE antibodies in the infected subjects. The question of whether any cross-reactivity between house dust mites and helminth antigens occurs, as reported in an earlier study [57], needs to be investigated in our samples. Considering SPT reactivity, none of the microfilaria-positive individuals were positive in mite SPT. Although the results do not reach statistical significance ($p = 0.10$), most probably due to the small number of microfilaria-positive subjects, the data suggest that the presence of microfilariae in blood has a protective effect on mite SPT reactivity. This is in agreement with the findings that schistosomes and intestinal helminths reduce the risk of atopy [8, 30], asthma [58] or wheeze [9]. However, when antifilarial IgG4 was used as an indicator of active infection, no effect of infection on SPT positivity was observed. The discrepancy between the results when using microfilarial positivity versus positivity for antifilarial IgG4 may be due to the fact that only high infection levels (those with detectable microfilaria) are able to protect from SPT positivity. Indeed antifilarial IgG4 is believed to detect low level infections or amicrofilareemics who have cryptic or single sex infections [59, 60].

Contribution of Mite IgE and Total IgE to SPT Positivity

As in studies carried out in western populations, the high level of mite IgE in this study population was associated with a 6-fold increase in the risk of responding to

mite extracts in skin prick testing. However, this risk factor is relatively low compared with industrialized countries where having specific IgE generally translates into positivity in SPT [61, 62]. In the Netherlands the presence of mite IgE was associated with a 39-fold increase in the risk of SPT to mite (van der Zee, unpubl. data). Several reasons may be responsible for this difference.

Total IgE. Some investigators have attributed the negative association between helminth infections and allergic disorders to the presence of high levels of total IgE in helminth-infected subjects; nonspecific IgE may be able to saturate Fcε receptors and therefore prevent the binding of allergen-specific IgE [63]. However, more recent studies using appropriate statistical methods have found no effect of total IgE on the risk of skin prick reactivity or wheeze [41, 64] supporting our findings here.

Specific IgE. It is possible that the affinity of IgE antibodies to mite is lower in individuals infected with helminths. The lower affinity of mite IgE would then lead to lower mast cell degranulation [26]. This possibility has yet to be investigated in detail.

Immunoregulation. In helminth-infected subjects, there are increased regulatory responses characterized by high IL-10 production [30, 65] and the presence of regulatory T cells [66]. These suppressor responses may contribute to lower allergic outcomes in the face of elevated IgE antibodies [30].

Clustering of Allergic Phenotypes

From studies carried out in the west, it is clear that genetic and household factors influence predisposition to allergic disorders. In our study carried out in a rural area in a developing country, we evaluated the influence of family and household factors on mite SPT, mite IgE and total IgE in an area endemic for filariasis. We used a test statistic that screens data sets for different correlation structures [48]. Significant clustering of mite SPT, mite IgE and total IgE was seen in the study population. It was tested whether clustering existed due to genetic or household factors. The genetic model indicates that a phenotype is more similar between genetically related subjects who may belong to different households whereas the household model indicates that the phenotype is clustered within a household due to factors that govern the lifestyle within a household.

When the test was applied to allergic responses, namely mite SPT reactivity, mite IgE sensitization and total IgE, it showed different results. When IgE response to mite was considered, both genetic and household factors could significantly explain clustering of this trait. Thus

both household factors and genetic factors may contribute to mite IgE and total IgE responses in this population. Interestingly when the test is used to examine the aggregation of mite SPT, only the household effect was found to be significant, indicating that household factors affect this outcome. The evidence for a genetic effect on SPT has been observed in some studies in the west [reviewed in 12]. The weak genetic effect on SPT reactivity in our population may result from the ability of helminth infections to modulate skin reactivity to allergens. When using microfilaria positivity as a measure of infection, it was clear that indeed there may be an effect of filarial infection on SPT; however, when antifilarial IgG4 was used to also detect low level infections or exposure to larval stages of parasites, we did not see any effect of infection on SPT outcomes. These results suggest that high level infections, but not light infections, modulate the SPT. The specificity of antifilarial IgG4 has been estimated to be 95% using sera from Europeans infected with other nematode or trematode infections [47]. The assay may therefore also indicate the minor presence of some other helminth infections. Considering this, one would interpret the result that low level filarial infections and other helminths such as intestinal nematodes are less effective in modulating SPT. To substantiate this, studies need to be undertaken in another area where microfilaria prevalence is higher in order to be able to determine with certainty the modulating effect of SPT by microfilariae.

This study has shown that some allergic determinants such as IgE antibodies are under genetic control in populations residing in areas where pathogens such as helminths can be encountered. However, not all allergic outcomes are under genetic control. For skin reactivity to house dust mites, which in areas of low pathogen exposure has been reported to be under genetic control, may be also affected by environmental effects. These data raise the question of whether frequent exposure to pathogens such as to helminths can override the genetic influences of allergic traits, indicating a situation of a gene by environment interaction.

Acknowledgements

This work was supported by The Netherlands Foundation for the Advancement of Tropical Research (W93-325), Van Deventer-Maas Stichting, Netherlands Organization for International Cooperation in Higher Education (NUFFIC) and European Commission, INCO program, contract No. ICA4-CT-2001-10081.

References

- Teeratakulpisarn J, Pairajkul S, Heng S: Survey of the prevalence of asthma, allergic rhinitis and eczema in schoolchildren from Khon Kaen, Northeast Thailand. An ISAAC study. International Study of Asthma and Allergies in Childhood. Asian Pac J Allergy Immunol 2000;18:187-194.
- Norzila MZ, Haifa AL, Deng CT, Azizi BH: Prevalence of childhood asthma and allergy in an inner city Malaysian community: Intra-observer reliability of two translated international questionnaires. Med J Malaysia 2000;55:33-39.
- Hasan MM, Gofin R, Bar-Yishay E: Urbanization and the risk of asthma among schoolchildren in the Palestinian Authority. J Asthma 2000;37:353-360.
- Hijazi N, Abalkhail B, Seaton A: Asthma and respiratory symptoms in urban and rural Saudi Arabia. Eur Respir J 1998;12:41-44.
- Sudhir P, Prasad CE: Prevalence of exercise-induced bronchospasm in schoolchildren: An urban-rural comparison. J Trop Pediatr 2003;49:104-108.
- Perzanowski MS, Ng'ang'a LW, Carter MC, Odhiambo J, Ngari P, Vaughan JW, Chapman MD, Kennedy MW, Platts-Mills TA: Atopy, asthma, and antibodies to *Ascaris* among rural and urban children in Kenya. J Pediatr 2002;140:582-588.
- Odhiambo JA, Ng'ang'a LW, Mungai MW, Gicheha CM, Nyamwaya JK, Karimi F, Macklem PT, Becklake MR: Urban-rural differences in questionnaire-derived markers of asthma in Kenyan school children. Eur Respir J 1998;12:1105-1112.
- Nyan OA, Walraven GE, Banya WA, Milligan P, Van Der Sande M, Ceesay SM, Del Prete G, McAdam KP: Atopy, intestinal helminth infection and total serum IgE in rural and urban adult Gambian communities. Clin Exp Allergy 2001;31:1672-1678.
- Dagoye D, Bekele Z, Woldemichael K, Nida H, Yimam M, Hall A, Venn AJ, Britton JR, Hubbard R, Lewis SA: Wheezing, allergy, and parasite infection in children in urban and rural Ethiopia. Am J Respir Crit Care Med 2003;167:1369-1373.
- Weiss ST: Gene by environment interaction and asthma. Clin Exp Allergy 1999;29(suppl 2):96-99.
- Martinez FD: Gene by environment interactions in the development of asthma. Clin Exp Allergy 1998;28(suppl 5):21-28.
- Wills-Karp M, Ewart SL: Time to draw breath: Asthma-susceptibility genes are identified. Nat Rev Genet 2004;5:376-387.
- Xu J, Postma DS, Howard TD, Koppelman GH, Zheng SL, Stine OC, Bleeker ER, Meyers DA: Major genes regulating total serum immunoglobulin E levels in families with asthma. Am J Hum Genet 2000;67:1163-1173.
- Liu X, Beaty TH, Deindl P, Huang SK, Lau S, Sommerfeld C, Fallin MD, Kao WH, Wahn U, Nickel R: Associations between specific serum IgE response and 6 variants within the genes IL4, IL13, and IL4RA in German children: The German Multicenter Atopy Study. J Allergy Clin Immunol 2004;113:489-495.
- Karihaloo C, Tovey ER, Mitakakis TZ, Duffy DL, Britton WJ: Evidence for the genetic control of immunoglobulin E reactivity to the allergens of *Alternaria alternata*. Clin Exp Allergy 2002;32:1316-1322.
- Koppelman GH, Stine OC, Xu J, Howard TD, Zheng SL, Kauffman HF, Bleeker ER, Meyers DA, Postma DS: Genome-wide search for atopy susceptibility genes in Dutch families with asthma. Allergy Clin Immunol 2002;109:498-506.
- Fryer AA, Bianco A, Hepple M, Jones PW, Strange RC, Spiteri MA: Polymorphism at the glutathione S-transferase GSTP1 locus. A new marker for bronchial hyperresponsiveness and asthma. Am J Respir Crit Care Med 2000;161:1437-1442.
- Ulbrecht M, Hergeth MT, Wjst M, Heinrich J, Bickeboller H, Wichmann HE, Weiss EH: Association of beta(2)-adrenoreceptor variants with bronchial hyperresponsiveness. Am J Respir Crit Care Med 2000;161:469-474.
- Wong GW, Ko FW, Lau TS, Li ST, Hui D, Pang SW, Leung R, Fok TF, Lai CK: Temporal relationship between air pollution and hospital admissions for asthmatic children in Hong Kong. Clin Exp Allergy 2001;31:565-569.
- Garty BZ, Kosman E, Ganor E, Berger V, Garty L, Wietzen T, Waisman Y, Mimouni M, Waisel Y: Emergency room visits of asthmatic children, relation to air pollution, weather, and airborne allergens. Ann Allergy Asthma Immunol 1998;81:563-570.
- Morgan WJ, Crain EF, Gruchalla RS, O'Connor GT, Kattan M, Evans R 3rd, Stout J, Malindzak G, Smartt E, Plaut M, Walter M, Vaughn B, Mitchell H; Inner-City Asthma Study Group: Results of a home-based environmental intervention among urban children with asthma. N Engl J Med 2004;351:1068-1080.
- Hodge L, Salome CM, Hughes JM, Liu-Brennan D, Rimmer J, Allman M, Pang D, Armour C, Woolcock AJ: Effect of dietary intake of omega-3 and omega-6 fatty acids on severity of asthma in children. Eur Respir J 1999;11:361-365.
- Chen R, Hu Z, Seaton A: Eating more vegetables might explain reduced asthma symptoms. BMJ 2004;328:1052-1053.
- Matricardi PM, Rosmini F, Ferrigno L, Nisini R, Rapietta M, Chionne P, Stroffolini T, Pasquini P, D'Amelio R: Cross-sectional retrospective study of prevalence of atopy among Italian military students with antibodies against hepatitis A virus. BMJ 1997;314:999-1003.
- von Mutius E: Infection: Friend or foe in the development of atopy and asthma? The epidemiological evidence. Eur Respir J 2001;18:872-881.
- Yazdanbakhsh M, Kreamsner PG, van Ree R: Allergy, parasites, and the hygiene hypothesis. Science 2002;296:490-494.
- Cooper PJ: The potential impact of early exposures to geohelminth infections on the development of atopy. Clin Rev Allergy Immunol 2004;26:5-14.
- Dold S, Heinrich J, Wichmann HE, Wjst M: *Ascaris*-specific IgE and allergic sensitization in a cohort of school children in the former East Germany. J Allergy Clin Immunol 1998;102:414-420.
- Wilson MS, Maizels RM: Regulation of allergy and autoimmunity in helminth infection. Clin Rev Allergy Immunol 2004;26:35-50.
- van den Biggelaar AH, van Ree R, Rodrigues LC, Lell B, Deelder AM, Kreamsner PG, Yazdanbakhsh M: Decreased atopy in children infected with *Schistosoma haematobium*: A role for parasite-induced interleukin-10. Lancet 2000;356:1723-1727.
- Elias D, Wolday D, Akuffo H, Petros B, Bronner U, Britton S: Effect of deworming on human T cell responses to mycobacterial antigens in helminth-exposed individuals before and after bacille Calmette-Guerin (BCG) vaccination. Clin Exp Immunol 2001;123:219-225.
- Marquet S, Abel L, Hillaire D, Dessein H, Kallil J, Feingold J, Weissenbach J, Dessein AJ: Genetic localization of a locus controlling the intensity of infection by *Schistosoma mansoni* on chromosome 5q31-q33. Nat Genet 1996;14:181-184.
- Dessein AJ, Hillaire D, Elwali NE, Marquet S, Mohamed-Ali Q, Mirghani A, Henri S, Abdelhameed AA, Saeed OK, Magzoub MM, Abel L: Severe hepatic fibrosis in *Schistosoma mansoni* infection is controlled by a major locus that is closely linked to the interferon-gamma receptor gene. Am J Hum Genet 1999;65:709-721.
- Terhell AJ, Houwing-Duistermaat JJ, Ruiterman Y, Haarbrink M, Abadi K, Yazdanbakhsh M: Clustering of *Brugia malayi* infection in a community in South-Sulawesi, Indonesia. Parasitology 2000;120:23-29.
- Wahyuni S, Houwing JJ, Syafruddin, Supali T, Yazdanbakhsh M, Sartono E: Clustering of filarial infection in age graded study: Genetic, household and environmental influences. Parasitology 2004;128:315-321.
- Williams-Blangero S, VandeBerg JL, Subedi J, Aivaliotis MJ, Rai DR, Upadhyay RP, Jha B, Blangero J: Genes on chromosomes 1 and 13 have significant effects on *Ascaris* infection. Proc Natl Acad Sci USA 2002;99:5533-5538.
- Oryszczyn MP, Annesi I, Neukirch F, Dore MF, Kauffmann F: Relationships of total IgE level, skin prick test response, and smoking habits. Ann Allergy 1991;67:355-358.

- 38 Suliaman FA, Holmes WF, Kwick S, Khouri F, Ratard R: Pattern of immediate type hypersensitivity reactions in the Eastern Province, Saudi Arabia. *Ann Allergy Asthma Immunol* 1997;78:415–418.
- 39 Saracevic E, Omercahic-Dizdarevic A, Hasanbegovic S: Allergy tests in children with seasonal allergic rhinitis (in Croatian). *Med Arh* 2002;56(3 suppl 1):23–25.
- 40 Smith PK, Krohn RI, Hermanson GT, Mallia AK, Gartner FH, Provenzano MD, Fujimoto EK, Goke NM, Olson BJ, Klenk DC: Measurement of protein using bicinchoninic acid. *Anal Biochem* 1985;150:76–85.
- 41 van den Biggelaar AH, Lopuhaa C, van Ree R, van der Zee JS, Jans J, Hoek A, Migombet B, Borrmann S, Luckner D, Kremsner PG, Yazdanbakhsh M: The prevalence of parasite infestation and house dust mite sensitization in Gabonese schoolchildren. *Int Arch Allergy Immunol* 2001;126:231–238.
- 42 Aalberse RC, Koshte V, Clemens JG: Immunoglobulin E antibodies that crossreact with vegetable foods, pollen, and Hymenoptera venom. *J Allergy Clin Immunol* 1981;68:356–364.
- 43 Schuurman J, Perdok GJ, Lourens TE, Parren PW, Chapman MD, Aalberse RC: Production of a mouse/human chimeric IgE monoclonal antibody to the house dust mite allergen Der p 2 and its use for the absolute quantification of allergen-specific IgE. *J Allergy Clin Immunol* 1997;99:545–550.
- 44 Haarbrink M, Terhell A, Abadi K, van Beers S, Asri M, Yazdanbakhsh M: IgG4 antibody assay in the detection of filariasis. *Lancet* 1995;346:853–854.
- 45 Wamae CN, Roberts JM, Eberhard ML, Lammie PJ: Kinetics of circulating human IgG4 after diethylcarbamazine and ivermectin treatment of bancroftian filariasis. *J Infect Dis* 1992;165:1158–1160.
- 46 Terhell AJ, Haarbrink M, van den Biggelaar A, Mangali A, Sartono E, Yazdanbakhsh M: Long-term follow-up of treatment with diethylcarbamazine on anti-filarial IgG4: Dosage, compliance, and differential patterns in adults and children. *Am J Trop Med Hyg* 2003;68:33–39.
- 47 Polderman AM, Verweij JJ: Recente ontwikkelingen in de diagnostiek van parasitaire infecties; in van't Wout JW (ed): *Importziekten*. Den Haag, CIP-Gegevens Koninklijke Bibliotheek, 1991.
- 48 Houwing-Duistermaat JJ, van Houwelingen HC, Terhell A: Modelling the cause of dependency with application to filaria infection. *Stat Med* 1998;17:2939–2954.
- 49 Yemaneberhan H, Bekele Z, Venn A, Lewis S, Parry E, Britton J: Prevalence of wheeze and asthma and relation to atopy in urban and rural Ethiopia. *Lancet* 1997;350:85–90.
- 50 Shaheen SO, Aaby P, Hall AJ, Barker DJ, Heyes CB, Shiell AW, Goudiaby A: Measles and atopy in Guinea-Bissau. *Lancet* 1996;347:1792–1796.
- 51 Akerman M, Valentine-Maher S, Rao M, Tanningco G, Khan R, Tuysugoglu G, Joks R: Allergen sensitivity and asthma severity at an inner city asthma center. *J Asthma* 2003;40:55–62.
- 52 Lynch NR, Hagel I, Perez M, Di Prisco MC, Lopez R, Alvarez N: Effect of anthelmintic treatment on the allergic reactivity of children in a tropical slum. *J Allergy Clin Immunol* 1993;92:404–411.
- 53 Kerkhof M, Droste JH, de Monchy JG, Schouten JP, Rijcken B: Distribution of total serum IgE and specific IgE to common aeroallergens by sex and age, and their relationship to each other in a random sample of the Dutch general population aged 20–70 years. Dutch ECRHS Group, European Community Respiratory Health Study. *Allergy* 1996;51:770–776.
- 54 Court CS, Cook DG, Strachan DP: The descriptive epidemiology of house dust mite-specific and total immunoglobulin E in England using a nationally representative sample. *Clin Exp Allergy* 2002;32:1033–1041.
- 55 Jedrychowski W, Flak E, Mroz E, Lis G: Sensitization to common airborne allergens in school children with bronchial asthma. *Przegl Lek* 2000;57:441–445.
- 56 Eriksson NE, Holmen A: Skin prick tests with standardized extracts of inhalant allergens in 7,099 adult patients with asthma or rhinitis: Cross-sensitizations and relationships to age, sex, month of birth and year of testing. *J Investig Allergol Clin Immunol* 1996;6:36–46.
- 57 Santiago ML, Hafalla JC, Kurtis JD, Aligui GL, Wiest PM, Olveda RM, Olds GR, Dunne DW, Ramirez BL: Identification of the *Schistosoma japonicum* 22.6-kDa antigen as a major target of the human IgE response: Similarity of IgE-binding epitopes to allergen peptides. *Int Arch Allergy Immunol* 1998;117:94–104.
- 58 Selassie FG, Stevens RH, Cullinan P, Pritchard D, Jones M, Harris J, Ayres JG, Newman Taylor AJ: Total and specific IgE (house dust mite and intestinal helminths) in asthmatics and controls from Gondar, Ethiopia. *Clin Exp Allergy* 2000;30:356–358.
- 59 Kwan-Lim GE, Forsyth KP, Maizels RM: Filarial-specific IgG4 response correlates with active *Wuchereria bancrofti* infection. *J Immunol* 1990;145:4298–4305.
- 60 Kurniawan A, Yazdanbakhsh M, van Ree R, Aalberse R, Selkirk ME, Partono F, Maizels RM: Differential expression of IgE and IgG4 specific antibody responses in asymptomatic and chronic human filariasis. *J Immunol* 1993;150:3941–3950.
- 61 Dreborg S, Frew A: Allergen standardization and skin test. EAACI position paper. *Allergy* 1993;48/S14:49–82.
- 62 Schuetze G, Storm van's Gravesande K, Sparhold S, Frischer T, Kuehr J: Comparison between serial skin-prick tests and specific serum immunoglobulin E to mite allergens. *Pediatr Allergy Immunol* 1999;10:138–142.
- 63 Lynch NR, Goldblatt J, Le Souef PN: Parasite infections and the risk of asthma and atopy. *Thorax* 1999;54:659–660.
- 64 Scrivener S, Yemaneberhan H, Zebeignus M, Tilahun D, Girma S, Ali S, McElroy P, Custovic A, Woodcock A, Pritchard D, Venn A, Britton J: Independent effects of intestinal parasite infection and domestic allergen exposure on risk of wheeze in Ethiopia: A nested case-control study. *Lancet* 2001;358:1493–1499.
- 65 King CL, Medhat A, Malhotra I, Nafeh M, Helmy A, Khaudary J, Ibrahim S, El-Sherbiny M, Zaky S, Stupi RJ, Brustoski K, Shehata M, Shata MT: Cytokine control of parasite-specific anergy in human urinary schistosomiasis. IL-10 modulates lymphocyte reactivity. *J Immunol* 1996;156:4715–4721.
- 66 Doetze A, Satoguina J, Burchard G, Rau T, Loliger C, Fleischer B, Hoerauf A: Antigen-specific cellular hyporesponsiveness in a chronic human helminth infection is mediated by T(h)3/T(r)1-type cytokines IL-10 and transforming growth factor-beta but not by a T(h)1 to T(h)2 shift. *Int Immunol* 2000;12:623–630.