

ORIGINAL PAPER

Immunoglobulin constant heavy G chain genes as risk factors in childhood allergies

V.-A. Oxelius^{*†}, L. Bråbäck[‡], S. Ahlstedt^{§¶} and B. Björkstén[¶]

^{*}Department of Pediatrics, [†]Institute of Clinical Immunology, University Hospital, Lund University, Lund, Sweden, [‡]Department of Public Health and Research, Sundsvall Hospital, Sundsvall, Sweden, [§]Pharmacia Diagnostics AB, Uppsala, Sweden and [¶]National Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden

Clinical and Experimental Allergy

Summary

Background Several candidate genes have been found to be associated with the inflammatory response of IgE-mediated allergy, so also the immunoglobulin constant heavy G chain (*IGHG*) genes. The *IGHG* genes are situated close to the *IGHE* gene on chromosome 14q32, 5'μ, δ, γ3, γ1, α1, γ2, γ4, ε, α2, 3'. They are inherited in a Mendelian fashion and expressed randomly in allelic exclusion. The alternative and functionally different γ3, γ1 and γ2 gene variants are found in four *IGHG* haplotypes, coding four B cell variants.

Objective The aim of this study was to assess the frequency of different *IGHG* genes in relation to phenotypes associated with allergy, in a case-control study.

Methods We identified the constant heavy-chain genes of IgG in 198 allergic and non-allergic children participating in the Phase II of the International Study of Asthma and Allergy in Children. The *IGHG* genes were assessed by the alternative serum IgG subclass allotypes expressing the alternative alleles of γ3, γ1 and γ2 genes, using ELISA and double immunodiffusion.

Results The *IGHG*^{*}*bfn* haplotype (= B1 cells) and *IGHG2*^{*}*n* allele dominated (51% vs. 24%, *P* = 0.002) and the *IGHG*^{*}*bf-n* haplotype (= B2 cells) was infrequent (16% vs. 52%, *P* < 0.001) in allergic children with a family history of allergy, clinical manifest allergy and positive skin prick test (SPT). The frequency of *IGHG* genes was similar in children with maternal and paternal heredity and in children with wheezing, eczema or rhinitis, as well as in children with different positive SPT. The *IGHG*^{*}*bfn* haplotype with the *IGHG2*^{*}*n* allele was strongly associated with heredity for allergy. The *IGHG*^{*}*bf-n* haplotype was inversely related to allergy.

Conclusions IgG allotypes, immunochemical and functional variants of IgG molecules from *IGHG* genes are associated with atopy. The *IGHG*^{*}*bfn* haplotype (= B1 cells) with the *IGHG2*^{*}*n* allele dominates, associated with an increased risk for atopy. In contrast, the *IGHG*^{*}*bf-n* haplotype (= B2 cells) with the *IGHG2*^{*}*n* allele is associated with low risk.

Keywords allergic rhinitis, allergy heredity, alternative IgG subclass allotypes, eczema, *IGHG* alleles, *IGHG* diplotypes, *IGHG* genotypes, *IGHG* haplotypes (= B-cells), SPT, wheezing

Submitted 14 March 2006; revised 13 September 2006; accepted 19 September 2006

Correspondence:

Dr Vivi-Anne Oxelius, Department of Pediatrics, University Hospital, Lund University, Se-221 85 Lund, Sweden.
E-mail: vivi-anne.oxelius@med.lu.se

Introduction

Asthma and allergies may affect as many as one in four urban children [1]; it is associated with substantial morbidity and economic costs [2] and the factors that confer susceptibility are not well understood. Identifying genetic pathways that allow asthma and allergies to be expressed

is fundamental for developing diagnostic tools and designing therapies [3].

Asthma, eczema and hayfever are all manifestations of the atopic state and they are strongly familial on a multifactorial genetic basis [4]. The genetic codes for the heavy constant chains of both IgE and IgG molecules are found on chromosome 14q32.3, with the immunoglobulin heavy

constant chain (*IGH*) genes in the order 5'μ, δ, γ3, γ1, α1, γ2, γ4, ε, α2, 3' [5]. The IgG allotypes are genetic markers of IgG proteins and therefore products of structural IgG genes. The alternative IgG allotypes from the γ3, γ1 and γ2 genes have minor amino acid differences, which correlate with single base substitutions in the genes [5]. The IgG allotypes are inherited in a Mendelian fashion and are expressed randomly in allelic exclusion. The IgG allotypes are separate entities with distinct immunochemical and functional characteristics, such as fractionation rates, electrophoretic rates [6], half-life times [7] and maturation rate during childhood [8]. Four subsets of B cells have been identified, based on the alternative protein expressions from the *IGHG3*, *IGHG1* and *IGHG2* genes [9]: the *IGHG***bfn* haplotype (= B1 cells) is typed by IgG3**b*, IgG1**f* and IgG2**n*; the *IGHG***bf-n* haplotype (= B2 cells) is typed by IgG3**b*, IgG1**f* and IgG2*-*n*; the *IGHG***gan* haplotype (= B3 cells) is typed by IgG3**g*, IgG1**a* and IgG2**n*; and the *IGHG***ga-n* haplotype (= B4 cells) is typed by IgG3**g*, IgG1**a* and IgG2*-*n* allotypes in serum [10] (Table 1).

Restricted *IGHG* genes are associated with both atopic and non-atopic asthma, replicated in several studies [11, 12]. In childhood asthma, the *IGHG***bfn*/**bfn* diplotype (= B1/B1 cells) with the *IGHG2***n*/**n* genotype is

associated with an *IGHE* gene, inducing increased levels of serum IgE and allergen specific IgE antibodies, while the alternative *IGHG***ga-n*/**ga-n* diplotype (= B4/B4) with the alternative *IGHG2***-n*/**-n* is linked to a low responding *IGHE* gene. Two different pathways of immune regulation are associated with the B1/B1 cells and B4/B4 cells, respectively, in asthmatic children with possible influences on pathogenesis [13]. Those with the B1/B1 cells have an imbalanced class switch in the rearrangement of the *IGHG* genes, expressed as low serum IgG1**f* level and high levels of serum IgG2**n* and IgE [14] and those with B4/B4 express low levels of both IgG1**a*, IgG2*-*n*, IgG3**g* and IgE.

We tested the hypothesis that various expressions of allergy are influenced by different *IGHG* gene-coded B cells and thereby add diagnostic power to other procedures.

Materials and methods

Study subjects

The study group (198 children) was selected among 1390 twelve-year-old children from 25 schools in Östersund, Sweden, participating in phase 2 of the International

Table 1. Abbreviations and description of the immunoglobulin heavy G chain (*IGHG*) genes on chromosome 14q32, alternative IgG subclass allotypes, *IGHG* alleles, *IGHG* haplotypes = B cells and *IGHG* diplotypes (individual)

(A) IGH genes of chromosome 14q32					
5' μ, δ, γ3, γ1, α1, γ2, γ4, ε, α2 3'					
(B) Alternative alleles of IGHG subclass genes					
IGHG gene			Alternative alleles		
IGHG3 (γ3)			*b and *g		
IGHG1 (γ1)			*f and *a		
IGHG2 (γ2)			*n and *-n		
(C) IGH genes of chromosome 14q32. Four different IGHG haplotypes coding four different B cells					
IGH genes of chromosome 14q32		IGHG	IgG allotypes in serum		B cell variation
5' μ, δ, γ3, γ1, α1, γ2, γ4, ε, α2 3'		haplotypes	Alternative protein variants = genetic markers		
with alternative alleles inserted for IGHG3, IGHG1 and IGHG2			for heavy G subclass chains of IgG3, IgG1 and IgG2		
5' μ, δ, b, f, α1, n, γ4, ε, α2 3'		IGHG*bfn	IgG3*b, IgG1*f, IgG2*n		B1 cells
5' μ, δ, b, f, α1, -n, γ4, ε, α2 3'		IGHG*bf-n	IgG3*b, IgG1*f, IgG2*-n		B2 cells
5' μ, δ, g, a, α1, n, γ4, ε, α2 3'		IGHG*gan	IgG3*g, IgG1*a, IgG2*n		B3 cells
5' μ, δ, g, a, α1, -n, γ4, ε, α2 3'		IGHG*ga-n	IgG3*g, IgG1*a, IgG2*-n		B4 cells
(D) Serum IgG, IgG subclasses and alternative IgG subclass allotypes					
IgG		IgG1		IgG1*f	
				IgG1*a	
		IgG2		IgG2*n	
				IgG2*-n	
		IgG3		IgG3*b	
				IgG3*g	
		IgG4			
(E) The six most common B cell combinations, IGHG diplotypes and IGHG2 genotypes					
B1/B1 cells	B1/B2 cells	B1/B4 cells	B2/B2 cells	B2/B4 cells	B4/B4 cells
IGHG*bfn/*bfn	IGHG*bfn/*bf-n	IGHG*bfn/*ga-n	IGHG*bf-n/*bf-n	IGHG*bf-n/*ga-n	IGHG*ga-n/*ga-n
IGHG2*n/*n	IGHG2*n/*-n	IGHG2*n/*-n	IGHG2*-n/*-n	IGHG2*-n/*-n	IGHG2*-n/*-n

Study of Asthma and Allergy in Children (ISAAC). All children with current wheezing and a random sample of non-wheezers were invited to a case-control questionnaire study [15]. Skin prick tests (SPT) were performed with five inhalant allergens (tree pollen, grass pollen, cat, dog and horse), strictly adhering to the ISAAC Phase 2 manual. Eighty-two were SPT positive and 116 SPT negative. Based on the questionnaire responses, the children were divided into those with and without clinical symptoms ($n = 130$ and 68 , respectively) and those with and without a family history of allergy ($n = 144$ and 54 , respectively) (Table 2). Sixty-one children had both clinical symptoms of allergy, a family history of allergy and positive SPT, while 21 children had none of these. The *IGHG* genotypes of the study population were in Hardy-Weinberg equilibrium $P > 0.05$, compared with Caucasian populations from Sweden [16], Finland and the Netherlands [17].

Methods

Immunoglobulin constant heavy G chain genotyping by serum IgG allotypes. The *IGHG* genes are situated on the long arm q at band 32 on chromosome 14. They are found in order $\gamma 3$, $\gamma 1$, $\gamma 2$, as used in the following, and together with all other immunoglobulin constant heavy chains, also ϵ for IgE, in the *IGH* gene $5'\mu$, δ , $\gamma 3$, $\gamma 1$, $\alpha 1$, $\gamma 2$, $\gamma 4$, ϵ , $\alpha 2$ 3'. Two alternative gene variants are reported by studies of allotypes, from the $\gamma 3$, $\gamma 1$ and $\gamma 2$ gene loci, producing IgG subclass molecules with only minor amino acid epitope differences of the constant heavy G chains [5] (Table 1). The alternative protein variations from the $\gamma 3$ (*IGHG3*), $\gamma 1$ (*IGHG1*) and $\gamma 2$ (*IGHG2*) genes can easily be studied qualitatively and quantitatively in serum [10]. The alternative *IGHG* genes are inherited in the Mendelian way and expressed randomly in allelic exclusion. The secreted alternative IgG subclass proteins are for *IGHG3*: IgG3**b* and IgG3**g*, for *IGHG1*: IgG1**f* and IgG1**a* and for *IGHG2*: IgG2**n* and IgG2**-n* allotypes (Table 1).

The $\gamma 3$ and $\gamma 1$ genes are concordant because of their close localization (only 26 kb in between): *IGHG3**b** together with *IGHG1**f**, in the gene complex**bf* and *IGHG3**g** together with *IGHG1**a**, **ga* [5]. Additional variation on *IGHG2* with the **n* or **-n* genes, respectively, makes four different haplotypes [5], which also code four different B cells (Table 1) [9].

A competitive ELISA was used to quantitate serum IgG1**f*, IgG1**a*, IgG2**n* and IgG3**b* allotypes [10]. The following monoclonal antibodies: anti IgG1**f* clone 5F10, anti IgG1**a* clone 5E7, anti IgG3**b*1/u clone 12D9 (Janssen Biochimica, Beerse, Belgium) and anti IgG2**n* clone SH21 (Sigma, St Louis, MO, USA) and predetermined concentration of purified myeloma proteins of the following IgG allotypes: IgG1**f*, IgG1**a*, IgG2**n* and IgG3**b* were used. The sensitivity of the ELISA assay was

0.0008 g/L for IgG1**a*, 0.0003 g/L for IgG1**f*, 0.0006 for IgG2**n* and 0.0007 g/L for IgG3**b*.

Determination of homozygosity and heterozygosity for the IgG2**n* allotype was performed with a double immuno diffusion assay with the monoclonal reagents anti IgG2**n* clone SH21 and anti HP 6014 (Sigma). The control test for IgG3**g* was, besides the absence of IgG3**b*, also the presence of the concordant IgG1**a* allotype. In heterozygous individuals, the IgG3**g* and IgG3**b* are about 50% : 50%, respectively, of the IgG3 amount. The method is described in detail elsewhere [10, 8].

Frequencies of the *IGHG* alleles, and the four *IGHG* (B cell) haplotypes were identified, i.e. *IGHG**bfn** (B1), *IGHG**bf-n** (B2), *IGHG**gan** (B3) and *IGHG**ga-n** (B4), making up 10 diplotypes: B1/B1, B1/B2, B1/B3, B1/B4, B2/B2, B2/B3, B2/B4, B3/B3, B3/B4 and B4/B4, of which six are common and four rare ($< 1-3\%$) in the Caucasian population [5, 8].

The data were analysed by the statistical package, - SPSS 11.5, calculating the *IGHG* allele, *IGHG* genotype, *IGHG* haplotype (= B cell) and *IGHG* diplotype frequencies for different allergic and non-allergic phenotypes. An analysis of 2×2 tables using the 'Wald' approximation of the variance of the odds ratios (OR) was used similar to simple logistic regression. An estimate of the OR for the final model variables and associated 95% confidence intervals (CI) are presented in the tables and results. In addition, stepwise logistic regression was performed in order to assess multi-variate associations between different B cell haplotypes and heredity, clinical symptoms and positive SPT.

Results

Of the 198 children in the study, 177 had one or more of either a family history of allergy, clinical allergy symptoms and/or a positive SPT, while 21 did not (Table 2). In children with any of these allergy indicators, the *IGHG2**n** allele and the *IGHG**bfn** haplotype (= B1 cells) were more and the frequencies of the *IGHG2**-n** allele and the *IGHG**bf-n** haplotype (= B2 cells) less common than in children with neither a positive family history nor clinical symptoms of allergy, or a positive SPT (Tables 2 and 3). The findings were most prominent in children with a family history of allergy. The presence of two or three of the allergy variables strengthened the relation to *IGHG* genes namely, for heredity and symptoms (B1 cells: OR 3.3, 95% CI 1.6-6.9; $P < 0.001$; B2 cells: OR 0.2, 95% CI 0.1-0.3, $P < 0.001$), for heredity and positive SPT (B1 cells: OR 2.5, 95% CI 1.4-4.6; $P = 0.002$; B2 cells: OR 0.3, 95% CI 0.1-0.5; $P < 0.001$) and for allergy symptoms and positive SPT (B1 cells: OR 1.7, 95% CI 1.0-2.7; $P = 0.048$; B2 cells: OR 0.4, 95% CI 0.2-0.7; $P < 0.001$). The B1 cells were particularly common in sensitized children combined with a positive family history of allergy

Table 2. Frequency of immunoglobulin heavy G2 chain (*IGHG2*) genotypes, *IGHG2***n* and *IGHG2***-n* alleles in 198 children with and without allergy variables

Allergy variables	<i>IGHG2</i> genotypes, <i>N</i> (%)		<i>IGHG2</i> alleles, <i>N</i> (%)				<i>P</i> value	OR	95% CI
	Yes	No	Yes	No					
Family history of allergy	144	54	288	108					
* <i>n</i> /* <i>n</i>	35 (24)	5 (9)							
* <i>n</i> /* <i>-n</i>	65 (45)	20 (37)							
* <i>-n</i> /* <i>-n</i>	44 (31)	29 (54)							
			* <i>n</i>	* <i>-n</i>	* <i>n</i>	* <i>-n</i>			
			135 (47)	153 (53)	30 (28)	78 (72)	< 0.001	2.3	1.4–3.7
Clinical symptoms	130	68	260	136					
* <i>n</i> /* <i>n</i>	28 (22)	12 (18)							
* <i>n</i> /* <i>-n</i>	61 (47)	24 (35)							
* <i>-n</i> /* <i>-n</i>	41 (32)	32 (47)							
			* <i>n</i>	* <i>-n</i>	* <i>n</i>	* <i>-n</i>			
			117 (45)	143 (55)	48 (35)	88 (65)	NS	1.5	1.0–2.3
Sensitized (SPT+)	82	116	164	232					
* <i>n</i> /* <i>n</i>	19 (23)	21 (18)							
* <i>n</i> /* <i>-n</i>	38 (46)	47 (41)							
* <i>-n</i> /* <i>-n</i>	25 (30)	48 (41)							
			* <i>n</i>	* <i>-n</i>	* <i>n</i>	* <i>-n</i>			
			76 (46)	88 (54)	89 (38)	143 (62)	NS	1.4	0.9–2.1
Any of the above	177	21	354	42					
* <i>n</i> /* <i>n</i>	38 (21)	2 (10)							
* <i>n</i> /* <i>-n</i>	79 (45)	6 (29)							
* <i>-n</i> /* <i>-n</i>	60 (34)	13 (62)							
			* <i>n</i>	* <i>-n</i>	* <i>n</i>	* <i>-n</i>			
			155 (44)	199 (56)	10 (24)	32 (76)	0.013	2.4	1.2–5.2
All of the above	61	21	122	42					
* <i>n</i> /* <i>n</i>	17 (28)	2 (10)							
* <i>n</i> /* <i>-n</i>	28 (46)	6 (29)							
* <i>-n</i> /* <i>-n</i>	16 (26)	13 (62)							
			* <i>n</i>	* <i>-n</i>	* <i>n</i>	* <i>-n</i>			
			62 (51)	60 (49)	10 (24)	32 (76)	0.002	3.3	1.5–7.3

SPT, skin prick test.

(OR 4.4, 95% CI 1.7–11.6; $P=0.001$), whereas the B4-cells were uncommon (OR 0.4, 95% CI 0.2–0.8; $P=0.013$).

In 61 atopic children with a positive family history, clinical allergy and positive SPT, the *IGHG2***n* allele and the B1 cells dominated (OR 3.3, 95% CI 1.5–7.3; $P=0.002$) and the *IGHG2***-n* allele and the B2 cells (OR 0.4, 95% CI 0.2–0.7; $P<0.001$) were infrequent, as compared with the 21 children with none of these traits (Tables 2 and 3). The relationship between allergy and the *IGHG***bfn* haplotype (= B1 cells) and *IGHG2***n* allele and inverse relationship with the *IGHG***bf-n* haplotype (= B2 cells) and *IGHG2***-n* allele were confirmed in subgroups of children with various allergic manifestations, such as either paternal or maternal heredity, wheezing, rhinitis and eczema and positive SPT to various inhalant allergens (Tables 4 and 5).

At the diplotic level, a high frequency of the *IGHG***bfn*/**bfn* diplotype (= B1/B1 cells) was found in children with a positive family history of allergy (OR 3.9, 95% CI 1.3–11.5; $P=0.001$) while the *IGHG***bf-n*/**bf-n*

diplotype (= B2/B2 cells) was uncommon (OR 0.3, 95% CI 0.1–0.7; $P=0.007$).

The frequencies of the *IGHG***gan* haplotype (B3 cells) (rare in the Caucasian population) and the *IGHG***ga-n* haplotype (B4 cells) (Tables 3 and 5), as well as *IGHG1* and *IGHG3* genotypes (data not shown) were similar in allergic and non-allergic children, with the exception of a low number of the *IGHG***ga-n* haplotype in sensitized children with a family history of allergy.

Discussion

The frequencies of the *IGHG2***n* allele, the *IGHG***bfn* haplotype and homozygous *IGHG***bfn*/**bfn* diplotype were high in children with either a family history of allergy, clinical symptoms or a positive SPT, while the alternative *IGHG2***-n* allele on *IGHG2* locus, the *IGHG***bf-n* haplotype and the *IGHG***bf-n*/**bf-n* diplotype were uncommon. This is in agreement with earlier

Table 3. Frequency of B cells [immunoglobulin heavy G chain (*IGHG*) haplotypes], B1 ($B^{*bf/n}$), B2 (B^{*bf-n}), B3 (B^{*gan}) and B4 (B^{*ga-n}) in relation to a family history of allergy, clinical symptoms and sensitization of 198 children (396 haplotypes)

Allergy variables	B cells coded by <i>IGHG</i> genes, <i>n</i> (%)		<i>P</i> -value	OR	95% CI
	Yes	No			
Family history of allergy	288	108			
B1 ($B^{*bf/n}$)	133 (46)	28 (26)	< 0.001	2.5	1.5–4.0
B2 (B^{*bf-n})	48 (17)	38 (35)	< 0.001	0.4	0.2–0.6
B3 (B^{*gan})	2 (1)	2 (2)	NS		
B4 (B^{*ga-n})	105 (37)	40 (37)	NS		
Clinical symptoms	260	136			
B1 ($B^{*bf/n}$)	115 (44)	46 (34)	0.045	1.6	1.0–2.4
B2 (B^{*bf-n})	43 (17)	43 (32)	< 0.001	0.4	0.3–0.7
B3 (B^{*gan})	2 (1)	2 (2)	NS		
B4 (B^{*ga-n})	100 (39)	45 (33)	NS		
Sensitized (SPT+)	164	232			
B1 ($B^{*bf/n}$)	73 (45)	88 (38)	NS	1.3	0.9–2.0
B2 (B^{*bf-n})	27 (17)	59 (25)	0.033	0.6	0.4–1.0
B3 (B^{*gan})	3 (2)	1 (< 1)	NS		
B4 (B^{*ga-n})	61 (37)	84 (36)	NS		
Any of the above	354	42			
B1 ($B^{*bf/n}$)	151 (43)	10 (24)	0.019	2.4	1.1–5.0
B2 (B^{*bf-n})	64 (18)	22 (52)	< 0.001	0.2	0.1–0.4
B3 (B^{*gan})	3 (1)	1 (2)	NS		
B4 (B^{*ga-n})	136 (38)	9 (21)	0.031	2.3	1.1–4.9
All of above	122	42			
B1 ($B^{*bf/n}$)	62 (51)	10 (24)	0.002	3.3	1.5–7.3
B2 (B^{*bf-n})	20 (16)	22 (52)	< 0.001	0.4	0.2–0.7
B3 (B^{*gan})	1 (1)	1 (2)	NS		
B4 (B^{*ga-n})	39 (32)	9 (21)	NS		

findings that the IgG2**n* allotype is associated with an allergen-responding *IGHG* gene [11–13].

The strongest association was seen in children with a positive family history of allergy, while the association was less obvious for clinical symptoms and sensitization. The high frequency of the *IGHG*2**n* allele and the *IGHG***bf/n* haplotype (=B1 cells) and low frequency of the *IGHG*2**-n* allele and the *IGHG***bf-n* haplotype (=B2 cells) in atopic children were similar in maternal and paternal heredity, and in wheezing children, as compared with those with eczema and rhinitis, as well as in children sensitized against different inhalant allergens. Our findings underline the importance of *IGHG*2**n*, but not the *IGHG*2**-n*, as a risk factor with a possible role of serum IgG2**n* allotype antibody molecules in the pathogenesis of allergy.

IGHG genotyping by alternative serum IgG subclass allotypes is the best way to demonstrate both *IGHG*-alleles, -genotypes, -haplotypes and -diplotypes. RFLP and DNA analyses have been performed, but with only limited success [5]. The IgG allotypes are products of structural genes. They have been extensively studied in humans, rabbits, rats and in the mink [5]. The basis of IgG allotypes in humans is precisely understood at the mole-

cular level by amino acid and codon substitutions IgG1**a* with Asp 356, Leu 358 and IgG1**f* with Lys 214 [5]. The substitutions of the alternative IgG2 allotypes have not yet been elucidated, however. We have previously shown that the alternative IgG2 allotypes have different electrophoretic and fractionation-rates, different half-life times and developmental rates during childhood [6, 7]. There is an advantage to study directly gene-derived proteins quantitatively, as this also describes the activity from the allelic variant of the gene. A serum sample is used for determination of the gene products from six *IGHG* alleles, nine *IGHG* genotypes, four *IGHG* haplotypes (B-cell variants) and 10 *IGHG* diplotypes (individual expressions). The advantage of the technique used is its simplicity, allowing both patient studies and studies of large populations.

Children with atopic manifestations and especially those with a family history of allergy presented particular IgG molecules IgG3**b*, IgG1**f* and especially IgG2**n* from the *IGHG***bf/n* haplotype, indicating that IgG antibodies are also involved, besides IgE. The precise role of IgG antibodies in allergic inflammation is unknown. During the inflammatory process, the variable antigen-binding part of the antibody molecule is combined with different

Table 4. Frequency of immunoglobulin heavy G2 chain (*IGHG2*) genotypes and *IGHG2* alleles as risk factors in atopy subgroups from 61 children with a family history of allergy, clinical allergy and positive skin prick test compared with 21 controls without

Atopy subgroups	Number (%)		<i>P</i> -value	OR	95% CI
	<i>IGHG2</i> genotypes	<i>IGHG2</i> alleles			
Controls	21	42			
* <i>n</i> /* <i>n</i>	2 (10)				
* <i>n</i> /*- <i>n</i>	6 (29)				
- <i>n</i> /- <i>n</i>	13 (62)				
		* <i>n</i>			
		10 (24)			
		*- <i>n</i>			
		32 (76)			
Family history					
Paternal heredity	34	68			
* <i>n</i> /* <i>n</i>	10 (29)				
* <i>n</i> /*- <i>n</i>	15 (44)				
- <i>n</i> /- <i>n</i>	9 (26)				
		* <i>n</i>			
		35 (51)	0.004	3.4	1.4–8.0
		*- <i>n</i>			
		33 (49)			
Maternal heredity	34	68			
* <i>n</i> /* <i>n</i>	7 (21)				
* <i>n</i> /*- <i>n</i>	16 (47)				
- <i>n</i> /- <i>n</i>	11 (32)				
		* <i>n</i>			
		30 (44)	0.031	2.5	1.0–5.9
		*- <i>n</i>			
		38 (56)			
Clinical allergy					
Wheezing	47	94			
* <i>n</i> /* <i>n</i>	14 (30)				
* <i>n</i> /*- <i>n</i>	23 (49)				
- <i>n</i> /- <i>n</i>	10 (21)				
		* <i>n</i>			
		51 (54)	< 0.001	3.7	1.7–8.6
		*- <i>n</i>			
		43 (46)			
Eczema	40	80			
* <i>n</i> /* <i>n</i>	15 (38)				
* <i>n</i> /*- <i>n</i>	16 (40)				
- <i>n</i> /- <i>n</i>	9 (23)				
		* <i>n</i>			
		46 (58)	< 0.001	4.3	1.9–10.0
		*- <i>n</i>			
		34 (43)			
Rhinit	30	60			
* <i>n</i> /* <i>n</i>	10 (33)				
* <i>n</i> /*- <i>n</i>	14 (47)				
- <i>n</i> /- <i>n</i>	6 (20)				
		* <i>n</i>			
		34 (57)	< 0.001	4.2	1.7–10.0
		*- <i>n</i>			
		26 (43)			

constant functional heavy chains: both γ and ϵ chains. In childhood asthma, the IgG1**f* is depressed but the IgG2**n* levels doubled compared with healthy age-matched and *IGHG* gene-matched controls [14]. During childhood, there is a slow developmental rate of IgG2**n* antibodies [8] while in IgE-mediated asthma the opposite seems to be the case.

The antibody production from *IGHG4* and *IGHE*, next to *IGHG2* on chromosome 14q32.3, is activated by IL-4 and IL-13 [18, 19], connecting T helper type 2 (TH2) cell activation to B cell activation. Increased serum level of the IgG2**n* allotype in IgE-mediated childhood asthma in-

dicates a linkage between an activated *IGHG2***n* gene and sensitization. As the *IGHG2* gene, which is situated within 18 kb from *IGHG4* and another 23 kb from *IGHE* on the long arm of chromosome 14q32, is supposed to be in strong linkage disequilibrium [5], it is likely that the *IGHE* gene is influenced by the *IGHG2***n* gene located upstream. Both the *IGHG* genes and the *IGHE* gene on the chromosome 14q32 are found within a range of 250 kb in linkage disequilibrium. The *IGHG* haplotype influences the amounts expressed by single *IGHG* alleles [20].

The serum IgG2**n* allotype is a marker of the *IGHG2***n* allele, either homozygous or heterozygous *IGHG2***n*/*n*

Table 5. Frequency of B cells [immunoglobulin heavy G chain (*IGHG*) haplotypes] B1 ($B^{*bf/n}$), B2 (B^{*bf-n}), B3 (B^{*gan}) and B4 (B^{*ga-n}) as risk factors in atopy subgroups from 61 children (122 haplotypes) with a family history of allergy, clinical allergy and positive skin prick test (SPT) compared with 21 controls without (42 haplotypes)

Atopy subgroups	B cell (<i>IGHG</i>) haplotypes, N (%)	P-value	OR	95% CI
Controls (21)	42			
B1 ($B^{*bf/n}$)	10 (24)			
B2 (B^{*bf-n})	22 (52)			
B3 (B^{*gan})	1 (2)			
B4 (B^{*ga-n})	9 (21)			
Family history of allergy				
Paternal heredity (34)	68			
B1 ($B^{*bf/n}$)	35 (52)	0.004	3.4	1.4–8.0
B2 (B^{*bf-n})	13 (19)	< 0.001	0.2	0.1–0.5
B3 (B^{*gan})	0	NS		
B4 (B^{*ga-n})	20 (29)	NS		
Maternal heredity (34)	68			
B1 ($B^{*bf/n}$)	30 (44)	0.03	2.5	1.1–5.9
B2 (B^{*bf-n})	14 (21)	< 0.001	0.2	0.1–0.5
B3 (B^{*gan})	0	NS		
B4 (B^{*ga-n})	24 (35)	NS		
Clinical allergy				
Wheezing	94			
B1 ($B^{*bf/n}$)	51 (54)	< 0.001	3.8	1.7–8.6
B2 (B^{*bf-n})	12 (12)	< 0.001	0.1	0.05–0.3
B3 (B^{*gan})	0	NS		
B4 (B^{*ga-n})	31 (33)	NS		
Eczema	80			
B1 ($B^{*bf/n}$)	46 (58)	< 0.001	4.3	1.9–10.0
B2 (B^{*bf-n})	12 (15)	< 0.001	0.2	0.1–0.4
B3 (B^{*gan})	0	NS		
B4 (B^{*ga-n})	22 (28)	NS		
Rhinit	60			
B1 ($B^{*bf/n}$)	34 (57)	< 0.001	4.2	1.7–10.0
B2 (B^{*bf-n})	7 (12)	< 0.001	0.1	0.04–0.3
B3 (B^{*gan})	0	NS		
B4 (B^{*ga-n})	19 (32)	NS		
Sensitized (SPT+)				
Tree	42			
B1 ($B^{*bf/n}$)	19 (45)	0.039	2.6	1.0–6.7
B2 (B^{*bf-n})	8 (19)	0.001	0.2	0.1–0.6
B3 (B^{*gan})	0	NS		
B4 (B^{*ga-n})	15 (36)	NS		
Grass	52			
B1 ($B^{*bf/n}$)	29 (47)	0.018	2.8	1.2–6.7
B2 (B^{*bf-n})	11 (18)	< 0.001	0.2	0.1–0.4
B3 (B^{*gan})	1 (2)	NS		
B4 (B^{*ga-n})	21 (34)	NS		
Cat	84			
B1 ($B^{*bf/n}$)	45 (54)	0.001	3.7	1.6–8.5
B2 (B^{*bf-n})	13 (16)	< 0.001	0.2	0.1–0.4
B3 (B^{*gan})	1 (2)	NS		
B4 (B^{*ga-n})	25 (30)	NS		
Dog	44			
B1 ($B^{*bf/n}$)	20 (46)	0.035	2.6	1.1–6.7
B2 (B^{*bf-n})	8 (18)	< 0.001	0.2	0.1–0.5
B3 (B^{*gan})	1 (2)	NS		
B4 (B^{*ga-n})	15 (34)	NS		
Horse	52			

B1 (B ^{*bfn})	27 (52)	0.006	3.5	1.4–8.5
B2 (B ^{*bf-n})	10 (19)	< 0.001	0.2	0.1–0.5
B3 (B ^{*gan})	1 (2)	NS		
B4 (B ^{*ga-n})	14 (27)	NS		

SPT, skin prick test.

(B1/B2-, B1/B4 cells). The *IGHG2*^{*n} allele is also found in the *IGHG*^{*gan} (B3) haplotype, but only very infrequently so (<1–2%). [5, 8]. Thus, the IgG2^{*n} allotype is mainly a marker for the *IGHG*^{*bfn} haplotype (= B1 cells).

Both the *IGHG*^{*bfn} haplotype and the homozygous *IGHG*^{*bfn/bfn} diplotype were frequent in a family history of allergy, but less so in environmental-dependent sensitization and sensitization-dependent clinical symptoms.

In infectious diseases, patients with the *IGHG*^{*bfn/bfn} diplotype (= B1/B1 cells) have a better outcome of bacterial infections as a consequence of a high production of specific bacterial antibodies especially to polysaccharide antigens [21, 22] and a more favourable outcome in autoimmune juvenile chronic arthritis [23]. Furthermore, the *IGHG*^{*bf-n/bf-n} diplotype (= B2/B2 cells), which was infrequent in children with a family history of allergy and atopy, is common in IgG2 deficiency and in common variable immunodeficiency [20, 24]. Infants hospitalized with the most severe respiratory syncytial virus (RSV) exhibited an increased frequency of the B2 cells [25]. The *IGHG*^{*ga-n/ga-n} diplotype (= B4/B4 cells) is associated with symptoms as bronchial reactivity to non-specific stimuli and to a history of recurrent upper and lower respiratory tract infections. It is also found in most patients with IgG3 deficiency, in whom recurrent or chronic bronchitis was the most common finding [13, 20, 26].

There is no single gene in allergy genetics but it is interesting to note that the most replicated genes *IL-4*, *IL-13*, *IL-4RA* and *FCERB1* in asthma genetics are associated with the activation of *IGH* genes, both *IGHG* and *IGHE* [27].

In conclusion, a family history, sensitization and clinical manifestations of allergy are associated with *IGHG* genes, from the *IGHG*^{*bfn} haplotype, expressing IgG3^{*b}, IgG1^{*f} and especially IgG2^{*n} molecules. The strongest association was observed for a family history of allergy. In contrast, there was an inverse relationship between allergy manifestations and the presence of the *IGHG2*^{*-n} allele and the *IGHG*^{*bf-n} haplotype. The *IGHG2*^{*-n} is associated with an increased risk, while the *IGHG2*^{*-n} is associated with a low risk of atopy. As there is no genetic marker for an activated or allergen-responding *IGHE* gene, the serum IgG2^{*n} allotype, from the closely located *IGHG2*^{*n} allele, is a possible genetic marker.

Contributors

V-A Oxelius had the idea for the project, designed the study, conducted the laboratory work and wrote the first version of the report. L Bråbäck was responsible for the clinical part of the study including diagnosis and presenting serum samples from the children. V-A Oxelius, B Björkstén and S Ahlstedt jointly interpreted the data and wrote the report.

Acknowledgements

We thank Ann-Margreth Carlsson for skilful laboratory assistance and Håkan Lövkvist for statistical advice. We thank Professor Juha Kere for valuable discussions and for reviewing the manuscript. The investigation was supported by grants of Lund University, University Hospital of Lund, Asthma and Allergy Association in Sweden and Vårdal Foundation in Sweden. The study was approved by the Institutional Review Board at Umeå University, Sweden.

Conflict of interest statement: None.

References

- 1 Asher MI, Keil U, Anderson HR *et al.* International Study of Asthma and Allergies in Childhood (ISAAC): rationale and methods. *Eur Respir J* 1995; **8**:483–91.
- 2 Weiss KB, Sullivan SD. The health economics of asthma and rhinitis. I. Assessing the economic impact. *J Allergy Clin Immunol* 2001; **107**:3–8.
- 3 Palmer LJ, Cookson WO. Genomic approaches to understanding asthma. *Genome Res* 2000; **10**:1280–7.
- 4 Cookson W. Genetics and genomics of asthma and allergic diseases. *Immunol Rev* 2002; **190**:195–206.
- 5 Grubb R. Human immunoglobulin allotypes and Mendelian polymorphism of the human immunoglobulin genes. In: Oss CJ, Regenmortel MHV., eds. *Immunochemistry*. New York: Marcel Dekker, 1994; 47–68.
- 6 Oxelius V-A. Preparation of IgG subclass allotypes from polyclonal IgG. *Scand J Immunol* 1999; **49**:395–8.
- 7 Oxelius V-A, Eibl MM. Different Gm allotype levels in human intravenous immunoglobulin (IVIg) preparations, survival of foreign Gm allotypes in immunodeficient patients. *Clin Exp Immunol* 1996; **106**:203–7.

- 8 Oxelius V-A, Aurivillius M, Carlsson A-M, Musil K. Serum Gm allotype development during childhood. *Scand J Immunol* 1999; 50:440-6.
- 9 Oxelius V-A. Genetic B cell variation based on immunoglobulin heavy G chain (Gm) genes. *Scand J Immunol* 1999; 49:345-6.
- 10 Oxelius V-A, Carlsson A-M. Quantitation of Gm allotypes. *Scand J Immunol* 1993; 37:143-8.
- 11 Oxelius V-A, Hultquist C, Husby S. Gm allotypes as indicators of non-atopic and atopic bronchial asthma. *Int Arch Allergy Appl Immunol* 1993; 101:66-71.
- 12 Oxelius V-A, Sjöstedt L, Willers S, Löw B. Development of allergy to laboratory animals is associated to particular Gm and HLA genes. *Int Arch Allergy Immunol* 1996; 110:73-8.
- 13 Oxelius V-A, Carlsson A-M, Aurivillius M. Alternative G1m, G2m and G3m allotypes of IGHG genes correlate with atopic and nonatopic pathways of immune regulation in children with bronchial asthma. *Int Arch Allergy Immunol* 1998; 115:215-9.
- 14 Oxelius V-A. Imbalanced switch of the IGHG (immunoglobulin constant heavy G chain) Gm(bfn) genes in atopic childhood asthma. *Allergy* 2000; 55:1063-8.
- 15 Annus T, Björkstén B, Mai X-M *et al*. Wheezing in relation to atopy and environmental factors in Estonian and Swedish schoolchildren. *Clin Exp Allergy* 2001; 31:1846-53.
- 16 Oxelius V-A. Serum IgG and IgG subclass contents in different Gm phenotypes. *Scand J Immunol* 1993; 37:149-53.
- 17 Sarvas H, Rautonen N, Mäkelä O. Allotype associated differences in concentrations of human IgG subclasses. *J Clin Immunol* 1991; 11:39-45.
- 18 Finkelman FD *et al*. IL-4 is required to generate and sustain in vivo IgE responses. *J Immunol* 1988; 141:2331-41.
- 19 Minty A *et al*. Interleukin 13 is a new human lymphokine regulating inflammatory and immune responses. *Nature* 1993; 362:248-50.
- 20 Oxelius V-A. Lack of G2m(n) allotype in IgG subclass deficiency, in IgG2 deficiency together with lack of G1m(a) and G3m(g) and in IgG3 deficiency together with lack of G1m(f) and G3m(b). *Scand J Immunol* 1990; 31:243-7.
- 21 Pandey JP, Fudenberg H, Virella G *et al*. Association between immunoglobulin allotypes and immune responses to Haemophilus influenzae and meningococcus polysaccharides. *Lancet* 1979; i:190-2.
- 22 Ambrosino D, Schiffman G, Gotschlich E *et al*. Correlation between G2m(n) immunoglobulin allotype and human antibody response and susceptibility to polysaccharide encapsulated bacteria. *J Clin Invest* 1985; 75:1935-42.
- 23 Oxelius V-A, Svantesson H, Carlsson A-M Gm phenotype linkage to subsets of juvenile chronic arthritis (JCA). *Scand J Rheum* 1993; 101:66-71.
- 24 Oxelius V-A, Ochs HD. Serum Gm allotype levels in common variable immunodeficiency: Preponderance of homozygous G2m(') on IGHG2. *Exp Clin Immunogenet* 1996; 13:70-7.
- 25 Aurivillius M, Oymar K, Oxelius V-A. Immunoglobulin heavy G2 chain (IGHG2) gene restriction in development of severe respiratory syncytial virus infection. *Acta Paediatr* 2005; 94:414-8.
- 26 Oxelius V-A, Hanson L-Å, Björkander J, Hammarström L, Sjöholm A. IgG3 deficiency: Common in obstructive lung disease. *Monogr Allergy* 1986; 20:106-15.
- 27 Ober C. Perspectives on the past decade of asthma genetics. *J Allergy Clin Immunol* 2005; 116:274-8.