Increasing Comorbidities Suggest that Atopic Dermatitis Is a Systemic Disorder

JID Open

Patrick M. Brunner[1], Jonathan I. Silverberg[2], Emma Guttman-Yassky[3], Amy S. Paller[2], Kenji Kabashima[4], Masayuki Amagai[5], Thomas A. Luger[6], Mette Deleuran[7], Thomas Werfel[8], Kilian Eyerich[9] and Georg Stingl[10], Councilors of the International Eczema Council

Atopic dermatitis comorbidities extend well beyond the march to allergic conditions (food allergy, asthma, allergic rhinitis, allergic conjunctivitis, and eosinophilic esophagitis), suggesting both cutaneous and systemic immune activation. In reviewing atopic dermatitis comorbidities, Councilors of the International Eczema Council found a strong pattern of immune activation in peripheral blood and the propensity to both skin and systemic infections. Associations with cardiovascular, neuropsychiatric, and malignant diseases were increasingly reported, but confirmation of their link with atopic dermatitis requires longitudinal studies. Given the possibility of atopic dermatitis-related systemic immune activation, future investigations of new interventions should concurrently examine the impact on these comorbidities.

Journal of Investigative Dermatology (2016) ■, ■–■; doi:10.1016/j.jid.2016.08.022

INTRODUCTION

The immune-mediated skin disease atopic dermatitis (AD) has long been recognized as closely associated with other atopic conditions. More recent studies have uncovered multiple previously unrecognized disease associations, further highlighting the possibility that AD is a systemic disorder. Councilors of the International Eczema Council, all leaders in AD research, gathered at the Annual European Society for Dermatological Research meeting in 2015 to review the increasingly recognized comorbidities, including controversial associations, that may contribute to the burden of AD.

Before the meeting, presenters searched for AD/eczema studies from 2000 through 2016 using PubMed, Google scholar, Science direct, and ClinicalTrials.gov using keywords such as *atopic dermatitis* or *eczema* combined with *infection, cardiovascular, neuro-psychiatric,* and *malignant.* Topics were then discussed by the group, and teams of Councilors further reviewed the literature from 2000–2016 literature to describe (i) pathogenic concepts of immune activation leading to both overt inflammation and increased susceptibility to infection in AD and (ii) associations with AD derived from epidemiologic studies, in which pathogenic mechanisms are still largely unclear. Selected effects sizes of disease associations for autoimmune, infectious, cardiovascular, malignant, and neuropsychiatric disorders are summarized in Supplementary Table S1 online.

ATOPIC DERMATITIS AND SYSTEMIC IMMUNE DEVIATION

The atopic march

AD usually begins early in life (Czarnowicki et al., 2015a; Eichenfield et al., 2012), and its onset is often followed by the serial occurrence of allergic diseases that represent the atopic march, especially food allergies, asthma, and allergic rhinoconjunctivitis (Schneider et al., 2016; Simpson et al., 2012). This sequence has recently been challenged, suggesting that developmental profiles might be much more heterogeneous (Belgrave et al., 2014). AD patients, often in adulthood, also suffer from increased frequencies of hand eczema and allergic contact dermatitis (Thyssen et al., 2010, 2014), although these conditions are not classically included as components of the atopic march.

Allergic sensitization may result from skin barrier dysfunction, facilitating cutaneous allergen penetration. Loss-of-function mutations in the gene encoding filaggrin, *FLG,* are associated with an increased risk of AD and asthma (Irvine et al., 2011; McAleer and Irvine, 2013) and are present in 15–46% of AD patients (Brown et al.,

[1]The Laboratory for Investigative Dermatology, The Rockefeller University, New York, New York, USA; [2]Department of Dermatology, Northwestern University Feinberg School of Medicine, Chicago, Illinois, USA; [3]Department of Dermatology and the Laboratory for Inflammatory Skin Diseases, Icahn School of Medicine at Mount Sinai, New York, New York, USA; [4]Department of Dermatology, Kyoto University Graduate School of Medicine, Kyoto, Japan; [5]Department of Dermatology, Keio University School of Medicine, Tokyo, Japan; [6]Department of Dermatology, University of Münster, Münster, Germany; [7]Department of Dermato-Venereology, Aarhus University Hospital, Aarhus C, Denmark; [8]Division of Immunodermatology and Allergy Research, Department of Dermatology and Allergy, Hannover Medical School, Hannover, Germany; [9]Department of Dermatology and Allergy, Technical University Munich, Munich, Germany; and [10]Division of Immunology, Allergy and Infectious Diseases, Department of Dermatology, Medical University of Vienna, Vienna, Austria

Correspondence: Amy S. Paller, Department of Dermatology, Northwestern University Feinberg School of Medicine, 676 North St. Clair Avenue, Suite 1600, Chicago, Illinois 60611, USA. E-mail: apaller@northwestern.edu

Abbreviations: AD, atopic dermatitis; CI, confidence interval; OR, odds ratio; Th, T helper

Received 7 April 2016; revised 4 August 2016; accepted 19 August 2016; corrected proof published online XXX

© 2016 The Authors. Published by Elsevier, Inc. on behalf of the Society for Investigative Dermatology. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

2008; Carson et al., 2012). However, T helper (Th) 2 (IL-4/IL-13/IL-25) and Th22 (IL-22) cytokines, which are increased in AD, suppress filaggrin expression in keratinocytes (Gutowska-Owsiak et al., 2011; Howell et al., 2007; Hvid et al., 2011; Pellerin et al., 2013) and may contribute to the barrier deficits in vivo. Thus, immune activation in AD may trigger barrier dysfunction and/or aggravate inherited barrier defects through a positive feedback loop (Leung and Guttman-Yassky, 2014). Subsequent migration of sensitized T cells and/or IgE into the nose and airways facilitates upper and lower airway disease (Kubo et al., 2012; Leung, 2013). The important role of barrier defects in AD has further justified vigorous application of emollients, which likely contribute more than barrier substitution, given that petrolatum application increases expression of barrier proteins and antimicrobial peptides and decreases proinflammatory cytokines (Czarnowicki et al., 2016). Furthermore, preliminary studies suggest that proactive application of emollients to infants at risk for AD, beginning as neonates, reduces AD development and/or severity (Horimukai et al., 2014; Simpson et al., 2014). Longitudinal studies are currently addressing the ability of early emollient application to decrease sensitization to antigens and development of the atopic march.

A recent mechanistic study suggested that Th2/Th22 skewing and epidermal hyperplasia, but not the deficiency of barrier proteins of adult AD, are seen in the skin of infants and toddlers with new-onset disease, suggesting an early role for T-cell skewing (Esaki et al., 2016). Nevertheless, the correlation of high transepidermal water loss at 2 days of age and the occurrence of AD at 1 and 2 years of age suggest an early barrier defect as an initial driver (Kelleher et al., 2015; Kelleher et al., 2016). Additional studies in infants and children are warranted to address potential differences between infant/childhood (i.e., early-onset) and adult (i.e., chronic) disease to better understand disease evolution with age and disease heterogeneity (Garmhausen et al., 2013).

Immune activation beyond the skin

Similar to psoriasis, AD skin lesions are accompanied by epidermal hyperplasia, large T-cell and dendritic cell infiltrates, and increased production of inflammatory products (Guttman-Yassky et al., 2011a, 2011b). Psoriasis has been associated with inflammation in several noncutaneous systems, particularly the joints, gastrointestinal, and cardiovascular systems, and elevations in C-reactive protein are associated with disease severity (Beygi et al., 2014). In contrast to psoriasis, however, AD immune abnormalities also profoundly involve nonlesional skin and the blood compartment (Leung and Guttman-Yassky, 2014; Suarez-Farinas et al., 2011; Tang et al., 2014), although acute phase reactants, such as C-reactive protein, do not tend to be elevated. These observations suggest that the mechanisms of the inflammatory component of AD and psoriasis are different. Indeed, blood from individuals with AD has more polar differentiation of central and effector memory T-cell subsets and significantly higher and more persistent T-cell activation than is seen in psoriasis (Czarnowicki et al., 2015b, 2015c). Ongoing blood T-cell activation is a feature of both adults and children with moderate to severe AD. In pediatric AD, Th2 imbalance is confined to skin homing/CLA$^+$ T cells and does not extend to CD8$^+$ T cells, whereas in adults, Th2 activation extends into systemic/CLA$^-$ CD4$^+$ T cells and CD8$^+$ T cells (Czarnowicki et al., 2015a). Consistent with elevated IgE levels in 80% of AD patients (Weidinger and Novak, 2016), AD blood is characterized by a unique peripheral B-cell signature, and subsets of memory B cells correlate with clinical AD severity (Scoring Atopic Dermatitis score/SCORAD) and IgE levels (Czarnowicki et al., 2016). In peripheral blood, several biomarkers were shown to correlate with disease activity, including systemic cytokine activation (Haeck et al., 2011; Hayashida et al., 2011; Jahnz-Rozyk et al., 2005; Raap et al., 2012; Waxweiler et al., 2011). Alarmins (especially IL-33, but also IL-25 and thymic stromal lymphopoietin) have recently been shown to drive the cutaneous expansion of innate lymphoid cells—particularly innate lymphoid cell-2, producing IL-5 and IL-13—in AD skin (Salimi et al., 2013; Soumelis et al., 2002); this recently described mechanism may play a key early role in systemic activation related to AD (Saunders et al., 2016).

AD has been shown to share susceptibility loci with other immune-mediated diseases such as psoriasis, inflammatory bowel disease, and alopecia areata (Ellinghaus et al., 2013; Weidinger et al., 2013). In a German cohort study, AD was associated with a slightly increased risk for rheumatoid arthritis and inflammatory bowel disease and a decreased risk for type 1 diabetes (relative risk = 0.72), independent of known risk alleles (see Supplementary Table S1) (Schmitt et al., 2016). Prominent Th1 and Th17 responses promoting autoimmunity are known to be present in inflammatory bowel disease and rheumatoid arthritis and are also implicated in the transition to chronic inflammation in AD (Gittler et al., 2012). Autoreactivity is most common in severe and persistent AD (Tang et al., 2012). In a Taiwanese study (Wu et al., 2014), AD was associated with lupus erythematosus (odds ratio [OR] = 1.94, 95% confidence interval [CI] = 1.48−2.54) (see Supplementary Table S1), another disease in which not only Th1 and Th2 but also Th17 responses may play a pathogenic role (Martin et al., 2014; Oh et al., 2011). Associations were more prominent in individuals younger than 18 years of age (OR = 3.02, 95% CI = 1.30−7.03) and in female patients (OR = 2.05, 95% CI = 1.53−2.76) (Wu et al., 2014). Future studies will need to identify potential subsets of AD patients at highest risk and elucidate the mechanism underlying these associations.

Infection

The increased association of AD with more frequent and severe bacterial and viral infections is a minor diagnostic criterion (Hanifin and Rajka, 1980). *Staphylococcus aureus* is cultured from up to 90% of AD lesions (Leyden et al., 1974), and methicillin-resistant *S. aureus* predominates in some patients. Skin microbiome and virome research has expanded our view on the role of infections in AD (Belkaid and Segre, 2014; Kong et al., 2012). Bacterial gene sequencing studies found that

AD flares are associated with increased *S. aureus* and decreased microbial diversity (Kong et al., 2012). Moreover, AD treatment (including intermittent bleach baths and anti-inflammatory medications) decreases *S. aureus* and increases microbial diversity (Kong et al., 2012).

The relative role of *S. aureus* dysbiosis as a result versus instigator of AD is unclear. Dysbiosis (microbial imbalance) may directly affect AD skin inflammation (Kobayashi et al., 2015). *S. aureus* has been shown to induce IL-4/IL-13 and IL-22 cytokines, promoting Th2/Th22 inflammation in AD (Brandt and Sivaprasad, 2011; Niebuhr et al., 2010) and might even affect systemic immune deviation such as food allergies (Jones et al., 2016). Experimentally, occurrence of pruritic AD-like dermatitis in keratinocyte-specific Adam17-deficient mice depends on the presence of *S. aureus* and *Corynebacterium bovis* (Blaydon et al., 2011; Kobayashi et al., 2015), suggesting that bacteria might trigger AD in humans as well. Further, sodium hypochlorite (bleach) baths are not just antiseptic but also reduce AD severity (Huang et al., 2009; Huang et al., 2011), although perhaps in part by direct anti-inflammatory effects (Leung et al., 2013). Increasing microbial diversity improves AD severity in mice (Volz et al., 2014), and *Staphylococcus epidermidis* products have anti-*S. aureus* activity (Cogen et al., 2010; Lai et al., 2010; Naik et al., 2012). There is evidence that cutaneous application of extracts from nonpathogenic bacteria (*Vitreoscilla filiformis*) (Gueniche et al., 2008) and even emollients (Seite et al., 2014) restores skin dysbiosis and thereby may reduce AD severity. Greater severity is associated with higher risk of infection, and both topical and systemic anti-inflammatory interventions reduce staphylococcal colonization (Bath-Hextall et al., 2010; Boguniewicz and Leung, 2010).

Eczema herpeticum, a well-recognized infectious complication, appears in a subset of patients with severe Th2-polarized AD, greater allergen sensitization, and more commonly a history of food allergy and/or asthma (Beck et al., 2009). Eczema coxsackium is a newly recognized complication of coxsackie diseases associated with AD, which can be confused with eczema herpeticum (Mathes et al., 2013).

Children with AD also have a higher prevalence of warts and extracutaneous infections, including streptococcal pharyngitis, upper and lower respiratory infections, recurrent otitis, chickenpox, and urinary tract infections (Silverberg and Silverberg, 2014) (see Supplementary Table S1). Recent reports also suggest a relationship between AD and infective endocarditis (Fukunaga et al., 2013; Patel and Jahnke, 2015) (see Supplementary Table S1).

ATOPIC DERMATITIS AND EPIDEMIOLOGIC ASSOCIATIONS

Cardiovascular risks

Analogous to psoriasis, recent studies have suggested an association between AD, obesity, and cardiovascular disease in both adults and pediatric patients. A meta-analysis of 30 studies showed higher odds of overweight and/or obesity in children and adults with AD in studies from North America and Asia but not Europe (Zhang and Silverberg, 2015) (see Supplementary Table S1). In a study using the 2007–2008 National Survey of Children's Health with 45,897 U.S. adolescents, an increased prevalence of overweight and/or obesity was found in adolescents with AD versus healthy control subjects (Silverberg and Simpson, 2014) (see Supplementary Table S1). Increased waist circumference percentiles (85th percentile or greater: OR = 3.92, 95% CI, 1.50–10.26) and waist-to-height ratios (ratio of 0.5 or greater: OR = 2.22, 95% CI = 1.10–4.50) (see Supplementary Table S1) have also been associated with moderate to severe AD in children (Silverberg et al., 2015). The recognition by practitioners of obesity in many patients with psoriasis, but not with AD, may reflect the higher overall body mass indexes among psoriasis versus AD patients with obesity.

Moderate to severe pediatric AD has significantly higher systolic (OR = 2.94, 95% CI = 1.04–8.36) and diastolic (OR = 3.68, 95% CI = 1.19–11.37) blood pressures and significantly higher odds of systolic blood pressure at or above the 90th percentile (OR = 2.06, 95% CI = 1.09–3.90) (Silverberg et al., 2015) (see Supplementary Table S1), unrelated to excess adiposity. Adults with AD have significantly higher odds of a self-reported history of hypertension, adult-onset diabetes, and hypercholesterolemia, even after controlling for body mass index and/or comorbid allergic disease (Silverberg and Greenland, 2015) (see Supplementary Table S1). Cardiovascular risk is further increased when fatigue and sleep disturbances are present.

Patients with AD may have poor health behaviors that increase their cardiovascular risk. In adults, the odds are higher for smoking at least 100 cigarettes in their lifetime, currently smoking, initiating smoking at a younger age, drinking alcohol, and having reduced vigorous physical activity (Silverberg and Greenland, 2015). Children with severe AD have less vigorous physical activity and sports participation and increased hours of daily television and/or video game use (Strom and Silverberg, 2016) (see Supplementary Table S1). The increased cardiovascular risk factors in AD likely play a prominent role in the development of cardiovascular disease.

Hjuler et al. (2015) found evidence of increased coronary artery calcium score in the cardiac computed tomography angiography 18-segment model of the coronary tree, and mild single-vessel disease, in adults with AD compared with both healthy subjects and patients with psoriasis (see Supplementary Table S1). Patients with psoriasis, however, have more coronary stenosis and three-vessel or left main artery disease compared with patients with AD. Cardiovascular events have also been evaluated using the 2005–2006 National Health and Nutrition Examination Survey and the 2010 and 2012 National Health Interview Surveys. Overall, adults with AD in all three cohorts had significantly higher odds of coronary disease, myocardial infarction, and congestive heart disease; odds of angina, stroke, and peripheral vascular disease were also increased in the National Health Interview Surveys (Silverberg, 2015) (see Supplementary Table S1). AD has been reported to be an independent risk factor for ischemic stroke in a large Taiwanese study (Su et al., 2014). After

multivariate adjustment, patients with AD had a 1.33-fold increased incidence of ischemic stroke, and adjusted hazard ratios in patients with mild, moderate, and severe AD were 1.20, 1.64, and 1.71, respectively (see Supplementary Table S1). However, the increased risk of cardiovascular disease in a Danish cohort was no longer present after adjustment for socioeconomic status, smoking, comorbidities, and medication use (Andersen et al., 2016) (see Supplementary Table S1). In addition, nonfatal stroke was significantly increased in female U.S.-based nurses with history of ever having AD when controlling for age (Drucker et al., 2016) but not after controlling for hypertension, hypercholesterolemia, and diabetes; nonfatal myocardial infarction was not increased. These studies suggest that poor health behaviors and increased cardiovascular risk factors are the major drivers of cardiovascular disease and events in patients with AD, rather than systemic inflammation (see Supplementary Table S1). Longitudinal studies are warranted to assess the potential burden of cardiovascular disease in AD.

Cancer

The relationship between AD and cancer is complex and remains controversial. A recent meta-analysis found small but significantly increased odds of any lymphoma in AD in cohort (relative risk = 1.43; 95% CI = 1.12−1.81), but not case-control (OR = 1.18; 95% CI = 0.94−1.47) studies (Legendre et al., 2015) (see Supplementary Table S1). AD severity was a significant risk factor. Although some studies suggest an association of cutaneous T-cell lymphoma with AD (Arellano et al., 2007, 2009) (see Supplementary Table S1), misdiagnosis of cutaneous T-cell lymphoma as AD in many adults may lead to the erroneous overestimation of the association (Tennis et al., 2011). The mechanisms underlying the possible association of AD and lymphoma are poorly understood, but chronic inflammation, perhaps driven by genetic factors, may promote carcinogenesis, and inappropriate Th2 skewing might divert immunity from tumor-protective Th1 responses (Josephs et al., 2013). Systemic immunosuppressive agents used to treat moderate to severe AD, such as cyclosporine and azathioprine, increase the risk of malignancy (Chockalingam et al., 2015; Simon and Bieber, 2014). Although a regional health record database study found an association between topical calcineurin inhibitors and increased risk of lymphoma (Hui et al., 2009), a nested case-control study with a cohort of almost 300,000 AD patients found no association (Arellano et al., 2007, 2009) (see Supplementary Table S1). Furthermore, a longitudinal cohort of 7,457 children treated with pimecrolimus had no increased malignancy (Margolis et al., 2015). A recent consensus report concluded that use of topical calcineurin inhibitors is effective and safe, even in infants as young as 3 months of age, suggesting that warnings of possible carcinogenicity and restrictions regarding their use under 2 years of age are no longer justified (Luger et al., 2015).

In contrast to a possible increased risk of lymphoma, a meta-analysis found an inverse association between AD and acute lymphoblastic leukemia in six studies and no association with acute myelogenous leukemia in two studies (Linabery et al., 2010) (see Supplementary Table S1). No association was found between AD and bladder cancer (Kim et al., 2000). Recent systematic reviews showed either inverse or no association between AD and pancreatic, brain, and skin cancers (Deckert et al., 2014). A prospective adult Dutch cohort study found an inverse association between AD and actinic keratosis but no association with basal cell carcinoma or squamous cell carcinoma (Hajdarbegovic et al., 2016). A recent population-based case-control study found an inverse association between a history of allergy and early-onset basal cell carcinoma but not with squamous cell carcinoma or melanoma (Cheng et al., 2015) (see Supplementary Table S1). Reasons for such inverse associations between AD and cancer may include increased immune surveillance and eradication of dysregulated cells by systemic inflammation (Fu et al., 2008), as well as the reported IgE-facilitated cross-presentation of tumor antigens by dendritic cells (Platzer et al., 2015).

Neuropsychiatric disorders

AD is associated with higher rates of psychological diseases such as depression, anxiety, suicidal ideation, attention deficit/hyperactivity disorder, and autism spectrum disorder (Arima et al., 2005; Buske-Kirschbaum et al., 2013; Dalgard et al., 2015; Dieris-Hirche et al., 2009; Riis et al., 2016; Sanna et al., 2014; Schmitt et al., 2010; Schut et al., 2014; Yaghmaie et al., 2013) (see Supplementary Table S1). Although most studies do not allow conclusions on temporal relationships and thus causality, a longitudinal Taiwanese study found that the presence of any atopic disease in early childhood (i.e., before 3 years of age) increased the risk of developing attention deficit/hyperactivity disorder (hazard ratio = 1.97) and autism spectrum disorder (hazard ratio = 3.40) later in life, with increasing risks with higher numbers of atopic diseases (Chen et al., 2014). The individual contribution of AD, however, was not reported.

The underlying mechanisms of neuropsychiatric disorders in AD are still unclear, but associations are potentially driven by the harmful effects of AD on quality of life (Chrostowska-Plak et al., 2013; Senra and Wollenberg, 2014), unbearable chronic itch, and sleeplessness (Weisshaar et al., 2008). Clinical investigations have associated depression, anxiety, and autism with elevated levels of proinflammatory (although not Th2) cytokines (Kiecolt-Glaser et al., 2011; Pollak and Yirmiya, 2002; Zimmerman et al., 2005), which theoretically could penetrate the blood-brain barrier (Yarlagadda et al., 2009) and modulate behavior and emotions (Buske-Kirschbaum et al., 2013; Ishiuji et al., 2009; Raison et al., 2006; Rosenkranz et al., 2005). Other chronic health conditions of childhood also increase the risk of emotional and behavioral problems, including attention deficit/hyperactivity disorder (Blackman et al., 2011).

Itch is central to the burden of AD and is significantly correlated with sleeplessness. Itch is also associated with coping behavior and poor quality of life in parents of children with AD (Weisshaar et al., 2008), which should be monitored when evaluating the impact of AD treatment (Metz et al.,

2013; Senra and Wollenberg, 2014). Itch mechanisms in AD are complex and involve cutaneous nerves as well as inflammatory cells and keratinocytes (Kim, 2012; Suarez et al., 2012). There is evidence for a direct correlation between the density of peripheral nerves and AD severity (Sugiura et al., 1997; Tominaga et al., 2009). Furthermore, central mechanisms are speculated to contribute to central sensitization, resulting in a loss of inhibitory control and synaptic recruiting of excitatory interneurons (Stander et al., 2015; Suarez et al., 2012).

OUTLOOK

The growing list of potential cutaneous and systemic AD comorbidities reflects the increasing appreciation of its high burden, both on affected individuals and their families, making the development of better interventions even more urgent. Pathophysiological interconnections are complex, because infections and allergies often develop during active disease, but malignancy or cardiovascular disease may be related to longstanding chronic inflammation or may even occur years after disease activity had abated. Although epidemiologic studies aim to reduce discernible bias and confounder effects (Nijsten and Wakkee, 2009; Schmitt and Weidinger, 2014), statistical associations cannot prove causality. As new treatment approaches for AD are being developed, monitoring of patients longitudinally to determine the impact of these approaches on AD-associated comorbidities, in addition to skin manifestations, will likely reveal mechanisms of disease associations and result in the development of more targeted, or even personalized, treatment approaches.

CONFLICT OF INTEREST

JIS is a consultant for AbbVie, GlaxoSmithKline, Lilly, Medimmune, Proctor & Gamble, Regeneron Sanofi, and Anacor. EGY is a board member for Sanofi Aventis, Regeneron, Stiefel/GlaxoSmithKline, MedImmune, Celgene, Anacor, AnaptysBio, Celsus, Dermira, Galderma, Glenmark, Novartis, Pfizer, Vitae, and Leo Pharma; has received consultancy fees from Regeneron, Sanofi, MedImmune, Celgene, Stiefel/GlaxoSmithKline, Celsus, BMS, Amgen, Drais, AbbVie, Anacor, AnaptysBio, Dermira, Galderma, Glenmark, LEO Pharma, Novartis, Pfizer, Vitae, Mitsubishi Tanabe, and Eli Lilly; and has received research support from Janssen, Regeneron, Celgene, Bristol-Myers Squibb, Novartis, Merck, LEO Pharma, and Dermira. ASP is an investigator for Anacor, Astellas, and LEO Pharma and has received honoraria related to atopic dermatitis consulting from Anacor, Galderma, GlaxoSmithKline-Stiefel, Novartis, Regeneron, and Vitae Pharmaceuticals. KK is in an advisory board for Chugai. MD is an advisor for Meda Pharma, Regeneron, Sanofi, and Pierre Fabre and an investigator for AbbVie, Novartis, Pierre Fabre, and Regeneron. The other authors state no conflict of interest.

ACKNOWLEDGMENTS

We acknowledge other Councilors of the International Eczema Council who participated in the original discussion for their valuable input: Martine Bagot, Dirk Jan Hijnen, Michael Ardern-Jones, Nick Reynolds, Phyllis Spuls, and Alain Taieb.

SUPPLEMENTARY MATERIAL

Supplementary material is linked to the online version of the paper at www.jidonline.org, and at http://dx.doi.org/10.1016/j.jid.2016.08.022.

REFERENCES

Andersen YM, Egeberg A, Gislason GH, Hansen PR, Skov L, Thyssen JP. Risk of myocardial infarction, ischemic stroke, and cardiovascular death in patients with atopic dermatitis. J Allergy Clin Immunol 2016;138:310–2.

Arellano FM, Arana A, Wentworth CE, Fernandez-Vidaurre C, Schlienger RG, Conde E. Lymphoma among patients with atopic dermatitis and/or treated with topical immunosuppressants in the United Kingdom. J Allergy Clin Immunol 2009;123:1111–6.

Arellano FM, Wentworth CE, Arana A, Fernandez C, Paul CF. Risk of lymphoma following exposure to calcineurin inhibitors and topical steroids in patients with atopic dermatitis. J Invest Dermatol 2007;127:808–16.

Arima M, Shimizu Y, Sowa J, Narita T, Nishi I, Iwata N, et al. Psychosomatic analysis of atopic dermatitis using a psychological test. J Dermatol 2005;32:160–8.

Bath-Hextall FJ, Birnie AJ, Ravenscroft JC, Williams HC. Interventions to reduce Staphylococcus aureus in the management of atopic eczema: an updated Cochrane review. Br J Dermatol 2010;163:12–26.

Beck LA, Boguniewicz M, Hata T, Schneider LC, Hanifin J, Gallo R, et al. Phenotype of atopic dermatitis subjects with a history of eczema herpeticum. J Allergy Clin Immunol 2009;124:260–9.

Belgrave DC, Granell R, Simpson A, Guiver J, Bishop C, Buchan I, et al. Developmental profiles of eczema, wheeze, and rhinitis: two population-based birth cohort studies. PLoS Med 2014;11:e1001748.

Belkaid Y, Segre JA. Dialogue between skin microbiota and immunity. Science 2014;346:954–9.

Beygi S, Lajevardi V, Abedini R. C-reactive protein in psoriasis: a review of the literature. J Eur Acad Dermatol Venereol 2014;28:700–11.

Blackman JA, Gurka MJ, Gurka KK, Oliver MN. Emotional, developmental and behavioural co-morbidities of children with chronic health conditions. J Paediatr Child Health 2011;47:742–7.

Blaydon DC, Biancheri P, Di WL, Plagnol V, Cabral RM, Brooke MA, et al. Inflammatory skin and bowel disease linked to ADAM17 deletion. N Engl J Med 2011;365:1502–8.

Boguniewicz M, Leung DY. Recent insights into atopic dermatitis and implications for management of infectious complications. J Allergy Clin Immunol 2010;125:4–13.

Brandt EB, Sivaprasad U. Th2 cytokines and atopic dermatitis. J Clin Cell Immunol 2011;2:110.

Brown SJ, Sandilands A, Zhao Y, Liao H, Relton CL, Meggitt SJ, et al. Prevalent and low-frequency null mutations in the filaggrin gene are associated with early-onset and persistent atopic eczema. J Invest Dermatol 2008;128:1591–4.

Buske-Kirschbaum A, Schmitt J, Plessow F, Romanos M, Weidinger S, Roessner V. Psychoendocrine and psychoneuroimmunological mechanisms in the comorbidity of atopic eczema and attention deficit/hyperactivity disorder. Psychoneuroendocrinology 2013;38:12–23.

Carson CG, Rasmussen MA, Thyssen JP, Menne T, Bisgaard H. Clinical presentation of atopic dermatitis by filaggrin gene mutation status during the first 7 years of life in a prospective cohort study. PLoS One 2012;7:e48678.

Chen MH, Su TP, Chen YS, Hsu JW, Huang KL, Chang WH, et al. Is atopy in early childhood a risk factor for ADHD and ASD? A longitudinal study. J Psychosom Res 2014;77:316–21.

Cheng J, Zens MS, Duell E, Perry AE, Chapman MS, Karagas MR. History of allergy and atopic dermatitis in relation to squamous cell and Basal cell carcinoma of the skin. Cancer Epidemiol Biomarkers Prev 2015;24:749–54.

Chockalingam R, Downing C, Tyring SK. Cutaneous squamous cell carcinomas in organ transplant recipients. J Clin Med 2015;4:1229–39.

Chrostowska-Plak D, Reich A, Szepietowski JC. Relationship between itch and psychological status of patients with atopic dermatitis. J Eur Acad Dermatol Venereol 2013;27:e239–42.

Cogen AL, Yamasaki K, Muto J, Sanchez KM, Crotty Alexander L, Tanios J, et al. Staphylococcus epidermidis antimicrobial delta-toxin (phenol-soluble modulin-gamma) cooperates with host antimicrobial peptides to kill group A *Streptococcus*. PLoS One 2010;5:e8557.

Czarnowicki T, Esaki H, Gonzalez J, Malajian D, Shemer A, Noda S, et al. Early pediatric atopic dermatitis shows only a cutaneous lymphocyte antigen$^+$ TH2/TH1 cell imbalance, whereas adults acquire cutaneous lymphocyte antigen$^+$ TH22/TC22 cell subsets. J Allergy Clin Immunol 2015a;136:941–51.

Czarnowicki T, Gonzalez J, Shemer A, Malajian D, Xu H, Zheng X, et al. Severe atopic dermatitis is characterized by selective expansion of circulating TH2/TC2 and TH22/TC22, but not TH17/TC17, cells within the skin-homing T-cell population. J Allergy Clin Immunol 2015b;136:104–15.

Czarnowicki T, Malajian D, Khattri S, da Rosa JC, Dutt R, Finney R, et al. Petrolatum: barrier

repair and antimicrobial responses underlying this "inert" moisturizer. J Allergy Clin Immunol 2016;137:1091–102.

Czarnowicki T, Malajian D, Shemer A, Fuentes-Duculan J, Gonzalez J, Suarez-Farinas M, et al. Skin-homing and systemic T-cell subsets show higher activation in atopic dermatitis versus psoriasis. J Allergy Clin Immunol 2015c;136:208–11.

Dalgard FJ, Gieler U, Tomas-Aragones L, Lien L, Poot F, Jemec GB, et al. The psychological burden of skin diseases: a cross-sectional multicenter study among dermatological outpatients in 13 European countries. J Invest Dermatol 2015;135:984–91.

Deckert S, Kopkow C, Schmitt J. Nonallergic comorbidities of atopic eczema: an overview of systematic reviews. Allergy 2014;69:37–45.

Dieris-Hirche J, Gieler U, Kupfer JP, Milch WE. Suicidal ideation, anxiety and depression in adult patients with atopic dermatitis. Hautarzt 2009;60:641–6.

Drucker AM, Li WQ, Cho E, Li T, Sun Q, Camargo CA, Jr., et al. Atopic dermatitis is not independently associated with non-fatal myocardial infarction or stroke among US women [e-pub ahead of print]. Allergy 2016;71:1496–500.

Eichenfield LF, Ellis CN, Mancini AJ, Paller AS, Simpson EL. Atopic dermatitis: epidemiology and pathogenesis update. Semin Cutan Med Surg 2012;31:S3–5.

Ellinghaus D, Baurecht H, Esparza-Gordillo J, Rodriguez E, Matanovic A, Marenholz I, et al. High-density genotyping study identifies four new susceptibility loci for atopic dermatitis. Nat Genet 2013;45:808–12.

Esaki H, Brunner PM, Renert-Yuval Y, Czarnowicki T, Huynh T, Tran G, et al. Early onset pediatric atopic dermatitis is Th2, but also Th17 polarized in skin. J Allergy Clin Immunol 2016;136(5 Suppl. 1):S7.

Fu SL, Pierre J, Smith-Norowitz TA, Hagler M, Bowne W, Pincus MR, et al. Immunoglobulin E antibodies from pancreatic cancer patients mediate antibody-dependent cell-mediated cytotoxicity against pancreatic cancer cells. Clin Exp Immunol 2008;153:401–9.

Fukunaga N, Okada Y, Konishi Y, Murashita T, Koyama T. Pay attention to valvular disease in the presence of atopic dermatitis. Circ J 2013;77:1862–6.

Garmhausen D, Hagemann T, Bieber T, Dimitriou I, Fimmers R, Diepgen T, et al. Characterization of different courses of atopic dermatitis in adolescent and adult patients. Allergy 2013;68:498–506.

Gittler JK, Shemer A, Suarez-Farinas M, Fuentes-Duculan J, Gulewicz KJ, Wang CQ, et al. Progressive activation of T(H)2/T(H)22 cytokines and selective epidermal proteins characterizes acute and chronic atopic dermatitis. J Allergy Clin Immunol 2012;130:1344–54.

Gueniche A, Knaudt B, Schuck E, Volz T, Bastien P, Martin R, et al. Effects of nonpathogenic gram-negative bacterium Vitreoscilla filiformis lysate on atopic dermatitis: a prospective, randomized, double-blind, placebo-controlled clinical study. Br J Dermatol 2008;159:1357–63.

Gutowska-Owsiak D, Schaupp AL, Salimi M, Taylor S, Ogg GS. Interleukin-22 downregulates filaggrin expression and affects expression of profilaggrin processing enzymes. Br J Dermatol 2011;165:492–8.

Guttman-Yassky E, Nograles KE, Krueger JG. Contrasting pathogenesis of atopic dermatitis and psoriasis—part I: clinical and pathologic concepts. J Allergy Clin Immunol 2011a;127:1110–8.

Guttman-Yassky E, Nograles KE, Krueger JG. Contrasting pathogenesis of atopic dermatitis and psoriasis—part II: immune cell subsets and therapeutic concepts. J Allergy Clin Immunol 2011b;127:1420–32.

Haeck IM, Knol MJ, Ten Berge O, van Velsen SG, de Bruin-Weller MS, Bruijnzeel-Koomen CA. Enteric-coated mycophenolate sodium versus cyclosporin A as long-term treatment in adult patients with severe atopic dermatitis: a randomized controlled trial. J Am Acad Dermatol 2011;64:1074–84.

Hajdarbegovic E, Blom H, Verkouteren JA, Hofman A, Hollestein L, Nijsten T. Atopic dermatitis is not associated with actinic keratosis: cross-sectional results from the Rotterdam study. Br J Dermatol 2016;175:89–94.

Hanifin JM, Rajka G. Diagnostic features of atopic dermatitis. Acta Derm Venereol (Stockh) 1980;92:44–7.

Hayashida S, Uchi H, Takeuchi S, Esaki H, Moroi Y, Furue M. Significant correlation of serum IL-22 levels with CCL17 levels in atopic dermatitis. J Dermatol Sci 2011;61:78–9.

Hjuler KF, Bottcher M, Vestergaard C, Deleuran M, Raaby L, Botker HE, et al. Increased prevalence of coronary artery disease in severe psoriasis and severe atopic dermatitis. Am J Med 2015;128:1325–34.

Horimukai K, Morita K, Narita M, Kondo M, Kitazawa H, Nozaki M, et al. Application of moisturizer to neonates prevents development of atopic dermatitis. J Allergy Clin Immunol 2014;134:824–30.

Howell MD, Kim BE, Gao P, Grant AV, Boguniewicz M, Debenedetto A, et al. Cytokine modulation of atopic dermatitis filaggrin skin expression. J Allergy Clin Immunol 2007;120:150–5.

Huang JT, Abrams M, Tlougan B, Rademaker A, Paller AS. Treatment of *Staphylococcus aureus* colonization in atopic dermatitis decreases disease severity. Pediatrics 2009;123:e808–14.

Huang JT, Rademaker A, Paller AS. Dilute bleach baths for Staphylococcus aureus colonization in atopic dermatitis to decrease disease severity. Arch Dermatol 2011;147:246–7.

Hui RL, Lide W, Chan J, Schottinger J, Yoshinaga M, Millares M. Association between exposure to topical tacrolimus or pimecrolimus and cancers. Ann Pharmacother 2009;43:1956–63.

Hvid M, Vestergaard C, Kemp K, Christensen GB, Deleuran B, Deleuran M. IL-25 in atopic dermatitis: a possible link between inflammation and skin barrier dysfunction? J Invest Dermatol 2011;131:150–7.

Irvine AD, McLean WH, Leung DY. Filaggrin mutations associated with skin and allergic diseases. N Engl J Med 2011;365:1315–27.

Ishiuji Y, Coghill RC, Patel TS, Oshiro Y, Kraft RA, Yosipovitch G. Distinct patterns of brain activity evoked by histamine-induced itch reveal an association with itch intensity and disease severity in atopic dermatitis. Br J Dermatol 2009;161:1072–80.

Jahnz-Rozyk K, Targowski T, Paluchowska E, Owczarek W, Kucharczyk A. Serum thymus and activation-regulated chemokine, macrophage-derived chemokine and eotaxin as markers of severity of atopic dermatitis. Allergy 2005;60:685–8.

Jones AL, Curran-Everett D, Leung DY. Food allergy is associated with Staphylococcus aureus colonization in children with atopic dermatitis. J Allergy Clin Immunol 2016;137:1247–8.

Josephs DH, Spicer JF, Corrigan CJ, Gould HJ, Karagiannis SN. Epidemiological associations of allergy, IgE and cancer. Clin Exp Allergy 2013;43:1110–23.

Kelleher M, Dunn-Galvin A, Hourihane JO, Murray D, Campbell LE, McLean WH, et al. Skin barrier dysfunction measured by transepidermal water loss at 2 days and 2 months predates and predicts atopic dermatitis at 1 year. J Allergy Clin Immunol 2015;135:930–5.

Kelleher MM, Dunn-Galvin A, Gray C, Murray DM, Kiely M, Kenny L, et al. Skin barrier impairment at birth predicts food allergy at 2 years of age. J Allergy Clin Immunol 2016;137:1111–6.

Kiecolt-Glaser JK, Belury MA, Andridge R, Malarkey WB, Glaser R. Omega-3 supplementation lowers inflammation and anxiety in medical students: a randomized controlled trial. Brain Behav Immun 2011;25:1725–34.

Kim K. Neuroimmunological mechanism of pruritus in atopic dermatitis focused on the role of serotonin. Biomol Ther (Seoul) 2012;20:506–12.

Kim WJ, Lee HL, Lee SC, Kim YT, Kim H. Polymorphisms of *N*-acetyltransferase 2, glutathione S-transferase mu and theta genes as risk factors of bladder cancer in relation to asthma and tuberculosis. J Urol 2000;164:209–13.

Kobayashi T, Glatz M, Horiuchi K, Kawasaki H, Akiyama H, Kaplan DH, et al. Dysbiosis and *Staphylococcus aureus* colonization drives inflammation in atopic dermatitis. Immunity 2015;42:756–66.

Kong HH, Oh J, Deming C, Conlan S, Grice EA, Beatson MA, et al. Temporal shifts in the skin microbiome associated with disease flares and treatment in children with atopic dermatitis. Genome Res 2012;22:850–9.

Kubo A, Nagao K, Amagai M. Epidermal barrier dysfunction and cutaneous sensitization in atopic diseases. J Clin Invest 2012;122:440–7.

Lai Y, Cogen AL, Radek KA, Park HJ, Macleod DT, Leichtle A, et al. Activation of TLR2 by a small molecule produced by *Staphylococcus epidermidis* increases antimicrobial defense against bacterial skin infections. J Invest Dermatol 2010;130:2211–21.

Legendre L, Barnetche T, Mazereeuw-Hautier J, Meyer N, Murrell D, Paul C. Risk of lymphoma in patients with atopic dermatitis and the role of topical treatment: a systematic review and meta-analysis. J Am Acad Dermatol 2015;72:992–1002.

Leung DY. New insights into atopic dermatitis: role of skin barrier and immune dysregulation. Allergol Int 2013;62:151–61.

Leung DY, Guttman-Yassky E. Deciphering the complexities of atopic dermatitis: shifting paradigms in treatment approaches. J Allergy Clin Immunol 2014;134:769–79.

Leung TH, Zhang LF, Wang J, Ning S, Knox SJ, Kim SK. Topical hypochlorite ameliorates NF-kappaB-mediated skin diseases in mice. J Clin Invest 2013;123:5361–70.

Leyden JJ, Marples RR, Kligman AM. *Staphylococcus aureus* in the lesions of atopic dermatitis. Br J Dermatol 1974;90:525–30.

Linabery AM, Jurek AM, Duval S, Ross JA. The association between atopy and childhood/adolescent leukemia: a meta-analysis. Am J Epidemiol 2010;171:749–64.

Luger T, Boguniewicz M, Carr W, Cork M, Deleuran M, Eichenfield L, et al. Pimecrolimus in atopic dermatitis: consensus on safety and the need to allow use in infants. Pediatr Allergy Immunol 2015;26:306–15.

Margolis DJ, Abuabara K, Hoffstad OJ, Wan J, Raimondo D, Bilker WB. Association between malignancy and topical use of pimecrolimus. JAMA Dermatol 2015;151:594–9.

Martin JC, Baeten DL, Josien R. Emerging role of IL-17 and Th17 cells in systemic lupus erythematosus. Clin Immunol 2014;154:1–12.

Mathes EF, Oza V, Frieden IJ, Cordoro KM, Yagi S, Howard R, et al. "Eczema coxsackium" and unusual cutaneous findings in an enterovirus outbreak. Pediatrics 2013;132:e149–57.

McAleer MA, Irvine AD. The multifunctional role of filaggrin in allergic skin disease. J Allergy Clin Immunol 2013;131:280–91.

Metz M, Wahn U, Gieler U, Stock P, Schmitt J, Blume-Peytavi U. Chronic pruritus associated with dermatologic disease in infancy and childhood: update from an interdisciplinary group of dermatologists and pediatricians. Pediatr Allergy Immunol 2013;24:527–39.

Naik S, Bouladoux N, Wilhelm C, Molloy MJ, Salcedo R, Kastenmuller W, et al. Compartmentalized control of skin immunity by resident commensals. Science 2012;337:1115–9.

Niebuhr M, Scharonow H, Gathmann M, Mamerow D, Werfel T. Staphylococcal exotoxins are strong inducers of IL-22: a potential role in atopic dermatitis. J Allergy Clin Immunol 2010;126:1176–83.

Nijsten T, Wakkee M. Complexity of the association between psoriasis and comorbidities. J Invest Dermatol 2009;129:1601–3.

Oh SH, Roh HJ, Kwon JE, Lee SH, Kim JY, Choi HJ, et al. Expression of interleukin-17 is correlated with interferon-alpha expression in cutaneous lesions of lupus erythematosus. Clin Exp Dermatol 2011;36:512–20.

Patel D, Jahnke MN. Serious complications from *Staphylococcal aureus* in atopic dermatitis. Pediatr Dermatol 2015;32:792–6.

Pellerin L, Henry J, Hsu CY, Balica S, Jean-Decoster C, Mechin MC, et al. Defects of filaggrin-like proteins in both lesional and nonlesional atopic skin. J Allergy Clin Immunol 2013;131:1094–102.

Platzer B, Elpek KG, Cremasco V, Baker K, Stout MM, Schultz C, et al. IgE/FcεRI-mediated antigen cross-presentation by dendritic cells enhances anti-tumor immune responses [e-pub ahead of print] Cell Rep 2015. http://dx.doi.org/10.1016/j.celrep.2015.02.015 (accessed September 21, 2016).

Pollak Y, Yirmiya R. Cytokine-induced changes in mood and behaviour: implications for 'depression due to a general medical condition', immunotherapy and antidepressive treatment. Int J Neuropsychopharmacol 2002;5:389–99.

Raap U, Weissmantel S, Gehring M, Eisenberg AM, Kapp A, Folster-Holst R. IL-31 significantly correlates with disease activity and Th2 cytokine levels in children with atopic dermatitis. Pediatr Allergy Immunol 2012;23:285–8.

Raison CL, Capuron L, Miller AH. Cytokines sing the blues: inflammation and the pathogenesis of depression. Trends Immunol 2006;27:24–31.

Riis JL, Vestergaard C, Deleuran MS, Olsen M. Childhood atopic dermatitis and risk of attention deficit/hyperactivity disorder: a cohort study. J Allergy Clin Immunol 2016;138:608–10.

Rosenkranz MA, Busse WW, Johnstone T, Swenson CA, Crisafi GM, Jackson MM, et al. Neural circuitry underlying the interaction between emotion and asthma symptom exacerbation. Proc Natl Acad Sci USA 2005;102:13319–24.

Salimi M, Barlow JL, Saunders SP, Xue L, Gutowska-Owsiak D, Wang X, et al. A role for IL-25 and IL-33-driven type-2 innate lymphoid cells in atopic dermatitis. J Exp Med 2013;210:2939–50.

Sanna L, Stuart AL, Pasco JA, Jacka FN, Berk M, Maes M, et al. Atopic disorders and depression: findings from a large, population-based study. J Affect Disord 2014;155:261–5.

Saunders SP, Moran T, Floudas A, Wurlod F, Kaszlikowska A, Salimi M, et al. Spontaneous atopic dermatitis is mediated by innate immunity, with the secondary lung inflammation of the atopic march requiring adaptive immunity. J Allergy Clin Immunol 2016;137:482–91.

Schmitt J, Buske-Kirschbaum A, Roessner V. Is atopic disease a risk factor for attention-deficit/hyperactivity disorder? A systematic review. Allergy 2010;65:1506–24.

Schmitt J, Schwarz K, Baurecht H, Hotze M, Folster-Holst R, Rodriguez E, et al. Atopic dermatitis is associated with an increased risk for rheumatoid arthritis and inflammatory bowel disease, and a decreased risk for type 1 diabetes. J Allergy Clin Immunol 2016;137:130–6.

Schmitt J, Weidinger S. Alternative models of comorbidity: a framework for the interpretation of epidemiological association studies. J Invest Dermatol 2014;134:303–7.

Schneider L, Hanifin J, Boguniewicz M, Eichenfield LF, Spergel JM, Dakovic R, et al. Study of the atopic march: development of atopic co-morbidities. Pediatr Dermatol 2016;33:388–98.

Schut C, Bosbach S, Gieler U, Kupfer J. Personality traits, depression and itch in patients with atopic dermatitis in an experimental setting: a regression analysis. Acta Derm Venereol 2014;94:20–5.

Seite S, Flores GE, Henley JB, Martin R, Zelenkova H, Aguilar L, et al. Microbiome of affected and unaffected skin of patients with atopic dermatitis before and after emollient treatment. J Drugs Dermatol 2014;13:1365–72.

Senra MS, Wollenberg A. Psychodermatological aspects of atopic dermatitis. Br J Dermatol 2014;170(Suppl. 1):38–43.

Silverberg JI. Association between adult atopic dermatitis, cardiovascular disease, and increased heart attacks in three population-based studies. Allergy 2015;70:1300–8.

Silverberg JI, Becker L, Kwasny M, Menter A, Cordoro KM, Paller AS. Central obesity and high blood pressure in pediatric patients with atopic dermatitis. JAMA Dermatol 2015;151:144–52.

Silverberg JI, Greenland P. Eczema and cardiovascular risk factors in 2 US adult population studies. J Allergy Clin Immunol 2015;135:721–8.

Silverberg JI, Silverberg NB. Childhood atopic dermatitis and warts are associated with increased risk of infection: a US population-based study. J Allergy Clin Immunol 2014;133:1041–7.

Silverberg JI, Simpson EL. Association between obesity and eczema prevalence, severity and poorer health in US adolescents. Dermatitis 2014;25:172–81.

Simon D, Bieber T. Systemic therapy for atopic dermatitis. Allergy 2014;69:46–55.

Simpson EL, Chalmers JR, Hanifin JM, Thomas KS, Cork MJ, McLean WH, et al. Emollient enhancement of the skin barrier from birth offers effective atopic dermatitis prevention. J Allergy Clin Immunol 2014;134:818–23.

Simpson EL, Eichenfield LF, Ellis CN, Mancini AJ, Paller AS. Current issues in atopic comorbidities and preventing the atopic march. Semin Cutan Med Surg 2012;31:S6–9.

Soumelis V, Reche PA, Kanzler H, Yuan W, Edward G, Homey B, et al. Human epithelial cells trigger dendritic cell mediated allergic inflammation by producing TSLP. Nat Immunol 2002;3:673–80.

Stander S, Pogatzki-Zahn E, Stumpf A, Fritz F, Pfleiderer B, Ritzkat A, et al. Facing the challenges of chronic pruritus: a report from a multidisciplinary medical itch centre in Germany. Acta Derm Venereol 2015;95:266–71.

Strom M, Silverberg JI. Associations of physical activity and sedentary behavior with atopic disease in US children. J Pediatr 2016;174:247–53.

Su VY, Chen TJ, Yeh CM, Chou KT, Hung MH, Chu SY, et al. Atopic dermatitis and risk of ischemic stroke: a nationwide population-based study. Ann Med 2014;46:84–9.

Suarez AL, Feramisco JD, Koo J, Steinhoff M. Psychoneuroimmunology of psychological stress and atopic dermatitis: pathophysiologic and therapeutic updates. Acta Derm Venereol 2012;92:7–15.

Suarez-Farinas M, Tintle SJ, Shemer A, Chiricozzi A, Nograles K, Cardinale I, et al. Nonlesional atopic dermatitis skin is characterized by broad terminal differentiation defects and variable immune abnormalities. J Allergy Clin Immunol 2011;127:954–64.

Sugiura H, Omoto M, Hirota Y, Danno K, Uehara M. Density and fine structure of peripheral nerves in various skin lesions of atopic dermatitis. Arch Dermatol Res 1997;289:125–31.

Tang TS, Bieber T, Williams HC. Does "autoreactivity" play a role in atopic dermatitis? J Allergy Clin Immunol 2012;129:1209–15.

Tang TS, Bieber T, Williams HC. Are the concepts of induction of remission and treatment of subclinical inflammation in atopic dermatitis clinically useful? J Allergy Clin Immunol 2014;133:1615–25.

Tennis P, Gelfand JM, Rothman KJ. Evaluation of cancer risk related to atopic dermatitis and use of topical calcineurin inhibitors. Br J Dermatol 2011;165:465–73.

Thyssen JP, Johansen JD, Linneberg A, Menne T. The epidemiology of hand eczema in the general population–prevalence and main findings. Contact Dermatitis 2010;62:75–87.

Thyssen JP, McFadden JP, Kimber I. The multiple factors affecting the association between atopic dermatitis and contact sensitization. Allergy 2014;69:28–36.

Tominaga M, Ogawa H, Takamori K. Histological characterization of cutaneous nerve fibers containing gastrin-releasing peptide in NC/Nga mice: an atopic dermatitis model. J Invest Dermatol 2009;129:2901–5.

Volz T, Skabytska Y, Guenova E, Chen KM, Frick JS, Kirschning CJ, et al. Nonpathogenic bacteria alleviating atopic dermatitis inflammation induce IL-10-producing dendritic cells and regulatory Tr1 cells. J Invest Dermatol 2014;134:96–104.

Waxweiler WT, Agans R, Morrell DS. Systemic treatment of pediatric atopic dermatitis with azathioprine and mycophenolate mofetil. Pediatr Dermatol 2011;28:689–94.

Weidinger S, Novak N. Atopic dermatitis. Lancet 2016;387:1109–22.

Weidinger S, Willis-Owen SA, Kamatani Y, Baurecht H, Morar N, Liang L, et al. A genome-wide association study of atopic dermatitis identifies loci with overlapping effects on asthma and psoriasis. Hum Mol Genet 2013;22:4841–56.

Weisshaar E, Diepgen TL, Bruckner T, Fartasch M, Kupfer J, Lob-Corzilius T, et al. Itch intensity evaluated in the German Atopic Dermatitis Intervention Study (GADIS): correlations with quality of life, coping behaviour and Scoring Atopic Dermatitis severity in 823 children. Acta Derm Venereol 2008;88:234–9.

Wu LC, Hwang CY, Chung PI, Hua TC, Chen YD, Chu SY, et al. Autoimmune disease comorbidities in patients with atopic dermatitis: a nationwide case-control study in Taiwan. Pediatr Allergy Immunol 2014;25:586–92.

Yaghmaie P, Koudelka CW, Simpson EL. Mental health comorbidity in patients with atopic dermatitis. J Allergy Clin Immunol 2013;131:428–33.

Yarlagadda A, Alfson E, Clayton AH. The blood brain barrier and the role of cytokines in neuropsychiatry. Psychiatry (Edgmont) 2009;6:18–22.

Zhang A, Silverberg JI. Association of atopic dermatitis with being overweight and obese: a systematic review and metaanalysis. J Am Acad Dermatol 2015;72:606–16.

Zimmerman AW, Jyonouchi H, Comi AM, Connors SL, Milstien S, Varsou A, et al. Cerebrospinal fluid and serum markers of inflammation in autism. Pediatr Neurol 2005;33:195–201.

This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. To view a copy of this license, visit http://creativecommons.org/licenses/by-nc-nd/4.0/

Supplementary Table 1. Effect sizes of observed evidence of potential comorbidities.

First author, year	Potential comorbidity	Data source	Study design	Age group	Definition of AD	Definition of comorbidity	Findings
(Schmitt *et al.*, 2016)	Rheumatoid arthritis	Allgemeine Ortskrankenkasse Saxony database	German population-based administrative health care database	0-40 years	ICD-10 codes	ICD-10 codes	Increased risk of RA in AD (risk ratio [95% CI]: 1.72 [1.25-2.37]).
	Inflammatory bowel disease						Increased risk of Crohn's disease (1.34 [1.11-1.61]) and ulcerative colitis (1.25 [1.03-1.53]) in AD.
	Type 1 diabetes						Decreased risk of type 1 diabetes (0.72 [0.53-0.998]) in AD after adjusting for health care utilization.
(Wu *et al.*, 2014)	Autoimmune disease	National Health Insurance Research Database	Taiwan population-based administrative health care database	All ages	ICD-9 codes	ICD-9 codes	Higher odds of systemic lupus erythematosus in AD (adjusted OR [95% CI]: 1.94 [1.48–2.54]). Odds were higher in females and children. No associations with other autoimmune diseases.
(Silverberg and Silverberg, 2014)	Warts	National Health Interview Survey	US population-based cohort study	0-17 years	Caregiver report	Caregiver report	AD *without* other atopic disease was associated with lower odds of warts (0.91 [0.90-0.92]). AD *with* other atopic disease was associated with higher odds of warts (1.83 [1.82-1.84])
	Streptococcal pharyngitis						AD with or without other atopic disease was associated with higher odds of strep throat (adjusted OR [95% CI]: 1.47 [1.46-1.47] and 1.56 [1.54-1.57]).
	Upper respiratory infection						AD with or without other atopic disease was associated with higher odds of head or chest cold

							(1.47 [1.46-1.47] and 1.48 [1.47-1.48]).
	Influenza or pneumonia						AD with or without other atopic disease was associated with higher odds of influenza or pneumonia (1.24 [1.23-1.24] and 1.67 [1.66-1.67]).
	Sinusitis						AD with or without other atopic disease was associated with higher odds of sinus infections (2.18 [2.18-2.19] and 2.59 [2.58-2.60]).
	Recurrent otitis media						AD with or without other atopic disease was associated with higher odds of recurrent ear infections (1.41 [1.40-1.42] and 1.33 [1.32-1.34]).
	Chickenpox/ varicella						AD with or without other atopic disease was associated with higher odds of chickenpox (1.33 [1.32-1.33] and 1.19 [1.19-1.20]).
	Urinary tract infections						AD with or without other atopic disease was associated with higher odds of urinary tract infections (1.72 [1.70-1.73] and 3.12 [10-3.14]).
(Fukunaga *et al.*, 2013)	Infective endocarditis	Kobe City Medical Center General Hospital	Retrospective review	All ages	Japanese Dermatological Association criteria	Not specified	Staphylococcal aureus more prevalent in infective endocarditis with AD than without AD (88% vs. 14%).
(Zhang and Silverberg, 2015)	Obesity	Meta-analysis	Observational studies (n=30)	All ages	Multiple	BMI	Increased odds of obesity in AD (random-effects OR [95% CI]: 1.47 [1.21-1.79]).

(Silverberg and Simpson, 2014)	Obesity	National Survey of Children's Health	Cohort study	0-17 years	Caregiver report of healthcare diagnosed AD	Caregiver report of height, weight	Increased odds of BMI ≥50% (OR [95% CI]: 1.61 [1.32-1.97]) and ≥95% (1.46 [1.15-1.86]) in AD. Even higher odds of BMI ≥50% (2.46 [1.73-3.51]) and ≥95% (2.95 [1.73-3.51]) in moderate-severe AD.
(Silverberg *et al.*, 2015)	Obesity	Multicenter pediatric dermatology practices	Case-control study	4-17 years	Hanifin & Rajka criteria	BMI and waist circumference Blood pressure	Increased odds of BMI ≥97% (adjusted OR [95% CI]: 2.64 [1.15-6.06]), International Obesity Task Force obesity cutoffs (2.38 [1.06-5.34]), waist circumference ≥85% (3.92 [1.50-10.26]) and waist to height ratio ≥0.5 (2.22 [1.10-4.50]) in moderate-severe AD.
	Hypertension						Higher systolic (2.94 [1.04-8.36]) and diastolic BP (3.68 [1.19-11.37]) in moderate-severe AD, particularly systolic BP ≥90% (2.06 [1.09-3.90]).
(Silverberg and Greenland, 2015)	Hypertension	National Health Interview Survey	US population-based cohort study	18-85 years	Self-report	Self-report	Increased odds of hypertension in AD (1.48 [1.18-1.85]).
	Prediabetes						Increased odds of prediabetes in AD (1.71 [1.19-2.45]).
	Adult onset diabetes						Increased odds of adult onset diabetes in AD in unadjusted models (1.41 [1.18-1.68]) but not models controlling for socio-demographics (1.41 [1.18-1.68]).
	Obesity (BMI>35)						Increased odds of obesity in AD (1.54 [1.16-2.05])
	Hypercholesterolemia						Increased odds of high cholesterol in AD in unadjusted models (1.36 [1.21-1.52]) but not models controlling for socio-demographics (1.14 [0.92 -

							1.43]).
(Strom and Silverberg, 2016)	Sedentary behavior	National Survey of Children's Health	US Population-based cohort study	0-17 years	Caregiver report	Caregiver report	Lower odds of ≥1 days of vigorous physical activity in severe AD (0.39 [0.19-0.78]) Lower odds of sports participation in the past year in moderate (0.76 [0.57-0.99]) and severe (0.45 [0.28-0.73]) AD.
(Hjuler *et al.*, 2015)	Coronary artery disease	Aarhus University Hospital	Case-control study	≥18 years	Hanifin & Rajka criteria	Cardiac computed tomography	Higher rates of coronary artery calcium score >0 in AD than psoriasis or controls (45.2% vs. 29.8% or 15.2%). Higher rates of coronary plaques in AD than psoriasis or controls (48.1% vs. 38.2% or 21.2%).
(Silverberg, 2015)	Coronary artery disease	National Health and Nutrition Examination Survey 2005-2006, National Health Information Surveys 2010 and 2012	3 US population based surveys	≥18 years	Self-report	Self-report	Higher odds of CAD in AD in 3 out of 3 studies (OR: 1.36-2.57).
	Myocardial infarction						Higher odds of heart attack in AD in 3 out of 3 studies (OR: 1.33-2.59).
	Congestive heart failure						Higher odds of congestive heart failure in AD in NHANES 2005-2006 (2.37 [1.27-4.41]).
	Stroke						Higher odds of stroke in 2 out of 3 studies (OR: 1.61-1.63).
	Peripheral vascular disease						Higher odds of poor circulation in AD in NHIS 2012 (1.92 [1.67-2.21]).
(Su *et al.*, 2014)	Stroke	Taiwan's National Health Insurance program	Case-control study	≥20 years	ICD-9 codes	ICD-9 codes	Increased risk of stroke in AD (1.33 [1.12-1.59]), overall, and mild (1.20 [1.00–1.45]), moderate (1.64 [1.23–2.19]) and severe AD (1.71 [1.15 – 2.56]), in particular.
(Andersen *et al.*, 2016)	Myocardial infarction	Danish National Registry	Danish population-based	≥15 years	ICD-10 codes	ICD-10 codes	Marginally increased risk of MI in severe AD in age- and sex-adjusted models (1,39 [0.95-2.03]), but not in

			administrative health care database; severity based on use of systemic medication (severe) in hospital-based patients				fully adjusted models (1.06 [0.72-1.56].
	Stroke						Increased risk of stroke in severe AD in age- and sex-adjusted models (1,51 [1.08-2.10]), but not in fully adjusted models (1.19 [0.85-1.65].
	Cardiovascular death						Increased risk of cardiovascular death in severe AD in age- and sex-adjusted models (1,53 [1.23-1.91]), but not in fully adjusted models (1.17 [0.94-1.46].
(Drucker *et al.*, 2016)	Non-fatal myocardial infarction	Nurses Health Study	Cohort study; no assessment of AD current history or severity	≥18 years	Self-report of *ever* clinician diagnosed AD	Self-report and confirmed by medical record review	Non-fatal MI was not associated with ever history of AD (0.97 [0.69-1.32]).
	Non-fatal stroke						Non-fatal stroke was significantly increased in those with ever history of AD when controlling for age (adjusted OR [95% CI]: 1.38, [1.03-1.85]).
(Legendre *et al.*, 2015)	Lymphoma	Meta-analysis	Case-control and cohort studies (n=24)	All ages	Multiple	Multiple	Increased risk of lymphoma in AD (RR [95% CI]: 1.43 [1.12-1.81]). More severe AD and highly potent topical corticosteroids were associated with an increased risk of lymphoma.
(Arellano *et al.*, 2007)	Lymphoma	Pharmetrics – managed care database	Case-control	0-61+	ICD-9 codes	ICD-9 codes	Higher odds of lymphoma with more severe AD (adjusted OR [95% CI]: 2.4 [1.5-3.8]) and oral corticosteroids (1.5 [1.0-2.4]).
(Arellano *et al.*, 2009)	Lymphoma	The Health Improvement Network (THIN) database	Nested case-control study from UK practice	All ages	READ codes, dermatologist referral	READ codes	Increased odds of lymphoma in AD (1.83 [1.41-2.36] and AD referred to a dermatologist (3.72 [1.40-9.87]).

			network				
(Linabery *et al.*, 2010)	Leukemia	Meta-analysis	Case-control studies (n=6 for AD)	All ages	Multiple	Multiple	Decreased odds of acute lymphoblastic leukemia (random-effects OR [95% CI]: 0.74 [0.58-0.96]) and acute myelogenous leukemia (0.78 [0.53-1.15]) in AD.
(Cheng *et al.*, 2015)	Basal cell carcinoma of the skin	New Hampshire Skin Cancer Study	Population-based case-control study	≥25 years	Self-report of physician diagnosed AD	Clinical diagnosis and histological review	No increased odds of early-onset BCC in AD (adjusted OR [95% CI]: 1.52 [0.77-3.01]).
	Squamous cell carcinoma of the skin						Marginally significant increased odds of SCC in AD (1.83 [0.97-3.45]). Significantly increased odds of SCC in men (3.50 [1.39-8.82]) but not women (0.95 [0.35-2.57]) with AD.
(Dalgard *et al.*, 2015)	Depression	Multicenter cohort from 13 European countries	Cross-sectional	>18 years	Dermatologist examination	HADS	Higher odds of depression in AD (3.27 [1.61–6.62]).
	Anxiety					HADS	Higher odds of anxiety in AD (2.01 [1.10–3.68]).
	Suicidal ideation					Questionnaire	Higher odds of suicidal ideation in AD in unadjusted models (2.03 [1.20–3.45]), but not adjusted models (1.32 (0.75–2.33)).
(Riis *et al.*, 2016)	ADHD	Danish National Registry of Patients	Danish population-based administrative health care database	Children	ICD-10 codes	ICD-10 code and ADHD medication requirement	Higher risk of clinical encounter (adjusted OR [95% CI]: 1.3 [1.1-1.6]) and medication (1.3 [1.2-1.5]) for ADHD in AD.
(Schmitt *et al.*, 2010)	Attention deficit (hyperactivity) disorder	Systematic review	Cross-sectional, case-control and cohort	<18 years	Multiple	Multiple	AD was associated with ADD/ADHD in 4 out of 6 studies.

			(n=6 for AD)				
(Yaghmaie *et al.*, 2013)	Depression	National Survey of Children's Health	US population-based cohort study	0-17 years	Caregiver report	Caregiver report	Increased odds of depression in AD (1.81 [1.33-2.46]). Prevalence increased with more severe AD.
	Anxiety						Increased odds of anxiety in AD (1.77 [1.36-2.29]). Prevalence increased with more severe AD.
	ADD/ADHD						Increased odds of ADD/ADHD in AD (1.87 [1.54-2.27). Prevalence increased with more severe AD.
	Conduct disorder						Increased odds of conduct disorder in AD (1.87 [1.46-2.39]). Prevalence increased with more severe AD.
	Autism spectrum disorder						Increased odds of conduct disorder in AD (3.04 [2.13-4.34]). Prevalence increased with more severe AD.

*Although this review was comprehensive, it was not a systematic review and it is possible that some papers were not included. ADD/ADHD = attention deficit disorder/ attention deficit-hyperactivity disorder; BCC = basal cell carcinoma; BMI = body mass index; BP = blood pressure; CAD = coronary artery disease; CI = confidence interval; HADS = Hospital Anxiety and Depression Score; ICD = International Statistical Classification; OR = odds ratio; NHANES = National Health and Nutrition Examination Survey; NHIS = National Health Interview Survey; RA = rheumatoid arthritis; SCC = squamous cell carcinoma; SLE = systemic lupus erythematosus.