

Allergic Angina Syndrome in Anesthesia and Diabetes

Sir,

We have read with great interest the report published in *Indian Journal of Critical Care Medicine*^[1] concerning a 55-year-old male diabetic patient who developed tachycardia, bronchospasm, high airway pressure, increased blood pressure, reduced oxygen saturation, and reduced left ventricular ejection fraction with electrocardiographic ST elevation in the

anterior leads following induction anesthesia with rocuronium, propofol, and fentanyl intravenous administration. The patient was diagnosed as rocuronium-induced Kounis syndrome and recovered with intravenous nitroglycerin, hydrocortisone, budesonide, and salbutamol nebulization. Subsequent coronary arteriography did not reveal obstructive coronary artery disease. The authors correctly diagnosed and treated the

patient. However, this report raises the following issues as far as perioperative anesthesia-induced allergy in diabetic patients.

Intravenous administration of rocuronium, propofol, and fentanyl is a routine practice for induction anesthesia, and the authors correctly used this combination. However, these three agents have been incriminated as inducing mild or severe hypersensitivity reactions.^[2] Clinical studies have shown that patients, prone to allergy, have more symptoms than monosensitized individuals when simultaneously exposed to several anesthetics, as in the described case. This is based on findings that subthreshold numbers of immunoglobulin E antibodies with different specificities may have additive effects and can join forces to trigger mast cells to release their mediators. This can happen when the patient is simultaneously exposed to the corresponding antigens.^[3]

The described patient's blood pressure was increased during the allergic reaction (190/72 mmHg), and his clinical picture and clinical course were mild that is a bit unusual in rocuronium-induced allergic reactions. Indeed, the patient was diabetic and such patients are refractory to allergic inflammatory diseases. In alloxan-induced diabetes, there is a clear downregulation of local allergic inflammatory responses.

Experiments in diabetic rats have shown that skin mast cells' and histamine release is reduced. Administration of insulin in diabetic rats restores basal mast cell numbers.^[4] Furthermore, patients with diabetes present markedly reduced lethality when they suffer severe allergic reactions. This refractoriness is associated with mast cell depletion and overexpression of endogenous corticosteroids.^[5]

In this patient, the development of allergic reaction was not associated with signs of shock, the left ventricular ejection fraction was dropped only to 40%, and he recovered with intravenous nitroglycerin, hydrocortisone, budesonide, and salbutamol nebulization. Neither epinephrine nor intravenous fluid replacement had been necessary. This is another example denoting that the human heart and especially the coronary arteries can be the primary site and the target of an allergic reaction resulting in the development of Kounis syndrome. The view, therefore, that the registered cardiac damage during severe anaphylactic reactions might be due to peripheral vasodilation cannot be supported in this case.^[6]

Therefore, detailed medical history, regarding the coronary arteries as the primary target of allergic reactions and considering, in medicine and especially in anesthesia, "the fewer the better" could be of paramount clinical importance for the patient's safety.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

Nicholas G. Kounis, Ioanna Koniari¹, George Tzanis², George D. Soufras³, George Hahalis

Department of Cardiology, University of Patras Medical School, ³Department of Cardiology, "Saint Andrews" State General Hospital, Patras, ²Department of Cardiology, "Laiko" General Hospital, Athens, Greece, ¹Department of Electrophysiology, Royal Brompton Hospital, London, United Kingdom

Address for correspondence: Prof. Nicholas G. Kounis, Department of Cardiology, University of Patras Medical School, Queen Olgas Square, 7 Aratou Street, Patras 26221, Greece. E-mail: ngkounis@otenet.gr

REFERENCES

1. Raut MS, Kar S, Maheshwari A. Awareness of allergic angina syndrome. *Indian J Crit Care Med* 2017;21:412-3.
2. Kounis N, Kounis G. Anaphylactic cardiovascular collapse during anesthesia: The kounis acute hypersensitivity syndrome seems to be the most likely cause. *J Korean Med Sci* 2013;28:638-9.
3. Kounis NG, Mazarakis A, Alimpanis G, Gkouias K, Kounis GN, Tsigkas G, *et al.* The more allergens an atopic patient is exposed to, the easier and quicker anaphylactic shock and Kounis syndrome appear: Clinical and therapeutic paradoxes. *J Nat Sci Biol Med* 2014;5:240-4.
4. de F Carvalho V, Campos LV, Farias-Filho FA, Florim LT, Barreto EO, Pirmez C, *et al.* Suppression of allergic inflammatory response in the skin of alloxan-diabetic rats: Relationship with reduced local mast cell numbers. *Int Arch Allergy Immunol* 2008;147:246-54.
5. Carvalho VF, Barreto EO, Diaz BL, Serra MF, Azevedo V, Cordeiro RS, *et al.* Systemic anaphylaxis is prevented in alloxan-diabetic rats by a mechanism dependent on glucocorticoids. *Eur J Pharmacol* 2003;472:221-7.
6. Felix SB, Baumann G, Berdel WE. Systemic anaphylaxis – Separation of cardiac reactions from respiratory and peripheral vascular events. *Res Exp Med (Berl)* 1990;190:239-52.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Access this article online

Quick Response Code:



Website:

www.ijccm.org

DOI:

10.4103/ijccm.IJCCM_363_17

How to cite this article: Kounis NG, Koniari I, Tzanis G, Soufras GD, Hahalis G. Allergic angina syndrome in anesthesia and diabetes. *Indian J Crit Care Med* 2018;22:128-9.

© 2018 Indian Journal of Critical Care Medicine | Published by Wolters Kluwer - Medknow