

Confirmed causal effect of obesity on asthma and new insights on potential underlying shared genetic mechanisms

Raquel Granell, PhD,^{a,b} and Santiago Rodriguez, PhD^{a,b} *Bristol, United Kingdom*

Key words: Obesity, asthma, cross-trait genome-wide association study, shared genetics, gene expression

Asthma is a highly heterogeneous chronic respiratory disease characterized by symptoms of wheeze, cough, and breathlessness. In the United States, 250,000 new asthma cases per year are related to obesity and characterized by severity of symptoms, poor prognosis, poor asthma control, and resistance to current treatments and this applies to both children and adults.¹ Most of the observational and experimental evidence in the literature points toward obesity modifying asthma risk in children and adults. Results from Mendelian randomization studies have provided causal evidence for obesity increasing the risk of asthma, whereas little evidence has been found to support the opposite effect.² However, the shared biological mechanisms remain unclear and the heterogeneity of asthma and the obese-asthma phenotypes adds complexity and inconsistency in the results.

It is well recognized that there are different asthma phenotypes, and different obese-asthma phenotypes are starting to emerge.^{1,3} In this issue of the journal, Zhu et al⁴ explored the shared genetic associations between obesity and obesity-related metabolic traits and early- versus later-onset asthma in an adult population. Early-onset asthma (often characterized as atopic) and late-onset asthma (nonatopic, more often in females) are recognized as distinct asthma phenotypes that differ clinically and genetically.⁵ The authors have also explored the shared genetics between obesity traits and atopic versus nonatopic asthma, and results were consistent with previous reports, showing stronger associations for nonatopic compared with atopic asthma.

Zhu et al reported, for the first time, positive genome-wide genetic correlations between body mass index (BMI) and later-onset asthma (0.25 ; $P = 9.8 \times 10^{-22}$), nonatopic asthma (0.24 ; $P = 5.8 \times 10^{-17}$), and atopic asthma (0.08 ; $P = .002$), implying that higher BMI single nucleotide polymorphisms correlated with higher asthma risk single nucleotide polymorphisms. Little evidence of shared genetic correlation between BMI and early-onset asthma was reported. Furthermore, they were able to confirm causal effects of BMI on later-onset (odds ratio [OR], 1.21 ; $P = 6.3 \times 10^{-7}$), atopic (OR, 1.20 ; $P = 8.4 \times 10^{-6}$),

and nonatopic (OR, 1.10; $P = .04$) asthma and identified 32 independent shared loci between these traits (via cross-trait genome-wide association study and functional analyses) including the *HLA* region (key role in immune system), *ERBB3* (regulation of bronchial epithelial repair and remodeling), and *SMAD3* (regulation of inflammatory response).

Two genes (*ACOXL* in chromosome 2 and *MYL6* in chromosome 12) were differentially expressed in mouse lung tissues when comparing obese db/db or high-fat-diet-fed mice to controls. The *ACOXL* gene encodes an important enzyme involved in fatty acid beta-oxidation using acyl-CoA oxidase. The differential expression of *ACOXL* gene shown by Zhu et al⁴ supports a role of fatty acid oxidation caused by obesity in the inflammatory state of asthma. The *MYL6* gene encodes a motor protein (that converts chemical energy into mechanical work) expressed in smooth and non-muscle tissues. This gene, shown by Zhu et al⁴ to be differentially expressed, is involved in airway smooth muscle contractile function, a contributing factor in the excessive airway narrowing in asthma. These 2 genes open potential new mechanisms underlying obesity and asthma identified by shared genetic analyses and further confirmed by experimental mouse models (Fig 1).

The study by Zhu et al has several limitations. Although different traits were used to define obesity, these were all BMI based. BMI does not distinguish between fat mass and lean mass nor capture information on the location of body fat. Assessing body composition is needed to investigate the development and progression of obese patients throughout life because these components might better reflect differing physiology and metabolic health,⁶ and this, in turn, might be relevant in obese patients with asthma.

They found more evidence for positive genetic correlation between obesity and later-onset asthma compared with early-onset asthma using a cutoff of 16 years. However, it is possible that early-onset asthma before 16 years was underreported in this adult population (age range at recruitment 40-69 years) because of recall bias. Similarly, atopic asthma is more common during childhood; therefore, the number of self-reported (early-onset) atopic asthma cases might also be underreported.

The role of sex remains unclear. Many studies have reported differing obesity-asthma associations by sex, with the obese patient more often characterized by being female, with less atopy, and late-onset asthma. Zhu et al found evidence of a higher genetic correlation between later-onset asthma and BMI (adjusted waist-to-hip ratio) in males compared with females, but not for other BMI traits.

The role of glucose metabolism was not discussed despite reporting significant genetic correlations of later-onset and non-atopic asthma with type 2 diabetes, fasting glucose, and fasting insulin and reporting the same region in chromosome 11 as a shared genetic region associated with fasting glucose and both

From ^aMRC Integrative Epidemiology Unit and ^bPopulation Health Sciences, Bristol Medical School, University of Bristol.

Disclosure of potential conflict of interest: The authors declare that they have no relevant conflicts of interest.

Received for publication October 17, 2019; revised November 22, 2019; accepted for publication November 27, 2019.

Corresponding author: Raquel Granell, PhD, MRC Integrative Epidemiology Unit, University of Bristol, Bristol, United Kingdom. E-mail: Raquel.Granell@bristol.ac.uk.

J Allergy Clin Immunol 2019;■■■:■■■-■■■.

0091-6749/\$36.00

© 2019 American Academy of Allergy, Asthma & Immunology

<https://doi.org/10.1016/j.jaci.2019.11.033>




FIG 1. Potential new mechanisms underlying obesity and asthma identified in shared-genetic and experimental analyses by Zhu et al.⁴ GWAS, Genome-wide association study.

later-onset and nonatopic asthma. These findings are in agreement with studies reporting that airway inflammation is not the only mechanism driving the association between obesity and asthma and the reason why some obese patients with asthma do not respond to anti-inflammatory drugs. Metabolic changes induced by altered glucose metabolism (insulin resistance) could mediate inflammatory T_H2 -independent mechanisms in obese patients with asthma.⁷ Zhu et al's results suggest that these mechanisms are particular to late-onset nonatopic asthma.

The strength of this study relies on performing the largest genome-wide association study ($N > 450,000$) of subphenotypes of asthma, providing summary results that will be used in future Mendelian randomization and genetic analyses. Their results have strengthened the evidence supporting a causal link between obesity and asthma and the identified shared loci support the involvement of inflammation (T_H2 - and non- T_H2 -dependent), airway repair and remodeling, and the immune system as having key roles in the shared etiologies between both traits. The question now for future research is whether there are any other factors, besides shared genetics, with a causal role.

Recently, Peters et al¹ have reviewed the many metabolic and immune function-related factors that contribute to the obese-asthma syndrome: changes in lung function associated with higher leptin, decreased adiponectin levels and adipose tissue inflammation, increased airway oxidative-stress in obese adults with late-onset asthma, innate immune responses (T_H17 pathways and innate lymphoid cells) implicated in both obesity and asthma,

CD4 cells skewed by obesity associated with worse asthma severity and control, and insulin resistance associated with asthma and poor lung function. Other factors proposed to mediate the obesity-asthma relationship include environmental exposures (smoking, air pollution), diet (vitamin D, high-fat), physical activity, and the microbiome. For instance, changes in the diet (low-fiber diet, antibiotic use) alters the gut microbiota and this, in turn, might affect the development of allergic airway disease.¹ A small study by Rastogi et al⁸ described differential epigenome-wide DNA methylation patterns in children according to their obesity and asthma status. The impact of indoor and outdoor pollution might also be of interest in the obese population with asthma.⁹

Obesity is associated with increased asthma risk and comorbidity. The large number and variety of inflammatory and noninflammatory processes that co-occur in the obese population with asthma and the fact that most asthma treatments only target a single pathway could explain the poor response to treatment observed in these patients. Zhu et al⁴ have highlighted that the obese-asthma syndrome is complex and multifactorial and the underlying biological mechanisms might differ between children and adults and males and females, reinforcing the idea of precision medicine and the need to identify and develop treatments that are effective for particular groups of patients.¹⁰ Future studies characterizing obesity beyond BMI and involving new areas of research such as epigenetics and the microbiome might suggest distinct novel pathways.

REFERENCES

1. Peters U, Dixon AE, Forno E. Obesity and asthma. *J Allergy Clin Immunol* 2018; 141:1169-79.
2. Skaaby T, Taylor AE, Thuesen BH, Jacobsen RK, Friedrich N, Mollehave LT, et al. Estimating the causal effect of body mass index on hay fever, asthma and lung function using Mendelian randomization. *Allergy* 2018;73: 153-64.
3. Holguin F, Bleecker ER, Busse WW, Calhoun WJ, Castro M, Erzurum SC, et al. Obesity and asthma: an association modified by age of asthma onset. *J Allergy Clin Immunol* 2011;127:1486-93.e2.
4. Zhu Z, Guo Y, Shi H, Liu CL, Panganiban RA, Chung W, et al. Shared genetic and experimental links between obesity-related traits and asthma subtypes in UK Biobank. *J Allergy Clin Immunol* 2020;145:XXX-XXX.
5. de Nijs SB, Venekamp LN, Bel EH. Adult-onset asthma: is it really different? *Eur Respir Rev* 2013;22:44-52.
6. Blundell JE, Dulloo AG, Salvador J, Fruhbeck G; EASO SAB Working Group on BMI. Beyond BMI—phenotyping the obesities. *Obes Facts* 2014;7: 322-8.
7. Leiria LO, Martins MA, Saad MJ. Obesity and asthma: beyond T(H)2 inflammation. *Metabolism* 2015;64:172-81.
8. Rastogi D, Fraser S, Oh J, Huber AM, Schulman Y, Bhagtani RH, et al. Inflammation, metabolic dysregulation, and pulmonary function among obese urban adolescents with asthma. *Am J Respir Crit Care Med* 2015; 191:149-60.
9. Lu KD, Breyse PN, Diette GB, Curtin-Brosnan J, Aloe C, Williams DL, et al. Being overweight increases susceptibility to indoor pollutants among urban children with asthma. *J Allergy Clin Immunol* 2013;131:1017-1023. 23. e1-3.
10. Canonica GW, Ferrando M, Baiardini I, Puggioni F, Racca F, Passalacqua G, et al. Asthma: personalized and precision medicine. *Curr Opin Allergy Clin Immunol* 2018;18:51-8.