

blood

2011 118: 6930-6938
Prepublished online October 14, 2011;
doi:10.1182/blood-2011-03-343962

Reduced mast cell and basophil numbers and function in *Cpa3-Cre; Mcl-1^{fl/fl}* mice

Jennifer N. Lilla, Ching-Cheng Chen, Kaori Mukai, Maya J. BenBarak, Christopher B. Franco, Janet Kalesnikoff, Mang Yu, Mindy Tsai, Adrian M. Piliponsky and Stephen J. Galli

Updated information and services can be found at:
<http://bloodjournal.hematologylibrary.org/content/118/26/6930.full.html>

Articles on similar topics can be found in the following Blood collections
[Phagocytes](#), [Granulocytes](#), and [Myelopoiesis](#) (293 articles)

Information about reproducing this article in parts or in its entirety may be found online at:
http://bloodjournal.hematologylibrary.org/site/misc/rights.xhtml#repub_requests

Information about ordering reprints may be found online at:
<http://bloodjournal.hematologylibrary.org/site/misc/rights.xhtml#reprints>

Information about subscriptions and ASH membership may be found online at:
<http://bloodjournal.hematologylibrary.org/site/subscriptions/index.xhtml>

Blood (print ISSN 0006-4971, online ISSN 1528-0020), is published weekly by the American Society of Hematology, 2021 L St, NW, Suite 900, Washington DC 20036.

Copyright 2011 by The American Society of Hematology; all rights reserved.



Reduced mast cell and basophil numbers and function in *Cpa3-Cre; Mcl-1^{fl/fl}* mice

Jennifer N. Lilla,¹ Ching-Cheng Chen,^{1,2} Kaori Mukai,¹ Maya J. BenBarak,¹ Christopher B. Franco,¹ Janet Kalesnikoff,¹ Mang Yu,¹ Mindy Tsai,¹ Adrian M. Piliponsky,^{1,3} and Stephen J. Galli¹

¹Department of Pathology, Stanford University School of Medicine, Stanford, CA; ²Division of Stem Cell and Leukemia Research, City of Hope Medical Center, Duarte, CA; and ³Center for Immunity and Immunotherapies, Seattle Children's Research Institute, Seattle, WA

It has been reported that the intracellular antiapoptotic factor myeloid cell leukemia sequence 1 (Mcl-1) is required for mast cell survival in vitro, and that genetic manipulation of Mcl-1 can be used to delete individual hematopoietic cell populations in vivo. In the present study, we report the generation of C57BL/6 mice in which Cre recombinase is expressed under the control of a segment of the carboxypeptidase A3 (*Cpa3*) promoter. C57BL/6-*Cpa3-Cre; Mcl-1^{fl/fl}* mice are se-

verely deficient in mast cells (92%-100% reduced in various tissues analyzed) and also have a marked deficiency in basophils (58%-78% reduced in the compartments analyzed), whereas the numbers of other hematopoietic cell populations exhibit little or no changes. Moreover, *Cpa3-Cre; Mcl-1^{fl/fl}* mice exhibited marked reductions in the tissue swelling and leukocyte infiltration that are associated with both mast cell- and IgE-dependent passive cutaneous anaphylaxis (except at sites en-

grafted with in vitro-derived mast cells) and a basophil- and IgE-dependent model of chronic allergic inflammation, and do not develop IgE-dependent passive systemic anaphylaxis. Our findings support the conclusion that Mcl-1 is required for normal mast cell and basophil development/survival in vivo in mice, and also suggest that *Cpa3-Cre; Mcl-1^{fl/fl}* mice may be useful in analyzing the roles of mast cells and basophils in health and disease. (*Blood*. 2011;118(26):6930-6938)

Introduction

Myeloid cell leukemia sequence 1 (Mcl-1)¹ has been identified as an intracellular antiapoptotic factor in a variety of hematopoietic cells, both in vitro and in vivo.²⁻⁶ Human mast cells express Mcl-1,^{7,8} and Mcl-1 can promote the survival of some populations of human neoplastic mast cells in vitro.⁷ Basophils, granulocytes with many characteristics and functions that partially overlap with those of tissue mast cells,⁹⁻¹² can also express Mcl-1.¹³ However, it is not clear to what extent Mcl-1 is important in the development and/or survival of mast cells or basophils in vivo.

Opferman et al showed that the genetic manipulation of Mcl-1 can be used to delete individual hematopoietic cell populations in mice.⁴ We therefore used this approach to examine the effects of reducing expression of Mcl-1 in the mast cell lineage in vivo. To attempt to delete Mcl-1 selectively in mast cells, we used the promoter for the peptidase carboxypeptidase A3 (CPA3; originally named mast cell carboxypeptidase A¹⁴). CPA3 is highly expressed in mast cells,¹⁵ but is also expressed in basophils¹⁶ and can be expressed in some populations of T-cell progenitors and thymic T cells^{17,18} and in certain hematopoietic progenitor cells.¹⁹

We generated C57BL/6 mice in which a segment of the *Cpa3* promoter drives expression of Cre recombinase, and then mated these *Cpa3-Cre* transgenic mice to mice bearing a floxed allele of Mcl-1.⁴ We found that C57BL/6-*Cpa3-Cre; Mcl-1^{fl/fl}* mice are severely deficient in mast cells and have a marked deficiency in basophils, and also exhibit striking impairment in mast cell- or basophil- and IgE-dependent biologic responses.

Methods

Mice

All animal experiments were carried out following protocols approved by the Stanford University Administrative Panel on Laboratory Animal Care. B6-Tg(*Cpa3-cre*)3Glli (*Cpa3-Cre*—transgenic mice) were generated by microinjecting the *Cpa3-Cre* transgene into embryonic stem cells in the B6 background (Stanford University). *Gt(ROSA)26Sor^{tm4}(ACTB-tdTomato-EGFP)^{Luo}/J(mT/mG)* mice, obtained from The Jackson Laboratory, were crossed to *Cpa3-Cre* mice for Cre expression analysis. *Mcl-1^{+/fl}* (B6;129-*Mcl1^{tm3sjk}*J) animals were obtained from The Jackson Laboratory. *Mcl-1^{+/fl}* mice were bred to progeny from 2 *Cpa3-Cre* founder lines (founder lines #4 and #5) to obtain *Cpa3-Cre; Mcl-1^{+/+}*, *Cpa3-Cre; Mcl-1^{+/fl}*, and *Cpa3-Cre; Mcl-1^{fl/fl}* animals, but only the *Cpa3-Cre; Mcl-1^{fl/fl}* mice derived from founder line #4 exhibited a substantial mast cell deficiency. Therefore, the mice used were derived from crosses between founder line #4 (subsequently referred to as *Cpa3-Cre* mice) and *Mcl-1^{fl}* animals, and these mice had been intercrossed a minimum of 6 generations into the C57BL/6 background. Heterozygous *Cpa3-Cre* mice were determined to have 5 copies of the *Cpa3-Cre* transgene by real-time PCR. To emphasize that *Cpa3-Cre; Mcl-1^{fl/fl}* mice have deficiencies in mast cells and basophils that are independent of mutations affecting Kit, we call them informally in our laboratory “Hello Kitty” mast cell- and basophil-deficient mice. C57BL/6J (B6)-*Kit^{W-sh/W-sh}* mast cell-deficient mice and B6-*Kit^{+/+}* wild-type control mice were from stocks bred in our laboratory, as described previously.²⁰ All experiments were performed at least 3 times unless otherwise indicated.

Abs, flow cytometry, and cell quantification

We used flow cytometry to identify and enumerate basophils (CD49b [DX5]⁺; FcεRIα⁺) in the BM, peripheral blood, and spleen; mast cells

Submitted March 22, 2011; accepted October 4, 2011. Prepublished online as *Blood* First Edition paper, October 14, 2011; DOI 10.1182/blood-2011-03-343962.

An Inside *Blood* analysis of this article appears at the front of this issue.

The online version of this article contains a data supplement.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked “advertisement” in accordance with 18 USC section 1734.

© 2011 by The American Society of Hematology

(c-Kit⁺; FcεRIα⁺) in peritoneal lavage fluid; B cells (B220⁺; CD3ε⁻) in peritoneal lavage fluid and spleen; eosinophils (CCR3⁺; SSC^{hi}) in the spleen and BM; macrophages/monocytes (F4/80⁺; Gr-1⁻) in BM, peritoneal lavage fluid, and spleen; neutrophils (Gr-1⁺) in BM, peritoneal lavage fluid, and spleen; T cells (CD3ε⁺; B220⁻) in the spleen and thymus; and erythroid cells (Ter119⁺) in the spleen. Briefly, RBCs were lysed with pH 7.3 ACK lysis buffer (0.15M NH₄Cl, 1mM KHCO₃, and 0.1mM EDTA, pH 8.0) for 5 minutes. Cells were blocked with unconjugated anti-CD16/CD32 on ice for 5 minutes and then stained with a combination of Abs on ice for 15 minutes. Abs used were: B220 (RA3-6B2; eBioscience), c-Kit (2B8; eBioscience), CCR3 (TG14/CCR3; BioLegend and R&D Systems), CD11b (M1/70; eBioscience), CD3ε (145-2C11; BioLegend), CD49b (DX5; BD Pharmingen and BioLegend), F4/80 (BM8; BioLegend), FcεRIα (MAR-1; BioLegend and eBioscience), and Ter119 (TER-119; BioLegend). The expression of cell-surface markers was analyzed on a FACSaria II (BD Biosciences) using FlowJo Version 8.8.6 software (Stanford University and TreeStar). Dead cells (identified with propidium iodide; Invitrogen) were not included in the analysis. Blood lymphocytes, neutrophils, monocytes, eosinophils, erythrocytes, and platelets were counted using the Abbott Cell-Dyn 3500 automated hematology analyzer.

Histology

Some tissue specimens were fixed with 10% neutral buffered formalin and embedded in paraffin. Four-micrometer sections were stained with 0.1% Toluidine blue or H&E for histologic examination and enumeration of mast cells or leukocytes, respectively. To examine mast cell morphology in ears subjected to passive cutaneous anaphylaxis (PCA), specimens of ear pinnae were fixed in 2% paraformaldehyde, 2.5% glutaraldehyde, 0.1M cacodylate buffer, and 0.025% CaCl₂, washed in 0.1M cacodylate buffer, and embedded in Epon (Electron Microscopy Sciences), and then 1-μm sections were stained with alkaline Giemsa (pH 8.2) as described previously.²¹ To examine mast cell numbers in mesenteric windows, samples of mesentery were spread on a glass slide, fixed in Carnoy solution and stained with Alcian blue/Safranin-O. For PCA and chronic allergic inflammation (CAI) experiments (see “PCA” and “CAI”), ear pinnae were collected 6 hours after challenge and fixed in 10% neutral buffered formalin for paraffin sections to be stained with H&E to enumerate leukocytes present in the ear dermis. In all histological assessments, cell numbers were enumerated by a single observer not aware of the identity (mouse group) of the individual sections. Cell numbers were based on counting 25 medium power fields (200×) for mast cells or high power fields (400×) for leukocytes from a minimum of 3 sections from individual tissues from each mouse, and the mean value was used as the number for that mouse. Mast cells were quantified according to area (per square millimeter) for all tissues except the forestomach, glandular stomach, duodenum, and ileum (per millimeter of mucosa and submucosa) and dorsal skin and ear pinna (per millimeter horizontal field length of dermis). Leukocytes also were quantified as per millimeter of horizontal field length of ear dermis. To visualize infiltrating basophils in CAI experiments, ear pinnae sections were pretreated as described previously,²² and incubated with anti-mMCP-8 (10 μg/mL) for 1 hour at room temperature, followed by HRP-conjugated goat anti-rat IgG (Santa Cruz Biotechnology) diluted 1:500. The sections were subsequently incubated in a DAB solution (Sigma-Aldrich) and counterstained with Giemsa. Images were captured with an Olympus BX60 microscope using a Retiga-2000R QImaging camera run by Image-Pro Plus Version 6.3 software (Media Cybernetics).

PCA

IgE-dependent PCA was induced in the ear pinna as described previously.²³ Briefly, mice under isoflurane anesthesia were sensitized passively with IgE by intradermal injection of 20 ng of dinitrophenol (DNP)-specific IgE (α-DNP clone e26²⁴) kindly provided by Dr Fu-Tong Liu (University of California-Davis) in 20 μL of HMEM-Pipes buffer (Sigma-Aldrich) in the right ear pinna; mice received 20 μL of vehicle intradermally in the left ear pinna as a control. The next day, mice were challenged intravenously with 100 μg of DNP₃₀₋₄₀-conjugated human serum albumin (DNP-HSA; Sigma-Aldrich) in 100 μL of NaCl. Immediately before and at intervals

after antigen challenge, ear thickness was measured with a dial thickness gauge (G-1A; Ozaki). Mice were killed 6 hours after antigen challenge and ear pinnae were collected for histological analysis.

CAI

IgE-dependent CAI of the ear skin was induced as described previously.²⁵ Briefly, mice under isoflurane anesthesia were sensitized IV with IgE (300 μg of trinitrophenol (TNP)-specific IgE mAb (IGELb4), kindly provided by Dr Hajime Karasuyama (Tokyo Medical and Dental University). A day later, 10 μg of TNP₁₄-conjugated OVA (TNP-OVA; Biosearch Technologies) in 10 μL of PBS was injected intradermally with a microsyringe into the left ear of mice under isoflurane anesthesia, and an equal amount of unconjugated OVA (Sigma-Aldrich) was injected into the right ear as a control. Ear thickness was measured with a dial thickness gauge (G-1A; Ozaki) at the indicated time points.

PSA

IgE-dependent passive systemic anaphylaxis (PSA) was induced as described previously.²⁶ Briefly, mice were sensitized passively with IgE by intraperitoneal injection of 20 ng of DNP-specific IgE in 100 μL of saline or were injected intraperitoneally with saline as a control and then challenged intraperitoneally the next day with 1 mg of DNP-HSA in 100 μL of saline. Immediately before and at intervals after antigen challenge, body temperature was measured with a rectal thermometer (Physitemp Instrument). Mice were killed 2 hours after antigen challenge.

Cell culture

Mouse femoral and tibial BM cells from C57BL/6J or various *Cpa3-Cre* mice were cultured for 6 weeks in 20% WEHI-3 cell-conditioned medium (as a source of IL-3) to generate BM-derived cultured mast cells (BMCMCs) or for 7-8 days to generate BM-derived basophils (BMBas). Mast cell differentiation was assessed by May-Grünwald-Giemsa staining of cytospin preparations for granule content and by flow cytometry for surface expression of c-Kit and FcεRIα (all BMCMC preparations were > 95% pure). Basophil differentiation was assessed by flow cytometry for surface expression of DX5 and FcεRIα.

Western blotting

BMCMCs were washed with DMEM and solubilized by boiling for 1 minute with Laemmli-SDS sample buffer (10⁶ cells/50 μL). Total cell lysates were sheared (1 cc syringe; 26-ga needle), and then separated by SDS-PAGE, electroblotted onto PVDF membranes (Invitrogen), and probed with polyclonal Abs against Mcl-1 (Rockland Immunochemicals) or GAPDH (Fitzgerald Industries).

Statistics

Unless otherwise specified, data were examined for statistical significance using the Student *t* test (2-tailed, unpaired). We used 2-way ANOVA to compare time courses of responses. Significance was attributed when *P* ≤ .05 was observed.

Results

Mcl-1 is expressed by mouse mast cells in vitro

We first tested whether BMCMCs express the Mcl-1 protein in vitro. After 6 weeks of culture in IL-3-supplemented medium, BMCMCs expressed Mcl-1 protein, and the amount of Mcl-1 expression decreased when BMCMCs were maintained for 24 hours in the absence of IL-3 (Figure 1A). This observation agrees with previous findings that these culture conditions result in decreased levels of Bcl-2 family antiapoptotic proteins in BMCMCs, which is concomitant with a loss of mast cell viability.²⁷




Figure 1. Mcl-1 expressed by mast cells in vitro and maps of the *Cpa3-Cre* and *Mcl-1^{fl}* transgenes. (A) Mcl-1 is expressed by mast cells in vitro. BMCMCs (5×10^5) from male wild-type C57BL/6J mice were subjected to electrophoresis and Western blotting using a polyclonal Ab against Mcl-1 (Rockland) and reprobed with anti-GAPDH Ab to show loading. Bands represent the long (35 kDa) and short (32 kDa) forms of Mcl-1. BMCMCs were cultured for 24 hours with 10% FCS alone (IL-3⁻) or FCS + IL-3 (IL-3⁺; 20% Wehi-3-conditioned medium). Expression of Mcl-1 decreased after 24 hours of culture in medium without added IL-3. Results are representative of those obtained in 6 independent experiments. (B) Map of the *Cpa3-Cre* transgene. A 780-bp fragment of the region just 5' to the *Cpa3* transcription start site was used to drive expression of Cre recombinase in vivo (*Cpa3-Cre*). These transgenic mice were bred with mice in which the first exon of *Mcl-1* has been flanked by *loxP* sites (triangles).

The *Cpa3* promoter can be used to express Cre recombinase in mast cells and basophils in vivo

Based on our in vitro findings, we hypothesized that it might be possible to reduce mast cell numbers in vivo by conditionally deleting Mcl-1 expression in these cells. To target mast cells, we constructed transgenic mice that express Cre recombinase under the control of a 780-bp fragment of the *Cpa3* gene (*Cpa3-Cre*) (Figure 1B). This region, which is just 5' to the start codon of *Cpa3*, contains GATA-binding domains²⁸ thought to regulate *Cpa3* transcription at early stages of mast cell differentiation.²⁹

To examine the extent of Cre expression in various hematopoietic lineages in *Cpa3-Cre* mice, we crossed these mice to *mT/mG* reporter mice³⁰; the latter mice express membrane-targeted red fluorescence in all cells except those in which Cre-mediated excision of the *mT* sequence (that drives expression of the membrane-targeted tandem dimer Tomato), and subsequent expres-

sion of *mG* (membrane-targeted enhanced green fluorescent protein [GFP]), results in green rather than red fluorescence. This analysis indicated that *Cpa3-Cre*; *mT/mG* mice exhibited high levels of Cre expression in peritoneal mast cells (Figure 2A). In contrast, we detected no evidence of Cre expression in *Cpa3-Cre*; *mT/mG* mice in peritoneal macrophages (Figure 2B), but a small population of splenic erythroid cells were GFP⁺ (Figure 2C). Voehringer et al identified *Cpa3* as a gene that is also highly expressed in mouse basophils,¹⁶ and we detected Cre expression in the spleen basophils of *Cpa3-Cre*; *mT/mG* mice (Figure 2D). However, similar percentages of splenic eosinophils (Figure 2E) and neutrophils (Figure 2F) also appeared to express Cre, and for all 3 granulocyte populations, the GFP⁺ and GFP⁻ cells from the spleen or BM (supplemental Figure 1, available on the *Blood* Web site; see the Supplemental Materials link at the top of the online article) expressed very similar levels of cell-specific markers and forward versus side scatter.




Figure 2. The *Cpa3* promoter drives Cre expression in mast cells and basophils in vivo. (A-I) Expression of Cre in *Cpa3-Cre* mice, representative of similar results that were obtained in the 3 independent experiments performed. Fluorescence intensity plots depict GFP in cells from *Cpa3-Cre*; *mT/mG* mice (green solid lines) that express Cre, but not in *mT/mG*-only mice (red dashed lines) or *Cpa3-Cre*-only mice (gray filled curves). Percentage of GFP⁺ cells are indicated on the plots for *Cpa3-Cre*; *mT/mG* mice for each of the following cell populations: (A) peritoneal mast cells, (B) peritoneal macrophages, (C) splenic Ter119⁺ (erythroid) cells, (D) splenic basophils, (E) splenic eosinophils, (F) splenic neutrophils, (G) thymic T cells, (H) splenic T cells, and (I) splenic B cells.

Small numbers of thymic and splenic T cells expressed mG in *Cpa3-Cre*; *mT/mG* mice (Figure 2G-H), a finding in agreement with reports indicating that *Cpa3* can be expressed in T-cell populations in mice^{17,18}; in contrast, splenic B cells were essentially negative (Figure 2I).

Evidence that *Mcl-1* is a survival factor for mast cells and basophils in vitro and in vivo

Our results with the *Cpa3-Cre*; *mT/mG* reporter mice indicated that levels of *Cpa3* that were high enough to drive substantial *Cpa3-Cre*-dependent loss of red fluorescence were restricted predominantly to mast cells, whereas lesser amounts of *Cpa3-Cre*-dependent loss of red fluorescence were detected in all 3 granulocyte populations. To test the extent to which mast cells and various granulocytes require *Mcl-1* for development and/or survival, we crossed mice carrying the *Cpa3-Cre* transgene with mice in which the first exon of *Mcl-1* is flanked by LoxP sites (*Mcl-1^{fl/fl}*)⁴ (Figure 1B). Because male mice carrying 2 floxed *Mcl-1* alleles exhibit reduced fertility (The Jackson Laboratory), heterozygous males (*Mcl-1^{+/-}*) were crossed with either heterozygous (*Mcl-1^{fl/fl}*) or homozygous (*Mcl-1^{fl/fl}*) females to generate mice for subsequent cell lineage and functional analyses.

Cpa3-Cre; *Mcl-1^{fl/fl}* mice appeared to be grossly normal, based on both external inspection and autopsy examination of their organs. However, *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice exhibited dramatic reductions in mast cell numbers in nearly all tissues and anatomic sites examined (Figure 3 and supplemental Figure 2). In control *Cpa3-Cre*; *Mcl-1^{+/+}* mice, mast cells were present in numbers very similar to those in wild-type C57BL/6J mice (eg, see Grimbaldston et al³¹). In contrast, 6-week- to 10-month-old *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice exhibited reductions in mast cell numbers that ranged from 92%-100% in all sites examined except for the spleen, where no differences were detected in the small numbers of mast cells (Figure 3B-E and supplemental Figure 2). We found that the peritoneal mast cells detected in *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice exhibited levels of surface expression of FcεRI that were similar to those in the peritoneal mast cells of control *Cpa3-Cre*; *Mcl-1^{+/+}* mice (Figure 3E). *Cpa3-Cre*; *Mcl-1^{fl/+}* mouse tissues were also assessed for mast cell numbers, but no significant differences were found between the corresponding mast cell populations in mice with 1 versus 2 functional copies of *Mcl-1* (data not shown).

In vitro evidence suggests that the few mast cells that persist in *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice may be those that have the highest levels of *Mcl-1*. Specifically, when we examined the generation of mast cells from adult BM cells derived from *Mcl-1^{fl/fl}* versus *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice, we found that BM cells from *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice generated far fewer BMCMCs than did those from *Mcl-1^{fl/fl}* mice, if any survived at all (supplemental Figure 3A). However, those rare mast cells that persisted in the cultures of *Cpa3-Cre*; *Mcl-1^{fl/fl}* mouse BM cells had levels of *Mcl-1* protein that were similar to those of the corresponding *Mcl-1^{fl/fl}* cells (supplemental Figure 3B), suggesting that these persisting *Cpa3-Cre*; *Mcl-1^{fl/fl}* cells expressed sufficient *Mcl-1* protein to evade death even after 6 weeks of in vitro culture. The *Mcl-1* expression in such cells may be the result of incomplete excision of the first exon of *Mcl-1* by *Cpa3*-driven Cre in these mast cells.

Cpa3-Cre; *Mcl-1^{fl/fl}* mice exhibit a marked reduction in basophils

Our findings in *Cpa3-Cre*; *mT/mG* reporter mice suggested that basophils, eosinophils, and neutrophils also may be affected in

Cpa3-Cre; *Mcl-1^{fl/fl}* mice. We therefore examined whether numbers of various hematopoietic cells are reduced in *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice. We observed substantial—and statistically significant—reductions in basophil numbers in the spleen, BM, and blood of *Cpa3-Cre*; *Mcl-1^{fl/fl}* versus control *Cpa3-Cre*; *Mcl-1^{+/+}* mice (Figure 4A). In 3- to 10-month-old mice, basophil numbers were reduced 58% in the spleen, 78% in the BM, and 74% in the blood. The basophils detected in *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice exhibited levels of surface expression of FcεRI that were similar to those in the basophils of the control *Cpa3-Cre*; *Mcl-1^{+/+}* mice (Figure 4A). *Cpa3-Cre*; *Mcl-1^{fl/fl}* mouse BM yielded on average 43.3% fewer BMBas after 8 days of in vitro culture than did *Cpa3-Cre*; *Mcl-1^{+/+}* BM cells. In the 3 independent experiments performed, each with cells from *n* = 3 mice/genotype, the yield of BMBas from *Cpa3-Cre*; *Mcl-1^{fl/fl}* mouse BM cells was reduced compared with that from *Cpa3-Cre*; *Mcl-1^{+/+}* cells by 31.2%, 45.9%, and 52.8%, with the differences between numbers of BMBas in the 2 groups in these 3 experiments at *P* = .052, *P* = .007, and *P* = .007, respectively, by Student unpaired 2-tailed *t* test.

However, the *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice exhibited no significant alterations compared with values in control mice in any of the other leukocyte populations tested (Figure 4 and supplemental Figure 4), including eosinophils in the blood, spleen, and BM (Figure 4E and supplemental Figure 4A) and neutrophils in the blood, BM, and peritoneal cavity (Figure 4B,D), except for a modest (56%) increase in numbers of neutrophils in the spleen (Figure 4B). We did not detect significant changes in numbers of T cells (Figure 4C) or B cells (supplemental Figure 4C) in *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice or in the absolute numbers of lymphocytes or myeloid cells analyzed in the blood (Figure 4D-E) or in the spleen, BM, or peritoneal fluid (supplemental Figure 4A-B). Although *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice had normal numbers of platelets (Figure 4F), they had a mild macrocytic anemia (Figure 4F and supplemental Figure 5), with an approximately 31% reduction in numbers of blood erythrocytes (Figure 4F) and a 24% reduction in hematocrit (supplemental Figure 5) compared with *Cpa3-Cre*; *Mcl-1^{+/+}* control mice. Whereas the physiologic significance of such a mild anemia is not clear, this observation does support recent findings suggesting that mast cell and megakaryocyte/erythroid progenitors may be more closely linked than was realized previously.¹⁹

Cpa3-Cre; *Mcl-1^{fl/fl}* mice exhibit markedly reduced PCA

To determine whether the marked deficiency in mast cells in *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice resulted in reduced mast cell function in vivo, we examined a classic example of a mast cell-dependent biologic response, IgE-dependent PCA.²³ *Cpa3-Cre*; *Mcl-1^{+/+}* mice developed a robust local inflammatory response, with tissue swelling that peaked approximately 30 minutes after antigen challenge (Figure 5A). In contrast, the tissue swelling was markedly reduced in identically treated *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice (Figure 5A).

As expected, mast cell numbers in the ear pinnae dermis of *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice were very low compared with those in *Cpa3-Cre*; *Mcl-1^{+/+}* mice, and were not significantly different in IgE- versus vehicle-treated ears 6 hours after antigen challenge (Figure 5B and 5D). In addition to having much higher numbers of dermal mast cells, *Cpa3-Cre*; *Mcl-1^{+/+}* control mice exhibited a reduction in mast cell numbers in ear pinnae at sites of PCA reactions versus sites of vehicle injection (Figure 5B), probably because of the difficulty in identifying extensively degranulated mast cells using a counting method that requires identification of the cells' cytoplasmic granules (Figure 5D top left). Some of the




Figure 3. *Cpa3-Cre; Mcl-1^{fl/fl}* mice have markedly reduced numbers of mast cells. (A) Toluidine blue staining for mast cells in 4-μm-thick paraffin sections of glandular stomach wall (top) and dorsal skin (bottom) in *Cpa3-Cre; Mcl-1^{+/+}* mice (left) and *Cpa3-Cre; Mcl-1^{fl/fl}* mice (right) shows a marked reduction in numbers of mast cells (purple) in mice in which Mcl-1 has been selectively deleted by *Cpa3-Cre*. Scale bar indicates 100 μm. (B-C) Numbers of mast cells in various tissues in *Cpa3-Cre; Mcl-1^{+/+}* (control) mice or *Cpa3-Cre; Mcl-1^{fl/fl}* mice. Numbers of mast cells are shown as means + SEM per millimeter or per square millimeter of 4-μm-thick paraffin sections stained with 0.1% Toluidine blue (see "Methods") in tissues from *Cpa3-Cre; Mcl-1^{+/+}* (n = 16) or *Cpa3-Cre; Mcl-1^{fl/fl}* (n = 14) mice, except for mesentery window (n = 12 per group) and mammary stroma (n = 6 per group). (D) Percentage of mast cells in the live-cell population isolated from peritoneal lavage fluid from *Cpa3-Cre; Mcl-1^{+/+}* (control, n = 10) or *Cpa3-Cre; Mcl-1^{fl/fl}* (n = 10) mice, analyzed by flow cytometry (FcεRIα⁺; c-Kit⁺). Peritoneal mast cell numbers are shown as means + SEM. (B-D) **P < .01 and ***P < .001 versus corresponding values for *Cpa3-Cre; Mcl-1^{+/+}* mice. (E) Representative flow cytometry plots show comparable expression of FcεRIα⁺ on peritoneal mast cells isolated from *Cpa3-Cre;*

rare mast cells observed in the dermis of specimens from *Cpa3-Cre; Mcl-1^{fl/fl}* mice also exhibited histological evidence of degranulation (Figure 5D top right). This finding suggests that activation of the rare mast cells present in the skin of *Cpa3-Cre; Mcl-1^{fl/fl}* mice by IgE and specific antigen probably accounted for the weak tissue-swelling response detected in these animals (Figure 5A). However, at 6 hours after antigen challenge, there was no significant difference in the numbers of leukocytes present in the IgE- versus vehicle-injected ear pinnae of *Cpa3-Cre; Mcl-1^{fl/fl}* mice, whereas the numbers of leukocytes (mainly consisting of neutrophils) were markedly increased in the dermis 6 hours after antigen challenge in IgE-dependent PCA reactions in *Cpa3-Cre; Mcl-1^{+/+}* control mice (Figure 5C-D).

These results clearly show that *Cpa3-Cre; Mcl-1^{fl/fl}* mice lack sufficient mast cells to orchestrate a robust IgE-dependent PCA response in vivo. Indeed, even though the few mast cells present in the ear dermis of these mice may have accounted for the slight and quickly resolved ear swelling observed in such animals (and the attenuated swelling response depicted in Figure 5A reflected mainly the responses which occurred in the 2 *Cpa3-Cre; Mcl-1^{fl/fl}* mice with the highest numbers of ear mast cells), the *Cpa3-Cre; Mcl-1^{fl/fl}* mice exhibited no detectable IgE-dependent increase in dermal leukocytes by 6 hours after antigen challenge (Figure 5C). In contrast, both increased tissue swelling (associated with local extravasation of Evan blue dye) and enhanced leukocyte recruitment occurred at PCA reaction sites elicited in the pinnae of *Cpa3-Cre; Mcl-1^{fl/fl}* mice that had been engrafted intradermally with mast cells 8 weeks earlier (supplemental Figure 6A-C and 6E-F). Therefore, *Cpa3-Cre*-mediated deletion of Mcl-1 results in a skin mast cell deficiency that largely eliminates the ability of such mice to develop tissue swelling or leukocyte infiltration at sites challenged to express IgE- and antigen-dependent PCA reactions, but the ability to express IgE-dependent PCA is enhanced when such mice undergo local engraftment with mast cells. *Cpa3-Cre; Mcl-1^{fl/fl}* mice may also be engrafted with BMCMCs in the peritoneal cavity (supplemental Figure 6D).

***Cpa3-Cre; Mcl-1^{fl/fl}* mice exhibit markedly reduced IgE-dependent CAI**

In light of the 74% reduction in blood basophils in *Cpa3-Cre; Mcl-1^{fl/fl}* mice (Figure 4A), we investigated whether *Cpa3-Cre; Mcl-1^{fl/fl}* mice exhibited attenuation of a basophil-dependent biologic response, namely IgE-dependent CAI of the skin.²⁵ We found that the tissue swelling associated with the IgE-dependent CAI response was essentially eliminated in *Cpa3-Cre; Mcl-1^{fl/fl}* mice (Figure 6A). Leukocyte numbers (mainly consisting of neutrophils and eosinophils, but with a prominent infiltrate of basophils) were noticeably increased in the dermis 3 days after antigen challenge in IgE-dependent CAI reactions in *Cpa3-Cre; Mcl-1^{+/+}* control mice, whereas *Cpa3-Cre; Mcl-1^{fl/fl}* mice virtually lacked leukocyte infiltration and exhibited essentially no mMCP8⁺ basophils by immunohistochemistry (Figure 6B).


***Cpa3-Cre; Mcl-1^{fl/fl}* mice exhibit markedly reduced PSA**

We assessed an IgE-dependent systemic immune response in *Cpa3-Cre; Mcl-1^{fl/fl}* mice by examining a model of IgE-dependent

Figure 3. (continued) *Mcl-1^{+/+}* or *Cpa3-Cre; Mcl-1^{fl/fl}* mice. In panels B-D, mice ranged from 3-10 months of age; mouse age did not influence mast cell numbers in each genotype (data not shown). Images were captured with an Olympus BX60 microscope with a 20× objective using a Retiga-2000R QImaging camera run by Image-Pro Plus Version 6.3 software (Media Cybernetics) and exported into Adobe Photoshop (CS3), in which images were white balanced and resized.

Figure 4. *Cpa3-Cre; Mcl-1^{fl/fl}* mice have reduced numbers of basophils but not other leukocytes.

(A-F) Percentages (A-C) or absolute numbers (D-F) of leukocytes (A-E) or platelets and erythrocytes (F) are depicted as values from individual mice, with bars indicating means $\pm$ SEM. * $P < .05$ and *** $P < .001$ versus corresponding values for *Cpa3-Cre; Mcl-1^{+/+}* mice (controls). All other comparisons are not significant ($P > .05$). (A) Left: Percentage of basophils (FceRI α ⁺; DX5⁺; c-Kit⁻) in the live-cell population isolated from *Cpa3-Cre* mouse spleen (*Mcl-1^{+/+}*, $n = 6$; *Mcl-1^{fl/fl}*, $n = 6$), BM (*Mcl-1^{+/+}*, $n = 9$; *Mcl-1^{fl/fl}*, $n = 8$), or blood (*Mcl-1^{+/+}*, $n = 8$; *Mcl-1^{fl/fl}*, $n = 8$), as analyzed by flow cytometry. Right: Representative flow cytometry plots show comparable expression of FceRI α on basophils isolated from *Cpa3-Cre; Mcl-1^{+/+}* or *Cpa3-Cre; Mcl-1^{fl/fl}* mouse blood. (B) Percentage of neutrophils (Gr-1⁺) in the live-cell population isolated from *Cpa3-Cre* mouse spleen, BM, or peritoneal lavage fluid (*Mcl-1^{+/+}*, $n = 11$; *Mcl-1^{fl/fl}*, $n = 10$), as analyzed by flow cytometry. (C) Percentage of T cells (CD3⁺) in the live-cell population isolated from *Cpa3-Cre* mouse spleen (*Mcl-1^{+/+}*, $n = 8$; *Mcl-1^{fl/fl}*, $n = 7$) or thymus (*Mcl-1^{+/+}*, $n = 6$; *Mcl-1^{fl/fl}*, $n = 6$), as analyzed by flow cytometry. (D) Numbers of lymphocytes and neutrophils in blood from *Cpa3-Cre* mice (*Mcl-1^{+/+}*, $n = 11$; *Mcl-1^{fl/fl}*, $n = 10$). (E) Numbers of monocytes (*Mcl-1^{+/+}*, $n = 8$; *Mcl-1^{fl/fl}*, $n = 8$) and eosinophils (*Mcl-1^{+/+}*, $n = 4$; *Mcl-1^{fl/fl}*, $n = 7$) in blood from *Cpa3-Cre* mice. (F) Numbers of platelets and erythrocytes in blood from *Cpa3-Cre* mice (*Mcl-1^{+/+}*, $n = 11$; *Mcl-1^{fl/fl}*, $n = 10$).



PSA. In IgE-sensitized *Cpa3-Cre; Mcl-1^{+/+}* mice, injection of antigen resulted in marked hypothermia and, in some mice, death (Figure 7). In contrast, IgE-sensitized *Mcl-1^{fl/fl}* mice were essentially unresponsive to identical antigen challenge, and none of the mice died (Figure 7).

Discussion

Our data support the conclusion that the hematopoietic survival factor Mcl-1 is required for normal levels of mouse mast cell and basophil development and/or survival in vitro or in vivo. These results are of particular interest given that whereas the majority (~89%) of peritoneal mast cells in our *Cpa3-Cre; Mcl-1^{fl/fl}* mice exhibited relatively high levels of *Cpa3* promoter-driven expression of Cre recombinase, as assessed in *Cpa3-Cre; mT/mG* mice (Figure 2A), populations of each of 3 types of granulocytes (basophils, eosinophils, and neutrophils) in such mice exhibited high levels of *Cpa3* promoter-driven expression of Cre recombinase in only a small proportion (~13%-15%) of the cells (Figure 2D-F). However, we found that numbers of basophils were markedly reduced in *Cpa3-Cre; Mcl-1^{fl/fl}* mice (Figure 4A), whereas numbers of blood, spleen, and BM eosinophils (Figure 4E and supplemental Figure 4A) and blood, BM, and peritoneal cavity neutrophils (Figure 4B and D) were not significantly affected (and there was a modest increase in the number of neutrophils in the spleen; Figure 4B). We do not know the reason for the differential

effects of the *Cpa3-Cre; Mcl-1^{fl/fl}* genotype on levels of basophils as opposed to eosinophils or neutrophils, but one possibility is that Mcl-1 is a more important antiapoptotic factor in basophils than in the other types of granulocytes.

In addition to exhibiting marked reductions in the numbers of mast cells and basophils, *Cpa3-Cre; Mcl-1^{fl/fl}* mice exhibited significant reductions in the features of IgE-dependent biologic responses that are known to require mast cells or basophils. Therefore, both the tissue swelling (Figure 5A) and leukocyte infiltration (Figure 5C) associated with IgE- and mast cell-dependent PCA reactions²³ were markedly and significantly reduced in *Cpa3-Cre; Mcl-1^{fl/fl}* mice, with any minor tissue inflammation at the PCA reaction sites probably reflecting the presence of small residual populations of mast cells in the skin of some of the *Cpa3-Cre; Mcl-1^{fl/fl}* mice (Figure 5D). The latter conclusion is supported by our observation that the tissue swelling and leukocyte infiltration associated with IgE-dependent PCA reactions in *Cpa3-Cre; Mcl-1^{fl/fl}* mice were significantly increased (versus values in contralateral control ears) in mice that had been engrafted in the ear pinnae with 0.5 or 2.0 million in vitro-derived mast cells 8 weeks before eliciting the PCA reaction (supplemental Figure 6A-C and E-F). The latter result, and the demonstration that mast cells can be engrafted into the peritoneal cavity of *Cpa3-Cre; Mcl-1^{fl/fl}* mice (supplemental Figure 6D), support the conclusion that *Cpa3-Cre; Mcl-1^{fl/fl}* mice will be useful for studies of mast cell function in vivo. *Cpa3-Cre; Mcl-1^{fl/fl}* mice also exhibited little or no hypothermia when tested for their ability to develop IgE-dependent PSA




Figure 5. *Cpa3-Cre; Mcl-1^{fl/fl}* mice exhibit markedly reduced mast cell-dependent tissue swelling and leukocyte recruitment in IgE-dependent PCA. The data shown in panels A through C were pooled from the 3 independent experiments performed, each of which gave similar results. (A) *Cpa3-Cre; Mcl-1^{fl/fl}* mice ($n = 9$) and *Cpa3-Cre; Mcl-1^{+/+}* control mice ($n = 12$) were sensitized by intradermal injection of 20 ng of anti-DNP IgE into the right ear pinna, with vehicle injection into the left ear pinna as a control. Mice were challenged by retro-orbital injection of 100 μg of DNP-HSA the next day. Ear swelling was measured at the indicated time points and data are shown as means $\pm$ SEM. $^{**}P < .01$ and $^{***}P < .001$ versus corresponding values for *Cpa3-Cre; Mcl-1^{+/+}* mice at the indicated time points. $^{+}P < .05$, $^{++}P < .01$, and $^{+++}P < .001$ versus corresponding values for contralateral vehicle control ears in mice of the same genotype at the indicated time points. $P < .0001$ by 2-way ANOVA for the swelling response in ears sensitized with anti-DNP IgE in *Cpa3-Cre; Mcl-1^{fl/fl}* versus *Cpa3-Cre; Mcl-1^{+/+}* mice. (B-C) Numbers of mast cells (B) and leukocytes (C) in the dermis of ear pinnae of *Cpa3-Cre; Mcl-1^{fl/fl}* mice ($n = 9$) and *Cpa3-Cre; Mcl-1^{+/+}* control mice ($n = 12$) 6 hours after induction of PCA (IgE) or after in vehicle-treated control ears (vehicle). Data are shown as means $\pm$ SEM for mast cells or leukocytes per millimeter of dermis. $^{***}P < .001$ versus corresponding values for *Cpa3-Cre; Mcl-1^{+/+}* mice; $^{+++}P < .001$ versus corresponding values for vehicle-treated mice in mice of the same genotype. (D) Giemsa staining to demonstrate mast cells (top, in 1-μm-thick, Epon-embedded sections) and H&E staining to demonstrate leukocytes (bottom, in 4-μm-thick, paraffin-embedded sections) in ear pinnae 6 hours after induction of PCA reactions. There are decreased numbers of mast cells (solid arrowheads in upper panels) and leukocytes (some indicated by solid arrows in lower panels) in sections from *Cpa3-Cre; Mcl-1^{fl/fl}* mice (right panels) versus *Cpa3-Cre; Mcl-1^{+/+}* mice (left panels) subjected to PCA. Top: Some mast cells (*) exhibit alterations of the staining or location of many cytoplasmic granules, changes that are indicative of degranulation, whereas other mast cells (+) exhibit few or no granules exhibiting such changes. Bottom: Black arrows indicate leukocytes (primarily neutrophils). Insets depict selected mast cells exhibiting morphological evidence of extensive (*) or minimal or no (+) degranulation. Scale bars indicate 50 μm (for insets, scale bars indicate 5 μm); e indicates epidermis. Images were captured with an Olympus BX60 microscope with a 20 $\times$ objective using a Retiga-2000R QImaging camera run by Image-Pro Plus Version 6.3 software (Media Cybernetics) and exported into Adobe Photoshop (CS3), in which images were white balanced and resized.

(Figure 7), as would be expected given that these mice have markedly reduced levels of the 2 key effector cells of such responses, mast cells and basophils.

In a form of IgE-dependent inflammation that has been shown to depend on basophils but not mast cells,²⁵ both the tissue swelling (Figure 6A) and the leukocyte infiltration (Figure 6B) associated with the IgE-dependent CAI responses in the skin were essentially ablated in the *Cpa3-Cre; Mcl-1^{fl/fl}* mice. Therefore, even though blood basophils are less severely depleted relative to wild-type numbers in *Cpa3-Cre; Mcl-1^{fl/fl}* mice than are ear skin mast cells (with reductions of 74% versus 97%, respectively), the basophil deficiency was sufficient to result in a marked reduction in the IgE-dependent CAI response. It is possible that the number of basophils required to orchestrate IgE-dependent CAI exceeds that

found in *Cpa3-Cre; Mcl-1^{fl/fl}* mice. Alternatively, perhaps the residual blood basophils in *Cpa3-Cre; Mcl-1^{fl/fl}* mice have impaired survival after recruitment to tissues and/or are functionally abnormal in ways that limit their ability to orchestrate IgE-dependent CAI.

It remains to be determined whether the induction of biologic responses other than the ones tested in the present study would reveal any significant abnormalities in hematopoietic lineages in addition to basophils and mast cells in *Cpa3-Cre; Mcl-1^{fl/fl}* mice. We detected only minor abnormalities in lineages other than mast cells and basophils when we analyzed the mice under baseline conditions. Moreover, when we performed pilot experiments with a "mast cell-independent" model of antigen-induced chronic allergic inflammation of the airways,³²⁻³⁴ examination of C57BL/6J-*Ki^{+/+}* and C57BL/6J-*Ki^{W-sh/W-sh}* mice showed that the recruitment of




Figure 6. *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice have markedly reduced basophil-dependent tissue swelling and leukocyte recruitment in IgE-dependent CAI. (A) *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice ($n = 10$), and *Cpa3-Cre*; *Mcl-1^{+/+}* control mice ($n = 14$) were sensitized passively by IV injection of 300 μ g of IgE anti-TNP. Mice were challenged the next day by intradermal injection of 10 μ g of TNP-OVA into the left ear pinna and OVA (as a control) into the right ear pinna. Ear swelling was measured daily and the results are shown as means $\pm$ SEM. The data shown were pooled from the 3 independent experiments performed, each of which gave similar results. * $P < .05$; ** $P < .01$; and *** $P < .001$ comparing results of ears of IgE-treated *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice versus corresponding values for *Cpa3-Cre*; *Mcl-1^{+/+}* mice at the indicated time points. $P < .0001$ by 2-way ANOVA comparing the ear swelling responses in IgE-injected *Cpa3-Cre*; *Mcl-1^{fl/fl}* versus *Cpa3-Cre*; *Mcl-1^{+/+}* mice. (B) Immunohistochemical visualization of basophils by staining with an anti-mMCP8 Ab (DAB substrate) and Giemsa counterstaining (top panel, in 4- μ m-thick, paraffin-embedded sections) and H&E staining to demonstrate leukocytes (bottom panel, in 4- μ m-thick, paraffin-embedded sections) in ear pinnae 3 days after induction of CAI reactions. There are markedly decreased numbers of basophils (brown) and leukocytes in sections from *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice (right) versus *Cpa3-Cre*; *Mcl-1^{+/+}* control mice (left) subjected to CAI. Scale bars indicate 100 μ m (for insets, scale bars indicate 25 μ m); *, ear cartilage; e, epidermis. Images were captured with an Olympus BX60 microscope with a 20 $\times$ objective using a Retiga-2000R QImaging camera run by Image-Pro Plus Version 6.3 software (Media Cybernetics) and exported into Adobe Photoshop (CS3), in which images were white balanced and resized.

leukocytes (including eosinophils and neutrophils, as assessed in bronchoalveolar lavage [BAL] fluid) and airway hyperreactivity to methacholine in this model can occur in *C57BL/6J-Kit^{W-sh/W-sh}* mice that virtually lack mast cells^{20,31,35,36} and that also have increased levels of neutrophils^{20,35,36} and basophils²⁰ in the blood and BM (supplemental Figure 7). We found that *Cpa3-Cre*;

Mcl-1^{fl/fl} mice developed levels of airway hyperreactivity to methacholine that were similar to those of the corresponding *Cpa3-Cre*; *Mcl-1^{+/+}* control mice (supplemental Figure 7). However, compared with *Cpa3-Cre*; *Mcl-1^{+/+}* mice, *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice developed lower levels of BAL fluid eosinophils and higher levels of BAL fluid neutrophils. Additional work will be necessary to assess whether such differences (which did not achieve statistical significance) are confirmed in experiments with larger groups of mice, as well as to assess the extent to which any reproducible differences in levels of eosinophils, neutrophils, or other leukocytes in the BAL fluid in this model (or in any other mouse model of host defense or disease) reflected intrinsic abnormalities affecting these cells in *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice versus downstream consequences of their mast cell and/or basophil deficiency. Indeed, in both guinea pigs and mice, attempts to deplete basophils selectively also resulted in secondary reductions in eosinophils at sites of tick infestation³⁷ or IgE-dependent CAI,³⁸ respectively.

Whereas it will be important to continue to assess the phenotype of *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice, our findings to date—particularly the marked impairment in mast cell- and IgE-dependent PCA, IgE-dependent PSA, and basophil- and IgE-dependent CAI of the skin—suggest that *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice will be useful for in vivo analyses of the functions of mast cells and basophils in health and disease. With the notable exception of the small mast cell population in the spleen, *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice have a mast cell deficiency that is similar in magnitude to that observed in the genetically mast cell-deficient mice most commonly used for studies of mast cell function in vivo, *WBB6F1-Kit^{W/W-v}* mice and *C57BL/6-Kit^{W-sh/W-sh}* mice.^{31,39} However, unlike *WBB6F1-Kit^{W/W-v}* or *C57BL/6-Kit^{W-sh/W-sh}* mice, *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice lack mutations affecting c-Kit structure or expression, and therefore the phenotypic consequences of these mutations that affect populations other than mast cells will not have to be taken into account when interpreting experiments using *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice. Moreover, we^{9,10} and others^{11,12} have pointed out that mast cells and basophils may perform overlapping or complementary functions in several models of host defense or disease. *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice




Figure 7. IgE-dependent PSA is markedly reduced in *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice. *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice ($n = 6$) and *Cpa3-Cre*; *Mcl-1^{+/+}* control mice ($n = 6$) were sensitized by IP injection of 20 ng of anti-DNP IgE. Mice were challenged by IP injection of 1 mg DNP-HSA the next day. Body temperature measured at the indicated time points is shown as mean $\pm$ SD. ** $P < .05$; *** $P < .001$ versus corresponding values for *Cpa3-Cre*; *Mcl-1^{+/+}* mice at the indicated time points. $P < .01$ by 2-way ANOVA for the change in temperature in *Cpa3-Cre*; *Mcl-1^{fl/fl}* mice versus *Cpa3-Cre*; *Mcl-1^{+/+}* mice. The data shown were pooled from the 2 independent experiments performed, each of which gave similar results. The death head symbols indicate when single mice in that group died.

will permit us to examine such models in mice that are markedly and functionally deficient in both mast cells and basophils.

Acknowledgments

The authors thank Chen Liu, Christine Chang, Mariola Liebersbach, and Justin Tang for excellent technical assistance, Dr Laurent Reber for critical advice regarding the manuscript, and Drs Hajime Karasuyama and Fu-Tong Liu for providing valuable reagents.

This study was supported by grants from the National Institutes of Health (AI070813, AI023990, and CA072074 to S.J.G.; HL087936 to C.-C.C.; and CA009151 to J.N.L.).

References

- Kozopas KM, Yang T, Buchan HL, Zhou P, Craig RW. MCL1, a gene expressed in programmed myeloid cell differentiation, has sequence similarity to BCL2. *Proc Natl Acad Sci U S A*. 1993;90(8):3516-3520.
- Dzhagalov I, Dunkle A, He YW. The anti-apoptotic Bcl-2 family member Mcl-1 promotes T lymphocyte survival at multiple stages. *J Immunol*. 2008;181(1):521-528.
- Dzhagalov I, St John A, He YW. The antiapoptotic protein Mcl-1 is essential for the survival of neutrophils but not macrophages. *Blood*. 2007;109(4):1620-1626.
- Opferman JT, Letai A, Beard C, Sorcinelli MD, Ong CC, Korsmeyer SJ. Development and maintenance of B and T lymphocytes requires antiapoptotic MCL-1. *Nature*. 2003;426(6967):671-676.
- Steimer DA, Boyd K, Takeuchi O, Fisher JK, Zambetti GP, Opferman JT. Selective roles for antiapoptotic MCL-1 during granulocyte development and macrophage effector function. *Blood*. 2009;113(12):2805-2815.
- Zhou P, Qian L, Bieszczyk CK, et al. Mcl-1 in transgenic mice promotes survival in a spectrum of hematopoietic cell types and immortalization in the myeloid lineage. *Blood*. 1998;92(9):3226-3239.
- Aichberger KJ, Herndlhofer S, Agis H, et al. Liposomal cytarabine for treatment of myeloid central nervous system relapse in chronic myeloid leukaemia occurring during imatinib therapy. *Eur J Clin Invest*. 2007;37(10):808-813.
- Berent-Maoz B, Salemi S, Mankuta D, Simon HU, Levi-Schaffer F. TRAIL mediated signaling in human mast cells: the influence of IgE-dependent activation. *Allergy*. 2008;63(3):333-340.
- Wedemeyer J, Tsai M, Galli SJ. Roles of mast cells and basophils in innate and acquired immunity. *Curr Opin Immunol*. 2000;12(6):624-631.
- Kawakami T, Galli SJ. Regulation of mast cell and basophil function and survival by IgE. *Nat Rev Immunol*. 2002;2(10):773-786.
- Marone G, Triggiani M, Genovese A, De Paulis A. Role of human mast cells and basophils in bronchial asthma. *Adv Immunol*. 2005;88:97-160.
- Crivellato E, Travan L, Ribatti D. Mast cells and basophils: a potential link in promoting angiogenesis during allergic inflammation. *Int Arch Allergy Immunol*. 2010;151(2):89-97.
- Didichenko SA, Spiegl N, Brunner T, Dahinden CA. IL-3 induces a Pim1-dependent antiapoptotic pathway in primary human basophils. *Blood*. 2008;112(10):3949-3958.
- Reynolds DS, Stevens RL, Gurley DS, Lane WS, Austen KF, Serafin WE. Isolation and molecular cloning of mast cell carboxypeptidase A. A novel member of the carboxypeptidase gene family. *J Biol Chem*. 1989;264(33):20094-20099.
- Serafin WE, Dayton ET, Gravalles PM, Austen KF, Stevens RL. Carboxypeptidase A in mouse mast cells. Identification, characterization, and use as a differentiation marker. *J Immunol*. 1987;139(11):3771-3776.
- Voehringer D, Shinkai K, Locksley RM. Type 2 immunity reflects orchestrated recruitment of cells committed to IL-4 production. *Immunity*. 2004;20(3):267-277.
- Taghon T, Yui MA, Rothenberg EV. Mast cell lineage diversion of T lineage precursors by the essential T cell transcription factor GATA-3. *Nat Immunol*. 2007;8(8):845-855.
- Feyerabend TB, Terszowski G, Tietz A, et al. Deletion of *Notch1* converts pro-T cells to dendritic cells and promotes thymic B cells by cell-extrinsic and cell-intrinsic mechanisms. *Immunity*. 2009;30(1):67-79.
- Franco CB, Chen CC, Drukker M, Weissman IL, Galli SJ. Distinguishing mast cell and granulocyte differentiation at the single-cell level. *Cell Stem Cell*. 2010;6(4):361-368.
- Piliponsky AM, Chen CC, Grimaldeston MA, et al. Mast cell-derived TNF can exacerbate mortality during severe bacterial infections in C57BL/6-*KiW-sh/W-sh* mice. *Am J Pathol*. 2010;176(2):926-938.
- Galli SJ, Hammel I. Unequivocal delayed hypersensitivity in mast cell-deficient and beige mice. *Science*. 1984;226(4675):710-713.
- Ugajin T, Kojima T, Mukai K, et al. Basophils preferentially express mouse mast cell protease 11 among the mast cell tryptase family in contrast to mast cells. *J Leukoc Biol*. 2009;86(6):1417-1425.
- Wershil BK, Wang ZS, Gordon JR, Galli SJ. Recruitment of neutrophils during IgE-dependent cutaneous late phase reactions in the mouse is mast cell-dependent. Partial inhibition of the reaction with antiserum against tumor necrosis factor- α . *J Clin Invest*. 1991;87(2):446-453.
- Liu FT, Bohn JW, Ferry EL, et al. Monoclonal dinitrophenyl-specific murine IgE antibody: preparation, isolation, and characterization. *J Immunol*. 1980;124(6):2728-2737.
- Mukai K, Matsuoka K, Taya C, et al. Basophils play a critical role in the development of IgE-mediated chronic allergic inflammation independently of T cells and mast cells. *Immunity*. 2005;23(2):191-202.
- Metz M, Schafer B, Tsai M, Maurer M, Galli SJ. Evidence that the endothelin A receptor can enhance IgE-dependent anaphylaxis in mice. *J Allergy Clin Immunol*. 2011;128(2):424-426 e421.
- Kalesnikoff J, Huber M, Lam V, et al. Monomeric IgE stimulates signaling pathways in mast cells that lead to cytokine production and cell survival. *Immunity*. 2001;14(6):801-811.
- Zon LI, Gurish MF, Stevens RL, et al. GATA-binding transcription factors in mast cells regulate the promoter of the mast cell carboxypeptidase A gene. *J Biol Chem*. 1991;266(34):22948-22953.
- Walsh JC, Dekoter RP, Lee HJ, et al. Cooperative and antagonistic interplay between PU.1 and GATA-2 in the specification of myeloid cell fates. *Immunity*. 2002;17(5):665-676.
- Muzumdar MD, Tasic B, Miyamichi K, Li L, Luo L. A global double-fluorescent Cre reporter mouse. *Genesis*. 2007;45(9):593-605.
- Grimbaldeston MA, Chen CC, Piliponsky AM, Tsai M, Tam SY, Galli SJ. Mast cell-deficient *W-shash c-kit* mutant *KiW-sh/W-sh* mice as a model for investigating mast cell biology in vivo. *Am J Pathol*. 2005;167(3):835-848.
- Kung TT, Stelts D, Zurcher JA, et al. Mast cells modulate allergic pulmonary eosinophilia in mice. *Am J Respir Cell Mol Biol*. 1995;12(4):404-409.
- Takeda K, Hamelmann E, Joetham A, et al. Development of eosinophilic airway inflammation and airway hyperresponsiveness in mast cell-deficient mice. *J Exp Med*. 1997;186(3):449-454.
- Williams CM, Galli SJ. Mast cells can amplify airway reactivity and features of chronic inflammation in an asthma model in mice. *J Exp Med*. 2000;192(3):455-462.
- Nigrovic PA, Gray DH, Jones T, et al. Genetic inversion in mast cell-deficient *Wsh* mice interrupts corin and manifests as hematopoietic and cardiac aberrancy. *Am J Pathol*. 2008;173(6):1693-1701.
- Zhou JS, Xing W, Friend DS, Austen KF, Katz HR. Mast cell deficiency in *KiW-sh* mice does not impair antibody-mediated arthritis. *J Exp Med*. 2007;204(12):2797-2802.
- Brown SJ, Galli SJ, Gleich GJ, Askenase PW. Ablation of immunity to *Amblyomma americanum* by anti-basophil serum: cooperation between basophils and eosinophils in expression of immunity to ectoparasites (ticks) in guinea pigs. *J Immunol*. 1982;129(2):790-796.
- Obata K, Mukai K, Tsujimura Y, et al. Basophils are essential initiators of a novel type of chronic allergic inflammation. *Blood*. 2007;110(3):913-920.
- Kitamura Y, Go S, Hatanaka K. Decrease of mast cells in W/W^v mice and their increase by bone marrow transplantation. *Blood*. 1978;52(2):447-452.

Authorship

Contribution: J.N.L., C.-C.C., K.M., M.J.B., C.B.F., J.K., M.Y., and A.M.P. performed the experiments; J.N.L., C.-C.C., K.M., M.J.B., C.B.F., J.K., M.Y., M.T., A.M.P., and S.J.G. designed the research and analyzed the data; J.N.L. and S.J.G. wrote the manuscript; and all authors reviewed and helped to edit the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Stephen J. Galli, MD, Department of Pathology, L-235, Stanford University School of Medicine, 300 Pasteur Dr, Stanford, CA 94305-5324; e-mail: sgalli@stanford.edu.

Figure S1. Comparison of GFP expressing versus non-expressing spleen and bone marrow granulocytes in *Cpa3-Cre; mT/mG* mice. Dot plots depicting GFP⁺ (green) and GFP⁻ (grey) populations of spleen (A-C) or bone marrow (D-F) basophils (A, D), eosinophils (B, E), and neutrophils (C, F) from *Cpa3-Cre; mT/mG* mice. In all three cases, GFP⁺ cells express similar levels of cell-specific markers (top plots) and forward versus side scatter (bottom plots). Results are representative of the similar results that were obtained in the three independent experiments performed.

Figure S1




Figure S2. *Cpa3-Cre; Mcl-1^{fl/fl}* mice have markedly reduced numbers of mast

cells. (A) Numbers of mast cells in various tissues in 3-10 month old *Cpa3-Cre; Mcl-1^{+/+}* control mice ($n=7-10$) or *Cpa3-Cre; Mcl-1^{fl/fl}* mice ($n=4-10$). Numbers of mast cells are shown as mean + SEM per mm or mm² of 4 μ m-thick paraffin sections stained with 0.1% Toluidine blue (see Methods). For quantification of dura mater mast cells, dura were excised intact and mounted on a glass slide, dried, and subsequently stained with 0.1% Toluidine blue. Numbers of mast cells were assessed for the entire tissue and shown as mean + SEM per mm². Mice ranged from 3 months to 10 months of age; mouse age did not influence mast cell numbers in each genotype (data not shown). (B-D) Numbers (B) or percentage (C) of mast cells and percentages of basophils (D) in various tissues in 6 week old *Cpa3-Cre; Mcl-1^{+/+}* control mice ($n=3$) or *Cpa3-Cre; Mcl-1^{fl/fl}* mice ($n=3$). (A-D) * = $P < 0.05$; ** = $P < 0.01$; *** = $P < 0.001$ versus corresponding values for *Cpa3-Cre; Mcl-1^{+/+}* mice.

Figure S2





Figure S3. Diminished production of mast cells *in vitro* from bone marrow

cells of *Cpa3-Cre; Mcl-1^{fl/fl}* mice. Femoral and tibial bone marrow cells from *Mcl-1^{fl/fl}* mice and *Cpa3-Cre; Mcl-1^{fl/fl}* mice were cultured for 6 weeks in IL-3-supplemented medium to generate bone marrow derived cultured mast cells (BMCMCs). Every week, total cell numbers were counted and cultures were reseeded at 2.5×10^5 cells/ml. As is usual for such cultures, the % of identifiable BMCMCs in the cultures increased over time, reaching > 95% by 6 weeks. (A) After the usual drop in cell numbers during the first two weeks of culture, the total number of bone marrow-derived cells increased over time in the *Mcl-1^{fl/fl}* cultures but there were few surviving cells in cultures from *Cpa3-Cre; Mcl-1^{fl/fl}* marrow. The results shown represent the best survival of cells from *Cpa3-Cre; Mcl-1^{fl/fl}* mice observed in the three independent experiments performed. In the other two experiments, none of the *Cpa3-Cre; Mcl-1^{fl/fl}* cells survived the 6 week differentiation period. (B) Western blotting for Mcl-1 protein in cultured cells from the experiment shown in (A). At 1 and 6 weeks *in vitro*, surviving *Cpa3-Cre; Mcl-1^{fl/fl}* cells continued to express Mcl-1. Methods: 5×10^5 bone marrow-derived cells were subjected to electrophoresis and Western blotting using a polyclonal antibody against Mcl-1 (Rockland) and reprobed with anti-GAPDH antibody to show loading. Bands represent the long (35 kDa) and short (32 kDa) forms of Mcl-1. Results are representative of the similar results that were obtained in the three Western blots performed.

Figure S3

A



B




Figure S4. *Cpa3-Cre; Mcl-1^{fl/fl}* mice do not have reduced numbers of eosinophils, monocytes, macrophages, or B cells. Percentages are depicted as values from individual mice with bars indicating mean $\pm$ SEM. All comparisons between values from mice of the two genotypes are not significant ($P > 0.05$). (A) Percentage of eosinophils (CCR3⁺; SSC^{hi}) in the live cell population isolated from *Cpa3-Cre* mouse spleen (*Mcl-1*^{+/+}, $n=7$; *Mcl-1*^{fl/fl}, $n=8$) and bone marrow (*Mcl-1*^{+/+}, $n=6$; *Mcl-1*^{fl/fl}, $n=6$), as analyzed by flow cytometry. (B) Percentage of monocytes and macrophages (F4/80⁺; Gr-1⁻) in the live cell population isolated from *Cpa3-Cre* mouse spleen (*Mcl-1*^{+/+}, $n=10$; *Mcl-1*^{fl/fl}, $n=11$), bone marrow (*Mcl-1*^{+/+}, $n=5$; *Mcl-1*^{fl/fl}, $n=4$), and peritoneal lavage fluid (*Mcl-1*^{+/+}, $n=10$; *Mcl-1*^{fl/fl}, $n=10$), as analyzed by flow cytometry. (C) Percentage of B cells (B220⁺) in the live cell population isolated from *Cpa3-Cre* mouse spleen (*Mcl-1*^{+/+}, $n=10$; *Mcl-1*^{fl/fl}, $n=11$) and peritoneal lavage fluid (*Mcl-1*^{+/+}, $n=12$; *Mcl-1*^{fl/fl}, $n=11$), as analyzed by flow cytometry.

Figure S4




Figure S5. *Cpa3-Cre; Mcl-1^{fl/fl}* mice exhibit a mild macrocytic anemia. Red

blood cell indices from blood collected from *Cpa3-Cre* mice (*Mcl-1^{+/+}*, *n*=11; *Mcl-1^{fl/fl}*, *n*=10)

are depicted as data from individual mice with bars indicating mean $\pm$ SEM. *** = $P < 0.001$

versus corresponding values from *Cpa3-Cre; Mcl-1^{+/+}* control mice. Hematocrit = volume of red

blood cells as a percentage of blood volume; HGB = concentration of hemoglobin of whole

blood in g/dL; MCV = mean red blood cell volume in dL; MCHC = average concentration of

hemoglobin in all red blood cells in g/dL. Gray boxes depict normal reference ranges, as reported

in “Hematology of the Laboratory Mouse”, The Mouse in Biomedical Research, 2007.

Figure S5




Figure S6. Enhancement of IgE-dependent passive cutaneous anaphylaxis

(PCA) in mast cell engrafted *Cpa3-Cre; Mcl-1^{fl/fl}* mice. For mice whose results are shown in A-C, BMCMCs derived from female WT (C57BL/6J) mice were transferred by intradermal injection (5×10^5 cells in 30 μ l PBS in a single injection into each ear pinna) into 6-week old female *Cpa3-Cre; Mcl-1^{fl/fl}* mice (BMCMCs $\rightarrow$ *Cpa3-Cre; Mcl-1^{fl/fl}* mice) 8 weeks before initiating the experiment. Mast cell engraftment was confirmed by histology in 4/4 *Cpa3-Cre; Mcl-1^{fl/fl}* mice injected with BMCMCs in this experiment. Age-matched *Cpa3-Cre; Mcl-1^{fl/fl}* ($n=4$) and BMCMCs $\rightarrow$ *Cpa3-Cre; Mcl-1^{fl/fl}* mice ($n=4$) were sensitized by intradermal injection of 20 ng anti-DNP IgE into the right ear pinna, with vehicle injection into the left ear pinna as a control. Mice were challenged by retro-orbital injection of 100 μ g DNP-HSA the next day. (A) Ear swelling was measured at the indicated time points and is shown as mean $\pm$ SD. * = $P < 0.05$ versus corresponding values for *Cpa3-Cre; Mcl-1^{fl/fl}* mice at the indicated time points. + = $P < 0.05$ and ++ = $P < 0.01$ versus corresponding values for contralateral vehicle control ears at the indicated time points. $P < 0.01$ by two-way ANOVA for the swelling response in IgE-sensitized ears of BMCMC-engrafted *Cpa3-Cre; Mcl-1^{fl/fl}* mice versus *Cpa3-Cre; Mcl-1^{fl/fl}* controls. (B and C) Numbers of mast cells (B) and leukocytes (C) in the dermis of ear pinnae were counted 6 h after antigen challenge. Data are shown as mean + SD for mast cells or leukocytes per mm of dermis. * = $P < 0.05$ versus corresponding values for *Cpa3-Cre; Mcl-1^{fl/fl}* mice and + = $P < 0.05$ versus corresponding values for contralateral vehicle control ears by the Mann-Whitney U test (two-tailed). (D) BMCMCs derived from female WT (C57BL/6J) mice were transferred by intraperitoneal injection (1×10^6 cells in 300 μ l PBS) into four 6-week old female *Cpa3-Cre; Mcl-1^{fl/fl}* mice (BMCMCs $\rightarrow$ *Cpa3-Cre; Mcl-1^{fl/fl}* mice). 8 weeks after engraftment, the percentage of mast cells in the live cell population isolated from peritoneal lavage fluid from *Cpa3-Cre;*

Mcl-1^{fl/fl} (n=4) or BMCMCs→*Cpa3-Cre; Mcl-1^{fl/fl}* (n=4) mice was analyzed by flow cytometry (FcεRI⁺; c-Kit⁺). Peritoneal mast cell numbers are shown as values from individual mice with bars indicating mean + SD. *P* = 0.029 versus corresponding values for *Cpa3-Cre; Mcl-1^{fl/fl}* mice by the Mann-Whitney U test (two-tailed). (E-F) BMCMCs derived from WT (C57BL/6J) mice were transferred by intradermal injection (each ear received two injections; 1 x 10⁶ cells/25 μl into each of two sites in each ear pinna) into 6-week old female *Cpa3-Cre; Mcl-1^{fl/fl}* mice (BMCMCs→*Cpa3-Cre; Mcl-1^{fl/fl}* mice) 8 weeks before initiating the experiment. Mast cell engraftment was confirmed by histology in 3/5 *Cpa3-Cre; Mcl-1^{fl/fl}* mice injected with BMCMCs in the two experiments whose results (including those for the 3 *Cpa3-Cre; Mcl-1^{fl/fl}* mice that were successfully engrafted with mast cells) are pooled in E. (E) Age-matched *Cpa3-Cre; Mcl-1^{+/+}* (n=5), *Cpa3-Cre; Mcl-1^{fl/fl}* (n=5) and BMCMCs→*Cpa3-Cre; Mcl-1^{fl/fl}* mice (n=3) were sensitized as in (A) and challenged by retro-orbital injection of 100 μg DNP-HSA in 0.25% Evan's blue. Ear swelling is shown as in (A). * *P* < 0.05 and ** *P* < 0.01 versus corresponding values for *Cpa3-Cre; Mcl-1^{fl/fl}* mice at the indicated time points. + *P* < 0.05 and ++ *P* < 0.01 versus corresponding values for contralateral vehicle control ears at the indicated time points. *P* = 0.0019 and *P* < 0.0001 by two-way ANOVA for the ear swelling responses in *Cpa3-Cre; Mcl-1^{fl/fl}* mice versus BMCMCs→*Cpa3-Cre; Mcl-1^{fl/fl}* mice and versus *Cpa3-Cre; Mcl-1^{+/+}* mice, respectively. (F) Representative visualization of Evan's blue extravasation at 30 minutes after antigen challenge from one of the two experiments depicted in (E). Right ear = IgE-treated and left ear = vehicle control.

Figure S6




Figure S7. *Cpa3-Cre; Mcl-1^{fl/fl}* mice sensitized with OVA and alum develop

features of OVA-induced chronic allergic inflammation of the airways. (A) OVA-treated B6-*Kit^{+/+}* control mice ($n=4$) and B6-*Kit^{W-sh/W-sh}* mice ($n=4$) and (C) OVA-treated *Cpa3-Cre; Mcl-1^{+/+}* control mice ($n=4$) and *Cpa3-Cre; Mcl-1^{fl/fl}* mice ($n=4$) exhibited similar changes in specific airway resistance (sRaw, measured as $\text{cmH}_2\text{O}\cdot\text{sec}$) to methacholine versus baseline sRaw (sRaw at 0 mg/ml of methacholine). As a measure of airway hyperresponsiveness, changes in sRaw induced by increasing amounts of aerosolized methacholine were measured 24 hours after the last exposure of mice to OVA or PBS aerosol using the FinePointTM NAM system (BUXCO Research System). $+ = P < 0.05$ and $++ = P < 0.01$ versus corresponding values for PBS-treated mice in mice of the same genotype at 40 mg/ml methacholine. Data are depicted as mean $\pm$ SD. (B, D) Numbers of leukocytes in BAL fluid from B6-*Kit^{+/+}*, B6-*Kit^{W-sh/W-sh}*, *Cpa3-Cre; Mcl-1^{+/+}*, or *Cpa3-Cre; Mcl-1^{fl/fl}* mice after OVA sensitization and OVA challenge or, in corresponding control mice, after treatment with PBS instead of OVA at the time of OVA sensitization and challenge. BAL was performed 24 hours after the last aerosol exposure to OVA or PBS. Data are depicted as values from individual mice with bars indicating mean $\pm$ SD. By the Mann-Whitney U test (two-tailed), no differences were observed between the results for similarly-treated B6-*Kit^{+/+}* versus B6-*Kit^{W-sh/W-sh}* mice or *Cpa3-Cre; Mcl-1^{+/+}* versus *Cpa3-Cre; Mcl-1^{fl/fl}* mice. Note the differences in the scales used in A versus C and in B versus D.

Figure S7

