



Diverse age-incidence patterns of atopic sensitization in an unselected Finnish population up to 12 years



Kaisa Pyrhönen, MD, PhD^{*}; Petri Kulmala, MD, PhD^{†,‡}; Simo Näyhä, MD, PhD[§]; Esa Läärä, Lic Soc Sc^{||}

^{*} Center for Life Course Health Research, University of Oulu, Oulu, Finland

[†] PEDEGO Research Unit and MRC Oulu, University of Oulu and Oulu University Hospital, Oulu, Finland

[‡] Biomedicine Research Unit, Medical Microbiology and Immunology, University of Oulu, Oulu, Finland

[§] Faculty of Medicine, University of Oulu, Oulu, Finland

^{||} Research Unit for Mathematical Sciences, University of Oulu, Oulu, Finland

ARTICLE INFO

Article history:

Received for publication December 10, 2018.

Received in revised form February 8, 2019.

Accepted for publication February 28, 2019.

ABSTRACT

Background: The temporal sequence in which allergic sensitization to different allergens emerges is not well characterized at the level of general population.

Objective: We describe the incidence patterns of atopic sensitization to different allergens from birth up to 12 years of age in an unselected Finnish population.

Methods: The study population comprised all children born between 2001 and 2006 identified from the nationwide population register as residents of the province of South Karelia, Finland (n = 5564). The results of allergy tests (22,380 results from skin prick tests, immunoglobulin E [IgE] antibodies, and open food challenges [OFCs], performed in 1827 children) were collected from patient records of all the health care units in the area.

Results: The incidence rates of positive results for food and animal allergens as well as positive OFCs for cow's milk showed prominent peaks at 5 months of age. Positive results for pollen allergens started to emerge after 1.5 years of age. The 12-year cumulative incidence of sensitization to food, animal, pollen, and any allergens was 12%, 8%, 10%, and 18%, respectively. The cumulative incidence of sensitization to house dust mites was 1% and to molds or latex less than 1%. Firstborn boys had the highest, and those who were not firstborn girls and children born in rural municipalities had the lowest early incidence of sensitization to inhalation allergens.

Conclusion: In the unselected population, the atopic sensitization against food and animal allergens began before 6 months of age and was followed by sensitization to pollen allergens before 2 years of age. Primary prevention of sensitization to food and inhalation allergens should therefore occur in early infancy.

© 2019 American College of Allergy, Asthma & Immunology. Published by Elsevier Inc. All rights reserved.

Introduction

The clinically observed progression of atopic diseases is often described in terms of an "atopic march", from atopic eczema to respiratory allergies (allergic rhinitis and/or asthma).^{1–5} The

incidence of atopic manifestations has previously been defined by using the first date of hospital admission for these conditions or dispensed prescription of specific medication⁶ or by the age at diagnoses of atopic manifestations.⁷

Reprints: Kaisa Pyrhönen, MD, PhD, Center for Life Course Health Research, University of Oulu, P.O. Box 5000, FIN-90014 Oulu, Finland; E-mail: kaisa.pyrhonen@oulu.fi.

Disclosures: None.

Funding Sources: The data collection and the work of the principal investigator (the first author) in years 2005 to 2006 was mainly funded by the Social Insurance Institution of Finland and partly by EVO grants from the hospital districts of South Karelia and Northern Ostrobothnia, by Lappeenranta City Council and personal grants from the Finnish Cultural Foundation, South Karelia Regional fund, Finland; the Viipuri Tuberculosis Foundation, Finland; the Väinö and Laina Kivi Foundation, Finland; the Tyyni Tani Foundation, Finland; Kymenlaakson Terveysturva ry, Finland; the Allergy Foundation, Finland; and the Medical Society of South Karelia,

Finland. The work of the principal investigator was funded in years 2012 to 2014 by University of Oulu and Lappeenranta University of Technology. In 2016 to 2019, the extended data collection and the work of the principal investigator was funded by the Finnish Cultural Foundation (Pekka and Jukka-Pekka Lylykari's Fund), Finland; South Karelia Regional Fund (Lauri and Lahja Hotinen Fund), Finland and by the Viipuri Tuberculosis Foundation, Finland, and the Finnish Pediatric Research Foundation, Finland. The work of the second author was supported by the research grants from the Alma and K.A. Snellman Foundation, Finland; the Finnish Medical Association, Finland; the Allergy Research Foundation, Finland and the Finnish Pediatric Research Foundation, Finland. None of the funding organizations was involved in the design or execution of the study.




Figure 1. Flow diagram showing the data sources and data linkages of the South Karelia Allergy Research Project (the SKARP).

The information on the occurrence of allergic sensitization and different allergies by age has been derived from population-based birth cohorts, mainly from the age of 1 year, in single or repeated cross-sectional settings,^{8–18} a laboratory database of positive specific immunoglobulin E (IgE) test results among tested subjects¹⁹ or clinical experience.²⁰ Allergen-specific IgEs were detected from birth in the Danish Allergy Research Centre (DARC)¹⁶ and the Protection Against Allergy-Study in Rural Environment (PASTURE)¹⁰ cohorts, the latter of which was partly carried out in Finland, with the aim of finding preventive environmental and genetic factors of atopy.²¹ However, the patterns of incidence of sensitization to different allergens from birth until early teenage years has been inadequately described at the level of the general population.

The South Karelian Allergy Research Project^{22–25} is a large observational population-based cohort study, which contains the information on exact dates and results of allergy testing during a long follow-up period. We report the age-incidence patterns of sensitization to different allergens from birth up to 12 years of age, using survival methods. From a public health perspective, our data add to previous descriptions of allergic sensitization and food allergies and will be useful for health care authorities, physicians, and primary prevention.

Methods

Definition of the Population

The target population comprised all children born between April 2001 and March 2006 who resided in the province of South Karelia in southeast Finland at the time they were invited to the questionnaire survey, which was conducted during the years 2005

and 2006.²² The target population (N = 5973) of our study was identified and their personal identity codes (PIC; including information on the date of birth) obtained from the Population Register Centre of Finland (Fig 1). The current paper focuses on the entire population, covering both the participants and the nonparticipants of the survey.

The parents of 53 children had refused the use of the PIC in the questionnaire, and their data were excluded. For the remaining 5920 children, information on vital status and migrations by September 2013 was available for 5902 children from the Population Register Centre. Children who had immigrated to the study area after 1 month of age (n = 356) were excluded, and those with missing information of migration history (n = 18) were included in the data analyses. Thus, the final study population comprised 5564 children (Fig 1).

Data Collection

The information on all allergy tests was collected from all of the health care units in the area, with the intention of covering the entire target population. These tests had been performed for diagnostic purposes between April 2001 and September 2013. The PIC was used to merge the individual test data with population register data.

Ethics and Permissions

The study protocol was reviewed and approved by the Ethical Committee of the Northern Ostrobothnia Hospital District (95/2003) and the South Karelia District of Social and Health Services (979/13.01.02/2014). The test data were collected with the

Table 1

Cumulative Incidences (% from Kaplan-Meier Analyses) and Number (n) of Children Tested^a for Different Allergies and Those with a Positive Test Result up to 12 Years of Age among the Entire SKARP Population (n = 5564)

Allergen	Tested children	Children with a positive test
	% (n)	% (n)
Food	27 (1371)	12 (594)
Essential food items (any) ^b	27 (1344)	11 (541)
Cow's milk	26 (1311)	8 (401)
Cow's milk supplementary formula ^c	16 (750)	2 (97)
Soya	15 (735)	2 (85)
Hen's egg	19 (908)	5 (249)
Cereal (wheat/rye/oat/barley/etc)	21 (1056)	4 (209)
Non-essential foods (any) ^d	15 (692)	3 (131)
Peanuts, tree nuts, or almond	13 (570)	2 (92)
Animal	25 (1231)	8 (378)
Dog	24 (1170)	7 (300)
Cat	24 (1150)	6 (266)
Horse	15 (822)	2 (97)
Pollen	23 (1081)	10 (409)
Birch	22 (1073)	7 (342)
Alder	14 (694)	4 (184)
Mugwort	21 (989)	3 (107)
Timothy	22 (1029)	6 (208)
Other		
House dust mite	19 (903)	1 (43)
Mold ^e	14 (602)	1 (31)
Latex	6 (310)	0 (2)
Other specific allergen eg, medicine, vaccine, poison of wasp or bee, etc.	1 (70)	0 (3)
Any test above	36 (1827)	18 (874)

^aSpecific immunoglobulin E antibodies or skin prick tests for the allergen(s) in question, or additionally an open food challenge regarding food items.

^bEssential food items (any) include either cow's milk, milk supplementary formula, egg, or cereals (wheat/barley/rye/etc).

^cIncluding extensively hydrolyzed and amino acid formulae, and soya.

^dNonessential food items (any) include any other food items than above.

^eInclude tests for the following mold allergens: *Cladosporium herbarum*, *Aspergillus fumigatus*, *Alternaria alternata* (*A tenuis*), *Penicillium chrysogenum* (*P notatum*), *Candida albicans*, and *Helminthosporium halodes*.

permission of the Finnish Ministry of Social Affairs and Health (49/07/2003) and the National Institute for Health and Welfare (THL/1490/5.05.01/2014 and THL/1519/5.05.00/2014).

Outcomes and Explanatory Variables

Skin prick tests (SPTs) were performed by trained local laboratory personnel or nurses, using standard commercial extracts and using negative controls and histamine as a positive control. The SPTs were often ordered and performed as panels including the most common allergens either for food allergens (cow's milk, hen's egg, wheat, barley, rye, gliadin, oat, fish, and often soya or nut), or inhalation allergens (dog, cat, horse, birch, alder, timothy, mugwort, meadow fescue, or cocksfoot grass, *Dermatophagoides pteronyssinus* or *D farinae*, *Cladosporium herbarum*), or both, or a contact allergen, latex, and sometimes for single allergens. Specific immunoglobulin E antibodies (sIgE) were used as an alternative method for SPTs depending on their availability in the health care center. Specific IgE had been performed for 34 and SPT for 28 individual allergens or group (eg, root vegetables, fruits, spices, legumes, molds, grasses, vaccines, or medicines) of allergens.

As the outcome variables, we considered the occurrence of 1) the first test, that is, SPT, sIgE, or open food challenge (OFC), and 2) the first positive result from any of these tests for any allergen, and separately for pollen, animal, and food allergens, and also separately for different food and other allergens. Cow's milk, milk supplementary formula (extensively hydrolyzed/amino acid formulae/soya), hen's egg, or cereals (wheat/barley/rye/oat, etc)

were considered as *essential food items* and other food items as *nonessential food items*.

The cutoff point for a positive sIgE result was 0.35 kU/L with RAST-CAP FEIA or Phadiatop Combi and 1.43 standardized units per milliliter with Magic Lite and for SPTs as the mean of 2 orthogonal diameters of the urticarial wheal of 3 mm, applied to both positive and negative controls. A wheal diameter of 3 mm or more was regarded as a positive SPT result. Most OFCs were supervised by a pediatrician in the hospitals of the study area; exceptionally, 3 OFCs in 2 children were performed in health centers. The OFC was considered "positive" when labeled as such in the patient records (the food allergy diagnosis made by the pediatrician, who supervised the OFC).

The information on sex, birth order, and the first place of residence were based on the data obtained from the Population Register Centre and used as explanatory variables.

Statistical Methods

For each allergen or a combination of them, the outcome event was either the first allergy test performed, or the first positive test, as appropriate. The risk time was primarily defined as the period between the child's date of birth and the date of the event. Where a pertinent event was not observed during the follow-up, the risk time ended either on the date of death (4 children died), date of emigration from the study area (to other regions in Finland or abroad) for a period longer than 1 year, or on the closing date of data collection (September 30, 2013), whichever occurred first. In analyses of the cumulative incidence of the first event among competing, that is, mutually exclusive alternative events, the risk time would also end on the date of the first of the competing events.

The R environment release 3.5.1 was used for computations and graphics.²⁶ The incidence rates for the outcomes by age were analyzed and presented as follows.^{27,28} For each outcome, the follow-up was organized and split into bimonthly intervals, or age bands, using functions *Lexis* and *splitLexis* of the *Epi* package in R.²⁹ In each age band, the incidence rate was calculated as the number of cases of the outcome in question occurring in the interval divided by the total person-time (summed risk times) accumulated in the same age band and expressed as cases per 100 child-years. Smoothed estimates of incidence rates were obtained as predictions from a Poisson regression model using function *glm*, in which the effect of age was smoothed by natural cubic splines,²⁸ with prespecified knots at the ages of 0, 0.25, 0.5, 1, 2, 4, 6, 8, 10, and 12 years. The smoothed age-incidence curves were graphically presented, also showing the empirical bimonthly incidence rates at the midpoint of each interval (eFigs 1 and 2).

The cumulative incidence proportions of each outcome event by age were first computed by the Kaplan-Meier method,^{30,31} in which possible competing events were treated as independent censorings. In the analyses addressing the cumulative incidence of the major combinations of allergen groups, the Aalen-Johansen method³² was applied to allow for competing events. Computations of cumulative incidence were performed by the *survfit* function of the *survival* package.³³ The Cox regression model, fitted by function *coxph* in R, was used to estimate the incidence rate ratios of the positive test for animal and pollen allergens associated with a history of previous sensitization (a positive sIgE/SPT) to food allergens and of a positive OFC.

Results

Number of Tests and Tested Children

Among the entire population (N = 5564), 22,380 individual allergy tests for specific IgE-mediated sensitization or allergy had been performed on altogether 1827 children (39,973 child-years of follow-up; median, 8.42 years), and 4120 tests had yielded a




Figure 2. Smoothed age-incidence curves of *testing* for food (any food test, food, or open food challenge, OFC), animal, pollen, and any allergens (A) and the curves of *positive test results* (B) among the SKARP population ($n = 5564$).

positive result in 874 children (47,242 child-years of follow-up; median, 9.23 years) (Table 1). At least 1 positive result in sIgE or SPT, that is, *sensitization* was observed in 803 children (3713 positive tests) out of 1822 children tested, the cumulative incidence being 17% by 12 years of age.

Altogether 706 OFCs had been performed on 295 children, of which 407 OFCs had been positive in 207 children. Most OFCs had been performed for cow's milk ($n = 245$ children). Other OFCs had been performed for cereals ($n = 95$), hen's egg ($n = 14$), soya or milk supplementary formulas ($n = 81$), fish ($n = 11$), pork meat ($n = 2$), potato ($n = 1$), and nut ($n = 1$).

Incidence Rates of Positive Test Results

The incidence rates of sensitization to food and animal allergens reached a prominent peak at approximately 5 months of age, whereas the incidence rates of sensitization to pollen allergens increased monotonically up to 12 years of age (Fig 2 and eFig 1). The incidence rates of the first positive test results for cow's milk allergens showed a peak at 5 months of age (Fig 3 and eFig 2). Other early incidence peaks of the positive tests were those for hen's egg and cereal allergens (wheat, barley, rye, oat, or other cereal).

The incidence rates of *any food allergy* (a positive OFC), cow's milk allergy, and cow's milk allergy with sensitization to cow's milk allergens peaked around the age of 6 months (Fig 4). Altogether 168 children had OFC-confirmed cow's milk allergy. Among them, 59 (35%) had sensitization to cow's milk allergens, 53 (32%) to some other allergens, 56 (33%) to none of the tested specific allergens, and for 1 child no results of SPT or sIgE tests were found. The incidence rates of cereal allergy, and cereal allergy with sensitization, peaked at the age of 10 months. Altogether 59 children had OFC-confirmed cereal allergy, and among them 32 children had cereal allergy with sensitization to cereal allergens.

Cumulative Incidence of Positive Test Results

The cumulative incidences for the first positive test were 12% for food, 8% for animal, 10% for pollen, and 18% for any allergens by 12 years of age (Table 1, Fig 5). Positive tests for food, most commonly for cow's milk (Table 1, Figs 5 and 6), started to emerge soon after the birth and those for animal and pollen allergens at 3 months and 1.5 years of age (Fig 5), respectively. The cumulative incidence of food allergy (a positive OFC) was 4%, cow's milk allergy 3%, and cow's milk allergy with sensitization 1%.

The cumulative incidence curves of sensitization to pollen and animal allergens crossed at 7 years of age (Fig 5), and those of birch and pet (dog and cat) allergens at approximately 5 years of age (eFig 3). Sensitization to tree pollen started to arise earlier than sensitization to grass pollen allergens.

The most common combination of the first positive test(s) was that for food allergens alone, whereas food allergens together with animal or pollen allergens, or for any of the latter allergens alone, were much more rarely found combinations of the first positive tests (Fig 5). The most common combination of the first positive food tests was that for cow's milk allergens alone (Fig 6).

Previous sensitization to food allergens (a positive sIgE/SPT) and food allergy (a positive OFC) was a strong predictor of subsequent sensitization to animal allergens (rate ratio, 37.5; 95% confidence interval, 28.9–48.5 and 29.1; 21.2–40.6, respectively) and of subsequent sensitization to pollen allergens (37.6; 29.3–48.2 and 30.1; 22.3–40.6, respectively).

Other Allergic Sensitizations

Altogether 903 children were tested for house dust mite allergy (SPTs performed on 539 children and sIgE tests on 466), but positive results were found in only 43 cases (Table 1). A positive result




Figure 3. Smoothed age-incidence curves of testing for cow's milk, hen's egg, cereals, and any essential allergens (A) and the curves of positive test results (B) among the SKARP population ($n = 5564$).

for mold allergens was found in only 31 children out of 602 tested subjects. Only 2 children had a positive test result for latex allergens, although 307 children were tested by SPT and 5 by sIgE for latex. Overall, 650 children were tested for fish or other seafood allergens, but only 23 children had positive results.

Incidence Rates by Sex, Birth Order, and First Residence

Incidence rates of positive test results for cow's milk and hen's egg allergens formed early incidence peaks in all subgroups according to the birth order and sex (Fig 7). Early incidence peaks also could be seen for sensitization to animal allergens among firstborn boys and girls, whereas such early peaks were not seen for not firstborn infants. The incidence curves of sensitization to pollen allergens started to arise after the first year of life in all subgroups, most steeply among the firstborn boys. Children with the first residence in rural municipalities had slightly lower incidence rates of sensitization to inhalation allergens than others, but within the first 2 years their incidence curves for cow's milk and hen's egg allergens formed at least as high early peaks as other children.

Discussion

The current paper describes the age-incidence patterns of sensitization for different allergens from birth up to 12 years of age in an unselected general pediatric population. The incidence rates of positive test results started to increase soon after birth, the peak

incidence for food and animal allergens being reached at the age of 5 months. In contrast, no unequivocal early peak was seen for sensitization to pollen allergens during the follow-up time. The cumulative incidence of sensitization to tree and grass pollen allergens started to rise after 1.5 year and 2.5 years of age, respectively.

Our quantitative estimates for the incidence of positive test results are in line with clinical experience⁵ on the "allergic march" and also with a recent questionnaire survey from Sweden,⁹ according to which food allergies (and atopic eczema) typically appear first and are followed by rhinoconjunctivitis and asthma. Previous information on the age-specific occurrence of sensitization is mainly based on point prevalence figures for sensitization after the first year of life obtained from cross-sectional studies and can be compared with our incidence figures only with caution.

According to previous studies, the occurrence of sensitization to food allergens varies depending on the food items studied, the test methods used, the duration of follow-up period (age), and participation rates.³⁴ In the Isle of Wight cohort,¹¹ the prevalence of food allergy at 4 years of age, based on SPTs only, was 3.5%. In primary care cohorts from the United States,⁷ the incidence of food allergy for any food item during the first 5 years of life was 8.2%, the peak incidence at 12 to 17 months of age, and we observed a similar figure for the cumulative incidence of sensitization to food allergens (8.6%), with the peak incidence occurring at approximately 6 months of age.




Figure 4. Smoothed age-incidence curves of testing for sensitization (sIgE or skin prick test, SPT, performed) to essential foods (A), the curves of sensitization (B), and the curves of open food challenge (OFC) confirmed food allergy to cow's milk and cereals, also with respective sensitization (C) among the SKARP population ($n = 5564$).

In our real-life setting, the first OFCs had been most commonly performed for essential rather than nonessential food items, usually before 2 years of age. By the age of 2 years, the incidence of challenge-proven cow's milk allergy was less than 1% in EuroPreval cohorts,³⁵ which is in line with the incidence of IgE-mediated OFC-confirmed cow's milk allergy in our study (1%). In the SchoolNuts study (in Australia),¹² the prevalence of OFC-confirmed food allergy was 4.5% and most common for peanuts among students aged 10 to 14 years, whereas the incidence of OFC-confirmed food allergies was similar in our population (4%), although in early childhood only, and most commonly for cow's milk. In these 2 studies from the United States and Australia,^{7,12} the prevalence of peanut allergy (2.6% and 2.7%, respectively) agree well with our cumulative incidence of sensitization (not necessarily allergy) to peanuts, tree nuts, or almond combined (2.4%, 92/5564).

In our study, the cumulative incidence of sensitization to inhalant allergens was lower than the prevalence of sensitization to inhalant allergens at the respective ages in the Multicenter Allergy Study from Germany (MAS)⁸ (1.5% and 26% at 1 and 6 years, respectively, although

the cutoff for sIgE was higher; ≥ 0.7 kU_A/L), the PARIS¹⁷ (2.5% at 1.5 years), and the Isle of Wight cohorts¹¹ (19.2% at 4 years). In the retrospective cohort study from the United States,⁷ the prevalence of rhinitis was also higher than sensitization to inhalant allergens in our study (5.6%, 13.7%, 21.7%, 23.3%, and 24.8% at ages of 0–3, 3–5, 6–10, 11–13, and 14–17 years, respectively). These differences might be explained by different study settings, variable definition of the outcomes, statistical methods used (point prevalence vs cumulative incidences with censoring), and genetic variations between populations in different countries.

Our population-based findings provide further support to the conclusion of the PASTURE cohort,¹⁰ according to which any potential preventive measures to curtail atopic sensitization should be timed at prenatal or early postnatal periods. In the DARC¹⁶ and the PASTURE cohorts,¹⁰ positive sIgEs for cat and dog allergens were detected at birth, whereas in our cohort the tests were performed for diagnostic purposes months or years after birth. In addition, we were able to show early incidence peaks for sensitization to animal and food allergens and their temporal order calculated as




Figure 5. Cumulative incidences by age of first testing (A, C) and positive tests (B, D) for food, animal, pollen, and any allergen shown by Kaplan-Meier curves (A, B) and their first combinations by stacked curves (Aalen-Johansen method; competing risks) (C, D) among the SKARP population ($n = 5564$).

competitive risk estimates in a general population. Preventing early sensitization is also justified because it can be considered as a primary prevention of allergic rhinitis and asthma.³⁶

According to our results, the sensitization to house dust mite allergens, molds, and latex were rarities; the occurrence of the former has been reported to be very low because of a cold and dry climate in other Nordic countries.³⁷ The prevalence of sensitization to house dust mite (based on sIgEs) was 1.2% at 18 months of age in the PARIS cohort¹⁷ and 12% (a positive SPT) at 4 years of age in the Isle of Wight cohort.¹¹ In a study population aged 7 years from the United States,³⁸ the prevalence of sensitization to mold allergens was 12% and 5% among children with and without asthma, respectively. Sensitization to natural rubber has recently decreased on account of the increased use of powder-free, low-latex gloves³⁹ and baby's dummies made of silicon replacing latex dummies. The development of allergic sensitization to house dust mites, mold, or latex also may require longer exposure periods than sensitization to

food or animal allergens. Because the available allergy tests mostly indicate IgE-mediated allergy, possible underlying non-IgE-mediated mechanisms may remain undiagnosed.

In our study population, the proportion of sensitization was 44% (803/1822) among children tested by either sIgE or SPT, and 40% (572/1444) if tested by sIgE only, whereas the proportion of children with at least 1 positive sIgE out of all repeatedly tested children at 4, 8, and 16 years of age was 51% (1328/2607) in the BAMSE cohort.¹³ Additionally, a Dutch study¹⁹ from a hospital laboratory reports 40% of sIgE tested children having a positive test result.

In accordance with previous reports,^{6,24,25,40,41} the incidences of positive test results for different allergens were higher in boys and firstborns than others. Data from the questionnaire survey are available and can be linked with these test results in the future. Because our focus was to show results among the general population, presenting occurrences of sensitization according to the questionnaire-based background variables would have restricted




Figure 6. Cumulative incidences by age of first testing (A, C) and positive test results (B, D) to cow's milk, cereals, hen's egg, and essential foods shown by Kaplan-Meier (A, B) and their first combinations by stacked curves (Aalen-Johansen method; competing risks) (C, D) among the SKARP population (n = 5564).

the study population to the questionnaire survey participants only. Additionally, the trajectories of sensitization in the same population may be possible to analyze in the future.

The main strength of this work is the population-based cohort identified from a nationwide population register, which could be individually linked with the results of all allergy tests conducted for the diagnostic purposes on cohort members from birth up to 12 years of age. The record linkage to the national population register also allowed us to take into account the migration histories of almost all cohort members, enabling us to properly allow for censoring during the follow-up time. The follow-up time of 12 years seems to be adequate to describe the early phases of sensitization to food, animal, tree pollen, and grass pollen allergens. However, the cumulative incidence of the first positive test result for animal, pollen, and nonessential food (eg, nut) allergens were still increasing in the end of the current follow-up period.

In our study, positive results in SPT or sIgE indicate allergic sensitization.⁴² Tests of molecular allergology (the third step of diagnostic workup) had been performed for very few children at the end of the follow-up only; therefore, results of these tests were not included here. Oral food challenges are considered a gold standard (an ideal method, a double-blind, placebo-controlled food challenge, was not available in our study area) and a cornerstone of food allergy diagnosis.^{43,44} In our population, positive OFCs can be considered to indicate the physician-diagnosed food allergies confirmed in a hospital, which probably underestimate the total

number of food allergies in the general population, because no data were available for OFCs carried out at home because of mild or delayed symptoms.

The main limitation of this study is a lack of information on specific indications of physician-ordered allergy tests such as specific symptoms of atopy (atopic eczema, rhinoconjunctivitis, or asthma) or parental concern only. In Finland, all children are followed up approximately 25 to 30 times in the child health clinics, scheduled to take place monthly during the first year of life, at 18 months of age, and thereafter annually until 16 years of age. During these visits, the public health nurses have the possibility of consulting a physician or pediatrician, who will order allergy tests if needed. Our data also cover allergy tests ordered by private doctors and doctors in the municipal health centers. However, the tests had probably not been performed for asymptomatic children among whom sensitization to allergens also could have occurred. Therefore, observational studies such as ours obviously underestimate the occurrences of sensitization compared with studies in which almost all recruited participants have been tested. However, because of possible false-positives, our sensitization rates probably overestimate the rates of clinical allergies.

Another limitation of our study is that the study population covered only 1 province of Finland, South Karelia. The collection of allergy test data would have been too expensive and even impossible to carry out in a larger area. South Karelia includes both rural (population density, 2.69 inhabitants/km²) and urban areas




Figure 7. Smoothed incidence curves (cases per 100 child-years) by age of first testing (greater plots) for cow's milk, hen's egg, animal, and pollen allergens and the respective curves for first positive test results (nested plots; note the different scaling) according to sex and birth order and first residence among the SKARP population (n = 5564).

(182/km²), the district of the Lake Saimaa, with several paper and pulp mills, and their surroundings. These characteristics of the area restrict the generalizability of our results to some extent.

Based on our real-world data, allergic sensitization was observed first for food and animal allergens, both reaching their incidence peak at the age of approximately 5 months, followed by gradually increasing allergic sensitization to pollen allergens after 18 months of age. General population-based information on the temporal order, in which sensitization to different allergens and physician-diagnosed allergies are observed by age, is useful for physicians when choosing appropriate allergy tests and explaining the natural course of allergic conditions to the parents. This information is useful in primary prevention. Obviously, preventive measures should be targeted at the age before the incidence of the allergic sensitization starts to arise in a general population.

Acknowledgments

We thank all the nurses in the child health clinics of South Karelia for their cooperation and the staff of the various health care units for their assistance and cooperation in collecting the test data. We also thank Mr. Markku Koiranen for his skillful technical assistance with the data management in the early phase of the study.

Supplementary Data

Supplementary data related to this article can be found online at <https://doi.org/10.1016/j.janai.2019.02.027>

References

- Ker J, Hartert TV. The atopic march: what's the evidence? *Ann Allergy Asthma Immunol*. 2009;103:282–289.
- Spergel JM. From atopic dermatitis to asthma: the atopic march. *Ann Allergy Asthma Immunol*. 2010;105:99–106.
- Wahn U. The significance of environmental exposure on the progression of allergic diseases. *Allergy*. 2011;66:7–9.
- Zheng T, Yu J, Oh MH, Zhu Z. The atopic march: progression from atopic dermatitis to allergic rhinitis and asthma. *Allergy Asthma Immunol Res*. 2011;3:67–73.
- Thomsen SF. Epidemiology and natural history of atopic diseases. *Eur Clin Respir J*. 2015;2. <https://doi.org/10.3402/ecrj.v2.24642>.
- Henriksen L, Simonsen J, Haerskjold A, et al. Incidence rates of atopic dermatitis, asthma, and allergic rhinoconjunctivitis in Danish and Swedish children. *J Allergy Clin Immunol*. 2015;136:360–366.
- Hill DA, Grundmeier RW, Ram G, Spergel JM. The epidemiologic characteristics of healthcare provider-diagnosed eczema, asthma, allergic rhinitis, and food allergy in children: a retrospective cohort study. *BMC Pediatr*. 2016;16:133.
- Kulig M, Bergmann R, Klettke U, et al. Natural course of sensitization to food and inhalant allergens during the first 6 years of life. *J Allergy Clin Immunol*. 1999;103:1173–1179.
- Goksör E, Loid P, Alm B, Åberg N, Wennergren G. The allergic march comprises the coexistence of related patterns of allergic disease not just the progressive development of one disease. *Acta Paediatr*. 2016;105:1472–1479.
- Depner M, Ege MJ, Genuneit J, et al. Atopic sensitization in the first year of life. *J Allergy Clin Immunol*. 2013;131:781–788.
- Arshad SH, Tariq SM, Matthews S, Hakim E. Sensitization to common allergens and its association with allergic disorders at age 4 years: a whole population birth cohort study. *Pediatrics*. 2001;108:E33.
- Sasaki M, Koplin JJ, Dharmage SC, et al. Prevalence of clinic-defined food allergy in early adolescence: the SchoolNuts study. *J Allergy Clin Immunol*. 2018;141:391–398.
- Ballardini N, Bergström A, Wahlgren C, et al. IgE antibodies in relation to prevalence and multimorbidity of eczema, asthma, and rhinitis from birth to adolescence. *Allergy*. 2016;71:342–349.
- Grabenhenrich LB, Keil T, Reich A, et al. Prediction and prevention of allergic rhinitis: a birth cohort study of 20 years. *J Allergy Clin Immunol*. 2015;136:940.
- Nissen SP, Kjær HF, Høst A, Nielsen J, Halken S. The natural course of sensitization and allergic diseases from childhood to adulthood. *Pediatr Allergy Immunol*. 2013;24:549–555.
- Kjaer HF, Eller E, Andersen KE, Høst A, Bindslev-Jensen C. The association between early sensitization patterns and subsequent allergic disease: the DARC birth cohort study. *Pediatr Allergy Immunol*. 2009;20:726–734.
- Gabet S, Just J, Couderc R, Bousquet J, Seta N, Momas I. Early polysensitization is associated with allergic multimorbidity in PARIS birth cohort infants. *Pediatr Allergy Immunol*. 2016;27:831–837.
- Goksör E, Alm B, Pettersson R, et al. Asthma and allergy from infancy into school age: the allergic march revisited. *Clin Transl Allergy*. 2014;4(suppl 1):P5.
- Baatenburg de Jong A, Dikkeschei LD, Brand PLP. Sensitization patterns to food and inhalant allergens in childhood: a comparison of non-sensitized, monosensitized, and polysensitized children. *Pediatr Allergy Immunol*. 2011;22:166–171.
- Barnetson R, Rogers M. Childhood atopic eczema. *Br Med J*. 2002;324:1376–1379.
- Mutius E, Schmid S. The PASTURE project. EU support for the improvement of knowledge about risk factors and preventive factors for atopy in Europe. *Allergy*. 2006;61:407–413.
- Pyrhönen K, Läärä E, Kaila M, Hiltunen L, Näyhä S. SKARP: a population-based cohort study of food associated symptoms and food allergies in childhood: design, methods, and participation. *Scand J Public Health*. 2011;39:194–202.
- Pyrhönen K, Näyhä S, Läärä E. Dog and cat exposure and respective pet allergy in early childhood. *Pediatr Allergy Immunol*. 2015;26:247–255.
- Pyrhönen K, Hiltunen L, Näyhä S, Läärä E, Kaila M. Real-life epidemiology of food allergy testing in Finnish children. *Pediatr Allergy Immunol*. 2011;22:361–368.
- Pyrhönen K, Hiltunen L, Kaila M, Näyhä S, Läärä E. Heredity of food allergies in an unselected child population: an epidemiological survey from Finland. *Pediatr Allergy Immunol*. 2011;22:e132.
- R Development Core Team. R: a language and environment for statistical computing. 2019. Available at: <https://CRAN.R-project.org/>. Accessed March 25, 2019.
- Carstensen B, Plummer M. Lexis: an R class for epidemiological studies with long-term follow-up. *J Stat Softw*. 2011;38:1–12.
- Harrell F. *Regression Modeling Strategies: With Applications to Linear Models, Logistic and Ordinal Regression, and Survival Analysis*. New York, NY: Springer-Verlag; 2015.
- Carstensen B, Plummer M, Läärä E, Hills M. Epi: a package for statistical analysis in epidemiology. R package version 2.35. 2019. Available at: <https://CRAN.R-project.org/package=Epi>. Accessed March 25, 2019.
- Clayton D, Hills M. Competing risks and selection. In: Clayton D, Hills M, eds. *Statistical Models in Epidemiology*. New York, NY: Oxford University Press; 1993:63–70.
- Greenland S. Applications of stratified analysis methods. In: Rothman KJ, Greenland S, Lash TL, eds. *Modern Epidemiology*. Philadelphia, PA: Lippincott Williams & Wilkins; 2008:283–302.
- Andersen PK, Geskus RB, de Witte T, Putter H. Competing risks in epidemiology: possibilities and pitfalls. *Int J Epidemiol*. 2012;41:861–870.
- Therneau T. A package for survival analysis in S package version 2.38. 2015. Available at: <https://cran.r-project.org/web/packages/survival/index.html>. Accessed March 25, 2019.
- Nwaru BI, Hickstein L, Panesar SS, et al. Prevalence of common food allergies in Europe: a systematic review and meta-analysis. *Allergy*. 2014;69:992–1007.
- Schoemaker AA, Sprinkelman AB, Grimshaw KE, et al. Incidence and natural history of challenge-proven cow's milk allergy in European children: Euro-Prevall birth cohort. *Allergy*. 2015;70:963–972.
- Alduraywish SA, Standl M, Lodge CJ, et al. Is there a march from early food sensitization to later childhood allergic airway disease? Results from two prospective birth cohort studies. *Pediatr Allergy Immunol*. 2017;28:30–37.
- Munir AKM. Mite sensitization in the Scandinavian countries and factors influencing exposure. *Allergy*. 1998;53:64–70.
- Reponen T, Lockey J, Bernstein DI, et al. Infant origins of childhood asthma associated with specific molds. *J Allergy Clin Immunol*. 2012;130:639–644.
- Niggemann B. IgE-mediated latex allergy: an exciting and instructive piece of allergy history. *Pediatr Allergy Immunol*. 2010;21:997–1001.
- Arshad SH, Kurukulaaratchy RJ, Fenn M, Matthews S. Early life risk factors for current wheeze; asthma; and bronchial hyperresponsiveness at 10 years of age. *Chest*. 2005;127:502–508.
- Tariq SM, Matthews SM, Hakim EA, Stevens M, Arshad SH, Hide DW. The prevalence of and risk factors for atopy in early childhood: a whole population birth cohort study. *J Allergy Clin Immunol*. 1998;101:587–593.
- Matricardi PM, Kleine-Tebbe J, Hoffmann HJ, et al. EAACI molecular allergology user's guide. *Pediatr Allergy Immunol*. 2016;27:1–250.
- Muraro A, Werfel T, Hoffmann-Sommergruber K, et al. EAACI food allergy and anaphylaxis guidelines: diagnosis and management of food allergy. *Allergy*. 2014;69:1008–1025.
- Eigenmann PA. Do we still need oral food challenges for the diagnosis of food allergy? *Pediatr Allergy Immunol*. 2018;29:239–242.




Figure 1. Incidence rates (number of cases per 100 child-years) by age of allergy testing and positive test results for different allergens estimated by piecewise constant rate approach and smoothed by cubic splines shown by lines (dashed line: testing; solid line: positive tests). Bimonthly point estimates of incidence rates are shown by circles (open circles: testing; filled circles: positive tests) in the SKARP population (n = 5564).





Figure 2. Incidence rates (number of cases per 100 child-years) by age of allergy testing and positive test results for different food allergens estimated by piecewise constant rate approach and smoothed by cubic splines shown by lines (dashed line: testing; solid line: positive tests). Bimonthly point estimates of incidence rates are shown by circles (open circles: testing; filled circles: positive tests) in the SKARP population ($n = 5564$).



eFigure 3. Cumulative incidences by age of allergy testing (dashed lines) and positive test results (solid lines) for different allergens in the SKARP population (n = 5564).