


Journal Pre-proof

Real life small airway outcomes in severe asthma patients receiving biologic therapies

Rory Chan, MBChB, Chris RuiWen Kuo, MBChB, Brian Lipworth, MD



PII: S2213-2198(21)00157-4

DOI: <https://doi.org/10.1016/j.jaip.2021.01.029>

Reference: JAIP 3406

To appear in: *The Journal of Allergy and Clinical Immunology: In Practice*

Received Date: 23 November 2020

Revised Date: 8 January 2021

Accepted Date: 15 January 2021

Please cite this article as: Chan R, RuiWen Kuo C, Lipworth B, Real life small airway outcomes in severe asthma patients receiving biologic therapies, *The Journal of Allergy and Clinical Immunology: In Practice* (2021), doi: <https://doi.org/10.1016/j.jaip.2021.01.029>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier Inc. on behalf of the American Academy of Allergy, Asthma & Immunology

1 Title: Real life small airway outcomes in severe asthma patients receiving biologic
2 therapies

3 Authors: Rory Chan, MBChB, r.chan@dundee.ac.uk
4 Chris RuiWen Kuo, MBChB, r.kuo@dundee.ac.uk
5 Brian Lipworth, MD, b.j.lipworth@dundee.ac.uk

6 Affiliation: Scottish Centre for Respiratory Research
7 School of Medicine,
8 University of Dundee
9 Ninewells Hospital, Dundee
10 DD1 9SY

11 Correspondence: Dr Brian Lipworth
12 Scottish Centre for Respiratory Research
13 Ninewells Hospital and Medical School
14 University of Dundee, DD1 9SY
15 Scotland, UK
16 Tel: +44 (0)1382 383188; Fax: +44 (0)1382 383259
17 b.j.lipworth@dundee.ac.uk
18

19 Conflict of Interest:

20 Dr. Chan has no relevant conflicts of interest.

21 Dr. Kuo reports personal fees (talks) from AstraZeneca, personal fees (advisory board) from
22 Circassia, personal fees (talks) in relation to the submitted work, personal fees (talks) and other
23 support from Chiesi (attending BTS) outside of the submitted work.

24 Dr. Lipworth reports non-financial support (equipment) from GSK; grants, personal fees (consulting,
25 talks and advisory board), other support (attending ATS and ERS) and from AstraZeneca, grants,
26 personal fees (consulting, talks, advisory board), other support (attending ERS) from Teva, personal
27 fees (consulting) from Sanofi, personal fees (consulting, talks and advisory board) from Circassia in
28 relation to the submitted work; personal fees (consulting) from Lupin, personal fees (consulting)
29 from Glenmark, personal fees (consulting) from Vectura, personal fees (consulting) from Dr Reddy,
30 personal fees (consulting) from Sandoz; grants, personal fees (consulting, talks, advisory board),
31 other support (attending BTS) from Boehringer Ingelheim, grants and personal fees (advisory board
32 and talks) from Mylan outside of the submitted work; and the son of BJL is presently an employee of
33 AstraZeneca.

34 Funding source: None
35

36 Word count: 1,059

37 Figures: 1

38 Tables: 1

39 Key words: small airways dysfunction, severe refractory asthma, omalizumab, mepolizumab,
40 benralizumab, impulse oscillometry, FEF25-75

Clinical implications

FEF₂₅₋₇₅ should be incorporated to assess improvement in small airways dysfunction (SAD) for severe asthma patients receiving biologics. Impulse oscillometry (IOS) can be useful as an adjunct, particularly in those with baseline IOS-defined SAD.

In Scotland, the current licensed biologic therapies for severe refractory asthma include omalizumab, mepolizumab and benralizumab. These agents have been shown to significantly reduce asthma exacerbations and oral corticosteroid (OCS) use.⁽¹⁾ Although the improvements observed in patient reported outcomes and lung function measured by forced expiratory volume in 1 second (FEV₁) were statistically significant, neither achieved the minimal clinically important difference (MCID).⁽¹⁾

Small airways dysfunction (SAD) can be measured using spirometry expressed as the forced expiratory flow between 25 and 75% of forced vital capacity (FEF₂₅₋₇₅) and using impulse oscillometry (IOS) as the difference between resistance at 5 and 20 Hz (R5-R20).^(2, 3) R5-R20 is related to GINA asthma severity,⁽⁴⁾ and the use of reliever therapy and long-term OCS is significantly higher in asthmatic patients with abnormal R5-R20 but preserved FEV₁.⁽⁵⁾

As SAD is directly related to asthma severity and type 2 inflammation,⁽⁶⁾ we performed a real-life retrospective cohort study to evaluate whether biologic therapies improved SAD in adult patients with severe refractory asthma treated in the National Health Service (NHS) Tayside specialist respiratory clinic. We primarily examined the differences in spirometry (FEV₁% and FEF₂₅₋₇₅%), IOS (R5-R20), peripheral blood eosinophils (PBE), asthma control questionnaire (ACQ) and OCS-requiring exacerbations pre- and post-biologic therapy.

56 severe refractory asthma patients who had previously been treated with or were presently receiving biologic therapy with omalizumab, mepolizumab or benralizumab in accordance with national guidelines were included. Pre- and post-biologic data on ACQ, OCS-requiring exacerbations,

spirometry, IOS, and PBE were collected. Spirometry (Micromedical, Chatham, UK) and IOS (Masterscreen™ IOS, Jaeger Ltd) measurements were performed in triplicate according to the European Respiratory Society guidelines with IOS always performed first.

Data were first analysed for normality. Paired Student's T-tests with alpha error set at 0.05 (2-tailed) were applied to compare outcomes pre- and post- biologic therapy. Caldicott Guardian approval was obtained prior to data collection. Values are given as means and SEM.

The mean baseline demographic data, prior to initiation of biologic therapy, were as follows: age 54 years; 14% with aspirin exacerbated respiratory disease; 21% with nasal polyps; ACQ 3.3; FEV₁ 81%; FEF₂₅₋₇₅ 43%; R5 0.50 kPa/L/s; R5-R20 0.12 kPa/L/s; AX 1.14 kPa/L; PBE 572 cells/μL and total IgE 347 kU/L. The mean daily beclomethasone dipropionate equivalent dose of inhaled corticosteroid was 1,889μg; 93% were taking LABA; 61% LAMA; 77% LTRA; 32% theophylline; and 55% oral antihistamine. 34% of patients were taking maintenance OCS prior to biologic initiation. 30% were ex-smokers with the remainder having never smoked. 20, 30 and 6 patients were treated with omalizumab, mepolizumab and benralizumab respectively with the mean duration of therapy between pre- and post-biologic lung function measurements being 10 months. All patients underwent spirometry with 23/56 (41%) also undergoing IOS. 17 patients from the IOS subgroup received anti-IL5 therapy with the remaining receiving omalizumab.

In the overall group (Table 1), FEF₂₅₋₇₅% but not FEV₁% improved significantly pre- vs post-biologic therapy. There were also significant improvements in ACQ, OCS-requiring exacerbations, and a significant drop in PBE count.

In the SAD subgroup defined by R5-R20≥0.08 kPa/L/s (n=15) (Figure 1), R5-R20 [0.17(0.02) vs 0.11(0.02)] but not AX improved significantly pre- vs post-biologic therapy. Of these, 11 patients received anti-IL5 therapy with the remaining receiving omalizumab. Differences in ACQ [3.7(0.3) vs 1.2(0.3)], OCS-requiring exacerbations [3.8(0.2) vs 0.5(0.2)] and PBE [798(129) vs 122(35)] were also significant in this subgroup. In patients with pre-biologic FEF₂₅₋₇₅<60% (n=43), there were significant

improvements in ACQ: 3.1(0.3) vs 1.7(0.3); OCS-requiring exacerbations: 4.4(0.3) vs 0.6(0.1); and PBE: 575(65) vs 168(27) cells/ μ L, but no differences were observed in FEV₁ or IOS. Notably, 10 patients with R5-R20 $\geq$ 0.08kPa/L/s also had a pre-biologic FEF₂₅₋₇₅<60%.

When comparing patients according to biologic therapy received (Table 1), FEF₂₅₋₇₅% increased significantly in the omalizumab subgroup (15/20 patients) but not significantly in the anti-IL5 (mepolizumab and benralizumab) subgroup (21/36 patients). In both the omalizumab and anti-IL5 subgroups, ACQ and OCS-requiring exacerbations significantly decreased; however, PBE count only decreased significantly in the anti-IL5 subgroup. In the omalizumab subgroup, there were insufficient evaluable data to perform a meaningful analysis of IOS. Although ACQ and PBE improved significantly in patients who were taking maintenance OCS (n=19) at baseline, there were no significant differences observed in spirometry pre- vs post-biologic therapy.

Our study has demonstrated that severe refractory asthma patients with relatively well-preserved baseline FEV₁% experienced significant improvements in small airways function measured by FEF₂₅₋₇₅%. This was accompanied by improvements in asthma control, exacerbation frequency and PBE in line with previous studies. FEV₁% improved numerically but was not statistically significant. As expected in the anti-IL5 subgroup, there was a significant reduction in PBE counts which was not seen in the anti-IgE subgroup. ACQ improvements exceeded MCID of 0.5(7) in all groups. Our study observed that in patients with IOS defined SAD (R5-R20 $\geq$ 0.08kPa/L/s)(4) R5-R20 but not R5, R20 or FEF₂₅₋₇₅ improved significantly with biologic therapy, in conjunction with ACQ, exacerbation frequency and PBE count (Figure 1).

Interestingly, a recent observational study demonstrated that oscillometry parameters improve earlier than spirometry after benralizumab therapy in severe asthma patients.(8) Although the change in R5 (total airway resistance) was significant after 4 weeks of benralizumab, R5-R20 (reflecting peripheral airway resistance)(3) was not statistically altered. Another observational study also showed that biologic therapy responders, defined using a composite of reduction in OCS-

requiring exacerbations and asthma symptoms, had significantly higher R5-R20 at baseline compared to partial or non-responders.(9) As small airway function is directly related to GINA asthma severity(4), this could theoretically propose an argument that FEF_{25-75} and R5-R20 are perhaps the missing link between asthma control, exacerbations and T2 biomarkers. We therefore hypothesise that incorporating IOS into the standard severe asthma workup prior to initiation of biologic therapy may help identify patients most likely to benefit.

The real-life observational aspect of this study was one of its main strengths. However, we recognise the limitations of our retrospective analysis. Firstly, our data was obtained from a single Scottish centre which limits the generalisability of the results. Secondly, our patient numbers are relatively small especially in relation to IOS results and therefore a larger prospective study is required to further validate our results.

In summary our findings have shown that biologic therapies result not only in improvements in asthma control and OCS exacerbation frequency but also in spirometry small airway function. Furthermore, in a subgroup of such patients the presence of abnormal peripheral airway resistance (R5-R20) on IOS was associated with a clinical response to biologic therapy.

127 REFERENCES

- 128 1. Chan R, RuiWen Kuo C, Lipworth B. Pragmatic Clinical Perspective on Biologics for Severe
129 Refractory Type 2 Asthma. *J Allergy Clin Immunol Pract*. 2020;8(10):3363-70.
- 130 2. Konstantinos Katsoulis K, Kostikas K, Kontakiotis T. Techniques for assessing small airways
131 function: Possible applications in asthma and COPD. *Respiratory medicine*. 2016;119:e2-e9.
- 132 3. Foy BH, Soares M, Bordas R, Richardson M, Bell A, Singapuri A, et al. Lung Computational
133 Models and the Role of the Small Airways in Asthma. *American journal of respiratory and critical*
134 *care medicine*. 2019;200(8):982-91.
- 135 4. Postma DS, Brightling C, Baldi S, Van den Berge M, Fabbri LM, Gagnatelli A, et al. Exploring
136 the relevance and extent of small airways dysfunction in asthma (ATLANTIS): baseline data from a
137 prospective cohort study. *Lancet Respir Med*. 2019;7(5):402-16.
- 138 5. Manoharan A, Anderson WJ, Lipworth J, Ibrahim I, Lipworth BJ. Small airway dysfunction is
139 associated with poorer asthma control. *Eur Respir J*. 2014;44(5):1353-5.
- 140 6. Kuo CR, Jabbal S, Lipworth B. Is small airways dysfunction related to asthma control and type
141 2 inflammation? *Ann Allergy Asthma Immunol*. 2018;121(5):631-2.
- 142 7. Juniper EF, Svensson K, Mork AC, Stahl E. Measurement properties and interpretation of
143 three shortened versions of the asthma control questionnaire. *Respir Med*. 2005;99(5):553-8.
- 144 8. Shirai T, Akamatsu T, Hirai K, Watanabe H, Tamura K, Kishimoto Y, et al. Oscillometry
145 improves earlier than spirometry after benralizumab initiation in severe asthma. *Allergy*.
146 2020;75(10):2678-80.
- 147 9. Abdo M, Watz H, Veith V, Kirsten A-M, Biller H, Pedersen F, et al. Small airway dysfunction as
148 predictor and marker for clinical response to biological therapy in severe eosinophilic asthma: a
149 longitudinal observational study. *Respiratory Research*. 2020;21(1):278.

150

151 **Table 1**

	Overall (n = 56)		Anti-IL5 (n = 36)		Omalizumab (n = 20)	
	Pre	Post	Pre	Post	Pre	Post
FEV ₁ %	81 (3)	85 (3)	81 (3)	87 (3)	80 (6)	81 (6)
FEF ₂₅₋₇₅ %	43 (3)	49 (4)*	46 (4)	52 (5)	37 (5)	44 (7)*
FVC%	100 (3)	104 (3)	100 (3)	104 (3)	101 (5)	103 (5)
FEV ₁ /FVC ratio	67 (2)	68 (2)	69 (2)	71 (2)	63 (3)	64 (3)
R5 (kPa/L/s)	0.50 (0.03)	0.48 (0.03)	0.46 (0.03)	0.48 (0.04)		
R5 – R20 (kPa/L/s)	0.12 (0.02)	0.12 (0.02)	0.10 (0.02)	0.11 (0.02)		
AX (kPa/L)	1.14 (0.24)	1.30 (0.25)	0.87 (0.17)	1.39 (0.32)		
PBE (cells/ μ L)	572 (55)	152 (22)***	728 (69)	71 (10)***	309 (53)	288 (41)
OCS exacerbations	4 (0.3)	1 (0.1)***	4 (0.2)	0.5 (0.1)***	4.7 (0.7)	0.6 (0.2)***
ACQ	3.3 (0.23)	1.6 (0.21)***	3.2 (0.3)	1.5 (0.3)***	3.7 (0.3)	1.8 (0.3)***

152

153 **Table 1 Legend**

154 Comparisons in spirometry, IOS, ACQ, OCS-requiring exacerbations and PBE count pre- and post-
 155 biologic therapy. Data presented as means (SEM). ACQ = 6-point asthma control questionnaire
 156 score; AX = area under reactance curve; FEV₁ = forced expiratory volume in 1 second; FEF₂₅₋₇₅ =
 157 forced expiratory flow between 25 and 75% of forced vital capacity (FVC); OCS = oral corticosteroid;
 158 PBE = peripheral blood eosinophils; R5 = resistance at 5 Hz; R5-R20 = difference between resistance
 159 at 5Hz and 20Hz; * denotes p<0.05 and ***p<0.001

160

161 **Figure 1 Legend**

162 Values shown as means and standard error of means for significant comparisons in the small airways
 163 dysfunction (R5-R20 $\geq$ 0.08 kPa/L/s) subgroup according to (a) R5-R20 (b) peripheral blood eosinophil
 164 count (c) ACQ and (d) OCS exacerbations pre- and post-biologic therapy.

