

Correspondence

Reply

To the Editor:

We agree with Calvani et al¹ that any death related to food immunotherapy should be reported. The death referred to in our letter² occurred during parenteral therapy and has been reported.³ We apologize to the readers of the *Journal* for this misstatement.

Richard L. Wasserman, MD, PhD^a

Robert W. Sugerman, MD^a

Lyndon Mansfield, MD^b

James W. Baker, MD^c

Disclosure of potential conflict of interest: The authors have declared that they have no conflict of interest.

REFERENCES

1. Calvani M, Sopo SM, Giorgio V. Oral immunotherapy in food allergy: how difficult to weigh its risks and its benefits? *J Allergy Clin Immunol* 2011; ■■■:■■■.
2. Wasserman RL, Sugerman RW, Mireku-Akomeah N, Mansfield L, Baker JW. Office-based oral immunotherapy for food allergy is safe and effective. *J Allergy Clin Immunol* 2011;127:290-1.
3. Oppenheimer JJ, Nelson HS, Bock SA, Christensen F, Leung DY. Treatment of peanut allergy with rush immunotherapy. *J Allergy Clin Immunol* 1992;90:256-62.

doi:10.1016/j.jaci.2011.03.038

From ^aDallasAllergyImmunology, Dallas, Tex; ^bWestern Sky Medical Research, El Paso, Tex; and ^cJames W. Baker, MD, LLC, Lake Oswego, Ore. E-mail: drichwasserman@gmail.com.