Check for updates

Uter Wolfgang (Orcid ID: 0000-0002-4498-3710)
Wilkinson Mark (Orcid ID: 0000-0001-7253-3461)
Borrego Leopoldo (Orcid ID: 0000-0002-0199-2756)
Buhl Timo (Orcid ID: 0000-0002-3139-129X)
Dickel Heinrich (Orcid ID: 0000-0001-6837-4402)
Mercader Pedro (Orcid ID: 0000-0002-8309-9725)
Giménez-Arnau Ana M (Orcid ID: 0000-0001-5434-7753)
Pesonen Maria (Orcid ID: 0000-0003-0356-7064)
Pónyai Györgyi (Orcid ID: 0000-0002-1695-6133)
Schubert Steffen (Orcid ID: 0000-0002-2359-2518)
Schuttelaar Marie Louise Anna (Orcid ID: 0000-0002-0766-4382)
Simon Dagmar (Orcid ID: 0000-0001-8965-9407)
Stingeni Luca (Orcid ID: 0000-0001-7919-8141)
Gonçalo Margarida (Orcid ID: 0000-0001-6842-1360)

Accepted Article

Patch test results with the European Baseline Series, 2019/20 – Joint European results of the ESSCA[A] and the EBS[B] working groups of the ESCD, and the GEIDAC[C]

Wolfgang Uter (1; ORCID-ID: 0000-0002-4498-3710), S. Mark Wilkinson (2; ORCID-ID: 0000-0001-7253-3461), Olivier Aerts (3; ORCID-ID: 0000-0002-0076-2887), Andrea Bauer (4; ORCID-ID: 0000-0002-4411-3088), Leopoldo Borrego (5; ORCID-ID: 0000-0002-0199-2756), Richard Brans (6; ORCID-ID: 0000-0002-1245-024X), Timo Buhl (7, ORCID 0000-0002-3139-129X), Heinrich Dickel (8; ORCID-ID: 0000-0001-6837-4402), Aleksandra Dugonik (9; ORCID-ID: 0000-0002-0199-2756), Francesca Larese Filon (10; ORCID-ID: 0000-0002-7717-0417), Pedro Mercader Garcìa (11; ORCID-iD: 0000-0002-8309-9725), Ana Giménez Arnau (12; ORCID-ID: 0000-0001-5434-7753), Cataldo Patruno (13; ORCID-ID: 0000-0001-8661-2822), Maria Pesonen (14; ORCID-ID: 0000-0003-0356-7064), Györgyi Pónyai (15; ORCID-ID: 0000-0002-1695-6133), Thomas Rustemeyer (16; ORCID-ID 0000-0001-7580-0684), Steffen Schubert (17; ORCID-ID: 0000-0002-2359-2518), Marie-L.A. Schuttelaar (18; ORCID-ID: 0000-0002-0766-4382), Dagmar Simon (19; ORCID-ID: 0000-0001-8965-9407), Luca Stingeni (20; ORCID-ID: 0000-0001-7919-8141), Skaidra Valiukevičienė (21; ORCID-ID: 0000-0003-1219-5928), Elke Weisshaar (22; ORCID-ID: 0000-0002-7016-0224), Thomas Werfel (23; ORCID-ID: 0000-0001-6830-7672), Margarida Gonçalo (24; ORCID-ID: 0000-0001-6842-1360), for the ESSCA[A] and EBS[B] ESCD working groups, and the GEIDAC[C]

1. Department of Medical Informatics, Biometry and Epidemiology, Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen, Germany
2. Dermatology Department, Leeds Teaching Hospitals NHS Trust, Leeds, UK
3. Department of Dermatology, University Hospital Antwerp (UZA) and Research group Immunology, Infla-Med Centre of Excellence, University of Antwerp, Antwerp, Belgium
4. Department of Dermatology, University Hospital Carl Gustav Carus, Technical University, Dresden, Germany
5. Complejo Hospitalario Universitario Insular Materno Infantil, Universidad de las Palmas de Gran Canaria, Spain

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/cod.14170

This article is protected by copyright. All rights reserved.

6. Institute for Interdisciplinary Dermatologic Prevention and Rehabilitation (iDerm) at the University of Osnabrück, Osnabrück, Germany
7. Department of Dermatology, Venereology and Allergology, University Medical Center Göttingen, Göttingen, Germany
8. Department of Dermatology, Venereology and Allergology, St. Josef Hospital, University Medical Center, Ruhr University Bochum, Bochum, Germany
9. Department of Dermatology, University Medical Centre Maribor, Maribor, Slovenia.
10. Department of Public Health, Occupational Medicine, University of Trieste, Trieste, Italy
11. Department of Dermatology, Hospital Morales Meseguer, Murcia, Spain
12. Department of Dermatology, Hospital del Mar, Institut Mar d´Investigacions Mèdiques, Universitat Pompeu Fabra de Barcelona, Barcelona, Spain
13. Department of Health Sciences, University Magna Graecia of Catanzaro, Catanzaro, Italy
14. Division Occupational Medicine, Finnish Institute of Occupational Health, Helsinki, Finland
15. Department of Dermatology, Venerology and Dermato-oncology, Semmelweis University, Budapest, Hungary
16. Department of Dermatology, Amsterdam University Medical Centers, Amsterdam, The Netherlands
17. Information Network of Departments of Dermatology (IVDK), Institute at the University Medical Centre Göttingen, Göttingen, Germany
18. Department of Dermatology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands
19. Department of Dermatology, Inselspital, Bern University Hospital, University of Bern, Bern, Switzerland
20. Dermatology Section, Department of Medicine and Surgery, University of Perugia, Italy
21. Department of Skin and Venereal Diseases, Lithuanian University of Health Sciences, Kaunas, Lithuania
22. Occupational Dermatology, Department of Dermatology, University Hospital Heidelberg, Heidelberg, Germany
23. Division of Immunodermatology and Allergy Research, Department of Dermatology and Allergy, Hannover Medical School, Hannover, Germany
24. Department of Dermatology, University Hospital and Faculty of Medicine, University of Coimbra, Coimbra, Portugal

Accepted Article

Accepted Article

Collaborators:

[A]ESSCA working group of the ESCD and [B]EBS working group of the ESCD (members contributing data)

Birger Kränke, Department of Dermatology, Medical University of Graz, Graz, Austria, Graz, Austria (birger.kraenke@medunigraz.at); Kathrin Scherer Hofreiter, Department of Dermatology, Allergy Unit, University Hospital Basel, Basel, Switzerland, Basel, Switzerland (Kathrin.scherer@usb.ch); Alexander Navarini, NA, Basel, Switzerland (alexander.navarini@usb.ch); Jürgen Grabbe, Department of Dermatology, Kantonsspital Aarau, Aarau, Switzerland, Aarau, Switzerland (juergen.grabbe@ksa.ch); Philip Spring, Dermatologie et vénéréologie FMH, Center Médical d'Epalinges, Epalinges, Switzerland, Lausanne/Epalignes, Switzerland (springphil@hotmail.com); Ulrike Beiteke, Department of Dermatology, Dortmund, Germany, Dortmund, Germany (ulrike.beiteke@klinikumdo.de); Cecilia Dietrich, Department of Dermatology, University of Schleswig-Holstein, Campus Kiel, Kiel, Germany, Kiel, Germany (cdietrich@dermatology.uni-kiel.de); Sibylle Schliemann, Department of Dermatology and Allergology, University Hospital Jena, Jena, Germany, Jena, Germany (schliemann@derma-jena.de); Detlef Becker, Department of Dermatology, University of Mainz, Mainz, Germany, Mainz, Germany (detlef.becker@unimedizin-mainz.de); Swen Malte John, Institute for Interdisciplinary Dermatologic Prevention and Rehabilitation (iDerm), University of Osnabrück, Lower Saxony Institute for Occupational Dermatology (NIB), Osnabrück, Germany, Osnabrück, Germany (sjohn@uni-osnabrueck.de); Nicola Wagner, Department of Dermatology, University of Erlangen-Nürnberg, Erlangen, Germany, Erlangen, Germany (nicola.wagner@uk-erlangen.de); Maddalena Napolitano, Department of Medicine and Health Sciences Vincenzo Tiberio, University of Molise, Campobasso, Italy (maddy.napolitano@gmail.com) Rosella Gallo, Clinica Dermatologica, IRCCS - AOU San Martino – IST and Department of Health Sciences, University of Genoa, 16132 Genoa, Italy, Genova, Italy (rs.gallo@unige.it); Maja Kalac Pandurovic, Outpatient clinic of dermatology, Maribor, Maribor, Slovenia, Maribor/Clinic, Slovenia (maja.kp@derma-mkp.si); Tanja Kmecl, Department of Dermatology, General Hospital Celje, Celje, Slovenia, Celje, Slovenia (tanja.kmecl@guest.arnes.si); Marko Vok, Outpatient clinic of dermatology, Izola, (please edit), Slovenia, Izola, Slovenia (marko.vok@siol.net); Mojca Simončič Godnič, Department of dermatology, Novo mesto, (please edit), Slovenia, Novo Mesto, Slovenia (mojcasg78@gmail.com); Nada Kecelj, Outpatient clinc of dermatology, Ljubljana (please edit), Slovenia, Ljubljana/Clinic, Slovenia (nada.kecelj@kclj.si); Susan M. Cooper, Churchill Hospital, Oxford, United Kingdom (sm_nc@btopenworld.com); Philippa Cousen, James Cook University Hospital, Middlesbrough, United Kingdom (Philippa.Cousen@stees.nhs.uk); Ella Dendooven, University Hospital Antwerp (UZA) and Research group Immunology, Infla-Med Centre of Excellence, University of Antwerp, Antwerp, Belgium (elladendooven@hotmail.com)

[C]GEIDAC

Members of the "Grupo Español de Investigación en Dermatitis de Contacto y Alergia Cutánea" (GEIDAC) comprise (also listing co-authors): Pedro Mercader-García (Servicio de

Accepted Article

Dermatología, Hospital General Universitario Jose M a Morales Meseguer, Murcia, Spain), Juan Francisco Silvestre Salvador (Servicio de Dermatología, Hospital General Universitario de Alicante, Alicante, Spain), Javier Sánchez Pérez (Servicio de Dermatología, Hospital Universitario de La Princesa, Madrid, Spain), Virginia Fernández Redondo (Servicio de Dermatología, Complejo Hospitalario Universitario de Santiago, Universidad de Santiago de Compostela, Santiago de Compostela, Spain), Francisco Javier Miquel Miquel (Servicio de Dermatología, Hospital Universitario Arnau de Vilanova, Valencia, Spain), Francisco Javier Ortiz de Frutos (Servicio de Dermatología, Hospital Universitario 12 de Octubre, Madrid, Spain), María Elena Gatica Ortega (Servicio de Dermatología, Complejo Hospitalario Universitario de Toledo, Toledo, Spain), José Manuel Carrascosa (Servicio de Dermatología, Hospital Universitario Germans Trias i Pujol, Badalona, Spain), Ana María Giménez Arnau (Servicio de Dermatología, Hospital del Mar, Instituto Municipal de Asistencia Sanitaria, Barcelona, Spain), Violeta Zaragoza Ninet (Servicio de Dermatología, Hospital General Universitario de Valencia, Valencia, Spain), Ricardo González Pérez (Servicio de Dermatología, Hospital Universitario Araba, Universidad del País Vasco, Vitoria, Spain), Tatiana Sanz Sánchez (Servicio de Dermatología, Hospital Universitario Infanta Sofía, Madrid, Spain), María Antonia Pastor Nieto (Servicio de Dermatología, Hospital Universitario de Guadalajara, Guadalajara, Spain), Esther Serra Baldrich (Servicio de Dermatología, Hospital de la Santa Creu i Sant Pau, Barcelona, Spain), Susana Córdoba Guijarro (Servicio de Dermatología, Hospital Universitario de Fuenlabrada, Madrid, Spain), Felipe Heras Mendaza (Servicio de Dermatología, Hospital Universitario Fundación Jiménez Díaz, Madrid, Spain), Inmaculada Ruiz González (Servicio de Dermatología, Complejo Asistencial Universitario de León, León, Spain), Mercedes Rodríguez Serna (Servicio de Dermatología, Hospital Universitario La Fe, Valencia, Spain), Marcos Hervella Garcés (Servicio de Dermatología, Complejo Hospitalario de Navarra, Pamplona, Spain), Leopoldo Borrego (Complejo Hospitalario Universitario Insular Materno Infantil, Universidad de las Palmas de Gran Canaria, Spain).

Correspondence

Prof. W. Uter, Department of Medical Informatics, Biometry and Epidemiology, University of Erlangen/Nürnberg, Erlangen, Germany wolfgang.uter@fau.de

Conflict of Interests

W.U. has accepted travel reimbursement and research funds from the cosmetic industry association IFRA and has received a lecture fee from dermatology-related sponsors for an educational lecture on contact allergy in 2018. O.A. is investigator, consultant and/or speaker for Leo Pharma and La Roche Posay/L'Oréal. M.G. has received honoraria for advisory boards and lectures from Novartis, Abbvie, Leo, Lilly, Pfizer, Sanofi and Takeda. C.P. is an advisor, consultant, speaker and / or investigator for AbbVie, Eli Lilly, Leo Pharma, Novartis, Pfizer, Pierre Fabre, Sanofi. M.-L.A.S. is an advisor, consultant, speaker and/or investigator for Abbvie, Pfizer, LEO Pharma, Regeneron, Sanofi-Genzyme, Eli Lilly and Galderma, and has received research grants from Regeneron, Sanofi-Genzyme, Novartis and Pfizer. The IVDK, maintained by the IVDK e.V., of which S.S. is an employee, is sponsored by the cosmetic and fragrance industry (associations) as well as by public funds. Th.W. is an advisor, consultant, speaker and/or investigator for Abbvie, Pfizer, LEO Pharma, Novartis, Regeneron, Sanofi-Genzyme, Eli Lilly and Galderma. S.M.W. has received travel

reimbursement to attend meetings with the cosmetic industry. The other authors have no pertinent conflict of interests to declare.

Funding

Partial funding by EADV Grant PPRC-2018-8. The REIDAC project is promoted by the Fundación Piel Sana Academia Española de Dermatología y Venereología, which has received financial support from the Spanish Medicines and Health Products Agency (Agencia Española de Medicamentos y Productos Sanitarios; available at https://www.boe.es/boe/dias/2021/12/15/pdfs/BOE-A-2021-20717.pdf), and from pharmaceutical companies (Sanofi, GSK and Novartis).

Accepted Article

Accepted Article

Summary

Background. Continual analyses of patch test results with the European baseline series (EBS) serve both contact allergy surveillance and auditing the value of included allergens.
Objectives. To present results of current EBS patch testing, obtained in 53 departments in 13 European countries during 2019 and 2020.
Methods. Anonymised or pseudonymised individual data, and partly aggregated data on demographic/clinical characteristics and patch test rest results with the EBS were prospectively collected and centrally pooled and analysed.
Results. In 2019 and 2020, 22581 patients were patch tested with the EBS. Sensitization to nickel remained most common (19.8 (19.2–20.4)% positivity (95% confidence interval)). Fragrance mix I and *Myroxylon pereirae* yielded very similar results with 6.80 (6.43–7.19)% and 6.62 (6.25–7.00)% positivity, respectively. Formaldehyde at 2% aq. yielded almost one percentage point more positive reactions than 1% concentration (2.49 (2.16–2.85)% vs. 1.59 (1.33–1.88)); methylchloroisothiazolinone/methylisothiazolinone (MCI/MI) and MI alone up to around 5% positives. Among the new additions, propolis was most commonly positive (3.48 (3.16–3.82)%), followed by 2-hydroxyethyl methacrylate (2.32 (2.0–2.68)%).
Conclusions. Ongoing surveillance on the prevalence of contact sensitization contributes to an up-to-date baseline series containing the most frequent and/or relevant contact sensitizers for routine patch testing in Europe.

Key words

baseline series, contact allergy, clinical epidemiology, surveillance, patch testing, RRID:SCR_001905

Accepted Article

1. Introduction

A "baseline" series should comprise those contact allergens which are of greatest importance and relevance for the majority of patients. It is regularly patch tested in all patients presenting with suspected allergic contact dermatitis. Its composition varies in time and with geographical region, reflecting changes and differences of exposure, respectively. Following fundamental conceptual thoughts delineating the objectives of a baseline series published two decades ago,[1] the criteria for inclusion have recently been revisited.[2] The 2019 version of the European Baseline Series (EBS)[3] has been used since then, partly with modifications by national contact dermatitis groups or single departments. The present paper summarizes results obtained in the years 2019/20 by members of the European Surveillance System on Contact Allergies (ESSCA), a working group of the European Society of Contact Dermatitis (ESCD; https://www.escd.org), by members of the EBS taskforce of the ESCD, and by the Spanish "Grupo Español de Investigación en Dermatitis de Contacto y Alergia Cutánea" (GEIDAC) / "Registro Español de Dermatitis de Contacto" (REIDAC) surveillance network described in.[4]) Moreover, several departments prospectively tested the recommended additions in their baseline series ("audit allergens"). Results with these have been published separately.[5] This article collates prospective audit data with the EBS from a wider spread of departments with a view to informing a further revision of the EBS. Thereby, similar reports on previous EBS results are continued; regarding the last two periods, see.[6,7]

2. Methods

Across Europe, different national contact dermatitis groups are active, many of these using electronic data collection of patch test results.[8] Moreover, a high degree of methodological standardization had been achieved, as documented by the ESCD patch test guidelines.[9] The value of networking has been multiply demonstrated.[10] The present analysis combines, for the first time, individual, anonymized/pseudonymized data and aggregated data (results) from three different sources, the methods of which are briefly outlined below, covering the period January 2019 to December 2020. All groups adhered to the ESCD patch test guideline.[9] As final reading the maximum reaction between day (D) 3 and D5 (inclusive) was used, following current ESSCA standards (see discussion).

2.1 ESSCA working group of the ESCD

The objective of the ESSCA is the clinical surveillance of contact allergy.[10,11] To this end, contributing departments (**Online supplemental Table S1**) submit either all patch test results, or just patch test results obtained with the European baseline series (or national or local adaptations thereof, as evident from **Online supplemental Table S2**), obtained following ESCD standards,[9] to the data centre in Erlangen. This is accompanied by important demographic and clinical information, ranging from "MOAHLFA index"[12] characteristics to a wider range of information according to the ESSCA "minimal dataset" definition.[13,10] Data from contributing departments are delivered in an anonymous format or partly, following national network standards, in a pseudonymized format, where the pseudonym cannot be related to actual personal data except in the contributing department

Accepted Article

itself. This difference is of importance, as only with pseudonymized data can re-investigations of patients be identified and eliminated, to avoid duplication of entries. However, in view of the short study period the effect of re-consultations appeared negligible. Data were quality checked, providing an "internal report" for each contributing department for scrutiny and approval before pooling of the respective data.[11] Data management and analysis was performed with the R software package (https://www.r-project.org; RRID:SCR_001905), version 4.0.3. For the calculation of 95% confidence intervals (CIs) to zero proportions an approximation to an exact CI was used.[14]

2.2 EBS working group of the ESCD

To enable contribution of individual, anonymized data, an online documentation was set up in 2018, based on a local academic implementation of a SoSci server (https://www.soscisurvey.de/). Subsequently, two departments, namely Coimbra, Portugal and Antwerp, Belgium, used this for data contribution. Budapest, Hungary contributed individual data using a spreadsheet, with a defined set of 12 of the EBS allergens related to the "audit allergens" tested in the same period.[5]

2.3 GEIDAC

In Spain, all the participating centers are members of the Spanish Contact Dermatitis and Skin Allergy Research Group (GEIDAC). Data were collected prospectively in the Spanish Contact Dermatitis Registry (REIDAC). This is an online based multicenter registry that uses the OpenClinica platform (OpenClinica LLC and collaborators). Data are anonymized at source and the registry complies with all ethical standards in terms of informed consent and data protection legislation. Clinical data match exactly with those set out in the minimal data set of the ESSCA, which allows them to be exported to other databases with identical categories. Ideally, the centers systematically upload clinical data and results of patch tests on the day of the last reading, thus providing epidemiological centralized data in real time.

The GEIDAC is responsible for the definition and publication of the Spanish Baseline Series, the use of which is recommended in all patients patch tested in Spain. During the study period, a particular form with an "Extended Spanish Baseline Series" which includes the European allergens not included in the Spanish series was temporally added to the registry (REIDAC).[4]

3. Results

In total, 22474 patients were patch tested with the EBS in 2019 to 2020 (inclusive). The individual contribution by country and department is shown in **Online supplemental Table S1**. Population characteristics according to the MOAHLFA index,[12] extended by the "P-measure", that is, the proportion of patients positive to at least one allergen from the baseline series used,[15] are illustrated in **Table 1**.

Patch test results with the EBS are shown in **Table 2**, grouped for allergen classes. A supplemental analysis stratified for the contributing countries can be found in **Online supplemental Table S2**. As the composition of the TRUE Test (panels 1 to 3) partly differs

from the EBS, and as concentrations (effectively: doses per area) are partly different, the results obtained in 12 Spanish and the one Dutch department are presented separately in **Table 3**, with a breakdown by country offered in **Online supplemental Table S3**.

Positive reactions to nickel were most common (19.8 (19.2–20.4)% positivity (95% CI in parentheses)), see **Table 2**; the higher share seen with the TRUE Test can be attributed to the Spanish results (**Table 3**), but also the Italian prevalences are high (**Online supplemental Table S2**) reflecting well-known regional differences.[16,17] Among the fragrances and related substances tested in the EBS, fragrance mix (FM) I and *Myroxylon pereirae* yielded very similar results with 6.80 (6.43–7.19)% and 6.62 (6.25–7.00)% pos., respectively.

Biocides (preservatives) comprise the largest number of allergens in the EBS (n=6). Formaldehyde at 2% aq. Yielded almost one percentage point more positive reactions than the 1% concentration (2.49 (2.16–2.85)% vs. 1.59 (1.33–1.88)); the pooled prevalence of both concentrations being 2.04 (95% CI: 1.83—2.26). However, the results are difficult to compare, as they were obtained in different patients. The higher concentration of methylchloroisothiazolinone/methylisothiazolinone (MCI/MI) and MI alone, respectively, elicited positive reactions more frequently than the lower concentration still in use (**Table 2**), namely, in 6.91 (6.39–7.46)% to MCI/MI and 7.57 (7.11–8.06)% to MI, respectively. MCI/MI tested in the TRUE Test yielded a very similar result with 6.91 (6.1–7.78)% positive reactions (**Table 3**). While positive reactions to paraben mix and quaternium 15 are (notoriously) rare, the prevalence seen of methyldibromo glutaronitrile (MDBGN) sensitization is still high, with 4.26 (3.73–4.84)% at 0.3% pet. and 5.64 (5.1–6.22)% at 0.5% pet., while only 0.65 (0.4–1.01)% positive reactions were observed when MDBGN was tested in Spain using the TRUE Test.

Contact allergy to rubber allergens is not common, the frequency of positive reactions being below 1% in the overall group of consecutively patch tested patients. An exception is thiuram mix, with positive reactions in 2.34 (2.12–2.58)% and 1.63 (1.24–2.1)%, respectively, when tested in the EBS and the TRUE Test, respectively. Positive patch test reactions to the 3 resins included in the EBS, namely, colophonium, epoxy resin, and *p-tert*-butylphenol formaldehyde resin, were seen in 3.32 (3.06–3.60)%, 1.29 (1.12–1.48)%, and 0.47 (0.35–0.61)%, respectively. Positive reactions to the corticosteroid screening allergens and to neomycin sulfate were rare in both EBS and TRUE Test results. Patch tests with both dye-related allergens, *p*-phenylenediamine (PPD) and textile dye mix, yielded very similar results, with 3.60 (3.26–3.96)% and 3.58 (3.23–3.96)% positive reactions, respectively (**Table 2**). Among 5403 patients simultaneously tested with both PPD and textile dye mix, coupled reactivity was marked, with n=110 patients reacting to both allergen preparations, n=86 only to PPD, and n=95 only to textile dye mix (OR: 68.8 (95% CI: 48.6–97.4)).

Among the substances added last to the EBS, propolis was most frequently positive (3.48 (3.16–3.82)%), followed by 2-hydroxyethyl methacrylate (HEMA; 2.32 (2.0–2.68)%), see **Table 2**. Coupled reactivity between propolis and *Myroxylon pereirae* resin was evaluated based on 7879 patients tested with both; 116 patients reacted to both, 480 only to *Myroxylon pereirae*, and 179 only to propolis (OR: 9.6 (95% CI: 7.5–12.3)). Co-reactivity between propolis and FM I was in a similar range, with 92 of 7821patients reacting to both allergens, 200 only to propolis, and 450 only to FM I (OR: 7.2 (95% CI: 5.5–9.4)).

Accepted Article

Concerning colophonium, 65 patients reacted to both natural mixtures, 228 only to propolis, and 237 only to colophonium (OR: 8.9 (95% CI: 6.6–12.1)). A positive patch test reaction to FM I, FM II, HICC, and/or *Myroxylon pereirae* resin was seen in 10.3% (95% CI: 9.8–10.7%) of patients tested with these (n=15071 for whom individual data were available for analysis). Furthermore, caine mix III 10% pet. elicited 1.56 (1.28–1.89)% pos.; in the TRUE Test, the frequency of positive reactions was 0.91 (0.63–1.28)%.

4. Discussion

This update of previous similar ESSCA reports on EBS patch test results[6,7] is novel in that it not only includes ESSCA data, but also additional data from the EBS working group and results obtained by the GEIDAC previously not included in the analyses. Before the results with different allergens are discussed, methodological issues shall be addressed.

4.1 Methodology

This study includes data collected 2019 and 2020. Owing to the coronavirus disease (COVID-19) pandemic, many dermatology departments experienced extreme variations of working conditions and patient selection, with the contact dermatitis units sometimes completely closed, and at other times trying to recover the patch testing backlog.[18] Therefore, the number of cases included is less than expected, also owing to the recommendations by authorities to use hospitals only when strictly necessary. At this time, it is unknown whether and how this feature has impacted on the global results. Considerable variation of the percentage of patients with positive reactions to at least one baseline series allergen (the "P-measure") has been noted before and to a similar extent,[15] thus likely not being indicative of COVID-19 related effects.

It has been repeatedly demonstrated that D7 readings improve the accuracy of patch test results, as a certain share of sensitized patients will exhibit a clear (be it weak) positive reaction only at a late reading, while having negative or doubtful reactions at earlier readings. The share of such late positive readings differs vastly between contact allergens and is highest in corticosteroids and e.g. neomycin sulfate.[19] Moreover, patient characteristics and patch test exposure time may have an impact on the share of reactions which are appearing as positive only on D7.[20] Accordingly, a D7 reading combined with an earlier reading (D2 and D3 or D4) is recommended as "optimum" schedule in the ESCD patch test guideline.[9] However, in practice many national networks deviate from this recommendation, usually owing to logistical reasons. In the part of the data documenting the full matrix of D1 to D7 readings (i.e., departments using the WinAlldat software)[13], 208 instances of positive reactions which had been non-positive at earlier readings were identified, ranging from single events to a maximum of 72 reactions; these were observed in Groningen, which appears to be the only department regularly reading D3 and D7. Three German departments assess 60 to 77% of their patients with a D7 reading, but in the remainder D7 readings are probably only used selectively. Thereby, some underestimation of true sensitization prevalences may be anticipated when relying on the conventional D3-D5 reading outcome (implemented some 20 years ago to cover most reading schedules), at least concerning those allergens which are known to cause late reactions.

Accepted Article

Some well-known characteristics of the TRUE Test, such as poor sensitivity of FM I,[21] and better detection of the MCI/MI preparation than with MCI/MI 100 ppm aq.[22] should be taken into account when comparing these results with other results obtained with investigator-loaded, pet.-/aq.-based allergens. Currently, the sensitization prevalence to MCI/MI 200 ppm aq. is very close to that obtained with the TRUE Test, possibly supporting the recommendation to use the 200 ppm concentration when patch testing MCI/MI aq. in investigator-loaded chambers.

4.2 Metals

Compared to the previous reporting period,[6] the share of positive patch test reactions was slightly higher in the present analysis. However – also for other allergens subsequently discussed – some impact of the limitations imposed by the COVID-19 pandemic in terms of potentially patch testing in a more selective way in 2020 could play a role. Previous analyses pointed to a decrease of nickel contact allergy in younger patients, while a "cohort effect" can be observed in those older females already sensitized before nickel regulation came into effect.[23] Moreover, the prevalence of nickel sensitization varies greatly between countries with e.g. higher crude estimates in Austria (30.32 (24.33-36.84)%) and Italy (26.79 (25.02-28.62)%), and a lower prevalence in Finland (14.77 (8.11-23.94)%) and the UK (13.1 (11.6-14.8)%), which can partially be explained by the late introduction of EU nickel regulation in Italy.[17]

Concerning chromium, a clear effect of regulating the chromium-VI content in leather to < 3 ppm/kg leather (EU 301/2014)[24] has probably yet to be demonstrated although a recent Danish analysis identified a significantly lower prevalence in recent years.[25] The clinical relevance of cobalt contact allergy is often elusive, e.g. in as much as 80% of consecutive patients testing positive.[26] Leather exposure seems to be increasingly important,[26] and the role of cobalt liberation from jewellery should probably not be underestimated,[27,28] although both exposures are not easy to ascertain in clinical practice, as the cobalt spot test is often negative, even if potentially relevant metal liberation is identified by chemical analysis along EN1811:2011+AC:2012.[28]

4.3 Fragrances and plant materials

The percentage of positive reactions to the fragrance markers FM I, FM II, and *Myroxylon pereirae* (balsam of Peru) is slightly higher than in 2015-2018.[6] However, the above-mentioned putative selection effects hamper a direct interpretation of time trends. Conversely, hydroxyisohexyl 3-cyclohexene carboxaldehyde (HICC) contact allergy has declined further, as compared to 2013/14 (1.7%; 95% CI: 1.5-1.8%).[7] This confirms a downward trend following a ban by the European Commission in 2017 during, or even before, a 4-year transition period ending in August 2021 owing to preemptive adaptation by industry.[29] New fragrance compounds introduced to the market, or generally changes in exposure, will evidently affect the screening abilities of the above-mentioned markers, as illustrated by considerable added reactivity to e.g. limonene or linalool hydroperoxides, which has been confirmed by the analysis of "audit allergens" during the same study period.[5] Moreover, the mixes themselves have been found to be insufficiently sensitive to diagnose contact allergy to their constituents, so that patch testing with individual components such as the 26 fragrances to be labelled in the EU has been recommended.[30]

Accepted Article

Propolis, a newly introduced part of the EBS, has been reported to often cross-react with *Myroxylon pereirae* resin.[31] In the present results, the OR of 9.6 (95% CI: 7.5–12.3) indicates only moderate concomitant reactivity, which is moreover asymmetrical: while 39.3% of propolis-positive patients also reacted to *Myroxylon pereirae*, only 19.5% of *Myroxylon pereirae*-positive patients also reacted to propolis. Interestingly, the increase of gross usage data of propolis in Germany has been found to concur with the increase of positive patch test reactions observed in that country,[32] providing supporting evidence of a real increase in sensitization prevalence. However, the prevalence of propolis contact allergy vastly differs between European countries, with a higher share of contact allergy in the centre and East, and a lower in the South(west), see **Online supplemental table S2**. Moreover, the different geographical origin (and thereby, different chemical composition) of propolis,[33] renders interpretation of patch test results complex. Co-reactivity between propolis and *Myroxylon pereirae*, FM I, and colophonium, respectively, which had been reported earlier as significant,[33] was significant in our results, too, albeit on a comparatively moderate level. The role of sesquiterpene lactone mix and the Compositae mixes in diagnosing contact allergy to this important plant family has been discussed in.[5]

4.4 Biocides (preservatives)

The current figures on MCI/MI and MI are slightly higher than those in the previous analysis,[6] despite a presumptive further decline after MI contact allergy has peaked in 2013/14. Again, the putatively different patient selection makes it difficult to interpret these data. Probably, pre-epidemic levels which have been reported by some groups,[34] but not by others,[35] will not be reached soon, in view of the vast number of MI-sensitised patients, who might present with other current problems, but still be MI-allergic. MI contact allergy might not be clinically relevant in many cases, as allergen-avoidance should be (relatively) easy owing to (i) cosmetic restrictions with a ban in leave-on cosmetics and a restriction to 15 ppm in leave-on cosmetics and (ii) classification as CLP/GHS "H 317: I A" contact allergen under REACH regulation in Europe, with a requirement to disclose the presence of MI in all products if this exceeds 1.5 ppm. However, less easily avoidable occupational or hidden exposures such as MI emanating from freshly painted walls, causing airborne allergic contact dermatitis, may still pose difficult to avoid problems for the MI sensitized.[36]

One puzzling finding in the present data is the frequency of positive reactions to methyldibromo glutaronitrile (MDBGN): the prevalences seen in 2019/20 are markedly higher – also considering some often-cited selection effect during the COVID-19 pandemic – than in the years 2015-18. In that preceding study period, MDBGN 0.3% pet. had caused 3.29% (95% CI: 3.09-3.51%) and MDBGN 0.5% pet. 3.89% (95% CI: 3.58-4.23%) positive reactions,[6] compared to 4.26 (3.73–4.84)% pos. to 0.3% pet. and particularly 5.64 (5.1–6.22)% pos. to MDBGN 0.5% pet. (**Table 2**). Of note, positive reactions to MDBGN tested as part of the TRUE Test in Spain are much lower, at 0.65 (0.4–1.01) % pos. (**Table 3** and **Online supplemental Table S3**). However, if the test concentration used in the TRUE Test (5 μg/cm^2) is compared to the dose/area of MDBGN 0.5% pet. (200 μg/cm^2),[37] a difference by a factor of 40 is apparent. Vehicle effects may differentially affect allergen bioavailability, but it appears unlikely that equivalence can be assumed. As it has been noted that the increase of the MDBGN patch test concentration from 0.2% pet. to 0.3% pet. – corresponding to a factor of just 1.5, albeit presumably in a different area of the dose-effect

Accepted Article

curve – has led to an almost doubled share of positive reactions,[38] it appears very likely that the seemingly low prevalence in Spain (when relying on TRUE Test data) is due to a presumably unduly low patch test dose. A suitable patch test concentration of 0.2% pet., and potentially false-positive reactions to 0.3% and particularly to 0.5% pet. have been identified in a previous study of the German Contact Dermatitis Research Group (DKG) gauging the diagnostic properties of 0.2, 0.3, and 0.5% pet. against repeated open application test (ROAT) results with MDBGN.[39] The concentration of 0.2% would correspond to a dose/area of 80 μg/cm^2; this could probably be regarded as target, or as lower floor, for an amendment of the TRUE Test MDBGN preparation, taking possible vehicle effects into account.

MDBGN has been banned from use in cosmetics more than 10 years ago, and other exposures to MDBGN are often elusive. Therefore, it has been suggested that MDBGN contact allergy nowadays is largely irrelevant and that MDBGN can safely be withdrawn from the baseline series.[40]. However, under the synonym 2-bromo-2-(bromomethyl)pentanedinitrile (DBDCB) MDBGN still is one of the permitted preservatives in the EU for "product type 6", that is, for preservation during storage of products other than foodstuffs, feeding stuffs, cosmetics or medicinal products or medical devices.[41] Examples of non-cosmetic exposures associated with MDBGN have been reviewed recently,[42] and hence it may be premature to regard the still high number of patients sensitised to MDBGN as an entirely "historical" problem. Hence, in addition to optimizing the MDBGN TRUE Test concentration, research efforts should be put into a better delineation of its currently most relevant exposure sources.

Contact allergy to formaldehyde and formaldehyde releasers, including quaternium 15 included in the EBS, has recently been analyzed in detail and presented in a separate paper[43] and will thus not be addressed here. Benzisothiazolinone and octyl isothiazolinones have been evaluated as "audit" or "candidate" allergens for inclusion in the EBS, as presented and discussed in.[5] Positive reactions to paraben mix are very infrequent, both in the EBS and in the TRUE Test. This may argue for a removal from the EBS.[44]

4.5 Rubber, plastic, and glues

With 2.34 (2.12–2.58)% positive reactions, thiuram mix (still) is the most important rubber allergen in the EBS. In the TRUE Test, the yield is just slightly lower, with 1.63 (1.24–2.1) % positive reactions. Importantly, the thiurams included in the mix also represent the corresponding dithiocarbamates, which are also used as accelerators in vulcanization, as both form a redox pair. Thiurams may actually be better screening substances for dithiocarbamate contact allergy than the carbamates themselves, as discussed previously.[45] By contrast, sensitization prevalences observed with other rubber compounds, i.e., the benzothiazoles and *N*-Isopropyl-*N*'-phenyl-*p*-phenylenediamine (IPPD) as representative of PPD-derivatives used as antioxidants and -ozonants, are low. Although difficult to compare, the yield with IPPD 0.1% pet. in the EBS, with 0.79 (0.66–0.94)% positive reactions, and with black rubber mix 75 μg/cm^2 eliciting 0.88 (0.61–1.25) % positive reactions is similar. However, previous results seem to indicate that patch testing with IPPD alone will under-diagnose contact allergy to this class of antidegradants.[45,46] The discussion of the best patch testing strategy for benzothiazoles – mercaptobenzothiazole (MBT) with either the four-component mercapto mix 2% pet. or, preferably, the three-component mercapto mix 1%

Accepted Article

pet.[45] – shall not be re-iterated here. All these allergens appear to be borderline for inclusion in the EBS, with an upper 95% CI between 0.5 and 1%; a discussion whether rubber allergens beyond thiuram mix qualify for inclusion for other reasons (such as "often unsuspected, but retrospectively often relevant") seems warranted.

Colophonium is a relatively common allergen with a multitude of sources of exposure. Co-reactivity to *Evernia furfuracea* (tree moss) is common,[5] owing to the (oxidised) resin acids present in the extract resulting from a rather robust collection process of this lichen from conifer bark.[47] Therefore colophonium-allergic patients should be advised to avoid fragrances containing this compound, and vice versa. Concerning epoxy resin, in terms of the diglycidyl ether of bisphenol A as tested in the EBS, it could be argued that (occupational) exposure can usually be traced relatively well. However, the sheer frequency of reactions along with the growing use of epoxy resin systems, also beyond occupational exposure (and thus perhaps often ill-recognized) argues in favor of this allergen being part of the EBS. Conversely, contact allergy to *p-tert*-butylphenol formaldehyde resin (PTBFR) appears to be a rare. It should, however, be noted that the performance of commercially available patch test preparations containing this allergen may vary.[48] Besides glues used in shoe manufacture, other relevant exposures to this complex mixture include, for example, textiles, leather and rubber items, adhesives, domestic glues and even medical devices.[49–52] Although it does not detect contact allergy to all resins based on phenol and formaldehyde,[53] PTBFR is currently the sole representative of this class of compounds routinely tested, and thus should probably maintained in the EBS for the time being. The recent addition of 2-HEMA has certainly proven to be well-justified, in terms of overall frequency of sensitisation, probably driven mainly by the current fashion trend of artificial nails, but also relevant in the field or medical devices, dental materials, and industrial applications.[54] In November 2020, the use of 2-HEMA in nail cosmetics has been restricted in the context of the EU Cosmetics Regulation (EC 1223/2009), permitting only professional use.[55] Time will tell whether this restriction has an effect on the on-going "nail cosmetics epidemic".

4.6 Medicaments, excipients, and dyes

In the last revision of the EBS in 2019, caine mix III 10% pet. has been included to replace benzocaine 5% pet. Comparing just the sensitization frequencies, this has resulted in a more than doubled detection: from 0.69 (0.58-0.82)% to benzocaine seen during 2015-18 to 1.56 (1.28–1.89)% positive reactions to caine mix III in the present analysis. While the sensitivity of this mix diagnosing benzocaine allergy appears acceptable, if concomitant reactivity is analysed, this is not the case regarding cinchocaine and tetracaine, the other two constituents of the mix.[6]

At present, two representatives of corticosteroid allergens are included in the EBS: tixocortol pivalate and budesonide. Both concentrations of the corticosteroid markers yielded low prevalences of contact allergy, the 95% CI (by far) not exceeding 1%, except for budesonide tested in the TRUE Test (**Table 3**). The vastly differing yield of positive reactions, even significant in case of budesonide, is difficult to interpret, as different patients (with a differing exposure background) have been tested. A previous study comparing budesonide 0.01 and 0.1% pet. by synchronous application found these to be equivalent.[56] The low sensitization prevalence to corticosteroids is striking, since exposure

Accepted Article

to topical corticosteroids in the treatment of many dermatoses is very high. The addition of a group 3 corticosteroid allergen (such as clobetasol propionate) has been suggested to improve the diagnosis of allergy to topical corticosteroids;[57] the usefulness seems to depend on (national) prescription habits and thus differences in exposure. Lanolin alcohols are a natural mixture of emulsifying alcohols of various chain lengths and structures. They are weak contact allergens contained in wool fat or wool wax (lanolin) produced from sheeps' wool.[58] Considering the broad use of lanolin (alcohols), a frequency of sensitization of 1.38 (1.20–1.58)% is relatively low. In contrast to the US, the importance of neomycin sulfate as a contact allergen seems to be diminishing, e.g. from 1.24 (1.11-1.37)%[58] in 2009-12 or 1.23 (1.11-1.35)%[6] in 2015-18 to 0.83 (0.67–1.01)% at present. This decline has been found to be linked to decreasing sales/prescriptions.[58]

The EBS includes two allergen preparations used to diagnose contact allergy to dyes (oxidative hair dyes or textile dyes), namely *p*-phenylenediamine (PPD) and textile dye mix (TDM) 6.6% pet. The latter includes Disperse Orange 3, which is chemically very similar to PPD and shows marked cross-reactivity, noted already during the initial study,[59] and certainly also in the present analysis, with an OR of 68.8 (95% CI: 48.6–97.4). Hence, a Swedish study evaluated a TDM without this dye and concluded that "results indicate that Dispers Orange 3 might probably be omitted from TDM", but further study is needed to finally decide whether this is actually sensible or not.[60] Sensitization to PPD itself is more prevalent in central and southern European patch test centers than in Scandinavian,[61] most likely owing to the common use of dark oxidative hair dye products towards the South. A corresponding geographical gradient is also reflected in differences of the share of PPD-containing oxidative hair dye products between countries, which, in addition, has changed over time.[62]

5. Conclusion

The surveillance results on the prevalence of contact sensitization to EBS allergens, combined with patch test results of possible candidate allergens for inclusion in the EBS,[5] contributes to maintaining an up-to-date baseline series comprising the most frequent and/or relevant contact sensitizers for routine patch testing in Europe. National adaptations and extensions based on typical geographical allergen exposures are of course possible and useful; these added allergens are often a fruitful starting point for co-operative evaluation on a broader, European scale. The EBS working group offers a forum for this important research aiming at a high level of diagnostic standards and patient care.

Acknowledgement

Partial funding by the EADV Grant PPRC-2018-8 is thankfully acknowledged. Since spring 2020, data for this study were collected during the pre-vaccine COVID-19 pandemic. We are indebted to all the doctors who were on the front lines of this war and especially to the dermatologists who supported other departments. Our deep appreciation goes to the colleagues infected during this period, our mourning to those who died.

Accepted Article

Table 1: Demographic and clinical characteristics according to the MOAHLFA index[12] with "P-measure,"[15] ie the proportion of patients with at least one positive reaction to a baseline series allergen per country.

Country	M	O	A	H	L	F	A(2)	P
AT	24.8	15.0	26.1	22.1	5.8	19.0	61.9	75.7
BE	32.3	13.0	38.4	19.7	10.9	20.6	58.4	66.1
CH	38.1	13.5	22.1	25.0	6.0	18.2	68.6	59.0
DE	38.0	31.8	28.2	43.8	5.3	9.6	73.1	54.9
ES	34.0	8.4	17.0	22.5	6.1	15.6	70.7	46.8
FI	49.5	76.3	23.7	80.4	1.0	9.3	44.3	49.5
HU	24.1	0.8	13.0	35.4	7.4	17.0	69.7	34.8
IT	30.8	7.2	16.6	19.9	3.5	11.6	62.2	40.1
LT	24.6	27.4	5.2	33.0	9.8	25.6	60.6	47.8
NL	33.3	16.3	38.8	13.5	0.9	4.5	59.8	69.9
PT	29.4	32.8	18.0	43.3	7.1	14.2	57.0	51.7
SI	29.4	3.9	7.9	47.1	16.7	21.0	61.5	44.4
UK	30.4	5.8	46.2	24.7	2.6	33.5	53.9	40.4

M, % male patients; O, % patients with occupational dermatitis; A, % patients with atopic dermatitis; H, % patients with hand dermatitis; L, % patients with leg dermatitis; F, % patients with face dermatitis; A(2), % patients age 40 and above; P, share of patients with at least one positive reaction to a baseline series allergen.

Accepted Article

Table 2: Patch test results (day 3 to day 5) with the European Baseline Series in consecutive patients in the 53 active departments of the European Surveillance System on Contact Allergies (ESSCA), additional contributors from the EBS working group, and the contributing "Grupo Español de Investigación en Dermatitis de Contacto y Alergia Cutánea" (GEIDAC) members (TRUE Test results see **Table 3**). Conc., concentration in %, tested in petrolatum, except where indicated otherwise: [a], aqua. For composition of mixes see.[3]

Allergen	Conc.	tested	+/++/+++	?+/IR	% pos. (95% CI)
Metals					
Potassium dichromate	0.5	16296	712	332	4.37 (4.06–4.69)
Cobalt chloride	1.0	16608	1027	396	6.18 (5.82–6.56)
Nickel sulfate	5.0	16540	3274	263	19.8 (19.2–20.4)
Fragrances					
Fragrance mix I	8.0	16928	1151	302	6.80 (6.43–7.19)
Fragrance mix II	14.0	17519	660	193	3.77 (3.49–4.06)
HICC	5.0	15191	201	43	1.32 (1.15–1.52)
Myroxylon pereirae (balsam of Peru)	25	16980	1124	370	6.62 (6.25–7.00)
Preservatives					
Formaldehyde	1.0[a]	8315	132	48	1.59 (1.33–1.88)
Formaldehyde	2.0[a]	8193	204	48	2.49 (2.16–2.85)
MCI/MI 3:1	0.01[a]	5627	192	25	3.41 (2.95–3.92)
MCI/MI 3:1	0.02[a]	8871	613	133	6.91 (6.39–7.46)
Methylisothiazolinone	0.05[a]	4863	192	33	3.95 (3.42–4.53)
Methylisothiazolinone	0.20[a]	11950	905	214	7.57 (7.11–8.06)
Paraben mix	16	13525	76	98	0.56 (0.44–0.70)
Quaternium-15	1.0	10156	62	12	0.61 (0.47–0.78)
Methyldibromo glutaronitrile	0.3	533125	227	165	4.26 (3.73–4.84)
Methyldibromo glutaronitrile	0.5	6596	372	102	5.64 (5.1–6.22)
Medicaments, excipients					
Caine mix III (Benzo-,Cincho-,Tetracaine)	10.0	6592	103	80	1.56 (1.28–1.89)
Budesonide	0.01	9438	46	43	0.49 (0.36–0.65)
Budesonide	0.1	1863	0	0	0 (0–0.19)
Tixocortol pivalate	0.1	7541	26	28	0.34 (0.23–0.5)
Tixocortol pivalate	1.0	3532	21	8	0.59 (0.37–0.91)

Accepted Article

Allergen	Conc.	tested	+/++/+++	?+/IR	% pos. (95% CI)
Neomycin sulfate	20	11737	97	18	0.83 (0.67–1.01)
Lanolin (wool alcohols)	30	14947	206	92	1.38 (1.20–1.58)
Rubber additives					
Thiuram mix	1.0	16632	389	49	2.34 (2.12–2.58)
N-Isopropyl-*N*'-phenyl-*p*-phenylenediamine	0.1	16866	133	47	0.79 (0.66–0.94)
Mercapto mix[(i)]	1.0	5454	39	27	0.72 (0.51–0.98)
Mercapto mix[(ii)]	2.0	9510	39	7	0.41 (0.29–0.57)
Mercaptobenzothiazole	2.0	16657	83	35	0.50 (0.40–0.62)
Resins/glues					
Colophonium	20.0	16994	564	75	3.32 (3.06–3.60)
Epoxy resin	1.0	16214	209	41	1.29 (1.12–1.48)
p-tert-Butylphenol formaldehyde resin	1.0	11753	55	14	0.47 (0.35–0.61)
2-Hydroxyethyl methacrylate	2.0	7675	178	15	2.32 (2.00–2.68)
Other					
p-Phenylenediamine	1.0	11173	402	28	3.60 (3.26–3.96)
Sesquiterpene lactone mix	0.1	8658	71	19	0.82 (0.64–1.03)
Propolis	10	11952	416	292	3.48 (3.16–3.82)
Textile dye mix	6.6	10323	370	82	3.58 (3.23–3.96)

Epoxy resin, diglycidyl ether of bisphenol A; HICC, hydroxyisohexyl 3-cyclohexene carboxaldehyde; mercapto mix[(i)], containing N-cyclohexylbenzothiazyl sulfenamide, dibenzothiazyl disulfide, and morpholinylmercaptobenzothiazole; mercapto mix[(ii)], containing N-cyclohexylbenzothiazyl sulfenamide, mercaptobenzothiazole, dibenzothiazyl disulfide, and morpholinylmercaptobenzothiazole; MCI, methylchloroisothiazolinone; MI, methylisothiazolinone.

Accepted Article

Table 3: Patch test results obtained with the TRUE Test used as baseline series (supplemented with other, pet.- or aq.-based allergens) in 12 Spanish departments and in Groningen, The Netherlands.

Allergen	mg/cm²	tested	+/++/+++	% pos. (95% CI)
Potassium dichromate	0.054	3620	114	3.15 (2.6–3.77)
Cobalt (II)-chloride, 6*H2O	0.02	3619	178	4.92 (4.24–5.67)
Nickel (II)-sulfate hexahydrate	0.2	3615	867	23.98 (22.6–25.41)
Gold sodium thiosulfate	0.075	3063	152	4.96 (4.22–5.79)
Fragrance mix	0.5	3620	123	3.4 (2.83–4.04)
Balsam of Peru (*Myroxolon pereirae*)	0.8	3619	60	1.66 (1.27–2.13)
Formaldehyde	0.18	3620	38	1.05 (0.74–1.44)
Methylchloroisothiazolinone Methylisothiazolinone (MCI/MI)	0.004	3620	250	6.91 (6.1–7.78)
Paraben mix	1	3620	11	0.3 (0.15–0.54)
Quaternium 15	0.1	3620	47	1.3 (0.96–1.72)
Diazolidinyl urea	0.55	3472	16	0.46 (0.26–0.75)
Imidazolidinyl Urea (Germall 115)	0.6	3472	17	0.49 (0.29–0.78)
Thiomersal (Thimerosal)	0.007	3620	113	3.12 (2.58–3.74)
2-Bromo-2-nitro-1,3-propanediol	0.25	3065	20	0.65 (0.4–1.01)
Methyldibromoglutaronitrile	0.005	3065	20	0.65 (0.4–1.01)
Caine Mix III (Benzo-,Cincho-,Tetracaine)	0.63	3620	33	0.91 (0.63–1.28)
Budesonide	0.001	3471	27	0.78 (0.51–1.13)
Tixocortol-pivalate	0.002	3471	13	0.37 (0.2–0.64)
Hydrocortisone-17-butyrate	0.02	3471	15	0.43 (0.24–0.71)
Neomycin sulfate	0.6	3619	28	0.77 (0.51–1.12)
Ethylene diamine-di-HCl	0.05	3619	32	0.88 (0.61–1.25)
Quinoline mix	0.19	3064	12	0.39 (0.2–0.68)
Bacitracine	0.6	3063	2	0.07 (0.01–0.24)
Lanolin (wool fat) alcohols	1	3619	25	0.69 (0.45–1.02)
Thiuram mix	0.027	3619	59	1.63 (1.24–2.1)
PPD (black rubber) mix	0.075	3619	32	0.88 (0.61–1.25)
Mercapto mix (CBS, MBTS, MOR)	0.075	3471	15	0.43 (0.24–0.71)
Mercaptobenzothiazole	0.075	3619	18	0.5 (0.3–0.78)
Carba mix	0.25	3619	81	2.24 (1.78–2.77)
Colophony (Rosin)	1.2	3619	79	2.18 (1.73–2.71)
Epoxy resin	0.05	3619	58	1.6 (1.22–2.07)
p-tert.-Butylphenol formaldehyde resin (PTBFR)	0.045	3618	81	2.24 (1.78–2.78)
p-Phenylenediamine (PPD)	0.08	3619	140	3.87 (3.26–4.55)
Parthenolide	0.003	3064	8	0.26 (0.11–0.51)
2-Bromo-2-nitro-1,3-propanediol	0.25	3065	20	0.65 (0.4–1.01)

Accepted Article

References

1. Bruze M, Condé-Salazar L, Goossens A, Kanerva L, White IR. Thoughts on sensitizers in a standard patch test series. The European Society of Contact Dermatitis. *Contact Dermatitis*. 1999;41(5):241-250.

2. Wilkinson SM, Badulici S, Giménez-Arnau A, et al. The European baseline series: Criteria for allergen inclusion (with reference to formaldehyde releasers). *Contact Dermatitis*. 2021;85:125-128. doi:10.1111/cod.13836

3. Wilkinson M, Gonçalo M, Aerts O, et al. The European baseline series and recommended additions: 2019. *Contact Dermatitis*. 2019;80(1):1-4. doi:10.1111/cod.13155

4. Hernández-Fernández CP, Mercader-García P, Silvestre Salvador JF, et al. Candidate Allergens for Inclusion in the Spanish Standard Series Based on Data from the Spanish Contact Dermatitis Registry. *Actas Dermosifiliogr*. 2021;112(9):798-805. doi:10.1016/j.adengl.2021.07.013

5. Uter W, Wilkinson SM, Aerts O, et al. European patch test results with audit allergens as candidates for inclusion in the European Baseline Series, 2019/20 - Joint results of the ESSCAA and the EBSB working groups of the ESCD, and the GEIDACC. *Contact Dermatitis*. 2022;86(5): 379-389. doi:10.1111/cod.14059

6. Uter W, Bauer A, Belloni Fortina A, et al. Patch test results with the European baseline series and additions thereof in the ESSCA network, 2015-2018. *Contact Dermatitis*. 2021;84(2):109-120. doi:10.1111/cod.13704

7. Uter W, Amario-Hita JC, Balato A, et al. European Surveillance System on Contact Allergies (ESSCA): Results with the European baseline series, 2013/14. *Journal of the European Academy of Dermatology and Venereology: JEADV*. 2017;31(9):1516-1525. doi:10.1111/jdv.14423

8. Uter W. Contact Dermatitis Research Groups. In: Johansen JD, Mahler V, Lepoittevin JP, Frosch PJ, eds. *Contact Dermatitis*. Sixth edition. Springer; 2020:1-4.

9. Johansen JD, Aalto-Korte K, Agner T, et al. European Society of Contact Dermatitis guideline for diagnostic patch testing recommendations on best practice. *Contact Derm*. 2015;73(4):195-221.

10. Uter W, Schnuch A, Giménez-Arnau A, Orton D, Statham B. Databases and networks: The benefit for research and quality assurance in patch testing. In: Johansen JD, Mahler V, Lepoittevin JP, Frosch PJ, eds. *Contact Dermatitis*. Sixth edition. Springer; 2020:1-16.

11. Uter W, Schnuch A, Wilkinson M, Dugonik A, Dugonik B, Ganslandt T. Registries in Clinical Epidemiology: The European Surveillance System on Contact Allergies (ESSCA). *Methods of information in medicine*. 2016;55(2):193-199.

12. Schnuch A, Geier J, Uter W, et al. National rates and regional differences in sensitization to allergens of the standard series. Population-adjusted frequencies of sensitization (PAFS) in 40,000 patients from a multicenter study (IVDK). *Contact Derm*. 1997;37(5):200-209.

13. Uter W, Arnold R, Wilkinson J, et al. A multilingual European patch test software concept: WinAlldat/ESSCA. *Contact Derm*. 2003;49(5):270-271.

14. Gefeller O, Pfahlberg AB, Uter W. What can be learnt from nothing? A statistical perspective. *Contact Derm*. 2013;69(6):350-354.

Accepted Article

15. Uter W, Schwitulla J, Thyssen JP, Frosch PJ, Statham B, Schnuch A. The 'overall yield' with the baseline series - a useful addition to the array of MOAHLFA factors describing departmental characteristics of patch tested patients. *Contact Dermatitis*. 2011;65(6):322-328.

16. García-Gavín J, Armario-Hita JC, Fernández-Redondo V, et al. Nickel allergy in Spain needs active intervention. *Contact Dermatitis*. 2011;64(5):289-291. doi:10.1111/j.1600-0536.2010.01865.x

17. Basso P, Mauro M, Miani A, Belloni Fortina A, Corradin MT, Larese Filon F. Sensitization to nickel in the Triveneto region: Temporal trend after European Union regulations. *Contact Dermatitis*. 2020;82(4):247-250. doi:10.1111/cod.13450

18. Carrascosa JM, Pastor-Nieto MA, Ruiz-González I, et al. Patch Testing During the COVID-19 Pandemic: Recommendations of the AEDV's Spanish Contact Dermatitis and Skin Allergy Research Group (GEIDAC). *Actas Dermo-Sifiliograficas*. 2020;111(8):650-654. doi:10.1016/j.ad.2020.06.003

19. van Amerongen CCA, Ofenloch R, Dittmar D, Schuttelaar MLA. New positive patch test reactions on day 7-The additional value of the day 7 patch test reading. *Contact Dermatitis*. 2019;81(4):280-287. doi:10.1111/cod.13322

20. Forkel S, Schubert S, Dickel H, et al. The benefit of late readings in patch testing depends both on allergen and patient characteristics. *Allergy*. 77(5):1477-1485. doi:10.1111/all.15149

21. Uter W. Fragrance mix I: TRUE Test() versus pet.-Based patch test. *Contact Dermatitis*. 2015;72(4):256-258. doi:10.1111/cod.12352

22. Dittmar D, Schuttelaar ML. Comparing patch test results of methylchloroisothiazolinone/methylisothiazolinone tested with both TRUE Test and 100 Ppm using investigator-loaded chambers. *Contact Dermatitis*. 2018;78(2):159-161. doi:10.1111/cod.12871

23. Uter W, Gefeller O, Mahler V, Geier J. Trends and current spectrum of contact allergy in Central Europe: Results of the Information Network of Departments of Dermatology (IVDK) 2007-2018. *The British Journal of Dermatology*. 2020;183(5):857-865. doi:10.1111/bjd.18946

24. Anonymous. Commission Regulation (EU) No 301/2014 of 25 March 2014 amending Annex XVII to Regulation (EC) No 1907/2006 of the European Parliament and of the Council on the Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH) as regards chromium VI compounds. *Official Journal*. 2014;90:1-3.

25. Alinaghi F, Zachariae C, Thyssen JP, Johansen JD. Temporal changes in chromium allergy in Denmark between 2002 and 2017. *Contact Dermatitis*. 2019;80(3):156-161. doi:10.1111/cod.13181

26. Alinaghi F, Zachariae C, Thyssen JP, Johansen JD. Causative exposures and temporal development of cobalt allergy in Denmark between 2002 and 2017. *Contact Dermatitis*. 2019;81(4):242-248. doi:10.1111/cod.13326

27. Uter W, Wolter J. Nickel and cobalt release from earrings and piercing jewellery - analytical results of a German survey in 2014. *Contact Dermatitis*. 2018;78(5):321-328. doi:10.1111/cod.12941

28. Wennervaldt M, Ahlström MG, Menné T, et al. Chromium and cobalt release from metallic earrings from the Danish market. *Contact Dermatitis*. 2021;85(5):523-530. doi:10.1111/cod.13917

29. Ahlström MG, Uter W, Ahlström MG, Johansen JD. Decrease of contact allergy to hydroxyisohexyl 3-cyclohexene carboxaldehyde in Europe prior to its ban and diagnostic value. *Contact Dermatitis*. 2021;84(6):419-422. doi:10.1111/cod.13786

Accepted Article

30. Bennike NH, Zachariae C, Johansen JD. Non-mix fragrances are top sensitizers in consecutive dermatitis patients – a cross-sectional study of the 26 EU-labelled fragrance allergens. *Contact Dermatitis* 2017;77(5):270-279. doi: 10.1111/cod.12822

31. Hausen BM, Evers P, Stüwe HT, König WA, Wollenweber E. Propolis allergy (IV). Studies with further sensitizers from propolis and constituents common to propolis, poplar buds and balsam of Peru. *Contact Dermatitis*. 1992;26(1):34-44. doi:10.1111/j.1600-0536.1992.tb00865.x

32. Schubert S, Geier J, Dickel H, et al. Contact sensitization to propolis in the Information Network of Departments of Dermatology (IVDK) 2013 to 2019 and market survey of propolis commerce in Germany. *Contact Dermatitis*. Published online August 2021. doi:10.1111/cod.13960

33. Nyman GSA, Giménez-Arnau AM, Grigaitiene J, Malinauskiene L, Paulsen E, Hagvall L. Patch Testing with Propolis of Different Geographical Origins in a Baseline Series. *Acta Dermato-Venereologica*. 2021;101(11):adv00591. doi:10.2340/actadv.v101.423

34. Herman A, Aerts O, Jacobs MC, et al. Evolution of methylisothiazolinone sensitization: A Belgian multicentric study from 2014 to 2019. *Contact Dermatitis*. Published online August 2021. doi:10.1111/cod.13956

35. Havmose M, Thyssen JP, Zachariae C, Menné T, Johansen JD. The epidemic of contact allergy to methylisothiazolinone-An analysis of Danish consecutive patients patch tested between 2005 and 2019. *Contact Dermatitis*. 2021;84(4):254-262. doi:10.1111/cod.13717

36. Amsler E, Aerts O, Raison-Peyron N, et al. Airborne allergic contact dermatitis caused by isothiazolinones in water-based paints: A retrospective study of 44 cases. *Contact Dermatitis*. 2017;77(3):163-170. doi:10.1111/cod.12795

37. Bruze M, Goossens A, Isaksson M. Recommendation to increase the test concentration of methylchloroisothiazolinone/methylisothiazolinone in the European baseline patch test series - on behalf of the European Society of Contact Dermatitis and the European Environmental and Contact Dermatitis Research Group. *Contact Dermatitis*. 2014;71(1):35-40. doi:10.1111/cod.12240

38. Schnuch A, Schubert S, Geier J, IVDK. Clinicians vs. Epidemiologists: Patch testing with methyldibromo glutaronitrile as a controversial issue. *Journal of the European Academy of Dermatology and Venereology: JEADV*. 2019;33(6):e242-e244. doi:10.1111/jdv.15505

39. Schnuch A, Kelterer D, Bauer A, et al. Quantitative patch and repeated open application testing in methyldibromo glutaronitrile-sensitive patients. *Contact Dermatitis*. 2005;52(4):197-206. doi:10.1111/j.0105-1873.2005.00529.x

40. Mercader-García P, Pastor-Nieto MA, Gonzalez-Perez R, et al. Arguments in favour of excluding MDBGN from the European baseline series. In response to "Comment on MDBGN/DBDCB, the European baseline series, and EU legislation". *Contact Dermatitis*. 2021;85(5):611-613. doi:10.1111/cod.13941

41. Anonymous. Commission Implementing Regulation (EU) 2016/1086 of 5 July 2016 approving 2-bromo-2-(bromomethyl)Pentanedinitrile (DBDCB) as an existing active substance for use in biocidal products of product-type 6 (Text with EEA relevance). *Official Journal L*. 2016;180:15-17.

42. Deza G, Giménez-Arnau AM. Allergic contact dermatitis in preservatives: Current standing and future options. *Current Opinion in Allergy and Clinical Immunology*. 2017;17(4):263-268. doi:10.1097/ACI.0000000000000373

Accepted Article

43. Whitehouse H, Uter W, Geier J, et al. Formaldehyde 2% is not a useful means of detecting allergy to formaldehyde releasers- results of the ESSCA network, 2015-2018. *Contact Dermatitis*. 2021;84(2):95-102. doi:10.1111/cod.13691

44. Giácaman-von der Weth MM, Ferrer-Guillén B, María Ortiz-Salvador J, et al. Is time to remove parabens from standard patch test batteries? Retrospective study of 10 461 patients. *Allergy*. 2020;75(11):2997-2999. doi:10.1111/all.14460

45. Warburton KL, Uter W, Geier J, et al. Patch testing with rubber series in Europe: A critical review and recommendation. *Contact Dermatitis*. 2017;76(4):195-203. doi:10.1111/cod.12736

46. Uter W, Warburton K, Weisshaar E, et al. Patch test results with rubber series in the European Surveillance System on Contact Allergies (ESSCA), 2013/14. *Contact Dermatitis*. 2016;75(6):345-352. doi:10.1111/cod.12651

47. Uter W, Schmidt E, Lessmann H, Schnuch A. Contact sensitization to tree moss (Evernia furfuracea extract, INCI) is heterogeneous. *Contact Dermatitis*. 2012;67(1):36-41.

48. Hamann CR, Zimerson E, Hamann D, et al. Concentration variability of potent allergens of p-tert-butylphenol-formaldehyde resin (PTBP-FR) in patch test preparations and commercially available PTBP-FR. *The British Journal of Dermatology*. 2012;166(4):761-770. doi:10.1111/j.1365-2133.2011.10781.x

49. Özkaya E. Linear eczema on the amputation stump as a clinical clue to the diagnosis. *Contact Dermatitis*. 2017;77(5):328-330. doi:10.1111/cod.12824

50. Herro EM, Jacob SE. Allergic contact dermatitis to p-tert butylphenol formaldehyde resin in a continuous positive airway pressure strap. *Dermatitis: Contact, Atopic, Occupational, Drug*. 23(3):125-126. doi:10.1097/DER.0b013e31824a63c6

51. Herro EM, Friedlander SF, Jacob SE. Bra-associated allergic contact dermatitis: P-tert-butylphenol formaldehyde resin as the culprit. *Pediatric Dermatology*. 29(4):540-541. doi:10.1111/j.1525-1470.2011.01533.x

52. Avenel-Audran M, Goossens A, Zimerson E, Bruze M. Contact dermatitis from electrocardiograph-monitoring electrodes: Role of p-tert-butylphenol-formaldehyde resin. *Contact Dermatitis*. 2003;48(2):108-111. doi:10.1034/j.1600-0536.2003.480210.x

53. Isaksson M, Ale I, Andersen K, et al. Multicenter Patch Testing With a Resol Resin Based on Phenol and Formaldehyde Within the International Contact Dermatitis Research Group. *Dermatitis: Contact, Atopic, Occupational, Drug*. 26(5):230-234. doi:10.1097/DER.0000000000000137

54. Uter W, Werfel T, Lepoittevin JP, White IR. Contact Allergy-Emerging Allergens and Public Health Impact. *International Journal of Environmental Research and Public Health*. 2020;17(7):E2404. doi:10.3390/ijerph17072404

55. Anonymous. COMMISSION REGULATION (EU) 2020/1682 of 12 November 2020 amending Annex III to Regulation (EC) No 1223/2009 of the European Parliament and of the Council on cosmetic products. *Official Journal L*. Published online 2020.

56. Isaksson M, Andersen KE, Brandão FM, et al. Patch testing with budesonide in serial dilutions. A multicentre study of the EECDRG. *Contact Dermatitis*. 2000 Jun;42(6):352-354

57. Mercader-García P, Pastor-Nieto MA, García-Doval I, et al. Are the Spanish baseline series markers sufficient to detect contact allergy to corticosteroids in Spain? A GEIDAC prospective study. *Contact Dermatitis*. 2018;78(1):76-82. doi:10.1111/cod.12874

Accepted Article

58. Uter W, Spiewak R, Cooper SM, et al. Contact allergy to ingredients of topical medications: Results of the European Surveillance System on Contact Allergies (ESSCA), 2009-2012. *Pharmacoepidemiology and Drug Safety*. 2016;25(11):1305-1312. doi:10.1002/pds.4064

59. Isaksson M, Ryberg K, Goossens A, Bruze M. Recommendation to include a textile dye mix in the European baseline series. *Contact Dermatitis*. 2015;73(1):15-20. doi:10.1111/cod.12400

60. Stenton J, Dahlin J, Antelmi A, et al. Patch testing with a textile dye mix with and without Disperse Orange 3. *Contact Dermatitis*. 2020;83(5):387-390. doi:10.1111/cod.13660

61. Thyssen JP, Andersen KE, Bruze M, et al. P-Phenylenediamine sensitization is more prevalent in central and southern European patch test centres than in Scandinavian: Results from a multicentre study. *Contact Dermatitis*. 2009;60(6):314-319. doi:10.1111/j.1600-0536.2009.01547.x

62. Kirchlechner S, Hübner A, Uter W. Survey of sensitizing components of oxidative hair dyes (retail and professional products) in Germany. *Journal der Deutschen Dermatologischen Gesellschaft = Journal of the German Society of Dermatology: JDDG*. 2016;14(7):707-715. doi:10.1111/ddg.13099

Accepted Article

Accepted Article



Online supplemental Table S1: Contribution by department and country, respectively. Additional contributions are from the "Grupo Español de Investigación en Dermatitis de Contacto y Alergia Cutánea" (GEIDAC):[4] (i) additional 1749 patients tested with investigator-loaded system, and (ii) 3065 using the TRUE Test in 2019/20.

Country	City	2019 tested	2020 tested
Austria	Graz	133	93
Belgium	Antwerp	529	395
Switzerland	Basel	175	156
Switzerland	Bern	332	271
Switzerland	Aarau	69	98
Switzerland	Lausanne/Epalignes	25	51
Germany	Dortmund	55	64
Germany	Göttingen	153	112
Germany	Kiel	163	187
Germany	Dresden	241	387
Germany	Jena	143	121
Germany	Mainz	126	124
Germany	Osnabrück	371	378
Germany	Erlangen	231	245
Germany	Heidelberg	116	66
Germany	Hannover	221	173
Germany	Bochum	222	201
Spain	Barcelona/IMAS	460	234
Finland	Helsinki	97	0
Hungary	Budapest	353	0
Italy	Napoli	614	289
Italy	Trieste	262	173
Italy	Genova	566	294
Italy	Perugia	422	187
Lithuania	Kaunas	331	169
The Netherlands	Groningen	310	255
The Netherlands	Amsterdam VU	887	716
Portugal	Coimbra	323	174
Slovenia	Maribor/Univ.	326	174
Slovenia	Maribor/Clinic	100	84
Slovenia	Celje	504	320
Slovenia	Izola	159	47

Country	City	2019 tested	2020 tested
Slovenia	Novo Mesto	221	185
Slovenia	Ljubljana/Clinic	32	31
United Kingdom	Middlesbrough		107
United Kingdom	Oxford	566	0
United Kingdom	Leeds	1112	0

Online supplemental Table S2: Patch test results (day 3 to day 5) with the European Baseline series, 2019–2020, stratified for country. For results from The Netherlands (Groningen), see also TRUE Test results. Conc., concentration in %, all tested in pet., except where indicated otherwise: [a], aqua

Allergen	Conc.	AT tested	AT % pos. (95% CI)	BE tested	BE % pos. (95% CI)	CH tested	CH % pos. (95% CI)	DE tested	DE % pos. (95% CI)
Metals									
Potassium dichromate	0.5	221	13.12 (8.97-18.3)	923	5.85 (4.43-7.56)	1035	5.7 (4.37-7.29)	3583	6.22 (5.45-7.07)
Cobalt chloride	1.0	219	19.63 (14.59-25.52)	923	7.58 (5.96-9.49)	1044	7.28 (5.78-9.03)	3611	6.65 (5.86-7.51)
Nickel sulfate	5.0	221	30.32 (24.33-36.84)	922	23.54 (20.83-26.41)	1040	23.17 (20.64-25.86)	3563	16.22 (15.03-17.47)
Fragrances									
Fragrance mix I	8.0	222	12.16 (8.17-17.2)	924	11.04 (9.09-13.24)	1045	9.38 (7.68-11.31)	3565	5.81 (5.06-6.63)
Fragrance mix II	14.0	222	6.31 (3.49-10.35)	924	5.74 (4.33-7.44)	1014	5.13 (3.85-6.67)	3626	4.05 (3.44-4.75)
HICC	5.0	221	1.36 (0.28-3.92)	924	1.19 (0.6-2.12)	1045	1.05 (0.53-1.88)	3540	2.03 (1.59-2.55)
Myroxylon pereirae (balsam of Peru)	25	221	12.67 (8.59-17.79)	924	16.99 (14.62-19.57)	1043	9.97 (8.22-11.95)	3629	6.45 (5.67-7.3)
Preservatives									
Formaldehyde	1.0[a]	221	1.81 (0.5-4.57)			1044	0.67 (0.27-1.38)	3625	1.05 (0.74-1.44)
Formaldehyde	2.0[a]			924	5.09 (3.76-6.71)				
MCI/MI 3:1	0.01[a]	172	4.07 (1.65-8.21)			619	2.75 (1.61-4.36)	2653	3.43 (2.77-4.19)
MCI/MI 3:1	0.02[a]			924	5.3 (3.95-6.95)				
Methylisothiazolinone	0.02[a]								
Methylisothiazolinone	0.05[a]	222	4.5 (2.18-8.13)			1045	2.97 (2.02-4.18)	3596	4.2 (3.57-4.91)
Methylisothiazolinone	0.20[a]			924	5.09 (3.76-6.71)				
Paraben mix	16	171	2.92 (0.96-6.69)	924	0.76 (0.31-1.55)	409	1.47 (0.54-3.17)	1498	0.67 (0.32-1.22)
Quaternium-15	1.0			924	0.97 (0.45-1.84)				
Methyldibromo glutaronitrile	0.3	171	7.6 (4.11-12.65)			409	5.38 (3.4-8.03)	1473	4.82 (3.78-6.04)
Methyldibromo glutaronitrile	0.5			924	5.41 (4.04-7.07)				
Medicaments, excipients									

Allergen	Conc.	AT tested	AT % pos. (95% CI)	BE tested	BE % pos. (95% CI)	CH tested	CH % pos. (95% CI)	DE tested	DE % pos. (95% CI)
Caine mix III (Benzo-,Cincho-,Tetracaine)	10.0	168	2.98 (0.97-6.81)	922	1.41 (0.75-2.4)	148	3.38 (1.11-7.71)	1336	1.05 (0.57-1.75)
Budesonide	0.01			924	0.76 (0.31-1.55)				
Budesonide	0.1								
Tixocortol pivalate	0.1			924	0.65 (0.24-1.41)				
Tixocortol pivalate	1.0								
Neomycin sulfate	20			924	0.65 (0.24-1.41)				
Lanolin (wool alcohols)	30	221	4.52 (2.19-8.16)	924	2.16 (1.33-3.32)	1044	1.82 (1.1-2.83)	3549	1.66 (1.27-2.14)
Rubber additives									
Thiuram mix	1.0	222	4.5 (2.18-8.13)	923	2.6 (1.67-3.84)	1044	2.87 (1.95-4.08)	3632	3.41 (2.85-4.06)
N-Isopropyl-*N'*-phenyl-*p*-phenylenediamine	0.1	221	2.26 (0.74-5.2)	923	1.19 (0.6-2.12)	1043	0.67 (0.27-1.38)	3602	1.17 (0.84-1.57)
Mercapto mix[(i)]	1.0	222	0 (0-1.65)			1043	0.38 (0.1-0.98)	3632	0.83 (0.56-1.18)
Mercapto mix[(ii)]	2.0			923	0.33 (0.07-0.95)				
Mercaptobenzothiazole	2.0	222	0 (0-1.65)	923	0.22 (0.03-0.78)	1043	1.05 (0.53-1.88)	3647	0.9 (0.62-1.27)
Resins/glues									
Colophonium	20.0	222	4.95 (2.5-8.69)	924	4 (2.83-5.48)	1045	3.64 (2.59-4.96)	3636	5.25 (4.55-6.03)
Epoxy resin	1.0	221	1.36 (0.28-3.92)	924	1.08 (0.52-1.98)	1043	1.92 (1.18-2.95)	3486	1.86 (1.44-2.37)
p-tert-Butylphenol formaldehyde resin	1.0			924	0.22 (0.03-0.78)				
2-Hydroxyethyl methacrylate	2.0	0		321	3.74 (1.95-6.44)				
Other									
p-Phenylenediamine	1.0			923	4.55 (3.3-6.1)				
Sesquiterpene lactone mix	0.1			924	0.97 (0.45-1.84)				
Propolis	10	222	10.36 (6.68-15.14)	924	4.44 (3.2-5.97)	1045	4.78 (3.57-6.26)	3518	4.43 (3.78-5.17)
Textile dye mix	6.6			923	3.9 (2.75-5.36)				

AT, Austria; BE, Belgium; CH, Switzerland; DE, Germany; epoxy resin, diglycidyl ether of bisphenol A; HICC, hydroxyisohexyl 3-cyclohexene carboxaldehyde; mercapto mix[(i)], containing N-cyclohexylbenzothiazyl sulfenamide, dibenzothiazyl disulfide, and morpholinylmercaptobenzothiazole; mercapto mix[(ii)], containing N-cyclohexylbenzothiazyl sulfenamide, mercaptobenzothiazole, dibenzothiazyl disulfide, and morpholinylmercaptobenzothiazole; MCI, methylchloroisothiazolinone; MI, methylisothiazolinone.

Allergen	Conc.	ES tested	ES % pos. (95% CI)	FI tested	FI % pos. (95% CI)	HU tested	HU % pos. (95% CI)	IT tested	IT % pos. (95% CI)
Metals									
Potassium dichromate	0.5	2163	3.88 (3.11-4.79)	89	1.12 (0.03-6.1)			2374	3.83 (3.1-4.69)
Cobalt chloride	1.0	2157	4.5 (3.66-5.46)	90	6.67 (2.49-13.95)			2373	6.49 (5.53-7.56)
Nickel sulfate	5.0	2158	19.51 (17.86-21.24)	88	14.77 (8.11-23.94)			2378	26.79 (25.02-28.62)
Fragrances									
Fragrance mix I	8.0	2166	6.69 (5.68-7.83)	90	4.44 (1.22-10.99)	353	13.03 (9.7-17)	2372	4.85 (4.02-5.79)
Fragrance mix II	14.0	2166	2.95 (2.28-3.76)	90	3.33 (0.69-9.43)	353	6.23 (3.95-9.28)	2372	2.36 (1.79-3.05)
HICC	5.0	2165	0.83 (0.49-1.31)			353	2.55 (1.17-4.78)	1450	0.97 (0.53-1.61)
Myroxylon pereirae (balsam of Peru)	25	2158	4.96 (4.08-5.96)	90	6.67 (2.49-13.95)	353	7.37 (4.87-10.61)	2372	3.75 (3.02-4.6)
Preservatives									
Formaldehyde	1.0[a]								
Formaldehyde	2.0[a]	2166	3.55 (2.82-4.42)	89	3.37 (0.7-9.54)	353	1.13 (0.31-2.88)	1937	2.17 (1.57-2.92)
MCI/MI 3:1	0.01[a]								
MCI/MI 3:1	0.02[a]	2159	6.67 (5.65-7.81)	90	8.89 (3.92-16.77)	353	5.95 (3.72-8.95)	1937	7.07 (5.97-8.31)
Methylisothiazolinone	0.02[a]								
Methylisothiazolinone	0.05[a]								
Methylisothiazolinone	0.20[a]	2165	7.9 (6.8-9.12)	90	6.67 (2.49-13.95)	353	13.6 (10.2-17.62)	1937	4.13 (3.29-5.11)
Paraben mix	16	2157	0.7 (0.39-1.14)	90	0 (0-4.02)			2080	0.38 (0.17-0.76)
Quaternium-15	1.0	2158	0.6 (0.32-1.03)	90	0 (0-4.02)	353	0.28 (0.01-1.57)	435	0.46 (0.06-1.65)
Methyldibromo glutaronitrile	0.3			89	1.12 (0.03-6.1)				
Methyldibromo glutaronitrile	0.5	2157	4.68 (3.83-5.66)						
Medicaments, excipients									
Caine mix III (Benzo-,Cincho-,Tetracaine)	10.0			90	2.22 (0.27-7.8)				

Allergen	Conc.	ES tested	ES % pos. (95% CI)	FI tested	FI % pos. (95% CI)	HU tested	HU % pos. (95% CI)	IT tested	IT % pos. (95% CI)
Budesonide	0.01	2158	0.46 (0.22-0.85)					1937	0.36 (0.15-0.74)
Budesonide	0.1			90	0 (0-4.02)				
Tixocortol pivalate	0.1	1607	0.12 (0.02-0.45)	90	0 (0-4.02)				
Tixocortol pivalate	1.0	549	0 (0-0.67)						
Neomycin sulfate	20	2157	1.07 (0.68-1.6)	90	0 (0-4.02)			2372	0.8 (0.48-1.25)
Lanolin (wool alcohols)	30	549	0.18 (0-1.01)	90	1.11 (0.03-6.04)			2378	0.71 (0.42-1.14)
Rubber additives									
Thiuram mix	1.0	2156	1.99 (1.45-2.68)	89	3.37 (0.7-9.54)			2372	2.49 (1.9-3.2)
N-Isopropyl-*N'*-phenyl-*p*-phenylenediamine	0.1	2159	0.65 (0.35-1.09)			353	0.28 (0.01-1.57)	2372	0.84 (0.52-1.3)
Mercapto mix[(i)]	1.0			90	0 (0-4.02)				
Mercapto mix[(ii)]	2.0	2157	0.46 (0.22-0.85)					1651	0.55 (0.25-1.03)
Mercaptobenzothiazole	2.0	2158	0.42 (0.19-0.79)	90	0 (0-4.02)			2378	0.5 (0.26-0.88)
Resins/glues									
Colophonium	20.0	2158	1.2 (0.79-1.76)	90	3.33 (0.69-9.43)	353	2.83 (1.37-5.15)	2372	1.52 (1.07-2.1)
Epoxy resin	1.0	2282	0.53 (0.27-0.92)	89	8.99 (3.96-16.95)			2372	0.76 (0.45-1.2)
p-tert-Butylphenol formaldehyde resin	1.0	2158	0.79 (0.46-1.26)	90	0 (0-4.02)			2387	0.29 (0.12-0.6)
2-Hydroxyethyl methacrylate	2.0	2440	1.84 (1.35-2.46)	90	4.44 (1.22-10.99)	342	0.88 (0.18-2.54)	1452	1.58 (1.01-2.37)
Other									
p-Phenylenediamine	1.0	2157	3.29 (2.58-4.13)	89	6.74 (2.51-14.1)			2372	4.72 (3.9-5.65)
Sesquiterpene lactone mix	0.1	549	0.18 (0-1.01)	90	1.11 (0.03-6.04)	353	0.85 (0.18-2.46)		
Propolis	10	2334	1.29 (0.87-1.83)			695	4.17 (2.81-5.94)		
Textile dye mix	6.6	2443	3.27 (2.6-4.06)	90	2.22 (0.27-7.8)			1450	4.34 (3.35-5.52)

ES, Spain; FI, Finland; HU, Hungary; IT, Italy; epoxy resin, diglycidyl ether of bisphenol A; HICC, hydroxyisohexyl 3-cyclohexene carboxaldehyde; mercapto mix[(i)], containing N-cyclohexylbenzothiazyl sulfenamide, dibenzothiazyl disulfide, and morpholinylmercaptobenzothiazole; mercapto mix[(ii)], containing N-cyclohexylbenzothiazyl sulfenamide, mercaptobenzothiazole, dibenzothiazyl disulfide, and morpholinylmercaptobenzothiazole; MCI, methylchloroisothiazolinone; MI, methylisothiazolinone.

Allergen	Conc.	LT tested	LT % pos. (95% CI)	NL tested	NL % pos. (95% CI)	PT tested	PT % pos. (95% CI)	SI tested	SI % pos. (95% CI)
Metals									
Potassium dichromate	0.5	214	5.14 (2.59-9.01)	1241	6.53 (5.22-8.05)	497	3.1 (1.49-5.62)	2183	2.34 (1.74-3.06)
Cobalt chloride	1.0	499	5.61 (3.76-8.01)	1242	12 (10.24-13.93)	497	6.81 (4.32-10.13)	2183	4.4 (3.58-5.34)
Nickel sulfate	5.0	499	16.63 (13.47-20.2)	1237	23.36 (21.03-25.82)	497	24.15 (19.58-29.2)	2183	17.45 (15.88-19.11)
Fragrances									
Fragrance mix I	8.0	499	2.2 (1.11-3.91)	1241	10.72 (9.05-12.57)	497	7.12 (4.57-10.49)	2183	6.05 (5.08-7.13)
Fragrance mix II	14.0	499	1.8 (0.83-3.4)	1801	5.89 (4.84-7.07)	497	3.1 (1.49-5.62)	2183	3.39 (2.67-4.24)
HICC	5.0	481	0.21 (0.01-1.15)	560	1.96 (0.98-3.49)	496	1.86 (0.69-4.01)	2183	1.51 (1.04-2.12)
Myroxylon pereirae (balsam of Peru)	25	499	4.21 (2.62-6.36)	1241	8.46 (6.97-10.15)	497	3.72 (1.93-6.4)	2183	6.41 (5.42-7.52)
Preservatives									
Formaldehyde	1.0[a]			1242	2.58 (1.77-3.62)			2183	2.34 (1.74-3.06)
Formaldehyde	2.0[a]	453	1.1 (0.36-2.56)			497	0.93 (0.19-2.69)		
MCI/MI 3:1	0.01[a]							2183	3.53 (2.79-4.39)
MCI/MI 3:1	0.02[a]	453	2.43 (1.22-4.3)	1240	14.84 (12.9-16.94)	497	7.43 (4.82-10.85)		
Methylisothiazolinone	0.02[a]								
Methylisothiazolinone	0.05[a]								
Methylisothiazolinone	0.20[a]	453	1.99 (0.91-3.74)	2130	18.4 (16.78-20.12)	497	8.67 (5.84-12.28)	2183	4.17 (3.37-5.09)
Paraben mix	16	499	0.4 (0.05-1.44)	1243	0.97 (0.5-1.68)	497	0.62 (0.08-2.22)	2183	0.27 (0.1-0.6)
Quaternium-15	1.0	499	0 (0-0.74)	1243	1.05 (0.56-1.78)	497	0.62 (0.08-2.22)	2183	0.73 (0.42-1.19)
Methyldibromo glutaronitrile	0.3			1243	8.77 (7.26-10.48)	172			
Methyldibromo glutaronitrile	0.5	453	3.09 (1.7-5.13)	556	7.19 (5.19-9.67)	323	0.62 (0.08-2.22)	2183	7.56 (6.48-8.75)
Medicaments, excipients									
Caine mix III (Benzo-,Cincho-,Tetracaine)	10.0	435	1.61 (0.65-3.29)	1237	3.07 (2.18-4.19)	482	0.32 (0.01-1.8)		
Budesonide	0.01	499	1.6 (0.69-3.13)	1242	0.16 (0.02-0.58)	495	0.31 (0.01-1.71)	2183	0.5 (0.25-0.9)
Budesonide	0.1								

Allergen	Conc.	LT tested	LT % pos. (95% CI)	NL tested	NL % pos. (95% CI)	PT tested	PT % pos. (95% CI)	SI tested	SI % pos. (95% CI)
Tixocortol pivalate	0.1	453	0.22 (0.01-1.22)	1803	0.67 (0.34-1.16)	481	0 (0-1.14)	2183	0.23 (0.07-0.53)
Tixocortol pivalate	1.0			1212	0.74 (0.34-1.4)				
Neomycin sulfate	20	499	1.4 (0.57-2.87)	1242	0.72 (0.33-1.37)	497	1.24 (0.34-3.14)	2183	1.05 (0.67-1.58)
Lanolin (wool alcohols)	30	499	1.2 (0.44-2.6)	1239	3.47 (2.52-4.65)	497	1.24 (0.34-3.14)	2183	0.82 (0.49-1.3)
Rubber additives									
Thiuram mix	1.0	499	0.4 (0.05-1.44)	1242	2.01 (1.31-2.96)	497	2.48 (1.08-4.82)	2183	1.88 (1.35-2.54)
N-Isopropyl-*N*'-phenyl-*p*-phenylenediamine	0.1	499	0.4 (0.05-1.44)	1242	0.97 (0.5-1.68)	495	0.31 (0.01-1.71)	2183	0.46 (0.22-0.84)
Mercapto mix[(i)]	1.0			467	1.07 (0.35-2.48)				
Mercapto mix[(ii)]	2.0	331	0.3 (0.01-1.67)			497	0.62 (0.08-2.22)	2183	0.41 (0.19-0.78)
Mercaptobenzothiazole	2.0	499	0.2 (0.01-1.11)	1243	0.32 (0.09-0.82)	497	0.93 (0.19-2.69)	2183	0.18 (0.05-0.47)
Resins/glues									
Colophonium	20.0	499	4.81 (3.11-7.07)	1241	3.55 (2.59-4.73)	497	0.62 (0.08-2.22)	2183	4.49 (3.66-5.44)
Epoxy resin	1.0	499	0.8 (0.22-2.04)	850	2.47 (1.54-3.75)	495	1.24 (0.34-3.14)	2183	1.65 (1.16-2.28)
p-tert-Butylphenol formaldehyde resin	1.0	499	0.2 (0.01-1.11)	1242	0.97 (0.5-1.68)	497	0.62 (0.08-2.22)	2183	0.37 (0.16-0.72)
2-Hydroxyethyl methacrylate	2.0	453	3.09 (1.7-5.13)	310	3.87 (2.02-6.66)	493	3.74 (1.95-6.44)		
Other									
p-Phenylenediamine	1.0	499	2.2 (1.11-3.91)	1239	3.55 (2.59-4.74)	497	3.1 (1.49-5.62)	2183	2.52 (1.9-3.27)
Sesquiterpene lactone mix	0.1	499	0.8 (0.22-2.04)	1790	1.56 (1.04-2.25)	496	0.31 (0.01-1.71)	2183	0.64 (0.35-1.07)
Propolis	10	48	6.25 (1.31-17.2)	1557	4.24 (3.29-5.36)	497	0 (0-1.14)		
Textile dye mix	6.6	453	3.97 (2.37-6.21)	1799	4.89 (3.94-5.99)	492	1.56 (0.51-3.61)	2183	2.98 (2.31-3.78)

LT, Lithuania; NL, The Netherlands; PT, Portugal; SI, Slovenia; epoxy resin, diglycidyl ether of bisphenol A; HICC, hydroxyisohexyl 3-cyclohexene carboxaldehyde; mercapto mix[(i)], containing N-cyclohexylbenzothiazyl sulfenamide, dibenzothiazyl disulfide, and morpholinylmercaptobenzothiazole; mercapto mix[(ii)], containing N-cyclohexylbenzothiazyl sulfenamide, mercaptobenzothiazole, dibenzothiazyl disulfide, and morpholinylmercaptobenzothiazole; MCI, methylchloroisothiazolinone; MI, methylisothiazolinone.

Allergen	Conc.	UK tested	UK % pos. (95% CI)
Metals			
Potassium dichromate	0.5	1773	0.90 (0.52-1.46)
Cobalt chloride	1.0	1770	2.03 (1.43-2.81)
Nickel sulfate	5.0	1744	13.1 (11.6-14.8)
Fragrances			
Fragrance mix I	8.0	1771	5.48 (4.46-6.64)
Fragrance mix II	14.0	1772	2.43 (1.76-3.26)
HICC	5.0	1773	0.56 (0.27-1.03)
Myroxylon pereirae (balsam of Peru)	25	1770	4.69 (3.75-5.78)
Preservatives			
Formaldehyde	1.0[a]		
Formaldehyde	2.0[a]	1774	0.90 (0.52-1.46)
MCI/MI 3:1	0.01[a]		
MCI/MI 3:1	0.02[a]	1218	1.89 (1.20-2.82)
Methylisothiazolinone	0.02[a]		
Methylisothiazolinone	0.05[a]		
Methylisothiazolinone	0.20[a]	1218	2.05 (1.33-3.02)
Paraben mix	16	1774	0.11 (0.01-0.41)
Quaternium-15	1.0	1774	0.23 (0.06-0.58)
Methyldibromo glutaronitrile	0.3	1774	0.51 (0.23-0.97)
Methyldibromo glutaronitrile	0.5		
Medicaments, excipients			
Caine mix III (Benzo-,Cincho-,Tetracaine)	10.0	1774	0.96 (0.56-1.53)
Budesonide	0.01		
Budesonide	0.1	1773	0 (0-0.21)

Allergen	Conc.	UK tested	UK % pos. (95% CI)
Tixocortol pivalate	0.1		
Tixocortol pivalate	1.0	1771	0.68 (0.35-1.18)
Neomycin sulfate	20	1773	0.28 (0.09-0.66)
Lanolin (wool alcohols)	30	1774	0.39 (0.16-0.81)
Rubber additives			
Thiuram mix	1.0	1773	0.96 (0.56-1.53)
N-Isopropyl-*N*'-phenyl-*p*-phenylenediamine	0.1	1774	0.28 (0.09-0.66)
Mercapto mix[(i)]	1.0		
Mercapto mix[(ii)]	2.0	1774	0.28 (0.09-0.66)
Mercaptobenzothiazole	2.0	1774	0.23 (0.06-0.58)
Resins/glues			
Colophonium	20.0	1774	2.31 (1.66-3.12)
Epoxy resin	1.0	1770	0.40 (0.16-0.81)
p-tert-Butylphenol formaldehyde resin	1.0	1773	0.34 (0.12-0.74)
2-Hydroxyethyl methacrylate	2.0	1774	2.54 (1.86-3.38)
Other			
p-Phenylenediamine	1.0	1214	3.38 (2.43-4.55)
Sesquiterpene lactone mix	0.1	1774	0.51 (0.23-0.96)
Propolis	10	1112	1.53 (0.89-2.44)
Textile dye mix	6.6	490	1.43 (0.58-2.92)

UK, United Kingdom; Epoxy resin, diglycidyl ether of bisphenol A; HICC, hydroxyisohexyl 3-cyclohexene carboxaldehyde; mercapto mix[(i)], containing N-cyclohexylbenzothiazyl sulfenamide, dibenzothiazyl disulfide, and morpholinylmercaptobenzothiazole; mercapto mix[(ii)], containing N-cyclohexylbenzothiazyl sulfenamide, mercaptobenzothiazole, dibenzothiazyl disulfide, and morpholinylmercaptobenzothiazole; MCI, methylchloroisothiazolinone; MI, methylisothiazolinone.

Online supplemental Table S3: Patch test results (day 3 to day 5) with the TRUE Test, 2019–2020, stratified for country (Spain and The Netherlands [Groningen]).

Allergen	Conc.	ES tested	ES % pos. (95% CI)	NL tested	NL % pos. (95% CI)
Potassium dichromate	0.054	3065	3.1 (2.5–3.8)	555	3.42 (2.07–5.29)
Cobalt (II)-chloride, 6*H2O	0.02	3064	4.7 (4–5.5)	555	5.95 (4.13–8.25)
Nickel (II)-sulfate hexahydrate	0.2	3064	25.3 (23.7–26.8)	551	16.88 (13.84–20.27)
Gold sodium thiosulfate	0.075	3063	5 (4.2–5.8)		
Fragrance mix	0.5	3065	3.4 (2.8–4.1)	555	3.24 (1.93–5.08)
Balsam of Peru (*Myroxolon pereirae*)	0.8	3064	1.7 (1.3–2.2)	555	1.44 (0.62–2.82)
Formaldehyde	0.18	3065	1 (0.7–1.4)	555	1.44 (0.62–2.82)
Methylchloroisothiazolinone Methylisothiazolinone (MCI/MI)	0.004	3065	7 (6.1–7.9)	555	6.49 (4.58–8.87)
Paraben mix	1	3065	0.4 (0.2–0.6)	555	0 (0–0.66)
Quaternium 15	0.1	3065	1.2 (0.9–1.7)	555	1.62 (0.74–3.06)
Diazolidinyl urea	0.55	3065	0.5 (0.2–0.8)	407	0.49 (0.06–1.76)
Imidazolidinyl Urea (Germall 115)	0.6	3065	0.4 (0.2–0.7)	407	0.98 (0.27–2.5)
Thiomersal (Thimerosal)	0.007	3065	3.5 (2.8–4.2)	555	1.26 (0.51–2.58)
2-Bromo-2-nitro-1,3-propanediol	0.25	3065	0.7 (0.4–1)		
Methyldibromoglutaronitrile	0.005	3065	0.7 (0.4–1)		
Caine Mix III (Benzo-,Cincho-,Tetracaine)	0.63	3065	1 (0.7–1.4)	555	0.54 (0.11–1.57)
Budesonide	0.001	3064	0.8 (0.6–1.2)	407	0.25 (0.01–1.36)
Tixocortol-pivalate	0.002	3064	0.4 (0.2–0.6)	407	0.49 (0.06–1.76)
Hydrocortisone-17-butyrate	0.02	3064	0.4 (0.2–0.7)	407	0.49 (0.06–1.76)
Neomycin sulfate	0.6	3064	0.9 (0.6–1.3)	555	0 (0–0.66)
Ethylene diamine-di-HCl	0.05	3064	1 (0.7–1.4)	555	0.18 (0–1)
Quinoline mix	0.19	3064	0.4 (0.2–0.7)		
Bacitracine	0.6	3063	0.1 (0–0.2)		
Lanolin (wool fat) alcohols	1	3064	0.7 (0.4–1)	555	0.9 (0.29–2.09)
Thiuram mix	0.027	3064	1.5 (1.1–2)	555	2.52 (1.39–4.2)

Allergen	Conc.	ES tested	ES % pos. (95% CI)	NL tested	NL % pos. (95% CI)
PPD (black rubber) mix	0.075	3064	0.8 (0.6–1.2)	555	1.08 (0.4–2.34)
Mercapto mix (CBS, MBTS, MOR)	0.075	3064	0.3 (0.2–0.6)	407	1.23 (0.4–2.84)
Mercaptobenzothiazole	0.075	3064	0.4 (0.2–0.7)	555	0.9 (0.29–2.09)
Carba mix	0.25	3064	1.8 (1.3–2.3)	555	4.86 (3.23–7)
Colophony (Rosin)	1.2	3064	1.7 (1.2–2.2)	555	5.05 (3.38–7.21)
Epoxy resin	0.05	3064	1.4 (1–1.9)	555	2.7 (1.52–4.42)
p-tert.-Butylphenol formaldehyde resin (PTBFR)	0.045	3064	2.1 (1.6–2.6)	554	3.25 (1.94–5.09)
p-Phenylenediamine (PPD)	0.08	3065	3.9 (3.3–4.7)	554	3.61 (2.22–5.52)
Parthenolide	0.003	3064	0.3 (0.1–0.5)		
2-Bromo-2-nitro-1,3-propanediol	0.25	3065	0.7 (0.4–1)		