

Airborne irritant contact dermatitis from benzalkonium chloride

NINA E. KROGSRUD AND ANDERS I. LARSEN

Occupational Health Service, Novo Nordisk A/S, Novo Allé, DK 2880 Bagsvaerd, Denmark

Key words: airborne; irritant contact dermatitis; benzalkonium chloride; laboratory worker; occupational. © Munksgaard, 1997.

Case Report

A laboratory technician, without previous skin diseases and non-atopic, developed urticaria on her face and neck 4–6 h after washing a laminar air flow (LAF) bench with a benzalkonium chloride (BAC) solution. The same evening, severe itching began. The skin became red and vesicles appeared on the chin and on the front of the neck. The symptoms lasted for 2 days. No symptoms were recognized on the hands or the arms, and she showed no symptoms of the airways and mucosa.

Environmental exposures: while bending halfway into the LAF bench, the patient sprayed a solution of BAC 2% mixed with ethanol 67% and distilled water inside the LAF bench and wiped up with a cloth. Following this procedure, she washed her hands. She did not use gloves while disinfecting the LAF bench. She works daily with industrial enzymes, bacteria and fungi.

RAST and scratch tests for relevant enzymes were negative, apart from a temporary reaction caused by dermographism. Patch test to BAC 0.1% aq. was negative (Scanpor® tape and Finn Chambers® were used, 2 days application, readings done at days 2 and 3).

The patient continues to work with BAC and enzymes, but she now pours the BAC directly onto the wipe cloth to avoid aerosol formation. So far no symptoms have reappeared.

Discussion

To the list of agents causing airborne dermatitis (1), we have hereby added BAC (2). Although the concentration of BAC in the application solution was low, evaporation of the volatile solvents (water and alcohol) may have resulted in an increased concentration of the irritant on the exposed area. In high concentrations, BAC is considered to be corrosive. Consequently, we think it is probable that BAC caused the eczema. Formation of and exposure to aerosols should be avoided when handling irritants.

Acknowledgement

We thank Professor Torkil Menné, Department of Dermatology, Gentofte Hospital, Denmark, for performing patch testing and for case discussion.

References

1. Dooms-Goossens A, Deleu H. Airborne contact dermatitis: an update. *Contact Dermatitis* 1991; 25: 211–217.
2. Rycroft R J G, Menné T, Frosch P J, Benezra C (eds): *Contact Dermatitis*, 1st edition. Berlin: Springer-Verlag, 1991: 38: 466.

Systemic contact dermatitis caused by nickel in a stainless steel orthodontic appliance

HEIDI KEROSUO¹ AND LASSE KANERVA²

¹Institute of Dentistry, Department of Orthodontics and Pedodontics, University of Helsinki, Helsinki, Finland

²Section of Dermatology, Finnish Institute of Occupational Health, Topeliuksenkatu 41aA, FIN-00250, Helsinki, Finland

Key words: systemic contact dermatitis; nickel; cobalt; stainless steel; orthodontic appliance; children; dental patients; allergic contact cheilitis. © Munksgaard, 1997.

Case Report

A 14-year-old atopic boy began orthodontic treatment with 2 permanent bands on the 1st maxillary molars, and an extraoral face bow (Fig. 1) to be worn at night. All appliances were stainless steel, usually 18% chromium and 8% nickel.

At the 1st follow-up visit 4 weeks later, he showed crusted yellowish vesicles on and around the lips (Fig. 1), and reddish eczema with scaling and fissures on the palms and soles. The dermatitis had developed soon after insertion of the appliance. The patient had seen a doctor and been prescribed local medication for the eczema. 5 weeks later, at the next visit, the eczema had

This document is a scanned copy of a printed document. No warranty is given about the accuracy of the copy. Users should refer to the original published version of the material.