

Efficacy of Waste Biomass Based Tablet Formulation for Cockroach Control

Megha Pant^{1,4} · Saurabh Dubey¹ · Eldho Varghese² · Neeraj Kumar³ · Satya Narayan Naik¹ · Phool Kumar Patanjali⁴

Received: 27 June 2016 / Accepted: 16 October 2017 / Published online: 10 November 2017
© Springer Science+Business Media B.V. 2017

Abstract The current paper has elaborated the efficient utilization of house hold botanical waste like OP (orange peel), UTL (used tea leaves) and industrial waste i.e., KCP and JCP (non-edible oil seed cakes of Karanja and jatropa) for developing a herbal tablet formulation for cockroach control. The percentage mortality and KD50 values recorded with and without KCP and JCP were 25.9597, 100% (after 96 h) and 50.6697, 90% (After 96 h). Analysis with ATR-FTIR also confirmed the presence of presence of OP; KCP& JCP in tablet composition. SEM analysis generated the

morphological details for the tablet containing active and other inert ingredients.

Keywords Jatropa cake powder (JCP) · Karanja cake powder (KCP) · Orange powder (OP) · Used tea leaves powder (UTL)

Introduction

Synthetic chemical insecticides have been considered as an important part of pest management for many years. Interest is however growing fast in discovering the possible role of plants and minerals as traditional protectants of stored products and as alternatives to the use of highly persistent synthetic chemicals with their attendant dangers and high cost. The use of natural products, which are cheap, effective, less toxic to mammals, easy to adopt and environmentally friendly have been investigated worldwide [1, 2]. Over 120 plants and plants products have been shown to have insecticidal or deterrent activity against stored product pests [3]. Currently, farmers in parts of Africa and Asia use some of these botanicals to protect their legumes from attack by bruchids, with varying degrees of success [4, 5]. However the number and quality of plants used by farmers is often limited by availability [5]. It was reported that powder peels of citrus fruits were effective in reducing the damage caused by *Callosobruchus maculatus* to cowpea seeds [6]. Also [7] reported that the crude extract of *Piper guineense* gave adequate protection to stored cowpea against *Callosobruchus maculatus*. Limonene, a monoterpene constituent in the extracts of citrus fruit peel was reported as a safe, natural pesticide for insect pests (Fig. 1a) [8]. The efficacy of orange peel has already been established [4].

A part of this paper has already been patented via patent application No. 2705/DEL/2012.

✉ Saurabh Dubey
drsaurabh_dubey@rediffmail.com

Megha Pant
mpmegha04@gmail.com

Eldho Varghese
eldhoiasri@gmail.com

Neeraj Kumar
neerajagrichem90@gmail.com

Satya Narayan Naik
snn@iitd.ac.in

Phool Kumar Patanjali
pkpatanjali@yahoo.com

¹ Centre for Rural Development and Technology(CRDT), Indian Institute of Technology, Delhi, New Delhi 110016, India

² Division of Design of Experiments, Indian Agricultural Statistics Research Institute, New Delhi 110012, India

³ Department of Agricultural Chemicals, Indian Agricultural Research Institute, New Delhi 110060, India

⁴ Formulation Division, Institute of Pesticide and Formulation Technology, Gurgaon, Haryana 122016, India




Fig. 1 **a** Structure of Limonene, **b** structure of Jatropa (Phorbol esters), **c** structure of Karinjin (3-methoxyflavone)

Considering the extraction of oil from the non-edible seeds, karanja (*Pongamia pinnata*) and jatropa (*Jatropha curcas*) have been selected as a major source of non-edible oil for production of biodiesel [9]. The technology of biodiesel production consumes only extracted vegetable oil from non-edible seeds and renders a large amount of unutilized biomass as seed cake. Disposal of cake after expelling oil from seed has become a major concern in the past few years. Due to its toxic nature, the cake neither can be used for animal feeding nor can be used in agricultural farming.

The non-edible seeds of jatropa (*Jatropha curcas*) and karanja (*Pongamia glabra*) contain some toxic components, i.e., phorbol esters (Fig. 1b) in *Jatropha curcas* and karanjin (Fig. 1c) in *P. glabra*, which can be exploited to function as bio-pesticides [10].

Literature survey does give some information about the insecticidal activity of neem and karanja oil and their cakes in combination against mosquito vectors [11–13].

Household pests include the various insect species that invade human habitation causing damage to structures, eating human foods, acting as disease vectors and

causing other threats. Cockroaches are one of the most commonly noted, unlikable household pests [14]. They have a widespread distribution with different species, eg. *Periplaneta americana*, *Blattella germanica*, *Blatta orientalis* etc. They produce odorous secretions that affect flavour of various food items. Disease producing organisms such as bacteria, protozoa and viruses have been found in their bodies. Different forms of gastroenteritis appear to be principal diseases transmitted by them. The disease causing organisms are carried on their legs and bodies and are deposited on food and utensils as they forage. Cockroach excrement and skin casts also contain a number of allergens to which people exhibit allergic responses such as skin rashes, asthma, etc [15].

In the present work, the knocking down and insecticidal activity of the tablet formulation containing bio-diesel by-product and orange peel powder based active ingredients individually and in combination has been compared with the efficacy of commercially available synthetic as well as herbal pesticide based product separately.

Materials and Methods

Karanja, and jatropha cakes were collected from the Centre for Rural Development Technology, I.I.T, and Delhi. Orange peel and used tea leaves collected as household waste. Magnesium stearate was purchased from Merck, Mumbai, India.

Preparation of Tablet

Botanical tablets (Fig. 2a) were prepared by mixing the botanical active ingredients and other components and compressing them into a solid mass using a Tablet Machine. They were kept in glass Petri dishes for some time at room temperature for drying and then used for Bio efficacy trials.

Test Insect & Contact Toxicity Bioassay

The efficacy of the Tablets with and without the karanja and jatropha deoiled cake was evaluated against the American cockroach, *Periplaneta americana*. The cockroaches were collected from the fields of Delhi University, India, which are maintained as pesticide free zones by the university authorities with an average life expectancy of 600 days i.e. females-440 days and males-200–300 days. Cultures of *Periplaneta Americana* were maintained at laboratory conditions, 60–70% RH, 27 ± 1 °C and darkness i.e. 12:12 (L:D) h photoperiod in specially designed cages. The cages were made of coated polyacrylic material with dimensions and windows of muslin cloth on two sides for proper aeration and a slider door to keep and remove the cockroaches out of the cages. 10 cockroaches were taken for each composition of tablet prepared and released into cages with 10 tablets as dose given to them. The experiment was carried in duplicates. Ten cockroaches were kept as control in cages with filter paper and moist bread provided to them as food with no other harbourage material in them. Observations were taken for knockdown and mortality characteristics for 96 h (Fig. 2b) [16].

Contact toxicity bioassay was performed by applying selected doses of the tablets (1 tablet for 1 cockroach) in the cages with the help of a micro applicator. 10 cockroaches were released in cages containing emulsion and mortality was observed after 24, 48, 72 and 96 h. Each assay for different doses of the emulsion was performed 3 times. A control set was also maintained [17].

Analysis of Variance (ANOVA) and DMRT

The results calculated here are means of three replicates. The standard deviation was calculated and reported as S.E. (Standard error) in the tables. The data was analysed by one-way analysis of variance (ANOVA) and significant effects were noted using SAS 9.4. along similar lines as in [18, 19]. Duncan's multiple range test (DMRT) 5% level was also employed to test for significant differences between treatments.

ATR-FTIR Analysis of Tablet

Infrared (IR) spectra were recorded on a Bruker alpha ATR-FTIR spectrophotometer using the attenuated total reflectance (ATR) technique, and values are expressed as gmax cm^{-1} [20].

Scanning Electron Microscopy (SEM)

SEM photographs were taken with JSM 5600 scanning electron microscopy, (JEOL limited Tokyo, Japan) to analyze surface morphology of tablet containing orange peel powder, karanja cake powder and jatropha cake powder as a active ingredient [21].

Fig. 2 a Tablet formulation, b cockroach mortality



Result and Discussion

The active component in tablet formulation is d-limonene, which is responsible for its insecticidal activity [18]. Confirmation of the active component in the tablet formulations prepared was done by using an ATR-FTIR technique. Limonene in general showed an infrared spectra in the following region of wave number; 2851 and C–H stretching spectra of limonene was found at 2900 cm^{-1} (Fig. 3) [22]. Phorbol esters [23] and karinjin are the major constituents of jatropha and karanja cake that were responsible for their bio-pesticidal properties [24, 25]. Their presence in the tablet was confirmed by ATR-FTIR analysis. ATR-FTIR technique gives confirmation of no chemical reaction between active and inert ingredient in tablet formulation.

Scanning electron microscopy (SEM) analysis helped to study the surface modification of orange peel powder before and after mixing with other ingredients for preparing tablets [26] Fig. 4a, b. The surface of orange peel powder showed irregular distributed pores and the pores became smooth and uniform when mixed with other inert ingredients like binders and lubricant with orange peel powder [27].

Table 1 shows the percentage mortality data with time of tablets containing using orange peel powder (OP), used tea leaves powder (UTLP) and glucose. The mortality results were also compared with commercially available synthetic as well as botanical pesticide based products i.e. gel and

tablets. Mortality values increased with time i.e. 10% in case of botanical (OP + UTLP + glucose) and 30% in case of market sample (synthetic and botanical) in 24 h. After 96 h values became similar i.e. 90% in both samples (botanical and market sample). Other compositions were also prepared by utilizing karanja cake powder (KCP) and jatropha cake powder (JCP) (Table 2) mixed with orange peel powder, used tea leaves powder and glucose along with magnesium stearate as a lubricant. In case of tablets containing KCP and JCP with combination of OP, UTLP and glucose, increase in the percentage mortality was 40–50% in 24 h as compared to market sample. After 96 h, Mortality rates reached to 100% as compared to market product i.e., synthetic (fipronil gel) as well as botanical (neemka paste) samples, where it was (90%).

Limonene [28] an active constituent of orange peel and also responsible for insecticidal activity [18], was used as an adsorbent in effluent treatment [29] process earlier. In the formulation described in the current paper, used tea leaves powder replaced the use of carbohydrate polymers like starch and cellulose. KCP and JCP containing karinjin and phorbol esters showed insecticidal activity against household as well as stored grain pest [10, 30], and the efficacy increased when combined with OP, UTLP and glucose. Toxicity of phorbol ester against cockroaches was reported [31] but the presence of limonene and karinjin enhanced the insecticidal activity and mortality was also found to




Fig. 3 ATR-FTIR spectra of tablet containing OP, KCP and JCP




Fig. 4 **a** SEM photographs of tablet containing OP, KCP, JCP and UTL, **b** SEM photographs of tablet containing OP and UTL

Table 1 Statistical evaluation of the percentage mortality data of tablets containing OP + UTL + glucose in comparison with commercially available synthetic and botanical products for American cockroach with mean and standard error

Sample name (tablets)	Replicates (N)	Mean% mortality with standard error (after 24 h)	Mean % mortality with standard error (after 48 h)	Mean % mortality with standard error (after 72 h)	Mean % mortality with standard error (after 96 h)
OP + UTL + glucose	3	6.14 ± 2.89 ^b	21.14 ± 2.89 ^c	46.92 ± 2.89 ^b	68.85 ± 2.89 ^a
Fipronil gel	3	35.21 ± 2.89 ^a	59.00 ± 2.89 ^a	61.21 ± 2.89 ^a	68.85 ± 2.89 ^a
Neemka paste	3	28.78 ± 2.89 ^a	35.21 ± 2.89 ^b	46.92 ± 2.89 ^b	54.78 ± 2.89 ^b
Control	3	0.00 ± 2.89 ^b	0.00 ± 2.89 ^d	0.00 ± 2.89 ^c	0.00 ± 2.89 ^c

Mean ± SE with same superscript letter are not significantly different at $p < 0.05$ by DMRT

Table 2 Statistical evaluation of the percentage mortality data of tablets containing OP + UTL + KCP + JCP + glucose in comparison with commercially available synthetic and botanical products for American cockroach with mean and standard error

Sample name (tablets)	Replicates (N)	Mean% mortality with standard error (after 24 h)	Mean % mortality with standard error (after 48 h)	Mean % mortality with standard error (after 72 h)	Mean % mortality with standard error (after 96 h)
OP + UTL + KCP + JCP + glucose	3	41.15 ± 2.89 ^a	52.77 ± 2.89 ^a	66.14 ± 2.89 ^a	90.00 ± 2.35 ^a
Fipronil gel	3	35.21 ± 2.89 ^{ab}	59.00 ± 2.89 ^a	61.21 ± 2.89 ^a	68.85 ± 2.35 ^b
Neemka paste	3	28.78 ± 2.89 ^b	35.21 ± 2.89 ^b	46.92 ± 2.89 ^b	54.78 ± 2.35 ^c
Control	3	0.00 ± 2.89 ^c	0.00 ± 2.89 ^c	0.00 ± 2.89 ^c	0.00 ± 2.35 ^d

Mean ± SE with same superscript letter are not significantly different at $p < 0.05$ by DMRT

be 100% within 96 h (Table 1). It was found KD50 values gradually decreased to 25.9597 min with the incorporation of KCP and JCP in tablet composition as compared to tablet containing OP and UTL (Table 2), where it was found to be 50.6697 min. The KD50 values of tablet composition with and without KCP and JCP were also compared with the KD50 value of commercially available synthetic and botanical products i.e. fipronil gel and neemka which was reported to be 29.971 and 63.5907 min respectively.

A Botanical tablet consists of ingredients derived from naturally occurring plant wastes. Various compositions

were obtained by mixing different botanical waste based active ingredients. In one of the compositions, feeding attractant, glucose was mixed with the insecticide to attract the pest. These botanical wastes when used in different combinations cause behavioural and physiological abnormalities disrupting the metabolism, digestion leading to starvation and ultimately death (Fig. 5).




Fig. 5 **a** Percentage mortality at different time intervals, **b** percentage mortality at different time intervals

Conclusion

Tablets prepared using these botanical by-products as active ingredients are less toxic, eco-friendly and economical as compared to commercially available synthetic as well as botanical products. The paper mainly focuses on the efficient use of orange peel and biodiesel product, having insecticidal activity, which would otherwise cause waste disposal problems.

Acknowledgements The authors gratefully acknowledge support from the University Grants Commission, India, Grant Number: F.15-42/SAII/2012, and from Dr. Rajesh, Senior Scientist, IARI, Agrochemical Division, New Delhi, India.

References

- Oparaeke, A.M., Dike, M.C.: *Monnodoramyristica* (Gaertn) *Dunalmyristicaceae* and *Allium cepa* L. (Liliaceae) as protectants against cowpea seed bruchid, *Callosobruchus maculatus* (F) infesting stored cowpea seeds. *Nig. J. Entomol.* **22**, 84–92 (2005)
- Oparaeke, A.M., Amodu, M.E.: Insecticidal efficacy of cashew products against *Callosobruchus maculatus* fab. on stored cowpea. *Span. J. Agric. Res.* **19**, 135–140 (2000)
- Dale, N.J.: A Review of Plant Materials Used for Controlling Insect Pests of Stored Products. *National Resource Insect bulletin*. Chatham, pp 36–50 (1996)
- Don-pedro, K.N.: Insecticidal activity of fatty acid constituents of fixed vegetable oils against *Callosobruchus maculatus* (F.) on cowpea. *Pest. Sci.* **30**(3), 295–302 (1990)
- Dharmasena, C. M. D., Simmonds, M. S. J., Blaney, W. M.: Insecticidal activity of eight plant species on egg laying, larval development and adult emergence of *Callosobruchus maculatus* (F.) in cowpea. *Trop. Agric. Res. Ext.* **1**, 67–69 (1998)
- Onu, L., Sulyman, A.: Effect of powdered peels of citrus fruits on damage by *Callosobruchus maculatus* (F) to cowpea seeds. *J. Sustain. Agric.* **9**(4), 85–92 (1997)
- Okonkwo, E. U., Okoye, W. I.: Effect of plant materials on damage and germination of cowpea seeds stored on small farms. *Nig. J. Entomol.* **19**, 90–98 (2002)
- Hilling, S. R.: Limonene, a citrus extract for control of mealy bugs and scale insects. *J. Econ. Entomol.* **98**(3), 772–779 (2005)
- Agarwal, A. K., Saravanan, I. B., Kumar, E. N.: Biodiesel as an alternative urban transport fuel in India. In: *Bio fuels in India: a new revolution first edition*, pp. 151e157. The ICFAI University Press, Hyderabad, (2007)
- Wakandigara, A., Nhamo, L. R. M., Kugara, J.: Chemistry of phorbol ester toxicity in *Jatropha curcas* seed—a review. *Int. J. Biochem. Res. Rev.* **3**(3), 146–161 (2013)
- Sharma, S., Verma, M., Prasad, R., Yadav, D.: Efficacy of non-edible oil seed cakes against termites *Odontotermes obesus*. *J. Sci. Indust. Res.* **70**, 1037–1041 (2011)
- Shanmugasundaram, R., Jeyalakshmi, T., Dutt, M.S., Murthy, P.B.: Larvicidal activity of neem and karanja oil cakes against mosquito vectors, *Culex quinquefasciatus* (Say), *Aedes aegypti* (L.) and *Anopheles stephensi* (L.). *J. Environ. Biol.* **29**, 43–50 (2008)
- Pant, M., Dubey, S., Raza, S. K., Patanjali, P. K.: Encapsulation of neem and karanja oil mixture for synergistic as well as larvicidal activity for mosquito control. *J. Sci. Indust. Res.* **71**, 348–352 (2012)
- Patel, S., Meher, B.R.: A review on emerging frontiers of house dust mites and cockroaches. *Allergol. Immunopathol.* **14**(6), 580–593 (2016)
- Pereira, R.M., Oi, D.H., Baggio, M.V., Kochler, P.G.: Microbial control of structural insect pest. *Microbial control of Insect and Mites Pest*, Chap. 29, pp. 431–442 (2017)
- Kumar, N., Pant, M., Singh, O.P., Dubey, S., Naik, S.N., Sharma, S., Patanjali, P.K.: Bioefficacy and stabilization of hydrolytically unstable pesticide in water based microemulsion against *Periplaneta americana*. *Int. J. Adv. Res. IT. Eng.* **5**(1), 9–25 (2016) ISSN: 2278–6244
- Kaakeh, W., Reid, B.L., Bennett, G.W.: Toxicity of fipronil to German and American cockroaches. *Entomol. Exp. Appl.* **84**, 229–237 (1997)
- Kumar, N., Mukherjee, I., Varghese, E.: Adsorption–desorption of tricyclazole: effect of soil types and organic matter. *Environ. Monit. Assess.* **187**, 61 (2015)
- Chatterjee, N.S., Gupta, S., Varghese, E.: Degradation of metafluzone in soil: impact of varying moisture, light, temperature, atmospheric CO₂ level, soil type and soil sterilization. *Chemosphere.* **90**, 729–736 (2013)
- Shi, Y., Zheng, T., Shang, Q.: Preparation of acrylic/acrylate copolymeric polymerization used in pesticide oil in water emulsions. *J. Appl. Polym. Sci.* **123**(5), 3117–3127 (2012)
- Lucia, A., Audino, P.G., Seccarini, E., Licastro, S., Zerba, E., Masuch, H.: Larvicidal effect of *Eucalyptus grandis* essential oil and turpentine and their major components on *Aedes aegypti* larvae. *J. Am. Mosq. Control. Assoc.* **23**, 299–303 (2007)

22. Palacios, M.S., Bertoni, A., Rossi, Y., Santander, R.: Efficacy of essential oils from edible plants as insecticides against the house fly, *Musca Domestica* L. J. Mol. Biol. **14**, 1938–1947 (2009)
23. Amanzadeh, Y., Ashrafi, M., Mohammadi, F.: New elaborated technique for isolation and purification of limonene from Orange Oil. Iran J. Pharm. Sci. **2**(2), 87–90 (2006)
24. Phasukarratchai, N., Tontayakom, V., Tongcumpou, C.: Reduction of phorbol esters in *Jatropha curcas* L. pressed meal by surfactant solutions extraction. Biomass Bioenerg. **45**, 48–56 (2012)
25. Baswa, M., Rath, C.C., Dash, S.K., Mishra, R.K.: Antibacterial activity of Karanj (*Pongamia pinnata*) and Neem (*Azadirachta indica*) seed oil: a preliminary report. Microbios. **105**, 183–189 (2000)
26. Verma, M., Pradhan, S., Sharma, S., Naik, S.N., Prasad, R.: Efficacy of karanjin and phorbol ester fraction against termites. *Odonotermesobesus*. Int. Biodeter. Biodegr. **65**, 877–882 (2011)
27. Liu, Y., Jehanathan, N., Yang, H., Laeng, J.: SEM observation of the “orange peel effect” of materials. Mater. Lett. **61**, 1433–1435 (2007)
28. Huang, H., Hu, X.Z., Liu, Y., Bush, M., Winchester, K., Musca, C., Dell, J., Faraone, L.: Characterization of mechanical properties of silicon nitride thin films for MEMS devices by nano-indentation. J. Mater. Sci. Technol. **21**, 13–16 (2005)
29. Phillips, A. K., Appel, A.G., Sims, S.R.: Topical toxicity of essential oils to the German cockroach (Dictyoptera: Blattellidae). J. Econ. Entomol. **103**, 448–459 (2010)
30. Tzeng, J.-H., Weng, C.-H., Huang, J.-W., Lin, Y.-H., Lai, C.-W., Lin, Y.-T.: Spent tea leaves: a new non-conventional and low-cost biosorbent for ethylene removal. Int. Biodeter. Biodegr. **104**, 67–73 (2015)
31. Pant, M., Dubey, S., Patanjali, P. K., Naik, S. N., Sharma, S.: Insecticidal activity of eucalyptus oil nanoemulsion with karanja and jatropha aqueous filtrates. Int. Biodeter. Biodegr. **91**, 119–127 (2014)