



Review

Soluble IgE receptors—Elements of the IgE network

Barbara Platzer^{a,b}, Floortje Ruiter^{a,b}, John van der Mee^{a,b}, Edda Fiebiger^{a,b,*}^a Division of Gastroenterology and Nutrition, Children's Hospital, Boston, MA 02115, United States^b Department of Pediatrics, Harvard Medical School, Boston, MA 02115, United States

ARTICLE INFO

Article history:

Received 22 July 2011

Received in revised form 15 August 2011

Accepted 27 August 2011

Available online 6 September 2011

Keywords:

IgE

Fc receptors

FcεRI

CD23

FcεRII

Galectin-3

ABSTRACT

Soluble isoforms of three human IgE Fc receptors, namely FcεRI, FcεRII, and galectin-3, can be found in serum. These soluble IgE receptors are a diverse family of proteins unified by the characteristic of interacting with IgE in the extracellular matrix. A truncated form of the alpha-chain of FcεRI, the high affinity IgE receptor, has recently been described as a soluble isoform (sFcεRI). Multiple soluble isoforms of CD23 (sCD23), the low affinity IgE receptor also known as FcεRII, are generated via different mechanisms of extracellular and intracellular proteolysis. The second low affinity IgE receptor, galectin-3, only exists as a secretory protein. We here discuss the physiological roles of these three soluble IgE receptors as elements of the human IgE network. Additionally, we review the potential and current use of sFcεRI, sCD23, and galectin-3 as biomarkers in human disease.

© 2011 Elsevier B.V. All rights reserved.

1. Introduction

Antibodies of the immunoglobulin E isotype (IgE) are key regulators of host defense against parasitic infections. Over the last three decades, IgE additionally gained undesirable fame as a central mediator of allergic responses. Allergic responses, however, are not regulated by IgE alone, but rather by a complex protein network including transmembrane and soluble IgE receptors and a variety of co-receptors that do not even bind IgE directly (for a detailed review on the human IgE network see Gould et al. [1]).

Soluble IgE receptors are constituents of the human IgE network and are part of feedback mechanisms that regulate IgE production. Therefore, the physiology of these serum components is highly interesting as they are potential *in vivo* modulators of allergic responses. Thus far, three human soluble IgE receptors have been described, namely, soluble FcεRI (sFcεRI), soluble CD23 (sCD23), and galectin-3 (Table 1 and Fig. 1). The focus of this review is to compare and contrast the role of these three soluble IgE receptors in the context of the human IgE network. We discuss the possible physiological roles of the soluble IgE receptors, clinical implications, and elaborate on the potential and current use of soluble IgE receptors as biomarkers of disease.

2. Generation of soluble IgE receptors

2.1. Soluble FcεRI, sFcεRI

sFcεRI is a single-chain receptor isoform of FcεRI, the high affinity IgE receptor. In humans and mice, robust levels of tetrameric FcεRIαβγ₂ are constitutively expressed on the cell surface of mast cells and basophils. This receptor isoform is well known for its function as a key regulator of allergic responses [2]. Under physiological conditions, FcεRIαβγ₂ is preloaded with IgE. When IgE-specific antigen crosslinks the receptor, the release of preformed inflammatory mediators and cytokines is triggered. Thus, IgE-FcεRI mediated activation of mast cells and basophils is considered a hallmark of immediate allergic reactions [2]. The following subunits assemble cotranslationally to form tetrameric FcεRI [3]: an IgE-binding α-chain, FcεRIα, and two signaling subunits, FcεRIβ and FcεRIγ; the latter is commonly referred to as the common Fcγ-chain and dimerizes. In addition to the tetrameric isoform, human antigen presenting cells (APCs), such as Langerhans cells of the skin and various other peripheral blood dendritic cell subpopulations, constitutively express a trimeric αγ₂ isoform of the FcεRI [2,4–6]. In contrast, murine APCs lack constitutive expression of the receptors, but an inducible version of FcεRIαγ₂ has been described in mice after viral infection or challenge with house dust mite [7,8]. Trimeric FcεRIαγ₂ is considered to be an antigen uptake receptor and has been shown to be involved in the regulation of Th2-type allergic tissue inflammation [5,9].

In allergic individuals, induction of FcεRI expression has also been described for many other cell types, including monocytes,

* Corresponding author at: 300 Longwood Avenue, EN630, Boston, MA 02115, United States. Tel.: +1 617 919 2549; fax: +1 617 730 0498.

E-mail address: edda.fiebiger@childrens.harvard.edu (E. Fiebiger).

Table 1
Soluble IgE receptors in human serum.

Soluble IgE receptor	Main source <i>in vivo</i>	Regulation of production
sFcεRI-soluble alpha-chain of FcεRI sCD23	Not defined B cells	IgE-mediated FcεRI activation Surface expression of membrane CD23 and accessibility of cleavage sites Expression and activity of shedding enzymes ADAM10, ADAM8, ADAM33, MMP9 Induction via IL-4 and IL-13 Inhibition by LPS and INF-γ
Galectin-3	Macrophages	

eosinophils, platelets, and gastrointestinal epithelial cell [10–14]. Immunoprecipitation studies from human serum show that sFcεRI consists of a smaller FcεRI alpha-chain with a molecular weight of ~40 kDa compared to the ~60 kDa full length protein [15]. The lower molecular weight is likely explained by the lack of the transmembrane and cytosolic domains [16]. This is also supported by the fact that FcεRIβ and FcεRIγ₂ require the alpha-chain transmembrane region to form a receptor complex and fail to co-immunoprecipitate with sFcεRI. Therefore, ultimately, mass spectrometric analysis is needed to precisely define the protein sequence of sFcεRI.

The alpha-chain of the multimeric FcεRI complex is a type I membrane protein that contains the receptor's IgE-binding site [17]. The soluble alpha-chain, sFcεRI, likewise contains an IgE binding site as it is precipitated with IgE and forms IgE-complexes in serum [15]. Dissociation studies [18] as well as subsequent analysis of the crystal structure [19] of FcεRI-α and IgE revealed an extraordinarily high affinity of this ligand–receptor interaction. As the crystals analyzed were generated with a recombinant soluble version of FcεRI-α, it is likely that serum sFcεRI has a high affinity for IgE consistent with reports in the literature [19].

It has not yet been characterized how the production of sFcεRI is induced *in vivo*. *In vitro* data show that sFcεRI can be generated after IgE-mediated crosslinking of surface-expressed FcεRI when the trimeric isoform of the receptor is expressed in MelJuso cells [15], which are a common model for non-professional

antigen presenting cells [20]. This set of data suggests that production of the soluble isoform is induced by FcεRI crosslinking-induced receptor activation. Since these data were generated with a stable cell line generated with full length FcεRI-α cDNA, sFcεRI could not be produced as a splice variant, but rather must be a product of a posttranslational modification such as cleavage by a protease. Nonetheless, several *in vivo* mechanisms for generating sFcεRI could be operating in parallel as discussed for the other sIgE receptors later in this review.

Currently, many more questions about sFcεRI remain open. For example, the cell types that release or shed this protein in humans remain to be defined. No *in vivo* modulators of sFcεRI production are as of yet known. Furthermore, experiments are needed to investigate whether activation of tetrameric FcεRI also induces the release of sFcεRI. Such experiments will answer the question as to whether sFcεRI exists in mice. If it does not, murine models might be inadequate for studying the potential physiological role of this receptor isoform. Initial experiments to detect sFcεRI from supernatants of IgE-activated RBL-2H3 cells, a rat basophilic leukemia cell line that expresses the tetrameric isoform of the receptor, and murine dendritic cells from a human FcεRIα-transgenic animal that express a chimeric form of FcεRIαγ₂ have failed (Fiebigler lab, unpublished observation and [9]). It is conceivable that expression of the sFcεRI isoform is different between humans and mice, comparable to the dissimilar expression patterns of trimeric sFcεRI between the two species [21]. However, more studies on the topic need to be performed before any conclusions are warranted.

2.2. Soluble FcεRII, sCD23

Soluble CD23 (sCD23) molecules result from proteolytic cleavage of the 45 kDa transmembrane form of the low affinity receptor for IgE, FcεRII (CD23). Unlike FcεRI, this IgE Fc receptor does not belong to the immunoglobulin receptor family. Its large extracellular globular C-type lectin domain places CD23 in the C-type lectin superfamily. Various cell types including B cells, T cells, NK cells, monocytes, macrophages, follicular dendritic cells, Langerhans cells, bone marrow stromal cells, neutrophils, eosinophils, platelets, and epithelial cells express CD23 at the cell surface. For a more detailed insight to the biology of transmembrane CD23, we refer the reader to Gould et al. and Acharya et al. who discussed this topic in their reviews in great detail [1,22].




Fig. 1. Human IgE Fc-receptors and their soluble isoforms. The high affinity IgE Fc receptor, FcεRI (FcεRI), has two transmembrane isoforms, FcεRIαβγ₂ and FcεRIαγ₂. The soluble isoform, sFcεRI, is a single chain receptor consisting of a truncated version of the IgE-binding FcεRIα subunit. Several different soluble isoforms of the transmembrane low affinity IgE Fc receptor, FcεRII or CD23, have been described. A detailed summary of soluble CD23 (sCD23) isoforms and their cleavage sites is provided in Table 2. Galectin-3 is a secretory IgE Fc receptor. After secretion, galectin-3 can attach to cell membranes via interacting with a large number of carbohydrate structures displayed by cell surface proteins. Additionally, an intracellular pool of galectin-3 can be found in the cytoplasm and the nucleus.

Table 2
Soluble CD23 isoforms.

Molecular weight (kDa)	Origin	Enzyme
37	Extracellular shedding of membrane CD23 at position Ala ⁸⁰	<i>in vivo</i> : ADAM10 and MMP9 <i>in vitro</i> : ADAM10, ADAM8 and ADAM 33
33	Extracellular shedding of membrane CD23 at position Arg ¹⁰¹	<i>in vivo</i> : ADAM10 and MMP9 <i>in vitro</i> : ADAM10, ADAM8 and ADAM 33
28–29	Intracellular processing of newly synthesized CD23 protein	Not defined
25–27	Proteolytic cleavage/degradation products of 33 kDa and 37 kDa sCD23	Not defined
16–17	Extracellular cleavage of membrane CD23 at Ser ¹⁵⁵ and Glu ²⁹⁸	Der p1

Based on distinct molecular weights, several sCD23 isoforms have been described (Table 2). The cleavage sites for all forms of sCD23 are located within the extracellular α -helical stalk region of the transmembrane protein. All cleavage products possess the globular lectin head domain, but retain stalk regions of different lengths. Since the IgE binding domain of CD23 resides in the globular lectin head domain, all isoforms of sCD23 have the capacity to interact with IgE.

sCD23 can be produced by intracellular proteolytic processing of newly synthesized CD23 molecules, which are subsequently secreted [23]. This intracellular cleavage event is responsible for the generation of sCD23 with a molecular weight of 28–29 kDa. Additionally, several enzymes have been described to generate sCD23 via extracellular processing. Metalloproteases of the ADAM family were identified as cleavage enzymes for the generation of 37 kDa as well as 33 kDa sCD23 fragments. *In vivo*, ADAM10 has also been found as the predominant sCD23-releasing enzyme [24,25]. B cell-specific deletion of ADAM10 reduces sCD23 levels by 70% [24], indicating that this cell type is the main source of sCD23 in serum. Additionally, ADAM8 and ADAM33 can generate sCD23 *in vitro* [26,27]. The contribution of both proteases for production of sCD23 *in vivo* is still debated, since experiments performed with knock-out animals showed that sCD23 production was not altered in the absence of either protease [25]. The finding that B cells express only very low levels of ADAM8 [28] and no ADAM33 [29] are in line with the observations seen in knock-out animals. Interestingly, ADAM8 is highly expressed on lung epithelial cells of asthmatic patients and upregulated in models of murine allergic airway inflammation [30–32]. Increased expression of ADAM33 in lung tissue has also been linked to the pathogenesis of asthma [33]. These findings imply that sCD23 production by ADAM8 and ADAM33 could be predominantly tissue-specific and restricted to the lung. As discussed later in this review in detail in Section 3.2, increased levels of sCD23 can foster IgE production as well as enhance production of inflammatory cytokines. As a result, sCD23 generated by ADAM8 or ADAM33 in the lung could be a local mediator that regulates asthma development or, more generally, the pathology of IgE-mediated asthmatic reactions. Additionally, smaller proteolytic cleavage products of sCD23 with molecular weights of 25–27 kDa are described. These isoforms of sCD23 are most likely extracellular degradation intermediates and can be found in cell culture supernatants [34,35], as well as in human serum of patients with chronic lymphocytic leukemia [36]. The house dust mite protease Der p1 has been shown to generate a 17 kDa sCD23 fragment [37,38]. It is speculated that, at least in part, the high allergenicity of the mites

is a result of the Der p1-mediated increase in serum sCD23 levels, which in turn modulates IgE production.

Recently, it was shown that membrane CD23 and ADAM10 can be found in exosomes of B cells [39]. The authors of this study suggest that membrane CD23 is internalized from the cell surface and traffics into endosomal compartments. Endosome-derived exosomes are then secreted and CD23 is cleaved extracellularly. Exosome-derived sCD23 could be an additional source of serum sCD23 in humans.

Production of sCD23 is dictated by the expression levels of transmembrane CD23, as well as the rate of proteolytic cleavage from the cell surface. Surface expression of CD23 is regulated by many cytokines including IL-4, IL-13, IL-5, IL-9, GM-CSF, IFN- γ , and CD40 (reviewed in [27]) and accordingly these immune mediators influence the generation of sCD23. Along this line, activation of murine splenic B cells and human peripheral blood B cells via lipopolysaccharides (LPS) induces release of sCD23. This LPS-induced increase in sCD23 production is mediated by both induction of de novo synthesis of CD23 and by enhanced CD23 cleavage by the matrix metalloproteinase MMP9 [40]. Supernatants from MMP9–/– splenocytes fail to induce sCD23 release from B cells and MMP9 expression itself is upregulated following LPS stimulation [40]. The cleavage sites and cleavage products of membrane CD23 by MMP9 are currently unknown. Recently, it was shown that glutamate, a major excitatory neurotransmitter in the central nervous system, also enhances sCD23 release by B cells. Activity of the glutamate-specific kainate receptor (KAR) mediates the glutamate-dependent release of sCD23 by increasing ADAM10 expression [41].

Membrane cleavage of sCD23 is primarily regulated via the accessibility of the cleavage sites for proteolysis. Ligation of membrane CD23 by IgE induces a conformational change that inhibits sCD23 shedding by ADAM10 [42]. Using a comparable mechanism, CD23-specific monoclonal antibodies (mAb) can either enhance or limit the release of sCD23 [43]. Binding of 19G5 mAb to murine sCD23 changes the topology of the α -helical coiled coil stalk region in favor of proteolytic cleavage [44]. Interestingly, the 19G5 mAb not only facilitates proteolytic cleavage of sCD23 from the cell surface but also induces internalization of CD23 into endosomal compartments [39]. This reagent could thus also induce production of sCD23 via the exosomal pathway as described earlier in the review. The anti-CD23 antibody Lumiliximab has been shown to decrease serum IgE, likely by inhibiting production of sCD23. It is speculated that Lumiliximab stabilizes surface CD23 by preventing its proteolytic cleavage [45].

2.3. Galectin-3

Galectin-3 is a secretory protein of ~30 kDa and belongs to a family of β -galactoside-binding animal lectins [46]. This protein was formerly named ϵ -binding protein because of its ability to interact with IgE as well as Fc ϵ RI [47]. In contrast to Fc ϵ RI and CD23, galectin-3 does not exist as a transmembrane protein. The structure of galectin-3 consists of a carbohydrate-recognition domain (CRD) linked to a non-lectin region of proline- and glycine-rich tandem repeats. Galectin-3 can form pentamers via the non-CRD domain which strongly resemble pentameric IgM; this is a unique structural feature of galectin-3 among the 15 members of the galectin family. Similar to all other galectin family members, galectin-3 lacks a classical signal for the secretory pathway. Therefore, the mechanism of galectin-3 secretion is currently poorly understood [46].

Galectin-3 resides in the cytosol or the nucleus, but has also been shown to associate with intracellular vesicles [48] and has been found in exosomes from dendritic cells [49]. Following secretion, galectin-3 is detected in the extracellular space from where it can attach to cell surfaces. IgE and Fc ϵ RI were among the first described binding partners at the cell surface of mast cells [50]. Interestingly,

differentially glycosylated IgE isoforms have been found that display distinct binding capacities for galectin-3 [51,52]. Additionally, galectin-3 appears to interact also with a large variety of other cell surface and extracellular matrix proteins. Several intracellular proteins were also described as galectin-3 binding partners (reviewed in detail in [46,48]). Because of the promiscuous binding pattern of galectin-3, it is important to note that the detection of this molecule at the cell surface does not indicate whether the cell type by itself produces the protein.

Galectin-3 has been found on the cell surface of eosinophils, neutrophils, mast cells, dendritic cells, macrophages, T cells, and B cells [46,47,53–57]. Macrophages have been shown to be a key source of extracellular soluble galectin-3. When monocytes differentiate into macrophages increased galectin-3 expression is observed [57]. Alveolar macrophages release galectin-3 into the alveolar space after infection with *Streptococcus pneumoniae* [58,59]. Furthermore, alternative macrophage activation with the Th2 response associated cytokines, IL-4 and IL-13, increase expression and release of galectin-3. In contrast, LPS or IFN- γ induced macrophage activation can inhibit galectin-3 release [60]. In line with the finding that Th2 cytokines regulate galectin-3 expression in macrophages, eosinophils from allergic donors show increased levels of galectin-3 [61]. Additionally, various fibrotic conditions in humans including liver cirrhosis and pulmonary fibrosis are characterized by increased levels of galectin-3 [62,63].

Interestingly, galectin-3 expression appears to be species specific and differences between humans and mice are described. Similar to Fc ϵ RI, galectin-3 was found on human but not murine eosinophils [64]. Galectin-3 might thus be another IgE Fc-receptors for which species-specific expression patterns might hamper interpretation of murine studies.

3. Functions of the soluble IgE receptors in the IgE network

3.1. sFc ϵ RI

Attempts to modulate IgE-mediated immune responses by recombinant Fc ϵ RI [65,66] well precede the description of sFc ϵ RI in human serum. Recombinant soluble forms of the extracellular domain of Fc ϵ RI alpha-chain that interact with the Fc-portion of IgE were described as inhibitors of cytokine release from human basophils [66] and RBL-2H3 release assays [65]. *In vivo* models of passive cutaneous anaphylaxis confirmed that such soluble IgE receptors indeed blunt IgE-mediated immune responses [65]. Omalizumab is an anti-IgE specific mAb that reacts with the Fc-portion of the immunoglobulin and is successfully used to clinically modulate serum IgE levels in several types of allergic diseases [67]. Based on its binding domains on IgE, a recombinant form of sFc ϵ RI could be used in a similar manner as Omalizumab to modulate IgE-mediated allergy without the disadvantages of antibody therapy.

In vivo, sFc ϵ RI has several potential binding partners in serum as well as on the surface of peripheral blood or tissue cells (Fig. 2). As a free IgE receptor, sFc ϵ RI can form a complex with IgE or form complexes of higher order that include antigen. Free sFc ϵ RI can additionally bind to membrane IgE (mIgE) expressed by B-cells. Based on a one-to-one ligand–receptor ratio, no crosslinking and activation of the B cell could occur. By blocking the Fc-region of IgE, free sFc ϵ RI could prevent IgE from interacting with other Fc ϵ RI receptors expressed on the cell surface [15]. On effector cells, such as basophils and mast cells, this would impair IgE-mediated degranulation and release of cytotoxic mediators. Thus, sFc ϵ RI has the potential to blunt the acute phase of an allergic response. If sFc ϵ RI–IgE–antigen complexes of higher order interact with surface-expressed Fc ϵ RI, however, such complexes could activate cells via Fc ϵ RI-crosslinking, provided the complexes contain

free IgE Fc-domains. Via such an activation pathway, sFc ϵ RI could exacerbate immediate type allergic responses. On dendritic cells, IgE-mediated antigen presentation might be down-regulated when free sFc ϵ RI blocks IgE from binding to surface expressed receptors, which are used for antigen-sampling. Consequently, the sensitization phase towards allergens as well as the Th2-type immune responses of chronic allergic reactions might be modified [9]. If sFc ϵ RI is internalized as part of a sFc ϵ RI–IgE–antigen complex by antigen presenting cells, the endogenous alpha-chain could be presented as an exogenous allergen. Such a process might provide a mechanistic explanation for how autoantibodies against Fc ϵ RI-alpha are generated [68,69]. Finally, it is conceivable that as of yet unidentified binding partners for sFc ϵ RI exist in human serum.

3.2. sCD23

IgE-mediated ligation of membrane CD23 inhibits IgE production in B cells via a negative feedback loop [1]. In contrast, sCD23 increases the production of IgE through co-ligation of CD21 and mIgE [70,71]. The size of clusters induced by sCD23-mediated crosslinking at the cell surface defines the strength of the IgE-inducing signal. Since the oligomerization capacity of sCD23 depends on the length of its stalk region, not all forms of sCD23 are equally potent at inducing IgE-production [72]. In this context, McCloskey et al. have reported that short recombinant sCD23 fragments corresponding to short sCD23 forms generated by Der p1 can form only small complexes, which were even inhibitory for IL-4 induced synthesis of IgE in their experimental settings [73]. This finding argues against the assumption that the high allergenicity of the house dust mite derives from increased production of sCD23 through Der p1. In addition to regulating IgE synthesis, sCD23 has been shown to promote B cell differentiation, survival of germinal center B cells as well as differentiation towards B cell blasts *in vitro* in the presence of IL-1 α [74–76]. Aside from acting on the B cell compartment, combinations of sCD23 and IL-1 α also promote proliferation of human myeloid bone marrow precursors, differentiation of thymic T cell precursors, and CD4 T cell responses [77–79].

sCD23 also magnifies allergic diseases by enhancing the production of inflammatory cytokines. On human monocytes, binding of sCD23 to CD11b/CD18 and CD11c/CD18 has been reported to activate nitric oxide synthase and to induce the production of inflammatory cytokines, including TNF- α , IL-6, IL-8, MIP1- α , and MIP1- β , as well as IL-1 β [80–83]. Similarly, murine monocytes and macrophages produce IL-6 after incubation with recombinant sCD23 [84]. Activation of vitronectin receptors by sCD23 is an additional pathway for the induction of pro-inflammatory cytokines [85]. Peripheral blood mononuclear cells of patients with hyper-IgE syndrome are particularly reactive to treatment with recombinant sCD23, producing high levels of IL-1 β and TNF- α [86].

3.3. Galectin-3

Galectin-3 is a versatile player of the immune system [46,87,88]. Secreted galectin-3 can activate cells directly by binding to cell surface receptors. Alternatively, galectin-3 is endocytosed, permitting it to modify intracellular signaling pathways. The large number of intracellular as well as extracellular galectin-3 binding partners, therefore, allows this protein to play a role in a large variety of inflammatory responses, including neutrophil activation, chemotraction of monocytes and macrophages, adhesion and migration of neutrophils and dendritic cells as well as regulation of apoptosis in immune cells [46,89–91]. Recently it was found that galectin-3 even displays antimicrobial functions against the fungus *Candida albicans* [92]. In the context of allergic reactions, it is important to note that extracellular galectin-3 is a potent activator of mast




Fig. 2. Possible interaction partners of soluble FcεRI (sFcεRI) *in vivo*. (A) FcεRI exists as a membrane bound as well as a soluble isoform in human blood. In serum, sFcεRI is found as a true soluble form or as a sFcεRI-IgE complex when bound to its natural ligand. Potentially, sFcεRI-IgE complexes can interact with antigens and form immune complexes of higher order. Free sFcεRI can additionally interact with membrane IgE expressed on B cells. sFcεRI-IgE complexes cannot bind to trimeric or tetrameric FcεRI expressed on the cell surface of peripheral blood cells, because the binding site of IgE and the cellular receptor is blocked by sFcεRI in solution. (B) In tissue, all interactions described for peripheral blood in (A) are theoretically possible. It remains to be defined whether the local concentration of sFcεRI in tissue is comparable to sFcεRI serum levels. Cells migrating from the periphery blood could also serve as vehicles to transport sFcεRI into tissue.

cells via crosslinking of IgE-loaded FcεRI or via crosslinking FcεRI directly in an IgE-independent manner. Galectin-3 was shown to induce inflammatory mediator release from IgE-sensitized as well as non-sensitized mast cells and human eosinophils [47,55]. Since membrane FcεRI is a binding partner, it is likely that galectin-3 can also interact with sFcεRI in serum. Such an interaction could potentially modulate the functions of sFcεRI *in vivo*.

4. Soluble IgE receptors as disease biomarkers

As detailed below, the soluble IgE receptors CD23 and galectin-3 are upregulated in a variety of diseases and have made their way into clinical practice as diagnostic biomarkers.

We speculate here that sFcεRI might also prove itself as a biomarker, although it should be noted that solid evidence for this role is so far lacking.

4.1. sFcεRI-a potential new biomarker?

Development of a standardized assay that allows for the comparative analysis of serum levels of sFcεRI in health and disease will facilitate an understanding of the clinical relevance of sFcεRI in serum. Currently, it is known only that serum levels of sFcεRI correlate with serum IgE in a pediatric patient cohort with elevated IgE [88,93]. Interestingly, sFcεRI was also found in serum of individuals with normal IgE levels. In fact, the highest levels of sFcεRI were actually described in this subpopulation [93]. More detailed

studies investigating the possible link between serum sFcεRI to clinical symptoms of allergy are urgently needed.

4.2. sCD23 as a biomarker

sCD23 levels appear to be upregulated in a plethora of diseases. Less frequently, a decrease of serum sCD23 has been described (Table 3). Elevated sCD23 levels have been found in association with allergic diseases, including asthma and atopic dermatitis [94–97]. There is some debate among allergists, however, as to whether sCD23 serum levels have a predictive value for the diagnosis of allergic reactions [98]. Clearly, further investigations are needed.

Chronic lymphocytic leukemia (CLL) is currently the only disease in which sCD23 is used as a clinical biomarker. CLL patients show significantly higher levels of serum sCD23 when compared to healthy controls [99,100]. Plasma levels of sCD23 correlate with disease outcome; high sCD23 concentrations indicate a more severe disease stage, a more rapid median progression time and shorter median survival time [99,101–105]. Other parameters of CLL which correspond to a negative prognosis, such as a short doubling time of lymphocyte and diffuse bone marrow histology, also correlate with increased levels of sCD23 [99,105].

In AIDS patients, sCD23 appears to be a predictive biomarker for the development of non-Hodgkin lymphoma. sCD23 is specifically elevated in HIV patients prior to the diagnosis of AIDS-associated non-Hodgkin lymphoma [106–108]. Recently, elevated serum levels of sCD23 were described in patients with pancreatic cancer

Table 3

Serum levels of sCD23 and galectin-3 are modulated in various pathologies.

Pathology	sCD23	Galectin-3
Cancer	Chronic lymphocytic leukemia ↑ ^a Pancreatic cancer ↑ AIDS-associated non-Hodgkin lymphoma ↑ Glioma ↓ ^b	Thyroid cancer ↑ [125–127] ^c Melanoma ↑ [128–130] Colorectal cancer ↑ [128] Head and neck squamous cell carcinomas ↑ [131] Bladder cancer ↑ [132] Breast and ovarian cancer ↑ [128] Non-Hodgkin lymphoma ↑ [128] Rheumatoid arthritis ↑ [136,137] Systemic lupus erythematosus ↑ [137] Behçet's disease ↑ [137] Inflammatory bowel disease ↑ [138]
Auto-immune diseases	Arthritis ↑ and ↓ Systemic lupus erythematosus ↑ [133] Primary Sjögren's disease ↑ [133] Autoimmune thyroiditis ↑ [134] Myasthenia gravis ↑ [134,135] Crohn's disease ↓ [134]	
Allergic diseases	Asthma ↑ Atopic dermatitis ↑	
Other		Chronic heart failure ↑ [123,139–141] Liver fibrosis ↑ Pulmonary fibrosis ↑

^a ↑ serum levels are found to be upregulated.^b ↓ serum levels are found to be downregulated.^c References are given only if not cited in the text.

[109], but additional studies are required to evaluate whether sCD23 can function as a biomarker in this disease. Several reports describe elevated sCD23 levels in serum and synovial fluid of patients with rheumatoid arthritis [110–117]. In contrast, Singh et al. showed recently that sCD23 levels were slightly decreased in patients with juvenile arthritis [118]. More extensive studies are needed to clarify this contradiction.

Low serum sCD23 levels are described for glioma patients [119]. The reason for this decrease of serum sCD23 is unknown as of yet. Glioma patients commonly do not have elevated serum IgE and suffer less often from allergies. Conversely, the survival time of glioma patients is prolonged if they show elevated serum IgE [120]. In these studies, the authors speculate that elevated IgE levels correlated with more effective anti-tumor response or less aggressive tumors. A study by Merrill et al. found that patients with allergy actually have a decreased risk for glioma [121]. More research is needed to understand how and if sCD23 and IgE have protective functions in this type of cancer. This topic is a focus of attention in the nascent field of allergooncology [122].

4.3. Galectin-3

Elevated serum levels of galectin-3 are found in many diseases including various types of autoimmune diseases and cancer (Table 3). So far, the only FDA approved application of galectin-3 as a biomarker is for the prognosis of chronic heart failure [123]. However, galectin-3 has recently emerged also as a promising diagnostic marker for thyroid cancers [124].

5. Summary and perspective

Soluble IgE receptors in human serum are a diverse group of proteins with the unifying characteristic of interacting with IgE. We here discussed the production and physiological roles of these

Table 4

Major physiological roles of transmembrane and soluble IgE Fc receptors.

IgE receptor	Transmembrane form	Soluble form
FcεRI	Mast cells, basophils: release of inflammatory mediators Dendritic cells: antigen uptake receptor	Not defined
CD23	Regulation of IgE synthesis Transport of IgE at mucosal surfaces	Promotion of IgE synthesis Induction of inflammatory cytokine production from monocytes and macrophages Promotion of T and B cell differentiation as well as B cell survival
Galectin-3 ^a	Induction of inflammatory mediator release from mast cells via binding of FcεRI and IgE Promotion of adhesion, migration and respiratory burst of neutrophils Chemoattraction of monocytes and macrophages Promotion of Th2 responses and alternative macrophage activation Regulation of apoptosis in immune cells Anti-microbial functions against <i>Candida albicans</i> Activation and growth induction of tissue fibroblasts	

^a No transmembrane form of galectin-3 exists. Next to the secreted pool, galectin-3 is found intracellularly in a non-secreted form.

receptors and compared them to their parental transmembranous receptor isoforms (Table 4). Further studies on how the generation of soluble IgE receptors is regulated, how these proteins function within the human IgE network and how they are connected to various disease pathologies will close large gaps in our understanding of IgE-mediated immune responses. In addition, a better understanding of the patho-physiology of soluble IgE receptors might point us towards novel intervention strategies for IgE-mediated allergies.

Acknowledgements

We apologize to colleagues whose work was not cited in this review due to space limitations and thank Bonny Dickinson and Michael Pardo for critically reading this manuscript. This work was supported by the Gerber Foundation and the National Institutes of Health grant AI075037 (both to E.F.).

References

- [1] Gould HJ, Sutton BJ. IgE in allergy and asthma today. *Nat Rev Immunol* 2008;8:205–17.
- [2] Kraft S, Kinet JP. New developments in FcεRI regulation, function, and inhibition. *Nat Rev Immunol* 2007;7:365–78.
- [3] Fiebigler E, Tortorella D, Jouvin MH, Kinet JP, Ploegh HL. Cotranslational endoplasmic reticulum assembly of FcεRI controls the formation of functional IgE-binding receptors. *J Exp Med* 2005;201:267–77.
- [4] Stone KD, Prussin C, Metcalfe DD. IgE, mast cells, basophils, and eosinophils. *J Allergy Clin Immunol* 2010;125:S73–80.
- [5] Maurer D, Fiebigler E, Ebner C, Reininger B, Fischer GF, Wichlas S, et al. Peripheral blood dendritic cells express FcεRI as a complex composed of FcεRIα- and FcεRIγ-chains and can use this receptor for IgE-mediated allergen presentation. *J Immunol* 1996;157:607–13.
- [6] Bieber T, de la Salle H, Wollenberg A, Hakimi J, Chizzonite R, Ring J, et al. Human epidermal Langerhans cells express the high affinity receptor for immunoglobulin E (FcεRI). *J Exp Med* 1992;175:1285–90.
- [7] Grayson MH, Cheung D, Rohlfing MM, Kitchens R, Spiegel DE, Tucker J, et al. Induction of high-affinity IgE receptor on lung dendritic cells during viral infection leads to mucous cell metaplasia. *J Exp Med* 2007;204:2759–69.
- [8] Hammad H, Plantinga M, Deswarte K, Pouliot P, Willart MA, Kool M, et al. Inflammatory dendritic cells—not basophils—are necessary and sufficient for induction of Th2 immunity to inhaled house dust mite allergen. *J Exp Med* 2010;207:2097–111.
- [9] Sallmann E, Reininger B, Brandt S, Duschek N, Hoflehner E, Garner-Spitzer E, et al. High-affinity IgE receptors on dendritic cells exacerbate Th2-dependent inflammation. *J Immunol* 2011;187:164–71.

- [10] Maurer D, Fiebiger E, Reininger B, Wolff-Winiski B, Jouvin M-H, Kilgus O, et al. Expression of functional high affinity immunoglobulin E receptors (FcεRI) on monocytes of atopic individuals. *J Exp Med* 1994;179:745–50.
- [11] Gounni AS, Lamkhioe B, Ochiai K, Tanaka Y, Delaporte E, Capron A, et al. High-affinity IgE receptor on eosinophils is involved in defence against parasites. *Nature* 1994;367:183–6.
- [12] Kayaba H, Dombrowicz D, Woerly G, Papin JP, Loiseau S, Capron M. Human eosinophils and human high affinity IgE receptor transgenic mouse eosinophils express low levels of high affinity IgE receptor, but release IL-10 upon receptor activation. *J Immunol* 2001;167:995–1003.
- [13] Joseph M, Gounni AS, Kusnier JP, Vorng H, Sarfati M, Kinet JP, et al. Expression and functions of the high-affinity IgE receptor on human platelets and megakaryocyte precursors. *Eur J Immunol* 1997;27:2212–8.
- [14] Untersmayr E, Bises G, Starkl P, Bevins CL, Scheiner O, Boltz-Nitulescu G, et al. The high affinity IgE receptor FcεRI is expressed by human intestinal epithelial cells. *PLoS ONE* 2010;5:e9023.
- [15] Dehlink E, Platzer B, Baker AH, Larosa J, Pardo M, Dwyer P, et al. A soluble form of the high affinity IgE receptor, FcεRI, circulates in human serum. *PLoS ONE* 2011;6:e19098.
- [16] Platzer B, Fiebiger E. The signal peptide of the IgE receptor alpha-chain prevents surface expression of an immunoreceptor tyrosine-based activation motif-free receptor pool. *J Biol Chem* 2010;285:15314–23.
- [17] Kinet JP. The high-affinity IgE receptor (FcεRI): from physiology to pathology. *Annu Rev Immunol* 1999;17:931–72.
- [18] Ortega E, Schweitzer-Stenner R, Pecht I. Kinetics of ligand binding to the type 1 Fcε receptor on mast cells. *Biochemistry* 1991;30:3473–83.
- [19] Garman SC, Kinet JP, Jardetzky TS. Crystal structure of the human high-affinity IgE receptor. *Cell* 1998;95:951–61.
- [20] Paul P, van den Hoorn T, Jongsma ML, Bakker MJ, Hengeveld R, Janssen L, et al. A genome-wide multidimensional RNA screen reveals pathways controlling MHC class II antigen presentation. *Cell* 2011;145:268–83.
- [21] Dehlink E, Fiebiger E. The role of the high-affinity IgE receptor, FcεRI, in eosinophilic gastrointestinal diseases. *Immunol Allergy Clin North Am* 2009;29:159–70, xii.
- [22] Acharya M, Borland G, Edkins AL, Maclellan LM, Matheson J, Ozanne BW, et al. CD23/FcεRII: molecular multi-tasking. *Clin Exp Immunol* 2010;162:12–23.
- [23] Lee BW, Simmons Jr CF, Wileman T, Geha RS. Intracellular cleavage of newly synthesized low affinity Fcε receptor (FcεR₂) provides a second pathway for the generation of the 28-kDa soluble FcεR₂ fragment. *J Immunol* 1989;142:1614–20.
- [24] Gibb DR, El Shikh M, Kang DJ, Rowe WJ, El Sayed R, Cichy J, et al. ADAM10 is essential for Notch2-dependent marginal zone B cell development and CD23 cleavage *in vivo*. *J Exp Med* 2010;207:623–35.
- [25] Weskamp G, Ford JW, Sturgill J, Martin S, Docherty AJ, Swendeman S, et al. ADAM10 is a principal 'shedase' of the low-affinity immunoglobulin E receptor CD23. *Nat Immunol* 2006;7:1293–8.
- [26] Fourie AM, Coles F, Moreno V, Karlsson L. Catalytic activity of ADAM8, ADAM15, and MDC-1 (ADAM28) on synthetic peptide substrates and in ectodomain cleavage of CD23. *J Biol Chem* 2003;278:30469–77.
- [27] Rosenwasser LJ, Meng J. Anti-CD23. *Clin Rev Allergy Immunol* 2005;29:61–72.
- [28] Ritchens J, Fairclough L, Ghaemmaghami AM, Mahdavi J, Shakib F, Sewell HF. The detection of ADAM8 protein on cells of the human immune system and the demonstration of its expression on peripheral blood B cells, dendritic cells, and monocyte subsets. *Immunobiology* 2007;212:29–38.
- [29] Umland SP, Garlisi CG, Shah H, Wan Y, Zou J, Devito KE, et al. Human ADAM33 messenger RNA expression profile and post-transcriptional regulation. *Am J Respir Cell Mol Biol* 2003;29:571–82.
- [30] Foley SC, Mogas AK, Olivenstein R, Fiset PO, Chakir J, Bourbeau J, et al. Increased expression of ADAM33 and ADAM8 with disease progression in asthma. *J Allergy Clin Immunol* 2007;119:863–71.
- [31] Knolle MD, Owen CA. ADAM8: a new therapeutic target for asthma. *Expert Opin Ther Targets* 2009;13:523–40.
- [32] Van Eerdewegh P, Little RD, Dupuis J, Del Mastro RG, Falls K, Simon J, et al. Association of the ADAM33 gene with asthma and bronchial hyperresponsiveness. *Nature* 2002;418:426–30.
- [33] Haitchi HM, Powell RM, Shaw TJ, Howarth PH, Wilson SJ, Wilson DI, et al. ADAM33 expression in asthmatic airways and human embryonic lungs. *Am J Respir Crit Care Med* 2005;171:958–65.
- [34] Letellier M, Nakajima T, Pulido-Cejudo G, Hofstetter H, Delespesse G. Mechanism of formation of human IgE-binding factors (soluble CD23): III. Evidence for a receptor (FcεRII)-associated proteolytic activity. *J Exp Med* 1990;172:693–700.
- [35] Letellier M, Sarfati M, Delespesse G. Mechanisms of formation of IgE-binding factors (soluble CD23)—I. FcεRII bearing B cells generate IgE-binding factors of different molecular weights. *Mol Immunol* 1989;26:1105–12.
- [36] Sarfati M, Bron D, Lagneaux L, Fonteyn C, Frost H, Delespesse G. Elevation of IgE-binding factors in serum of patients with B cell-derived chronic lymphocytic leukemia. *Blood* 1988;71:94–8.
- [37] Schulz O, Laing P, Sewell HF, Shakib F. Der p 1, a major allergen of the house dust mite, proteolytically cleaves the low-affinity receptor for human IgE (CD23). *Eur J Immunol* 1995;25:3191–4.
- [38] Schulz O, Sutton BJ, Beavil RL, Shi J, Sewell HF, Gould HJ, et al. Cleavage of the low-affinity receptor for human IgE (CD23) by a mite cysteine protease: nature of the cleaved fragment in relation to the structure and function of CD23. *Eur J Immunol* 1997;27:584–8.
- [39] Mathews JA, Gibb DR, Chen BH, Scherle P, Conrad DH, Sheddase A. CD23 disintegrin and metalloproteinase 10 (ADAM10) is also required for CD23 sorting into B cell-derived exosomes. *J Biol Chem* 2010;285:37531–41.
- [40] Jackson L, Cady CT, Cambier JC. TLR4-mediated signaling induces MMP9-dependent cleavage of B cell surface CD23. *J Immunol* 2009;183:2585–92.
- [41] Sturgill JL, Mathews J, Scherle P, Conrad DH. Glutamate signaling through the kainate receptor enhances human immunoglobulin production. *J Neuroimmunol* 2011;233:80–9.
- [42] Conrad DH, Ford JW, Sturgill JL, Gibb DR. CD23: an overlooked regulator of allergic disease. *Curr Allergy Asthma Rep* 2007;7:331–7.
- [43] Kilmon MA, Shelburne AE, Chan-Li Y, Holmes KL, Conrad DH. CD23 trimers are preassociated on the cell surface even in the absence of its ligand, IgE. *J Immunol* 2004;172:1065–73.
- [44] Kilmon MA, Ghirlando R, Strub MP, Beavil RL, Gould HJ, Conrad DH. Regulation of IgE production requires oligomerization of CD23. *J Immunol* 2001;167:3139–45.
- [45] Rosenwasser LJ, Busse WW, Lizambri RG, Olejnik TA, Totoritis MC. Allergic asthma and an anti-CD23 mAb (IDEC-152): results of a phase I, single-dose, dose-escalating clinical trial. *J Allergy Clin Immunol* 2003;112:563–70.
- [46] Liu FT, Rabinovich GA. Galectins: regulators of acute and chronic inflammation. *Ann N Y Acad Sci* 2010;1183:158–82.
- [47] Frigeri LG, Zuberi RI, Liu FT. εBP, a beta-galactoside-binding animal lectin, recognizes IgE receptor (FcεRI) and activates mast cells. *Biochemistry* 1993;32:7644–9.
- [48] Dumic J, Dabelic S, Flogel M. Galectin-3: an open-ended story. *Biochim Biophys Acta* 2006;1760:616–35.
- [49] Thery C, Bousset M, Veron P, Ricciardi-Castagnoli P, Raposo G, Garin J, et al. Proteomic analysis of dendritic cell-derived exosomes: a secreted subcellular compartment distinct from apoptotic vesicles. *J Immunol* 2001;166:7309–18.
- [50] Liu FT, Frigeri LG, Gritzmacher CA, Hsu DK, Robertson MW, Zuberi RI. Expression and function of an IgE-binding animal lectin (epsilon BP) in mast cells. *Immunopharmacology* 1993;26:187–95.
- [51] Robertson MW, Albrandt K, Keller D, Liu FT. Human IgE-binding protein: a soluble lectin exhibiting a highly conserved interspecies sequence and differential recognition of IgE glycoforms. *Biochemistry* 1990;29:8093–100.
- [52] Robertson MW, Liu FT. Heterogeneous IgE glycoforms characterized by differential recognition of an endogenous lectin (IgE-binding protein). *J Immunol* 1991;147:3024–30.
- [53] Truong MJ, Gruart V, Kusnier JP, Papin JP, Loiseau S, Capron A, et al. Human neutrophils express immunoglobulin E (IgE)-binding proteins (Mac-2/εBP) of the S-type lectin family: role in IgE-dependent activation. *J Exp Med* 1993;177:243–8.
- [54] Truong MJ, Gruart V, Liu FT, Prin L, Capron A, Capron M. IgE-binding molecules (Mac-2/εBP) expressed by human eosinophils. Implication in IgE-dependent eosinophil cytotoxicity. *Eur J Immunol* 1993;23:3230–5.
- [55] Frigeri LG, Liu FT. Surface expression of functional IgE binding protein, an endogenous lectin, on mast cells and macrophages. *J Immunol* 1992;148:861–7.
- [56] Wollenberg A, de la Salle H, Hanau D, Liu FT, Bieber T. Human keratinocytes release the endogenous beta-galactoside-binding soluble lectin immunoglobulin E (IgE-binding protein) which binds to Langerhans cells where it modulates their binding capacity for IgE glycoforms. *J Exp Med* 1993;178:777–85.
- [57] Liu FT, Hsu DK, Zuberi RI, Kuwabara I, Chi EY, Henderson Jr WR. Expression and function of galectin-3, a beta-galactoside-binding lectin, in human monocytes and macrophages. *Am J Pathol* 1995;147:1016–28.
- [58] Sato S, Hughes RC. Regulation of secretion and surface expression of Mac-2, a galactoside-binding protein of macrophages. *J Biol Chem* 1994;269:4424–30.
- [59] Sato S, Ouellet N, Pelletier I, Simard M, Rancourt A, Bergeron MG. Role of galectin-3 as an adhesion molecule for neutrophil extravasation during streptococcal pneumonia. *J Immunol* 2002;168:1813–22.
- [60] MacKinnon AC, Farnworth SL, Hodgkinson PS, Henderson NC, Atkinson KM, Leffler H, et al. Regulation of alternative macrophage activation by galectin-3. *J Immunol* 2008;180:2650–8.
- [61] Rao SP, Wang Z, Zuberi RI, Sikora L, Bahaie NS, Zuraw BL, et al. Galectin-3 functions as an adhesion molecule to support eosinophil rolling and adhesion under conditions of flow. *J Immunol* 2007;179:7800–7.
- [62] Henderson NC, MacKinnon AC, Farnworth SL, Poirier F, Russo FP, Iredale JP, et al. Galectin-3 regulates myofibroblast activation and hepatic fibrosis. *Proc Natl Acad Sci U S A* 2006;103:5060–5.
- [63] Nishi Y, Sano H, Kawashima T, Okada T, Kikkawa K, et al. Role of galectin-3 in human pulmonary fibrosis. *Allergol Int* 2007;56:57–65.
- [64] de Andres B, Rakasz E, Hagen M, McCormik ML, Mueller AL, Elliot D, et al. Lack of Fcε-epsilon receptors on murine eosinophils: implications for the functional significance of elevated IgE and eosinophils in parasitic infections. *Blood* 1997;89:3826–36.
- [65] Ra C, Kuromitsu S, Hirose T, Yasuda S, Furuichi K, Okumura K. Soluble human high-affinity receptor for IgE abrogates the IgE-mediated allergic reaction. *Int Immunol* 1993;5:47–54.
- [66] Vangelista L, Laffer S, Turek R, Gronlund H, Sperr WR, Valent P, et al. The immunoglobulin-like modules Cε3 and α2 are the minimal units necessary for human IgE-FcεRI interaction. *J Clin Invest* 1999;103:1571–8.
- [67] Kopp MV. Omalizumab: anti-IgE therapy in allergy. *Curr Allergy Asthma Rep* 2011;11:101–6.

- [68] Fiebigier E, Hammerschmid F, Stingl G, Maurer D. Anti-FcεRIα autoantibodies in autoimmune-mediated disorders. Identification of a structure–function relationship. *J Clin Invest* 1998;101:243–51.
- [69] Fiebigier E, Stingl G, Maurer D. Anti-IgE and anti-FcεRI autoantibodies in clinical allergy. *Curr Opin Immunol* 1996;8:784–9.
- [70] Bonnefoy JY, Gauchat JF, Life P, Graber P, Aubry JP, Lecoanet-Henchoz S. Regulation of IgE synthesis by CD23/CD21 interaction. *Int Arch Allergy Immunol* 1995;107:40–2.
- [71] Hibbert RG, Teriete P, Grundy GJ, Beavil RL, Reljic R, Holers VM, et al. The structure of human CD23 and its interactions with IgE and CD21. *J Exp Med* 2005;202:751–60.
- [72] Beavil RL, Graber P, Aubonne N, Bonnefoy JY, Gould HJ. CD23/Fc epsilon RII and its soluble fragments can form oligomers on the cell surface and in solution. *Immunology* 1995;84:202–6.
- [73] McCloskey N, Hunt J, Beavil RL, Juttan MR, Grundy GJ, Girardi E, et al. Soluble CD23 monomers inhibit and oligomers stimulate IGE synthesis in human B cells. *J Biol Chem* 2007;282:24083–91.
- [74] Kagan JM, Champlin RE, Saxon A. B-cell dysfunction following human bone marrow transplantation: functional–phenotypic dissociation in the early posttransplant period. *Blood* 1989;74:777–85.
- [75] Liu YJ, Mason DY, Johnson GD, Abbot S, Gregory CD, Hardie DL, et al. Germinal center cells express bcl-2 protein after activation by signals which prevent their entry into apoptosis. *Eur J Immunol* 1991;21:1905–10.
- [76] Liu YJ, Cairns JA, Holder MJ, Abbot SD, Jansen KU, Bonnefoy JY, et al. Recombinant 25-kDa CD23 and interleukin 1 alpha promote the survival of germinal center B cells: evidence for bifurcation in the development of centrocytes rescued from apoptosis. *Eur J Immunol* 1991;21:1107–14.
- [77] Mossalayi MD, Merle-Beral H, Dalloul AH, Arock M, Michel A, Hofstetter H, et al. Inhibition of interleukin-3-dependent growth of CD34⁺ acute myelogenous leukemia cells by recombinant soluble CD23. *Ann N Y Acad Sci* 1991;628:362–7.
- [78] Mossalayi MD, Arock M, Bertho JM, Blanc C, Dalloul AH, Hofstetter H, et al. Proliferation of early human myeloid precursors induced by interleukin-1 and recombinant soluble CD23. *Blood* 1990;75:1924–7.
- [79] Mossalayi MD, Lecron JC, Dalloul AH, Sarfati M, Bertho JM, Hofstetter H, et al. Soluble CD23 (FcεRII) and interleukin 1 synergistically induce early human thymocyte maturation. *J Exp Med* 1990;171:959–64.
- [80] Aubry JP, Dugas N, Lecoanet-Henchoz S, Ouaz F, Zhao H, Delfraissy JF, et al. The 25-kDa soluble CD23 activates type III constitutive nitric oxide-synthase activity via CD11b and CD11c expressed by human monocytes. *J Immunol* 1997;159:614–22.
- [81] Lecoanet-Henchoz S, Gauchat JF, Aubry JP, Graber P, Life P, Paul-Eugene N, et al. CD23 regulates monocyte activation through a novel interaction with the adhesion molecules CD11b-CD18 and CD11c-CD18. *Immunity* 1995;3:119–25.
- [82] Rezzonico R, Chicheportiche R, Imbert V, Dayer JM. Engagement of CD11b and CD11c beta2 integrin by antibodies or soluble CD23 induces IL-1beta production on primary human monocytes through mitogen-activated protein kinase-dependent pathways. *Blood* 2000;95:3868–77.
- [83] Rezzonico R, Imbert V, Chicheportiche R, Dayer JM. Ligation of CD11b and CD11c beta(2) integrins by antibodies or soluble CD23 induces macrophage inflammatory protein 1alpha (MIP-1alpha) and MIP-1beta production in primary human monocytes through a pathway dependent on nuclear factor-kappaB. *Blood* 2001;97:2932–40.
- [84] Lecoanet-Henchoz S, Plater-Zyberk C, Graber P, Gretener D, Aubry JP, Conrad DH, et al. Mouse CD23 regulates monocyte activation through an interaction with the adhesion molecule CD11b/CD18. *Eur J Immunol* 1997;27:2290–4.
- [85] Hermann P, Armant M, Brown E, Rubio M, Ishihara H, Ulrich D, et al. The vitronectin receptor and its associated CD47 molecule mediates proinflammatory cytokine synthesis in human monocytes by interaction with soluble CD23. *J Cell Biol* 1999;144:767–75.
- [86] Daniels BB, Askew SL, van de Venter M, Oosthuizen V. Production of biologically active recombinant human soluble CD23 and its effect on PBMCs isolated from hyper-IgE blood. *Cell Immunol* 2005;234:146–53.
- [87] Chen HY, Liu FT, Yang RY. Roles of galectin-3 in immune responses. *Arch Immunol Ther Exp (Warsz)* 2005;53:497–504.
- [88] Henderson NC, Sethi T. The regulation of inflammation by galectin-3. *Immunol Rev* 2009;230:160–71.
- [89] Farnworth SL, Henderson NC, Mackinnon AC, Atkinson KM, Wilkinson T, Dhaliwal K, et al. Galectin-3 reduces the severity of pneumococcal pneumonia by augmenting neutrophil function. *Am J Pathol* 2008;172:395–405.
- [90] Karlsson A, Follin P, Leffler H, Dahlgren C. Galectin-3 activates the NADPH-oxidase in exudated but not peripheral blood neutrophils. *Blood* 1998;91:3430–8.
- [91] Hsu DK, Chernyavsky AI, Chen HY, Yu L, Grando SA, Liu FT. Endogenous galectin-3 is localized in membrane lipid rafts and regulates migration of dendritic cells. *J Invest Dermatol* 2009;129:573–83.
- [92] Kohatsu L, Hsu DK, Jegalian AG, Liu FT, Baum LG. Galectin-3 induces death of *Candida* species expressing specific beta-1,2-linked mannans. *J Immunol* 2006;177:4718–26.
- [93] Dehlink E, Baker AH, Yen E, Nurko S, Fiebigier E. Relationships between levels of serum IgE, cell-bound IgE, and IgE-receptors on peripheral blood cells in a pediatric population. *PLoS ONE* 2010;5:e12204.
- [94] Di Lorenzo G, Drago A, Pellitteri ME, Candore G, Colombo A, Potestio M, et al. Serum levels of soluble CD23 in patients with asthma or rhinitis monosensitive to parietaria. Its relation to total serum IgE levels and eosinophil cationic protein during and out of the pollen season. *Allergy Asthma Proc* 1999;20:119–25.
- [95] Rogala B, Rymarczyk B. Soluble CD23 in allergic diseases. *Arch Immunol Ther Exp (Warsz)* 1999;47:251–5.
- [96] Takigawa M, Tamamori T, Horiguchi D, Sakamoto T, Yamada M, Yoshioka A, et al. Fcε receptor II/CD23-positive lymphocytes in atopic dermatitis. I. The proportion of FcεRII + lymphocytes correlates with the extent of skin lesion. *Clin Exp Immunol* 1991;84:275–82.
- [97] Tanaka A, Ohashi Y, Nakai Y. Decrease of serum levels of soluble CD23 during immunotherapy in patients with perennial allergic rhinitis. *Ann Otol Rhinol Laryngol* 1999;108:193–200.
- [98] Ott H, Wilke J, Baron JM, Hoger PH, Folster-Holst R. Soluble immune receptor serum levels are associated with age, but not with clinical phenotype or disease severity in childhood atopic dermatitis. *J Eur Acad Dermatol Venereol* 2010;24:395–402.
- [99] Saka B, Aktan M, Sami U, Oner D, Sanem O, Dincol G. Prognostic importance of soluble CD23 in B-cell chronic lymphocytic leukemia. *Clin Lab Haematol* 2006;28:30–5.
- [100] Sarfati M, Delespesse G. Possible role of human lymphocyte receptor for IgE (CD23) or its soluble fragments in the *in vitro* synthesis of human IgE. *J Immunol* 1988;141:2195–9.
- [101] Sarfati M, Chevret S, Chastang C, Biron G, Stryckmans P, Delespesse G, et al. Prognostic importance of serum soluble CD23 level in chronic lymphocytic leukemia. *Blood* 1996;88:4259–64.
- [102] Reinisch W, Willheim M, Hilgarth M, Gasche C, Mader R, Szeplafalusi S, et al. Soluble CD23 reliably reflects disease activity in B-cell chronic lymphocytic leukemia. *J Clin Oncol* 1994;12:2146–52.
- [103] Schwarzmeyer JD, Shehata M, Hilgarth M, Marschitz I, Louda N, Hubmann R, et al. The role of soluble CD23 in distinguishing stable and progressive forms of B-chronic lymphocytic leukemia. *Leuk Lymphoma* 2002;43:549–54.
- [104] Del Principe MI, Del Poeta G, Buccisano F, Maurillo L, Venditti A, Zucchetto A, et al. Clinical significance of ZAP-70 protein expression in B-cell chronic lymphocytic leukemia. *Blood* 2006;108:853–61.
- [105] Meuleman N, Stamatopoulos B, Dejenefte M, El Housni H, Lagneaux L, Bron D. Doubling time of soluble CD23: a powerful prognostic factor for newly diagnosed and untreated stage A chronic lymphocytic leukemia patients. *Leukemia* 2008;22:1882–90.
- [106] Yawetz S, Cumberland WG, van der Meyden M, Martinez-Maza O. Elevated serum levels of soluble CD23 (sCD23) precede the appearance of acquired immunodeficiency syndrome-associated non-Hodgkin's lymphoma. *Blood* 1995;85:1843–9.
- [107] Schroeder JR, Saah AJ, Hoover DR, Margolick JB, Ambinder RF, Martinez-Maza O, et al. Serum soluble CD23 level correlates with subsequent development of AIDS-related non-Hodgkin's lymphoma. *Cancer Epidemiol Biomarkers Prev* 1999;8:979–84.
- [108] Breen EC, Hussain SK, Magpantay L, Jacobson LP, Detels R, Rabkin CS, et al. B-cell stimulatory cytokines and markers of immune activation are elevated several years prior to the diagnosis of systemic AIDS-associated non-Hodgkin B-cell lymphoma. *Cancer Epidemiol Biomarkers Prev* 2011;20:1303–14.
- [109] Fu SL, Pierre J, Smith-Norowitz TA, Hagler M, Bowne W, Pincus MR, et al. Immunoglobulin E antibodies from pancreatic cancer patients mediate antibody-dependent cell-mediated cytotoxicity against pancreatic cancer cells. *Clin Exp Immunol* 2008;153:401–9.
- [110] Bansal AS, Haeney MR, Cochran S, Pumphrey RS, Green LM, Bhavnani M, et al. Serum soluble CD23 in patients with hypogammaglobulinaemia. *Clin Exp Immunol* 1994;97:239–41.
- [111] Chomarat P, Briolay J, Banchereau J, Miossec P. Increased production of soluble CD23 in rheumatoid arthritis, and its regulation by interleukin-4. *Arthritis Rheum* 1993;36:234–42.
- [112] Huissoon AP, Emery P, Bacon PA, Gordon J, Salmon M. Increased expression of CD23 in rheumatoid synovitis. *Scand J Rheumatol* 2000;29:154–9.
- [113] Kogure T, Itoh T, Shimada Y, Shintani T, Ochiai H, Terasawa K. Detection of serum soluble markers of immune activation in rheumatoid arthritis. *Mediators Inflamm* 1996;5:262–5.
- [114] Kutukculer N, Caglayan S. Plasma and synovial fluid soluble CD23 concentrations in children with juvenile chronic arthritis. *Autoimmunity* 1998;27:155–8.
- [115] Massa M, Pignatti P, Oliveri M, De Amici M, De Benedetti F, Martini A. Serum soluble CD23 levels and CD23 expression on peripheral blood mononuclear cells in juvenile chronic arthritis. *Clin Exp Rheumatol* 1998;16:611–6.
- [116] Rezonzew R, Newkirk MM. Impaired release of sCD23 by activated B-cells from RA patients. *Clin Immunol Immunopathol* 1994;71:156–63.
- [117] Ribbens C, Bonnet V, Kaiser MJ, Andre B, Kaye O, Franchimont N, et al. Increased synovial fluid levels of soluble CD23 are associated with an erosive status in rheumatoid arthritis (RA). *Clin Exp Immunol* 2000;120:194–9.
- [118] Singh A, Vastert SJ, Prakken BJ, Illges H. Decreased levels of sCD21 and sCD23 in blood of patients with systemic-juvenile arthritis, polyarticular-juvenile arthritis, and pauciarticular-juvenile arthritis. *Rheumatol Int* 2011, epub ahead of print.
- [119] Zhou M, Wiemels JL, Bracci PM, Wrensch MR, McCoy LS, Rice T, et al. Circulating levels of the innate and humoral immune regulators CD14 and CD23 are associated with adult glioma. *Cancer Res* 2011;70:7534–42.
- [120] Wrensch M, Wiencke JK, Wiemels J, Milke R, Patoka J, Moghaddasi M, et al. Serum IgE, tumor epidermal growth factor receptor expression,

- and inherited polymorphisms associated with glioma survival. *Cancer Res* 2006;66:4531–41.
- [121] Merrill RM, Isakson RT, Beck RE. The association between allergies and cancer: what is currently known? *Ann Allergy Asthma Immunol* 2007;99:102–16 [Quiz 117:109, 150].
- [122] Jensen-Jarolim E, Achatz G, Turner MC, Karagiannis S, Legrand F, Capron M, et al. Allergooncology: the role of IgE-mediated allergy in cancer. *Allergy* 2008;63:1255–66.
- [123] Lok DJ, Van Der Meer P, de la Porte PW, Lipsic E, Van Wijngaarden J, Hillege HL, et al. Prognostic value of galectin-3, a novel marker of fibrosis, in patients with chronic heart failure: data from the DEAL-HF study. *Clin Res Cardiol* 2010;99:323–8.
- [124] Chiu CG, Strugnell SS, Griffith OL, Jones SJ, Gown AM, Walker B, et al. Diagnostic utility of galectin-3 in thyroid cancer. *Am J Pathol* 2010;176:2067–81.
- [125] Saussez S, Glinier D, Chantrain G, Pattou F, Carnaille B, Andre S, et al. Serum galectin-1 and galectin-3 levels in benign and malignant nodular thyroid disease. *Thyroid* 2008;18:705–12.
- [126] Inohara H, Segawa T, Miyauchi A, Yoshii T, Nakahara S, Raz A, et al. Cytoplasmic and serum galectin-3 in diagnosis of thyroid malignancies. *Biochem Biophys Res Commun* 2008;376:605–10.
- [127] Sethi K, Sarkar S, Das S, Mohanty B, Mandal M. Biomarkers for the diagnosis of thyroid cancer. *J Exp Ther Oncol* 2010;8:341–52.
- [128] Iurisci I, Tinari N, Natoli C, Angelucci D, Cianchetti E, Iacobelli S. Concentrations of galectin-3 in the sera of normal controls and cancer patients. *Clin Cancer Res* 2000;6:1389–93.
- [129] Vereecken P, Awada A, Suciu S, Castro G, Morandini R, Litynska A, et al. Evaluation of the prognostic significance of serum galectin-3 in American Joint Committee on Cancer stage III and stage IV melanoma patients. *Melanoma Res* 2009;19:316–20.
- [130] Vereecken P, Zouaoui Boudjeltia K, Debray C, Awada A, Legssyer I, Sales F, et al. High serum galectin-3 in advanced melanoma: preliminary results. *Clin Exp Dermatol* 2006;31:105–9.
- [131] Saussez S, Lorfèvre F, Lequeux T, Laurent G, Chantrain G, Vertongen F, et al. The determination of the levels of circulating galectin-1 and -3 in HNSCC patients could be used to monitor tumor progression and/or responses to therapy. *Oral Oncol* 2008;44:86–93.
- [132] Sakaki M, Oka N, Nakanishi R, Yamaguchi K, Fukumori T, Kanayama HO. Serum level of galectin-3 in human bladder cancer. *J Med Invest* 2008;55:127–32.
- [133] Bansal A, Roberts T, Hay EM, Kay R, Pumphrey RS, Wilson PB. Soluble CD23 levels are elevated in the serum of patients with primary Sjogren's syndrome and systemic lupus erythematosus. *Clin Exp Immunol* 1992;89:452–5.
- [134] Bansal AS, Ollier W, Marsh MN, Pumphrey RS, Wilson PB. Variations in serum sCD23 in conditions with either enhanced humoral or cell-mediated immunity. *Immunology* 1993;79:285–9.
- [135] Itoh M, Uchimura K, Hayakawa N, Makino M, Hayashi R, Nagata M, et al. Surface expression and release of soluble forms of CD8 and CD23 in CD40- and IL-4-activated mononuclear cells from patients with Graves' disease (GD). *Clin Exp Immunol* 1998;113:309–14.
- [136] Ohshima S, Kuchen S, Seemayer CA, Kyburz D, Hirt A, Klinzing S, et al. Galectin 3 and its binding protein in rheumatoid arthritis. *Arthritis Rheum* 2003;48:2788–95.
- [137] Lee YJ, Kang SW, Song JK, Park JJ, Bae YD, Lee EY, et al. Serum galectin-3 and galectin-3 binding protein levels in Behcet's disease and their association with disease activity. *Clin Exp Rheumatol* 2007;25:S41–5.
- [138] Frol'ova L, Smetana Jr K, Borovska D, Kitanovicova A, Klimesova K, Janatkova I, et al. Detection of galectin-3 in patients with inflammatory bowel diseases: new serum marker of active forms of IBD? *Inflamm Res* 2009;58:503–12.
- [139] de Boer RA, Lok DJ, Jaarsma T, van der Meer P, Voors AA, Hillege HL, et al. Predictive value of plasma galectin-3 levels in heart failure with reduced and preserved ejection fraction. *Ann Med* 2011;43:60–8.
- [140] Lin YH, Lin LY, Wu YW, Chien KL, Lee CM, Hsu RB, et al. The relationship between serum galectin-3 and serum markers of cardiac extracellular matrix turnover in heart failure patients. *Clin Chim Acta* 2009;409:96–9.
- [141] van Kimmenade RR, Januzzi Jr JL, Ellinor PT, Sharma UC, Bakker JA, Low AF, et al. Utility of amino-terminal pro-brain natriuretic peptide, galectin-3, and apelin for the evaluation of patients with acute heart failure. *J Am Coll Cardiol* 2006;48:1217–24.