




Contents lists available at ScienceDirect



Association of neck circumference and pulmonary function in children

Onur Akin, MD^{*}; Mutluay Arslan, MD[†]; Cem Haymana, MD[‡]; Erdem Karabulut, PhD[§]; Bulent Hacıhamdioglu, MD^{||}; Suleyman Tolga Yavuz, MD^{¶, #}

^{*} Department of Pediatric Endocrinology, Gülhane Education and Research Hospital, Ankara, Turkey

[†] Department of Pediatrics, Gülhane Education and Research Hospital, Ankara, Turkey

[‡] Department of Endocrinology, Gülhane Education and Research Hospital, Ankara, Turkey

[§] Department of Biostatistics, Faculty of Medicine, Hacettepe University, Ankara, Turkey

^{||} Department of Pediatric Endocrinology, GATA Military School of Medicine, Ankara, Turkey

[¶] Department of Pediatric Allergy, GATA Military School of Medicine, Ankara, Turkey

[#] Department of Pediatric Allergy, Children's Hospital, University of Bonn, Bonn, Germany

ARTICLE INFO

Article history:

Received for publication March 16, 2017.

Received in revised form April 6, 2017.

Accepted for publication April 24, 2017.

ABSTRACT

Background: Childhood obesity leads to many complications including impaired respiratory function. There are various anthropometric parameters related to obesity.

Objective: To investigate the correlation between anthropometric indices and pulmonary function test results in children without asthma.

Methods: Children without any respiratory disorders were enrolled in this study. Anthropometric measurements, such as height, weight, neck circumference (NC), and waist circumference, were obtained from the enrollees and body mass index was calculated. Afterward, pulmonary function tests were performed using spirometry.

Results: A total of 178 children (106 boys, 59.5%) with a mean age of 9.7 years were included the study. NC was above the 90th percentile in 65 children. Importantly, pulmonary parameters, such as forced expiratory volume during the first second (FEV₁) and the ratio of FEV₁ to forced vital capacity (FVC), were lower in subjects with a large NC. Similarly, waist circumference was above the 90th percentile in 67 children, and FEV₁/FVC was significantly lower in children with a large waist circumference. Moreover, there was a statistically significant negative correlation among FEV₁, FEV₁/FVC, and body mass index SD score. Also, multivariable linear regression analysis showed that an NC above the 90th percentile was associated with lower FEV₁ and FEV₁/FVC values.

Conclusion: We identified NC as a novel anthropometric index that is strongly correlated with respiratory functions in children. Therefore, close monitoring of respiratory symptoms, particularly in children with obesity and a large NC, could help with early and prompt determination of respiratory complications of obesity.

© 2017 American College of Allergy, Asthma & Immunology. Published by Elsevier Inc. All rights reserved.

Introduction

Childhood obesity is a growing health problem that can lead to impairment of organ functions, including the respiratory system.¹ Respiratory impairments are mostly related to obstructive sleep apnea and obesity hypoventilation syndrome. In fact, numerous studies have reported a clear association of obesity with asthma in children.^{2–4}

Body mass index (BMI) is the most commonly used parameter to define obesity. However, BMI does not provide information on the body fat distribution that is particularly associated with

obesity-related respiratory complications.^{5,6} Therefore, anthropometric measurements that identify upper body obesity better than BMI are needed. Several studies have reported an association between waist circumference (WC) and pulmonary functions.^{7,8} Neck circumference (NC) is a new measurement to determine body fat distribution in children and adults.^{9,10} Previous studies have shown that NC is associated with respiratory disorders, such as obstructive sleep apnea, respiratory endurance, and perioperative adverse events.^{11–13} More recent studies have identified an association between NC and severity of asthma.^{14,15} Nevertheless, more clarity is required on the mechanisms that help explain the strong association between obesity and pulmonary function.

The present study investigated the correlation between anthropometric indices and pulmonary function test results in children without asthma.

Reprints: Suleyman Tolga Yavuz, MD, Associate Professor of Pediatric Allergy, Department of Pediatric Allergy, Children's Hospital, University of Bonn, 53113 Bonn, Germany; E-mail: styavuz@hotmail.com.

Disclosures: Authors have nothing to disclose.

<http://dx.doi.org/10.1016/j.anai.2017.04.018>

1081-1206/© 2017 American College of Allergy, Asthma & Immunology. Published by Elsevier Inc. All rights reserved.

Methods

One hundred seventy-eight children 5 to 15 years old who were admitted to pediatric allergy outpatient department for various symptoms from 2013 through 2015 were enrolled in this study. Patients with asthma, other lung diseases, and/or chronic systemic disorders were excluded from the study. Consent forms were obtained from the enrolled children and their parents. The institutional review board approved the study.

Different anthropometric indices such as weight, height, WC, and NC were measured under postabsorptive conditions by a physician. Height was measured to the nearest 0.5 cm on a standard height board, and weight (with no clothes or shoes on, except for underwear) was measured to the nearest 0.1 kg. Subsequently, BMI was calculated as weight (kilograms) divided by height (meters) squared. WC was measured at the end of normal expiration using a flexible tape at the circumference equidistant between the superior border of the iliac crest and the lowest rib margin. Hip circumference was measured at the level yielding the maximum circumference between the trochanters. NC was measured at the level of the cricoid cartilage. All measurements were performed by the same researcher. Obesity was defined as BMI above the 95th percentile for Turkish children. BMI z-scores, WC, and NC percentiles were calculated.¹⁶ Patients were graded according to WC and NC. Values above the 90th percentile were defined as “high.”^{17,18}

Pulmonary function tests were recorded in the sitting position by a physician using a spirometer (Zan 100, Nspire Health, Oberthulba, Germany) according to recommendations of the European Respiratory Society. Parameters, such as forced vital capacity (FVC), forced expiratory volume during the first second (FEV₁), FEV₁/FVC ratio, and forced expiratory flow between 25% and 75% (FEF_{25–75}), were measured and the predicted values were calculated. During the measurement, volume and flow curves were closely monitored to detect artifacts and to confirm enough effort was rendered by the participants. Overall, 3 appropriate measurements were recorded and the highest value was considered the basal value. All values were represented as percentages according to age, sex, weight, and height.

Calculations were performed using SPSS Statistics 21.0 (IBM, Armonk, New York). Descriptive data for categorical variables were expressed as frequencies. Grouped variables were compared using the Student *t* test for numerical values and the Pearson χ^2 test for categorical variables. Pearson correlations were calculated to analyze the relation between BMI SD score (SDS) and lung function test results. Normality of the variable distributions was evaluated by the Kolmogorov-Smirnov test. Multiple linear regression analysis used anthropometric parameters as predictors, and significant variables in the Student *t* test and Pearson correlation test were included as dependent variables. Significantly associated variables were determined by the backward elimination procedure. A *P* value less than .05 was set as the cutoff for statistical significance.

Results

A total of 178 children (106 boys, 59.5%) with a mean age of 9.7 years were included. Atopy was present in 73 patients (41%). Forty-one patients (23%) in the study population were obese and 73 (41%) were overweight (Table 1). NC was above the 90th percentile in 65 children. There was no significant difference in NC by age, sex, birthweight, atopy prevalence, FVC, peak expiratory flow (PEF), and FEF_{25–75}. However, FEV₁ and FEV₁/FVC were higher in subjects with a large NC (*P* = .04 and .01, respectively; Table 2). Similarly, WC was above the 90th percentile in 67 children. Atopy prevalence, birthweight, age, sex, FVC, FEV₁, PEF, and FEF_{25–75} were not significantly different between children with a large or small WC. Only FEV₁/FVC was significantly high in children with a WC higher than the 90th percentile (*P* = .01; Table 3). There was a statistically significant

Table 1

Demographic Features of Study Population (N = 178)^a

Boys	106 (59.5)
Age (y)	9.7 ± 2.6
Atopy	73 (41.0)
BMI z-score	0.94 ± 1.2
Obese	41 (23)
Overweight	73 (41)

Abbreviation: BMI, body mass index.

^aData are presented as frequency (percentage) or mean ± SD.

negative correlation among FEV₁, FEV₁/FVC, and BMI SDS. There were no significant correlations between other pulmonary function test parameters and BMI SDS.

Multiple linear regression analysis used pulmonary function test parameters, such as FEV₁ and FEV₁/FVC, as dependent variables that were significantly associated with obesity indices. Independent variables were determined as anthropometric parameters, such as NC, WC, and BMI SDS. The results showed a significant association among FEV₁, FEV₁/FVC, and NC.

Discussion

In the present study, we found an association between the pulmonary function variable FEV₁ and anthropometric indices of obesity. NC, a novel anthropometric measurement of obesity, was strongly associated with FEV₁ and FEV₁/FVC.

Many studies have shown an association between obesity using BMI as an index and respiratory functions. For instance, FEV₁/FVC was negatively associated with BMI in obese and overweight adults.⁸ In another adult study, a decrease in midexpiratory flow rate and FEV₁/FVC was shown in patients with morbid obesity.¹⁹ Similar findings were observed in cross-sectional studies performed in children with or without asthma.^{20,21} A different study showed that FEV₁ and FEV₁/FVC were significantly lower in children with obesity compared with those without obesity, whereas FVC did not differ by obesity status.²² In the present study, we found that FEV₁ and FEV₁/FVC were negatively correlated with BMI SDS. However, when the comparison was adjusted for other indices of obesity, BMI SDS was not related to FEV₁ and FEV₁/FVC. The regression model showed only the association between pulmonary functions and NC.

Various mechanisms have been suggested to explain the association between obesity and pulmonary functions. Mechanical changes due to fat deposition on the abdomen, chest wall, and upper airway have been implicated in changes in lung volumes,²³ leading to decreases in FVC and FEV₁.^{24–26} Conversely, other pediatric studies, including the present study, found no association between FVC and BMI and/or other obesity indices.²² Therefore, we suggest that, in addition to mechanical pressure, airway resistance

Table 2

Respiratory Function Test Results of Groups According to Neck Circumference^a

	<90th percentile (n = 113)	>90th percentile (n = 65)	<i>P</i> value
Boys	66 (58.4)	40 (61.5)	.28
Age (y)	9.5 ± 2.6	9.8 ± 2.4	.59
Atopy	48 (42.5)	25 (38.5)	.63
Birthweight (g)	3,050 ± 100	3,130 ± 110	.84
FVC (%)	97.5 ± 11.0	95.8 ± 11.3	.35
FEV ₁ (%)	100.9 ± 12.0	96.9 ± 11.3	.04
FEV ₁ /FVC	91.0 ± 5.7	87.0 ± 8.3	.01
PEF (%)	88.3 ± 15.3	87.8 ± 18.3	.35
FEF _{25–75} (%)	97.9 ± 21.7	98.1 ± 18.3	.94

Abbreviations: FEF_{25–75}, forced expiratory flow between 25% and 75%; FEV₁, forced expiratory volume during first second; FVC, forced vital capacity; PEF, peak expiratory flow.

^aData are presented as frequency (percentage) or mean ± SD.

Table 3
Respiratory Function Test Results of Groups According to Waist Circumference^a

	<90th percentile (n = 111)	>90th percentile (n = 67)	P value
Boys	63 (56.7)	43 (64.1)	.09
Age (y)	9.6 ± 2.4	9.7 ± 2.5	.72
Atopy	44 (39.6)	29 (43.2)	.25
Birthweight (g)	3,060 ± 120	3,100 ± 100	.85
FVC (%)	96.8 ± 11.8	95.2 ± 11.0	.78
FEV ₁ (%)	100.8 ± 12.0	97.2 ± 11.4	.06
FEV ₁ /FVC	90.6 ± 5.4	87.8 ± 8.8	.03
PEF (%)	87.3 ± 16.4	87.0 ± 16.7	.91
FEF _{25–75} (%)	98.6 ± 22.8	96.8 ± 15.6	.61

Abbreviations: FEF_{25–75}, forced expiratory flow between 25% and 75%; FEV₁, forced expiratory volume during first second; FVC, forced vital capacity; PEF, peak expiratory flow.

^aData are presented as frequency (percentage) or mean ± SD.

can be associated with obesity. Increased airway resistance observed in children with obesity can be mediated by inflammatory mediators derived from adipocytes, including adiponectin and leptin.^{27,28} Another possible mechanism is remodeling of the airway structure by fat deposition.²⁹ Airway lipid deposition also can affect surfactant production and function.^{30,31} In addition, hyperinsulinemia, a complication of obesity, can induce contraction of airway smooth muscles.³²

The impact of obesity on respiratory functions is affected by the quantity of fat mass and fat distribution.^{7,8} Clearly, BMI does not measure fat distribution. WC is widely used to measure fat distribution and define central obesity. In previous studies, WC was associated with decreased FEV₁ and FVC, but not with FEV₁/FVC.^{5,29} However, in the present study, FEV₁/FVC was decreased in subjects with a larger WC. Another study performed in children found that the FEV₁/FVC ratio was negatively associated with WC, whereas FEV₁ and FVC were independently and positively associated with WC. The investigators of that study suggested that WC is an important predictor of lung volume and FVC because a larger body size indicates a larger lung size in children.³³ However, in the present study, there was no clear association between FVC and WC. Unlike studies in adults, the association between WC and decreased FEV₁/FVC in children might be due to the relation between WC and lung volume. These conflicting results indicate the low reliability of WC as an indicator of central obesity. The abdomen is a changeable region; patients can voluntarily cause changes in WC, and WC depends on the period from the most recent meal to the time of measurement and on problems such as dyspepsia.

NC is a new measurement that could be a good indicator of central obesity. The neck is a sturdier region than the waist, and it does not vary in size during the day. Therefore, NC provides consistency between measurements made by different researchers, thus decreasing false positives.³⁴ Moreover, NC is simple to measure and is well tolerated in children and adolescents with obesity. The neck is anatomically related to the respiratory airways and excess neck fat can mechanically affect respiratory functions.³⁵ Others have suggested an association between NC and upper airway compliance.³⁶ Notably, a study showed that, rather than abdominal adiposity, neck adiposity was associated with respiratory muscle endurance, because NC was related to PEF.¹³ Notably, in the present study, we found that FEV₁ and FEV₁/FVC were lower in patients whose NC value was higher than the 90th percentile. Further, some studies have investigated the relation between NC and respiratory events, particularly in children. For instance, an association between NC and adverse respiratory events in children undergoing surgery has been reported.¹¹ Larger NC also has been associated with higher pCO₂ and lower pO₂ in adults with obesity and normal pulmonary function.³⁷ More recent studies have reported that NC is related to the severity of asthma; it is more closely

associated with and can serve as a better indicator than other indices of obesity.^{14,15} Not surprisingly, the present study found that NC has a stronger association with pulmonary functions compared with WC or BMI.

However, there were some limitations to this study. First, this is a cross-sectional study, and therefore studies with longer follow-up might better indicate the association between NC and obesity indices. Second, the frequency of overweight and obese children in this study was higher than the overweight and obesity prevalence for children, which could have been due to our institution's tertiary institutional structure.

In conclusion, this is probably the first study to demonstrate a possible association between NC and respiratory functions, particularly in children. We found that NC, a new anthropometric indicator of obesity, is closely associated with respiratory functions in children. Therefore, close monitoring of respiratory functions, particularly in children with obesity and a large NC, could help in the early and prompt diagnosis of respiratory complications owing to obesity.

References

- [1] Dietz WH, Robinson TN. Clinical practice. Overweight children and adolescents. *N Engl J Med*. 2005;352:2100–2109.
- [2] Castro-Rodriguez JA, Holberg CJ, Morgan WJ, Wright AL, Martinez FD. Increased incidence of asthmatic symptoms in girls who become overweight or obese during the school years. *Am J Respir Crit Care Med*. 2001;163:1344–1349.
- [3] Gilliland FD, Berhane K, Islam T, et al. Obesity and the risk of newly diagnosed asthma in school-age children. *Am J Epidemiol*. 2003;158:406–415.
- [4] Gold DR, Damokosh AI, Dockery DW, Berkey CS. Body-mass index as a predictor of incident asthma in a prospective cohort of children. *Pediatr Pulmonol*. 2003;36:514–521.
- [5] Canoy D, Luben R, Welch A, et al. Abdominal obesity and respiratory function in men and women in the EPIC-Norfolk Study, United Kingdom. *Am J Epidemiol*. 2004;159:1140–1149.
- [6] Chen R, Tunstall-Pedoe H, Bolton-Smith C, Hannah MK, Morrison C. Association of dietary antioxidants and waist circumference with pulmonary function and airway obstruction. *Am J Epidemiol*. 2001;153:157–163.
- [7] Collins LC, Hoberty PD, Walker JF, Fletcher EC, Peiris AN. The effect of body fat distribution on pulmonary function tests. *Chest*. 1995;107:1298–1302.
- [8] Lazarus R, Sparrow D, Weiss ST. Effects of obesity and fat distribution on ventilatory function: the Normative Aging Study. *Chest*. 1997;111:891–898.
- [9] Hatipoglu N, Mazicioglu MM, Kurtoglu S, Kendirci M. Neck circumference: an additional tool of screening overweight and obesity in childhood. *Eur J Pediatr*. 2010;169:733–739.
- [10] Ben-Noun L, Laor A. Relationship of neck circumference to cardiovascular risk factors. *Obes Res*. 2003;11:226–231.
- [11] Nafiu OO, Burke CC, Gupta R, Christensen R, Reynolds PI, Malviya S. Association of neck circumference with perioperative adverse respiratory events in children. *Pediatrics*. 2011;127:e1198–e1205.
- [12] Figueroa-Munoz JI, Chinn S, Rona RJ. Association between obesity and asthma in 4–11 year old children in the UK. *Thorax*. 2001;56:133–137.
- [13] Goncalves MJ, do Lago ST, Godoy Ede P, Fregonezi GA, Bruno SS. Influence of neck circumference on respiratory endurance and muscle strength in the morbidly obese. *Obes Surg*. 2011;21:1250–1256.
- [14] Hacıhamdioglu B, Arslan M, Yesilkaya E, Gok F, Yavuz ST. Wider neck circumference is related to severe asthma in children. *Pediatr Allergy Immunol*. 2015;26:456–460.
- [15] Akin O, Sari E, Arslan M, Yesilkaya E, Hacıhamdioglu B, Yavuz ST. Association of wider neck circumference and asthma in obese children. *Ann Allergy Asthma Immunol*. 2016;116:514–517.
- [16] Fredriks AM, van Buuren S, Jeurissen SE, Dekker FW, Verloove-Vanhorick SP, Wit JM. Height, weight, body mass index and pubertal development reference values for children of Turkish origin in the Netherlands. *Eur J Pediatr*. 2003;162:788–793.
- [17] Mazicioglu MM, Kurtoglu S, Ozturk A, Hatipoglu N, Cicek B, Ustunbas HB. Percentiles and mean values for neck circumference in Turkish children aged 6–18 years. *Acta Paediatr*. 2010;99:1847–1853.
- [18] Unalan D, Senol V, Bayat M, et al. Change in waist circumference over 3 years in Turkish children and adolescents. *Ann Hum Biol*. 2013;40:419–425.
- [19] Biring MS, Lewis MI, Liu JT, Mohsenifar Z. Pulmonary physiologic changes of morbid obesity. *Am J Med Sci*. 1999;318:293–297.
- [20] Vo P, Makker K, Matta-Arroyo E, Hall CB, Arens R, Rastogi D. The association of overweight and obesity with spirometric values in minority children referred for asthma evaluation. *J Asthma*. 2013;50:56–63.
- [21] Chu YT, Chen WY, Wang TN, Tseng HI, Wu JR, Ko YC. Extreme BMI predicts higher asthma prevalence and is associated with lung function impairment in school-aged children. *Pediatr Pulmonol*. 2009;44:472–479.

- [22] Strunk RC, Colvin R, Bacharier LB, et al. Airway obstruction worsens in young adults with asthma who become obese. *J Allergy Clin Immunol Pract.* 2015;3: 765–771.e762.
- [23] Brazzale DJ, Pretto JJ, Schachter LM. Optimizing respiratory function assessments to elucidate the impact of obesity on respiratory health. *Respirology.* 2015;20:715–721.
- [24] Jones RL, Nzekwu MM. The effects of body mass index on lung volumes. *Chest.* 2006;130:827–833.
- [25] Littleton SW. Impact of obesity on respiratory function. *Respirology.* 2012;17: 43–49.
- [26] Ulger Z, Demir E, Tanac R, et al. The effect of childhood obesity on respiratory function tests and airway hyperresponsiveness. *Turk J Pediatr.* 2006;48: 43–50.
- [27] Guerra S, Wright AL, Morgan WJ, Sherrill DL, Holberg CJ, Martinez FD. Persistence of asthma symptoms during adolescence: role of obesity and age at the onset of puberty. *Am J Respir Crit Care Med.* 2004;170: 78–85.
- [28] Lu Y, Van Bever HP, Lim TK, et al. Obesity, asthma prevalence and IL-4: roles of inflammatory cytokines, adiponectin and neuropeptide Y. *Pediatr Allergy Immunol.* 2015;26:530–536.
- [29] Salome CM, King GG, Berend N. Physiology of obesity and effects on lung function. *J Appl Physiol (1985).* 2010;108:206–211.
- [30] Inselman LS, Chander A, Spitzer AR. Diminished lung compliance and elevated surfactant lipids and proteins in nutritionally obese young rats. *Lung.* 2004; 182:101–117.
- [31] Inselman LS, Wapnir RA, Spencer H. Obesity-induced hyperplastic lung growth. *Am Rev Respir Dis.* 1987;135:613–616.
- [32] Gosens R, Nelemans SA, Hiemstra M, Grootte Bromhaar MM, Meurs H, Zaagsma J. Insulin induces a hypercontractile airway smooth muscle phenotype. *Eur J Pharmacol.* 2003;481:125–131.
- [33] Chen Y, Rennie D, Cormier Y, Dosman JA. Waist circumference associated with pulmonary function in children. *Pediatr Pulmonol.* 2009;44:216–221.
- [34] LaBerge RC, Vaccani JP, Gow RM, Gaboury I, Hoey L, Katz SL. Inter- and intra-rater reliability of neck circumference measurements in children. *Pediatr Pulmonol.* 2009;44:64–69.
- [35] Carpenter MA, Tockman MS, Hutchinson RG, Davis CE, Heiss G. Demographic and anthropometric correlates of maximum inspiratory pressure: the Atherosclerosis Risk in Communities Study. *Am J Respir Crit Care Med.* 1999;159:415–422.
- [36] Rowley JA, Sanders CS, Zahn BR, Badr MS. Gender differences in upper airway compliance during NREM sleep: role of neck circumference. *J Appl Physiol (1985).* 2002;92:2535–2541.
- [37] Gabrielsen AM, Lund MB, Kongerud J, Viken KE, Roislien J, Hjelmessaeth J. The relationship between anthropometric measures, blood gases, and lung function in morbidly obese white subjects. *Obes Surg.* 2011;21:485–491.