

Analytical Methods

A novel method using immuno-affinity chromatography for isolating β -conglycinin from soybean proteinsJinming You^a, Peng Sun^{b,1}, Defa Li^{b,*}, Shiyao Qiao^b, Mingren Qu^a, Zirui Wang^a, Guanhong Li^a, Ke Pan^a^a College of Animal Science and Technology, Jiangxi Agricultural University, No. 1101 Zhimin Road, Nanchang 330045, PR China^b National Key Laboratory of Animal Nutrition, China Agricultural University, No. 2 West Yuanmingyuan Road, Beijing 100193, PR China

ARTICLE INFO

Article history:

Received 29 November 2007

Received in revised form 17 March 2009

Accepted 29 March 2009

Keywords:

 β -Conglycinin

Monoclonal antibody (Mab)

Immuno-affinity chromatography

Purification

ABSTRACT

A monoclonal antibody (Mab) 6G₄ against soybean β -conglycinin has been prepared using a conjugate of chicken ovalbumin and a synthetic peptide that corresponded to one of the epitope sequences of β -conglycinin as the immunogen. An ELISA method for the quantification of β -conglycinin has also been developed. In the present study, we report a novel method for the purification of β -conglycinin by Mab 6G₄-based immuno-affinity chromatography. β -Conglycinin with a purity of 92.9% was successfully isolated from soybean proteins. Western blot assay was used to further identify its characteristics and the results demonstrated that the purified β -conglycinin maintains its biological activities. Therefore, the Mab-based immuno-affinity chromatography is an available method for purification of β -conglycinin. It also provides a new opportunity for future study on the mechanism of food allergy responses using high purity β -conglycinin as the experimental material.

© 2009 Elsevier Ltd. All rights reserved.

1. Introduction

Soybean is widely used as food ingredients because of its nutritional and functional properties (Astadi, Astuti, Santoso, & Nugraheni, 2008; Friedman & Brandon, 2001; Liu et al., 2008; Saito, Kohno, Tsumura, Kugimiya, & Kito, 2001). However, soybean is also considered as a major source of food allergies especially in industrialised countries (Guo, Piao, Cao, Ou, & Li, 2008; Herman, Helm, Jung, & Kinney, 2003; Huisman & Jansman, 1991). Soybean allergies are prevalently becoming “health hacker” in the world recently (Chandra, 2002; Song, Frias, Martinez-Villaluenga, Vidal-Valverde, & de Mejia, 2008; Zeiger, 2000).

As a major soybean allergen, β -conglycinin with α , α' and β subunits plays important roles in hypersensitive responses for children (Xiang, Beardslee, Zeece, Marwell, & Sarath, 2002) and young animals such as piglets (Li et al., 1990), calves (Lallès & Dreau, 1996) and mice (Guo et al., 2008). However, study on β -conglycinin has been limited due to the difficulty of separating β -conglycinin from other soybean proteins. Most previous studies have mainly used soybeans or their extracts as the experimental materials but not pure β -conglycinin because of the lack of an available isolation approach. Although some methods such as ammonium sulphate precipitation (Thanh & Shibasaki, 1976), pH adjustment isolation (Lovati et al., 1992), ultrafiltration membrane separation

(Wu, Murphy, Reuber, & Fratzke, 2000) and phytase treatment (Saito et al., 2001) can be conducted to isolate β -conglycinin from soybean proteins, it is difficult to obtain enough pure β -conglycinin for research purposes. Since monoclonal antibodies (Mabs) are powerful tools for many molecular immunology investigations (Nelson et al., 2000), they have frequently been applied in practical studies on recognising antigens and identifying new cell surface molecules. These binding concepts of allergen and antibody also provide a possibility for us to prepare an immuno-affinity column mediated by Mab and develop a new approach for purification of β -conglycinin which can be used for future research purposes.

2. Materials and methods

2.1. Preparation of monoclonal antibody against soybean β -conglycinin

The preparation of a stable murine-derived hybridoma cell line, named as 6G₄, has been described previously (You et al., 2008). The Mab against soybean β -conglycinin produced by 6G₄ is an IgG1 isotype. It shows high affinity for β -conglycinin with an association constant of $6.9 \times 10^9 \text{ M}^{-1}$ and can specifically bind to the α and α' subunits of β -conglycinin (You et al., 2008).

2.2. Production and purification of antibodies

A large amount of Mab was produced *in vivo*. The hybridoma cells from 6G₄ (1×10^6 cells/mouse) were injected

* Corresponding author. Tel.: +86 10 62733588; fax: +86 10 62733688.

E-mail address: defali@public2.bta.net.cn (D. Li).¹ Co-first author with equal contribution.

intraperitoneally into BLAB/c mice pre-treated 7 days before with 0.5 mL mineral oil. About two weeks later, the ascitic fluids from mice inoculated by hybridoma cells were collected and centrifuged for 15 min at $10000\times g$ to eliminate cells and dregs. The ascite supernatant was diluted with equal volume of 0.04 M barbital buffer (pH 7.0). An appropriate amount of SiO_2 powder was added into ascites dilution and gently stirred for 30 min. After centrifuging for 20 min at $2000\times g$, Mab 6G₄ was primarily isolated from supernatant by ammonium sulphate, and a commercial Protein G Affinity Column (Amersham Biosciences, Uppsala, Sweden) chromatography was then performed to further purify IgG1 from ammonium sulfate precipitation.

2.3. Preparation of immuno-affinity column

The CNBr-activated SepharoseTM 4B (Amersham Biosciences, Uppsala, Sweden) medium was prepared according to its production instructions. Briefly, 2.3 g of CNBr-activated SepharoseTM 4B freeze-dried powder was suspended in 1 mM HCl (pH 2.5). After swelling, the medium was washed for 15 min with 1 mM HCl on a sintered glass filter to remove all additives.

Fifty-six mg of Mab 6G₄ (IgG1) as ligand was dissolved in 8 mL of coupling buffer (0.1 M NaHCO_3 , pH 8.3 containing 0.5 M NaCl) and mixed with prepared Sepharose 4B gel in a 10 mL test tube fitted with a stopper. Then, the tube was tied on the orbital shaker and the mixture was gently shaken for 1 h at 20–25°C. Excessive ligand was flushed away with at least five volumes of coupling buffer. Tris–HCl buffer (0.1 M, pH 8.0) was added to block any remaining active groups for 2 h. The gel was washed for at least three cycles of alternating pH using five volumes of each buffer. Each cycle consisted of a wash with 0.1 M acetate buffer (pH 4.0 containing 0.5 M NaCl) followed by a wash with 0.1 M Tris–HCl (pH 8.0 containing 0.5 M NaCl).

The gel coupled with IgG1 was loaded into a column (10 $\times$ 100 mm) which was washed with 20 mM sodium phosphate buffer (pH 7.0) at 4 °C. The residual space of the column was immediately filled with buffer, and the top of the column was connected to a pump. The pump was set to run at a flow rate of 1.3 mL/min. The elution was maintained for 10 bed volumes after a constant bed height was reached.

2.4. Extraction of total soybean proteins

Soybean seeds were ground with a Cyclotec Sample Mill (Model 1093, Tecator Inc., Herndon, VA, USA) to pass through a 60-mesh sieve. The soybean flour was defatted by *n*-hexane. The fat-free flour was then suspended in 0.03 M Tris–HCl buffer (pH 8.0 containing 0.01 M β -mercaptoethanol) (100 g flour per litre buffer). Proteins were extracted from the flour for 1.5 h by vortexing. Extracts were centrifuged at $12,000\times g$ for 20 min at 4 °C using an OptimaTM L-80 XP Ultracentrifuge (Beckman Coulter Inc, Palo Alto, CA, USA). After being filtered through a 0.45 μm Millex GP filter (Millipore, Cork, Ireland), the supernatant was used as test sample solution and stored at –20 °C until analysis.

2.5. Immuno-affinity chromatography of β -conglycinin

The total proteins was loaded onto the immuno-affinity column. The outlet tubing of the column was connected to an HD21-1 detector and a protein auto-collector (Huxi instrument Co., Shanghai, China). Flow rate was maintained for 1.0 mL/min. The medium was washed with binding buffer until the base line was stable. Glycine–HCl elution buffer (0.1 M, pH 2.7) was used to elute the sample and β -conglycinin was collected using the auto-collector according to absorbance profile. The pH of collection

(β -conglycinin) was regulated immediately to neutral with 100 μL of 1 M Tris–HCl (pH 9.0).

2.6. Identification of β -conglycinin by Western blot

The proteins isolated from the immuno-affinity column were analysed by 12.5% sodium dodecyl sulphate–polyacrylamide gel electrophoresis (SDS–PAGE) according to the standard procedure (Laemmli, 1970). The separated proteins were then transferred electrophoretically from gel to a 0.45 μm nitrocellulose membrane in a mini Trans-Blot Cell Apparatus (BioRad) for 1.5 h at 100 V. The membrane was incubated in 3% bovine serum albumin (BSA) for 2 h at 37 °C. After a stringent wash with Tris-buffered saline (TBS) containing 0.05% Tween-20 (TBST), the membrane was incubated with 0.1 $\mu\text{g/mL}$ Mab 6G₄ for 2 h at 37 °C. The unbound primary antibody was removed by washing with TBST. The horseradish peroxidase-conjugated secondary antibody (goat anti-mouse IgG–HRP) was then added and the substrate 3,3'-diaminobenzidine (DAB) was used to show the specific protein bands.

The concentration of isolated β -conglycinin was determined with a Mab-based competitive ELISA method described previously (You et al., 2008) and the concentration of total proteins was tested by Bicinchoninic Acid Kit for Protein Determination (BCATM Kit, Pierce, Rockford, IL, USA). The purity of β -conglycinin is expressed as a percentage as the content of β -conglycinin compared with that of total proteins.

2.7. Evaluation of chromatography efficiency in the case of different Mab/gel ratio

There is a close relationship between the ligand (Mab)/gel ratio and the chromatography efficiency. In order to enhance the chromatography efficiency, it is necessary to obtain an optimal ligand (Mab)/gel ratio. In this study, five grade concentrations of Mab, 2.5, 5.0, 7.5, 10.0, 12.5 mg/mL gel, were conducted to prepare five immuno-affinity columns with the same size. The efficiency of the purification of β -conglycinin under identical conditions, including soybean protein extraction and isolation procedures, was then evaluated.

The concentrations and purities of isolated β -conglycinin were determined and the results are reported as mean $\pm$ SEM. All data were the average values from five replicate determinations. Statistical analysis was done using the Stat-View program (SAS Institute, Cary, NC, USA). Differences between the concentrations (or purities) of β -conglycinin obtained from five immuno-affinity columns were examined for statistical significance using the Dunnett test. A difference was considered as significant at $P < 0.05$.

3. Results and discussion

Fig. 1 shows the SDS–PAGE profile (Left) and Gary Density profile (Right) of β -conglycinin purified from soybean proteins. Lane b represents total soybean proteins before purification and lane c represents β -conglycinin purified from total proteins. The purified β -conglycinin fraction contained mainly α , α' , and β subunits of β -conglycinin. Lane d is a grey density profile obtained from track c of electrophoresis. Direct observation clearly indicated that β -conglycinin was successfully isolated from soybean proteins. A Mab-based competitive ELISA method was used to further test its content. The result indicated that the purity of β -conglycinin was 92.9% (Data not shown).

In previous reports, the purity of β -conglycinin acquired from soybeans were 68% by ultrafiltration membrane separation method (Wu et al., 2000) and no more than 79% by ethanol extraction method (Rickert, Johnson, & Murphy, 2004). In the present study,




Fig. 1. SDS–PAGE profile (Left), grey density profile and Western blot assay (Right) of β -conglycinin purified by Mab-based immuno-affinity chromatography method. Proteins were separated by a 12.5% SDS–PAGE and stained with Coomassie Blue. Lane: a. Protein molecular weight markers; b. Total protein before purification; c. Goal protein (β -conglycinin) separated from soybean proteins through the immuno-affinity column; d. Gray density profile obtained from track c of electrophoresis using Syngene Analysis Software (Version 3.03.03); e. The proteins separated by SDS–PAGE were transferred electrophoretically to a 0.45 μ m nitrocellulose membrane in a mini Trans-Blot Cell Apparatus. Wet blotting was performed at 100 V for 1.5 h. The blots were probed with 0.1 μ g/mL of primary antibody (Mab). The goat anti-mouse IgG–HRP was used as the second antibody and KC™ General Western Blot Detection Kit (Roche Diagnostics Ltd., Lewes, UK) was used to detect the bound antibody.

a higher purity (92.9%) β -conglycinin was obtained by a Mab-based immuno-affinity chromatography method. We attribute the purity improvement to two factors. The first one is the use of immuno-chemical approach based on a highly specific Mab with moderate affinity. Using the prepared immuno-affinity column, the target protein was captured by the specific antibody molecule that immobilized on Sepharose 4B gel whilst other unwanted proteins were flushed from the column. The second factor is the convenient and rapid isolation procedure. It is well known that a complex and time consuming isolation procedure has an increased probability of degradation or denaturation of the target protein. Our purification procedure is very convenient and the time consumed per processing cycle is 5–10 min, which is less than about 30 min by methods of Thanh and Shibasaki (1976), Wu et al. (2000), and Rickert et al. (2004). The method developed by Thanh and Shibasaki (1976) and Wu et al. (2000) can separate two main soybean protein fractions, glycinin and β -conglycinin, by pH adjustment. However, it is difficult to avoid the present of an intermediate protein mixture (Wu et al., 2000). Also, other proteins of which the isoelectric point close to that of β -conglycinin are easy to precipitate in β -conglycinin samples. For the method described by Rickert et al. (2004), extraction environments (pH, ethanol concentration, temperature, and water-to-flake ratio) are each varied during bench-scale optimisation. Optimised conditions may yield more β -conglycinin with higher isoflavone and saponin concentrations, but fraction purity was diminished by glycinin contamination. Therefore, the probability that the target protein is degraded or denatured can be greatly reduced.

In addition to purity, the bioactivity of β -conglycinin is also an important index to evaluate the feasibility of the immuno-affinity column. The bioactivity of isolated β -conglycinin was assessed by

the specific binding of β -conglycinin to Mab using Western blot assay. Fig. 1 (Lane e.) demonstrated that the Mab 6G₄ could recognise the α' and α subunits of the β -conglycinin molecule. This result indicated that the purified β -conglycinin maintained the natural bioactivity of protein molecule.

During the preparation of the immuno-affinity column, the coupling ratio of Mab and gel is one major factor impacting the isolation efficiency of β -conglycinin. The effects of Mab/gel ratio on isolation efficiency are presented in Table 1. No significant difference in terms of purity of isolated β -conglycinin was observed when the concentration of Mab coupled in Sepharose gel ranged from 2.5 to 12.5 mg/mL. However, different concentrations of Mab induced marked changes in yield of β -conglycinin. The concentrations of β -conglycinin in the 5.0, 7.5, and 10.0 mg/mL treatments were significantly higher than those in the 2.5 and 12.5 mg/mL treatments ($P < 0.05$). Interestingly, differences between 5.0, 7.5, and 10.0 mg/mL treatments were not significant ($P > 0.05$). Using this method, the yield of β -conglycinin separated from soybean proteins was 27.3–33.1 g/kg soybean. No differences were presented between 5.0, 7.5, and 10.0 mg/mL treatments ($P > 0.05$). Nevertheless, the yields in the 5.0, 7.5, and 10.0 mg/mL treatments were statistically superior to those in the 2.5 and 12.5 mg/mL treatments ($P < 0.05$). These results suggested that a low concentration of ligand (Mab) can reduce the isolation efficiency of β -conglycinin. Meanwhile, a very high concentration of ligand can have adverse effects on affinity chromatography. This may be because the less Mab couples in gel, the less likely goal protein in soybean solution binds to Mab. Namely, the more Mab couples in gel the more opportunity there is for goal protein to bind to Mab. However, too much Mab increases steric hindrance between the active sites and reduces the binding efficiency of the adsorbent. In

Table 1
Effects of Mab/gel ratio on efficiency of immuno-affinity chromatography for the isolation of soybean β -conglycinin.

	Concentration of Mab coupled in Sepharose 4B gel (mg/mL)				
	2.5	5.0	7.5	10.0	12.5
Concentration of isolated β -conglycinin (mg/mL)	0.48 $\pm$ 0.013 ^b	0.55 $\pm$ 0.010 ^a	0.56 $\pm$ 0.011 ^a	0.54 $\pm$ 0.008 ^a	0.50 $\pm$ 0.009 ^b
Yield in soybean (g/kg)	27.3 $\pm$ 1.9 ^b	31.6 $\pm$ 2.4 ^a	33.1 $\pm$ 2.1 ^a	32.4 $\pm$ 1.8 ^a	29.3 $\pm$ 1.5 ^b
Purity of isolated β -conglycinin (%)	93.2 $\pm$ 0.122	92.8 $\pm$ 0.292	93.1 $\pm$ 0.250	92.9 $\pm$ 0.203	92.7 $\pm$ 0.240

All data are means $\pm$ SEM ($n = 5$). Means within the same row that have no common superscript letters are significantly different ($P < 0.05$).

addition, substances bind more strongly to the immobilized ligand, which may result in difficult elution. On the other hands, the extent of non-specific binding increases at high ligand concentrations. For an efficient adsorbent, 5.0–10.0 mg ligand (Mab) per mL gel is suitable. In view of the above information, we considered that 92.7–93.2% purity of β -conglycinin can be successfully isolated from soybean proteins using the immuno-affinity column packed with 5.0–10.0 mg Mab per mL gel.

4. Conclusions

The Mab-based immuno-affinity chromatography developed in our study provides a novel method for the isolation of β -conglycinin. Under the optimized conditions, this technique can be used for collection of high purity β -conglycinin. In addition, the developed Mab-based immuno-affinity chromatography method also opens a new opportunity for the future study of the mechanism of food allergy responses using high purity β -conglycinin as an allergen protein material.

Acknowledgements

We thank the former research members, Dr. He P. L., Ou D. Y., & Dong B. (National Key Laboratory of Animal Nutrition, China Agricultural University), for the earlier stage works in the preparation of monoclonal antibody against soybean β -conglycinin and the development of competitive ELISA method for quantification of β -conglycinin. This work was supported in part by Grant from the National Natural Science Foundation of China (No. 30430520).

References

- Astadi, I. R., Astuti, M., Santoso, U., & Nugraheni, P. S. (2008). In vitro antioxidant activity of anthocyanins of black soybean seed coat in human low density lipoprotein (LDL). *Food Chemistry*, doi: 10.1016/j.foodchem.2008.06.034.
- Chandra, R. K. (2002). Food hypersensitivity and allergic diseases. *European Journal of Clinical Nutrition*, 56(Suppl. 3), 54–56.
- Friedman, M., & Brandon, D. L. (2001). Nutritional and health benefits of soy proteins. *Journal of Agricultural and Food Chemistry*, 49, 1069–1086.
- Guo, P., Piao, P., Cao, Y., Ou, D., & Li, D. (2008). Recombinant soybean protein β -conglycinin α' -subunit expression and induced hypersensitivity reaction in rats. *International Archives of Allergy and Immunology*, 145, 102–110.
- Herman, E. M., Helm, R. M., Jung, R., & Kinney, A. J. (2003). Genetic modification removes an immunodominant allergen from soybean. *Plant Physiology*, 132, 36–43.
- Huisman, J., & Jansman, A. J. M. (1991). Dietary effects and some analytical aspects of antinutritional factors in peas (*Pisum sativum*), common beans (*Phaseolus vulgaris*) and soyabeans (*Glycine max* L) in monogastric farm animals. A literature review. *Nutrition Abstract and Reviews (Series B)*, 61, 901–921.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T₄. *Nature*, 207, 680–685.
- Lallès, J. P., & Dreau, D. (1996). Identification of soyabean allergens and immune mechanisms of dietary sensitivities in preruminant calves. *Research in Veterinary Science*, 60, 111–116.
- Li, D. F., Nelssen, J. L., Reddy, P. G., Blecha, F., Hancock, J. D., Allee, G. L., et al. (1990). Transient hypersensitivity to soybean meal in the early weaned pig. *Journal of Animal Science*, 68, 1790–1799.
- Liu, C., Wang, X., Ma, H., Zhang, Z., Gao, W., & Xiao, L. (2008). Functional properties of protein isolates from soybeans stored under various conditions. *Food Chemistry*, 111(1), 29–37.
- Lovati, M. R., Manzoni, C., Corsini, A., Granata, A., Frattini, R., Fumagalli, R., et al. (1992). Low density lipoprotein receptor activity is modulated by soybean globulins in cell culture. *Journal of Nutrition*, 122, 1971–1978.
- Nelson, P. N., Reynolds, G. M., Waldron, E. E., Ward, E., Giannopoulos, K., & Murry, P. G. (2000). Monoclonal antibodies demystified. *Journal of Clinical Pathology*, 53, 111–118.
- Rickert, D. A., Johnson, L. A., & Murphy, P. A. (2004). Improved fractionation of glycinin and β -conglycinin and partitioning of phytochemicals. *Journal of Agriculture and Food Chemistry*, 52, 1726–1734.
- Saito, T., Kohno, M., Tsumura, K., Kugimiya, W., & Kito, M. (2001). Novel method using phytase for separating soybean β -conglycinin and glycinin. *Bioscience Biotechnology and Biochemistry*, 65, 884–887.
- Song, Y. S., Frias, J., Martinez-Villaluenga, C., Vidal-Valverde, C., & de Mejia, E. G. (2008). Immunoreactivity reduction of soybean meal by fermentation, effect on amino acid composition and antigenicity of commercial soy products. *Food Chemistry*, 108(2), 571–581.
- Thanh, V. H., & Shibasaki, K. (1976). Major proteins of soybean seed: A straightforward fractionation and their characterisation. *Journal of Agricultural and Food Chemistry*, 24, 1117–1121.
- Wu, S. W., Murphy, P. A., Reuber, M. A., & Fratzke, A. R. (2000). Simplified process for soybean glycinin and β -conglycinin fractionation. *Journal of Agriculture and Food Chemistry*, 48, 2702–2708.
- Xiang, P., Beardslee, T. A., Zeece, M. G., Marwell, J., & Sarath, G. (2002). Identification and analysis of a conserved immunoglobulin E-binding epitope in soybean G1a and G2a and peanut Ara h 3 glycinins. *Archives of Biochemistry and Biophysics*, 408, 51–57.
- You, J. M., Li, D. F., Qiao, S. Y., Wang, Z. R., He, P. L., Ou, D. Y., et al. (2008). Development of monoclonal antibody-based competitive ELISA for detection of β -conglycinin, an allergen from soybean. *Food Chemistry*, 106(1), 352–360.
- Zeiger, R. S. (2000). Dietary aspects of food allergy prevention in infants and children. *Journal of Pediatric Gastroenterology and Nutrition*, 30, 77–86.