


Accepted Manuscript

Disparities in mortality and morbidity in pediatric asthma hospitalizations, 2007 to 2011

Alexander F. Glick, MD, Suzy Tomopoulos, MD, Arthur H. Fierman, MD, Leonardo Trasande, MD, MPP



PII: S1876-2859(15)00387-3

DOI: [10.1016/j.acap.2015.12.014](https://doi.org/10.1016/j.acap.2015.12.014)

Reference: ACAP 796

To appear in: *Academic Pediatrics*

Received Date: 13 September 2015

Revised Date: 27 December 2015

Accepted Date: 30 December 2015

Please cite this article as: Glick AF, Tomopoulos S, Fierman AH, Trasande L, Disparities in mortality and morbidity in pediatric asthma hospitalizations, 2007 to 2011, *Academic Pediatrics* (2016), doi: 10.1016/j.acap.2015.12.014.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Disparities in mortality and morbidity in pediatric asthma hospitalizations, 2007 to 2011

Authors: Alexander F. Glick^{1*}, MD; Suzy Tomopoulos¹, MD; Arthur H. Fierman¹, MD; Leonardo Trasande^{1,2,3,4,5}, MD, MPP

- 1) Department of Pediatrics, New York University (NYU) School of Medicine/Bellevue Hospital Center, New York, NY, 10016, USA
- 2) Departments of Environmental Medicine and Population Health, NYU School of Medicine, New York, NY, 10016, USA
- 3) NYU Wagner School of Public Service, New York, NY, 10003, USA
- 4) NYU Steinhardt School of Culture, Education, and Human Development, Department of Nutrition, Food & Public Health, New York, NY, 10003, USA
- 5) NYU Global Institute of Public Health, New York, NY, 10003, USA

* Corresponding Author

Alexander F. Glick, MD
Bellevue Hospital Center
Department of Pediatrics
Administration Building, Room A314A
New York, NY 10016
Phone: 212-562-2455
Fax: 212-562-5518

Email addresses

Alexander F. Glick: alexander.glick@nyumc.org
Suzy Tomopoulos: soultana.tomopoulos@nyumc.org
Arthur H. Fierman: arthur.fierman@nyumc.org
Leonardo Trasande: leonardo.trasande@nyumc.org

Keywords: asthma, disparities, hospitalizations, mortality, morbidity

Running/Abbreviated Title: Disparities in pediatric asthma hospitalizations, 2007 to 2011

Funded by HRSA 1 T32HP22238 DHHS, HRSA, Primary Care (NRSA T32: Postdoctoral Primary Care Research Training Program, PI Arthur H. Fierman) and the NYU CTSA grant UL1TR000038 from the National Center for the Advancement of Translational Science (NCATS), NIH. The sponsor had no role in the study design, collection/analysis/interpretation of data, writing of the manuscript, or the decision to submit the manuscript for publication. We wish to confirm that there are no known conflicts of interest associated with this publication and there has been no significant financial support for this work that could have influenced its outcome.

Abstract Word Count: 249

Manuscript Word Count: 3437

Abstract**Objective**

Asthma is a leading cause of pediatric admissions. While several factors including race have been linked to increased overall asthma morbidity and mortality, few studies have explored factors associated with inpatient asthma outcomes. We examined factors associated with mortality and morbidity in children admitted for asthma.

Design/Methods

Data were obtained from the US Nationwide Inpatient Sample for 2007-2011. Patients 2-18 years old with a primary diagnosis of asthma were included. Predictor variables were sociodemographic and hospital factors and acute/chronic secondary diagnoses. Outcomes were mortality, intubation, length of stay (LOS), and costs. Weighted national estimates were calculated. Multivariable analyses were performed.

Results

There were 97,379 (478,546 weighted) asthma admissions. Most patients were male (60.6%); 30% were white, 28% black, and 18% Hispanic. Mortality rate was 0.03%. 0.3% were intubated. Median (IQR) LOS was 2 (1-3) days. Median (IQR) costs were \$2760 (\$1860-4320). Native American race, older age (13-18 years), and West region were significant independent predictors of mortality. Intubation rate was lower in Hispanic compared to white children ($p=0.028$). LOS was shorter in Asian compared to white children ($p=0.022$) but longer in children with public insurance and from low income areas ($p < 0.001$). Average costs were higher in black, Hispanic, and Asian compared to white children ($p < 0.05$).

Conclusions

With the exception of Native Americans, race/ethnicity is not associated with inpatient asthma mortality and has varied effects on morbidity. Recognition of factors associated with increased asthma mortality and morbidity may allow for earlier, more effective treatment and avoidance of complications.

What's New: Native American children have higher inpatient asthma mortality rates compared to white children. Race and ethnicity are otherwise not independent predictors of inpatient mortality and have varied effects on intubation, length of stay, and costs.

Introduction

Asthma is one of the most common conditions seen in pediatric patients. In 2009, 9.6% of children (7.1 million children) in the United States (US) were estimated to have asthma.¹ Asthma symptoms lead to 10.5 million missed school days per year.¹ More than 5% of all pediatric hospital admissions are for asthma,² making it a leading cause of pediatric hospitalizations.

It has been well established that racial and ethnic disparities exist in pediatric patients with asthma. Black and Hispanic children have a higher asthma prevalence and overall rates of mortality compared to white children.¹⁻³ Readmission rates are twice as high in black compared to white children.⁴ Black and Hispanic children have higher rates of emergency department (ED) use and hospitalizations for asthma.⁵⁻⁸ Black and Hispanic children often use the ED for their usual source of care⁹ and miss more school days due to their asthma compared to white children.¹⁰ These disparities can in part be explained by financial and social hardship,⁴ worse access to care including to primary care physicians,^{7,11} and improper controller and rescue medication use.^{6,10,12}

While black and Hispanic race and ethnicity have been shown to be predictors of mortality and morbidity from asthma in the outpatient setting, little is known on the national level about whether disparities persist for children after they have been admitted. One national study that examined inpatient asthma disparities in pediatric and adult patients ages five and older found no significant differences in hospital deaths from asthma by race or ethnicity in their multivariable analysis.¹³ The study did find that mortality was more likely in older adults, males, patients with more chronic conditions, and during winter months.¹³ While this study provides some evidence against racial and ethnic disparities associated with inpatient asthma mortality, it

did not specifically examine predictors of other more common adverse outcomes such as increased costs, longer length of stay (LOS), or other measures of more severe disease/morbidity which may have different predictors or in which disparities would be more easily identified given that mortality is rare. In addition, the study did not provide detail regarding the specific comorbidities that may lead to worse outcomes as has been done in studies in other settings.^{14–16} Another study of asthmatics at academic children's hospitals (from July 2002 to June 2003) found that black children had statistically shorter unadjusted LOS and lower charges (after adjusting for LOS) than white children although these differences may not be clinically significant;¹⁷ additional multivariable analyses were not presented. Furthermore, in the last decade there have been no published nationally representative studies on racial and ethnic disparities in inpatient asthma admissions and little work examining asthma outcomes in Native American or Asian children.

The purpose of this study was to determine the factors associated with mortality, intubation, LOS, and costs in children admitted for asthma exacerbations between 2007 and 2011 in a nationally representative sample in the US, including whether or not racial and ethnic disparities or the presence of specific diagnoses are independent predictors of these outcomes.

Methods

Data Source

We analyzed data from the 2007 to 2011 Nationwide Inpatient Sample (NIS), the largest all-payer publicly available inpatient data set, which is part of the Healthcare Cost and Utilization Project of the Agency for Healthcare Quality and Research. The NIS consists of a representative stratified sample of approximately 20% of US hospital admissions; the complex sampling design allows for the calculation of national estimates.¹⁸ Given the deidentified nature

of the data present in the NIS, this study did not meet the definition for human subjects research and was exempt from full review by the New York University School of Medicine Institutional Review Board.

Subjects

Patient records were included in the analysis if they had a primary discharge diagnosis of asthma by *International Classification of Diseases, Ninth Revision* (ICD-9) code 493.* and were 2 to 18 years old. Patients with asthma as a secondary diagnosis were not included. There were no additional exclusion criteria.

Measures

The primary predictor variable of interest was race/ethnicity (a single category in the NIS, with “Hispanic” defining anyone identified as being of Hispanic ethnicity in the data set). Data were also collected on additional possible indicators of income disparities (insurance type and median household income quartile [income ranges vary by year¹⁸; 2007 ranges used for simplicity] by patient zip code [patient location by zip code is not available; note that income]), demographic factors (sex, age [grouped as ages 2 to 5, 6 to 12, and 13 to 18 years]), hospital-level characteristics (region, rural/urban location, teaching status, and bed size [small, medium, or large; definition differs depending on location and teaching status as defined by HCUP¹⁸]), and hospitalization timing (weekend admission, year [as a continuous variable], and discharge quarter [January to March, April to June, July to September, October to December]). Data on presence of several acute (pneumonia, influenza, and Respiratory Syncytial Virus [RSV]) as well as chronic (obesity, obstructive sleep apnea [OSA], allergic rhinitis, eczema, mood disorders, gastroesophageal reflux disease, and tobacco addiction [coded in the patient]) secondary

diagnoses chosen based on prior studies showing associations with presence of asthma or worse asthma severity^{14–16} were also collected (see appendix for secondary diagnosis coding criteria).

The outcome variables of interest were mortality, intubation (defined by presence of ICD-9 procedural codes for intubation/mechanical intubation: 96.04, 96.70, 96.71, 96.72), LOS, and total costs. To determine estimated total costs of hospitalization, charges were converted to costs using NIS Cost-to-Charge data files.¹⁹ Group-weighted average cost-to-charge ratios were used in the analysis.

Analysis

Bivariate and multivariable regression analyses were performed to determine whether any of the above-mentioned predictor variables were associated with the outcomes of mortality, intubation, LOS, and costs. LOS was also included as a predictor in the multivariable regressions in which cost was the outcome. In addition, missing data for all predictor variables were reassigned to a separate missing variable category and analyzed to determine if missing data predicted any of the outcomes. Logistic and linear regressions were utilized for dichotomous (mortality and intubation) and continuous (LOS and costs) outcomes, respectively. Because LOS and cost distributions were skewed, the data were natural log-transformed for all bivariate and multivariable linear regression analyses. We then applied Duan's method to retransform all linear regression results from the log to the original scale.²⁰ A p value less than 0.05 was considered significant in all cases. All statistical methods conducted respected the complex survey design using sample weighting in Stata SE 12.1 (StataCorp, College Station, TX).

Results

Descriptive Results

Data from 97,379 admissions met inclusion criteria and were analyzed; this represents a weighted statistical sample of 478,546 hospitalizations. Information related to patient demographics, hospital-level factors, hospitalization timing, and patient outcomes are presented in Table 1. The majority of patients were male and had public insurance. White and black children each represented approximately 30% of the sample, while 18.1% of our sample was Hispanic. Over one third were in the lowest income quartile. Most of the admissions were to large, urban, teaching hospitals. A secondary diagnosis of pneumonia was present in 13.7% of hospitalizations. The frequency of other diagnoses of interest in asthmatic patients ranged from just under 1 to 3% of patients. Mortality rate was 0.03%, and 0.3% of patients required intubation. Median (Interquartile Range [IQR]) LOS was 2 (1 to 3) days. Median (IQR) costs were \$2,760 (\$1,860 to \$4,320).

Multivariable Analyses

Independent predictors of mortality from the multivariable regression analysis are presented in Table 2. After adjusting for all relevant predictor variables (see Table 2), Native American children had a higher odds of mortality than white children (Odds Ratio [OR] 11.15, 95% Confidence Interval [CI] 1.09 to 114.49, $p = 0.042$). Race/ethnicity was otherwise not an independent predictor of mortality. Older age group (13 to 18 years old) and West region were also independently associated with mortality (Table 2).

Independent predictors of intubation are presented in Table 3. Hispanic children were less likely than white children to be intubated (OR 0.59, 95% CI 0.36 to 0.94, $p = 0.028$). Race/ethnicity was otherwise not an independent predictor of intubation. Older age, several secondary diagnoses (pneumonia, influenza, RSV, OSA, tobacco addiction), West region,

urban/teaching hospital status, and large bed size were all independently associated with intubation (Table 3).

Multivariable analyses for LOS are presented in Table 4. On average, Asians/Pacific Islanders had 0.12 fewer hospital days than white children (95% CI -0.24 to -0.02 days, $p = 0.022$). Race/ethnicity was otherwise not independently associated with LOS. Children with private insurance had 0.17 fewer hospital days compared to those with public insurance (95% CI -0.20 to -0.14 days, $p < 0.001$). Children living in zip codes with the highest average income (\$63,000+) had 0.30 fewer hospital days compared to the lowest income (\$1 to \$38,999) group (95% CI -0.36 to -0.22 days, $p < 0.001$) (Table 4). Other independent predictors of longer LOS included older age group, female sex, presence of several secondary diagnoses (including pneumonia, influenza, RSV, and several chronic diagnoses), urban/nonteaching hospitals (compared to rural hospitals), and large bed size.

Multivariable analyses for costs (adjusted for LOS) are presented in Table 4. Significantly higher costs were seen in black (increment \$272, 95% CI \$128 to \$467, $p < 0.001$), Hispanic (increment \$316, 95% CI \$174 to \$504, $p < 0.001$), and Asian (increment \$218, 95% CI \$34 to \$471, $p = 0.016$) children compared to white children. Average cost of admission was \$291 higher in children living in the highest compared to the lowest income areas (95% CI \$102 to \$549, $p = 0.001$). There were no significant differences in costs between children with private and public insurance; however, total costs were lower in the “other” insurance group compared to those with public insurance (Table 4). Other independent predictors of higher costs included older age groups, female sex, several secondary diagnoses, and LOS. Independent predictors of lower costs included urban/non-teaching hospitals (compared to rural and medium and large bed

size (compared to small bed size). Finally, missing median income by zip code data was associated with lower costs, while missing sex data was associated with higher costs (Table 4).

Results of bivariate analyses for all outcomes are provided in an appendix.

Discussion

This large national study of disparities among pediatric inpatients with a primary diagnosis of asthma found that Native American children had higher odds of inpatient death compared to white children and that otherwise, race and ethnicity were not independent predictors of inpatient asthma mortality. In terms of measures of inpatient morbidity, Hispanic children were less likely to be intubated than white children. LOS was longer in children with public insurance and living in low income areas and shorter in Asian children than whites. Total costs were higher in black, Hispanic, and Asian children and in children living in high income areas. The results of this study support previous research from a decade ago showing that mortality rates in black and Hispanic children who are sick enough to be admitted for asthma exacerbations are not significantly different than rates in white children.^{13,17}

While this study did not find a difference in mortality rates for black and Hispanic compared to white children admitted for asthma exacerbations, there are a few important caveats. The low inpatient asthma mortality rates seen in this and other studies^{15,19} make it difficult to detect statistically significant black and Hispanic disparities in mortality when important differences may actually exist. In addition, white and black children each represented approximately 30% of the sample, despite the white population in the US being close to six times the size of the black population according to recent census data.²³ So while there were no differences in mortality rates among black compared to white children already admitted to the hospital for asthma, our results would support previous studies showing that combined inpatient

and outpatient asthma mortality rates in US children are higher in black compared to white children.^{2,3} It is also possible that these overall racial disparities in mortality may be due to an excess of deaths occurring in the outpatient and prehospital/ED settings.

Statistical disparities in inpatient mortality were identified among Native American children, a group that is less well-studied. Native Americans do experience substantial morbidity from asthma with higher prevalence and similar hospitalization rates in some subgroups compared to white children.²²⁻²⁴ Prior studies have indicated that many barriers exist to management of asthma in Native Americans, including poor access to care, poor understanding of asthma, improper use of controller medications, cultural factors, and difficulty communicating with providers,²⁴⁻²⁶ suggesting that these children have more severe disease, and supporting the higher inpatient mortality rate found in this study. Finally while higher levels of mortality in Native Americans would be consistent with outpatient barriers and disparities experienced in this group, the large confidence interval (given the small number of subjects and the rare outcome of mortality) and p value just below 0.05 limit the strength of this conclusion, so further study of Native American inpatient asthma outcomes is warranted.

Costs (adjusted for LOS) were close to \$300 higher in black and Hispanic children compared to white children in our multivariable analysis even though other adverse outcomes such as longer LOS and mortality were not higher in these minority groups. It has been well established that there are substantial disparities between these groups and white children in the outpatient setting with higher rates of death,^{2,3} more missed school days,¹⁰ worse access to care,^{7,11} and improper controller and rescue medication use.^{6,10,12} It is possible that these children require additional treatment and workup upon arrival to the ED to compensate for some of these outpatient disparities, which may lead to higher costs. Given the large numbers of black and

Hispanic children who are admitted for asthma each year,² these higher costs are likely to have substantial economic impact. Conversely, Hispanic children had significantly lower odds of intubation compared to white children. One could speculate that given the increased mortality seen in Hispanic children in the outpatient setting^{2,3} that these sicker children are not surviving to make it to the hospital for intubation. It is also possible that Hispanic children are presenting to hospitals that are less likely to intubate or that there are other factors not accounted for in the NIS.

Average adjusted costs were about \$200 higher in Asians/Pacific Islanders compared to white children. Other studies have established that there are cultural barriers affecting Asian families and that Asian families have less asthma knowledge compared to whites.^{27,28} Asthma may also be underdiagnosed in some Asians.²⁸ It is possible that Asians may be sicker in the outpatient setting and the higher costs may be associated with more treatment early on in the hospital course to compensate for these disparities. One study examining pediatric asthma admissions in California found that Asians were more likely to have experienced an adverse outcome (in this case defined by cardiopulmonary resuscitation, intubation, or death) compared to white children,²⁹ so the presence of disparities in our study was not surprising. Conversely, Asian children had a shorter LOS compared to white children in our study. It is possible that there are factors that are not being measured that account for the discrepancy between LOS and costs in Asian children. Given the limited number of studies focusing on asthma in Asian children, additional research is needed.

In addition to the racial and ethnic disparities observed, our study also found associations between the presence of several acute and chronic comorbidities and worse outcomes in children admitted with asthma exacerbations, adding to a prior national study that showed that a higher

number of chronic conditions was associated with increased mortality.¹³ In our study, a secondary diagnosis of pneumonia or presence of RSV was associated with a two to three fold increased odds of intubation, more than an additional day added on to LOS, and added additional costs. These data support previous studies showing that children with asthma admitted to California hospitals were more likely to experience adverse outcomes if they also had a diagnosis of pneumonia.²⁹ Diagnoses of influenza, OSA, tobacco addiction, allergic rhinitis, and gastroesophageal reflux disease all were associated increased rates of intubation, longer LOS, and/or higher costs, which supports prior studies showing worse asthma outcomes in people with these diagnoses.^{14–16} The effect of secondary diagnoses on mortality could not be assessed, as there were very few deaths, and coding of secondary diagnoses was also less likely among those patients who died. However secondary diagnosis data should be interpreted with caution. The absence of a diagnosis of pneumonia or a viral illness in an administrative data set signifies only that it was not coded for. As such, these diagnoses could be undercoded in children who are not as ill.

This study has limitations. Race/ethnicity data were missing for 16% of the subjects in this study; this is common in studies of large administrative data sets.³⁰ Conclusions based on race/ethnicity data are only applicable for those states that report race/ethnicity. Several states do not report race/ethnicity data to the NIS.¹⁸ It is possible that the race/ethnicity of children in these states could have affected our analysis; however, missing race/ethnicity was not an independent predictor of any of the outcomes studied when it was analyzed as a separate category. In addition, the variability in how data are collected and coded for in administrative data sets leads to some limitations in reliability of the data. Additional studies are therefore needed in which race/ethnicity as well as outcome data on mortality/morbidity are collected prospectively in a

standardized manner to minimize any bias presented by missing or retrospectively collected data.” Of note, when sex and income quartile by zip code data were missing, costs were respectively \$677 higher and \$502 lower on average. It is unclear why missing income and sex data would be associated with costs, indicating that sex and income data should be interpreted with caution. Costs were unexpectedly higher in patients living in zip codes with higher average income. Future studies could consider collecting socioeconomic status data prospectively. We also acknowledge that disease coded for as asthma in the 2 to 5-year-old age group may not represent the same clear diagnosis of asthma as seen in older children. Also other studies have established that asthma severity often differs within different subgroups of Asian, Native American, and Hispanic children; for example, outcomes are different for Mexican, Puerto Rican, and other Hispanic subgroups.^{24,31} This level of analysis is not available in administrative data sets. In addition, further research should examine additional factors associated with worse asthma outcomes in other settings that are not included in the NIS, such as the negative effects of air pollution^{32,33} or second-hand smoke and the association between poor outcomes and local viral activity patterns. It is possible that some of these factors may explain some of the differences in outcomes seen in different regions of the country (such as the higher odds of mortality and intubation seen in the West region compared to the Northeast). We also acknowledge that the selection of $p < 0.05$ as a cutoff or significance may have increased the chance of a type I error.

In conclusion, our study shows that black and Hispanic race/ethnicity are not independent predictors of inpatient asthma mortality, a finding consistent with studies of inpatient asthma mortality conducted one decade ago. The higher rate of combined outpatient and inpatient asthma mortality which previously has been reported in black and Hispanic children is therefore

likely to be due to factors occurring prior to hospitalization. A new finding in our study is the higher odds of inpatient death among Native Americans. Race and ethnicity had other varying effects on asthma morbidity, including intubation, LOS and costs. Several other factors, including the presence of several acute and chronic secondary diagnoses, also were associated with in-hospital morbidity. Recognition of risk factors predictive of mortality and morbidity from asthma exacerbations may allow for earlier and more effective treatment and should be the impetus for targeted interventions in groups with these risk factors; several studies have identified interventions that improve outcomes in Hispanic/black, low-income, and urban children.^{34,35} Future research should explore other factors, such as local air pollution data, not available for study in large data sets that may be associated with mortality and morbidity. In addition, the finding of apparent disparities in inpatient asthma outcomes suggests the need to expand the study of groups such as Native Americans and Asians.

Acknowledgements

This study was funded by HRSA 1 T32HP22238 DHHS, HRSA, Primary Care (NRSA T32: Postdoctoral Primary Care Research Training Program) and the NYU CTSA grant UL1TR000038 from the National Center for the Advancement of Translational Science (NCATS), NIH.

References

1. Akinbami LJ, Moorman JE, Liu X. Asthma prevalence, health care use, and mortality: United States, 2005-2009. *Natl Health Stat Report*. 2011;32:1-14.
2. Akinbami LJ, Moorman JE, Garbe PL, Sondik EJ. Status of childhood asthma in the United States, 1980-2007. *Pediatrics*. 2009;123 Suppl:S131-145.
3. Moorman JE, Zahran H, Truman BI, Molla MT. Current asthma prevalence - United States, 2006 - 2008. *Morb Mortal Wkly Rep CDC Heal Disparities Inequalities Rep - United States, 2011*. 2011;60:84-86.
4. Beck AF, Huang B, Simmons JM, et al. Role of financial and social hardships in asthma racial disparities. *Pediatrics*. 2014;133:431-9.
5. Boudreaux ED, Emond SD, Clark S, Camargo CA. Race/ethnicity and asthma among children presenting to the emergency department: differences in disease severity and management. *Pediatrics*. 2003;111:e615-e621.
6. Crocker D, Brown C, Moolenaar R, et al. Racial and ethnic disparities in asthma medication usage and health-care utilization: data from the National Asthma Survey. *Chest*. 2009;136:1063-1071.
7. Jones R, Lin S, Munsie JP, Radigan M, Hwang S-A. Racial/ethnic differences in asthma-related emergency department visits and hospitalizations among children with wheeze in Buffalo, New York. *J Asthma*. 2008;45:916-922.
8. Joseph C, Havstad S, Ownby D, Johnson C, Tilley B. Racial differences in emergency department use persist despite allergist visits and prescriptions filled for antiinflammatory medications. *J Allergy Clin Immunol*. 1998;101:484-490.
9. Flores G, Snowden-Bridon C, Torres S, et al. Urban minority children with asthma: substantial morbidity, compromised quality and access to specialists, and the importance of poverty and specialty care. *J Asthma*. 2009;46:392-8.
10. Lieu T, Lozano P, Finkelstein J. Racial/ethnic variation in asthma status and management practices among children in managed Medicaid. *Pediatrics*. 2002;109:857-865.
11. Piper CN, Glover S, Elder K, Baek J-D, Wilkinson L. Disparities in access to care among asthmatic children in relation to race and socioeconomic status. *J Child Health Care*. 2010;14:271-9.
12. Ortega A, Gergen P, Paltiel A. Impact of site of care, race, and Hispanic ethnicity on medication use for childhood asthma. *Pediatrics*. 2002;109:E1.
13. Krishnan V, Diette GB, Rand CS, et al. Mortality in patients hospitalized for asthma exacerbations in the United States. *Am J Respir Crit Care Med*. 2006;174:633-8.
14. Sumino K, O'Brian K, Bartle B, Al E. Coexisting chronic conditions associated with mortality and morbidity in adult patients with asthma. *J Asthma*. 2014;51:306-14.
15. Cazzola M, Calzetta L, Bettoncelli G, Al E. Asthma and comorbid medical illness. *Eur Respir J*. 2011;38:42-9.
16. Boulet L, Boulay M. Asthma-related comorbidities. *Expert Rev Respir Med*. 2011;5:377-393.
17. Fisher-Owens SA, Turenne WM, Chavanu K, Slonim AD. Racial disparities in children hospitalized with asthma at academic children's hospitals. *Pediatr Asthma Allergy Immunol*. 2006;19:162-171.
18. HCUP Nationwide Inpatient Sample (NIS). Healthcare Cost and Utilization Project (HCUP). Agency for Healthcare Research and Quality, Rockville, MD. 2007-2011.

- Available at: <https://www.hcup-us.ahrq.gov/nisoverview.jsp>.
19. HCUP Cost-to-Charge Ratio Files (CCR). Healthcare Cost and Utilization Project (HCUP). Agency for Healthcare Research and Quality, Rockville, MD. 2007-2011. Available at: <https://www.hcup-us.ahrq.gov/db/state/costtocharge.jsp>. Accessed March 2, 2015.
 20. Duan N. Smearing estimate: a nonparametric retransformation method. *J Am Stat Assoc*. 1983;78:605-610.
 21. United States Census Bureau. State & county quick facts. 2015. Available at: <http://quickfacts.census.gov/qfd/states/00000.html>.
 22. Singleton RJ, Holman RC, Cobb N, Al E. Asthma hospitalizations among American Indian and Alaska Native people and for the general US population. *Chest*. 2006;130:1554-1562.
 23. Mehal JM, Holman RC, Steiner CA, Bartholomew ML. Epidemiology of asthma hospitalizations among American Indian and Alaska Native people and the general United States population. *Chest*. 2015;146:624-632.
 24. Brim SN, Rudd RA, Funk RH, Callahan DB. Asthma prevalence among US children in underrepresented minority populations: American Indian/Alaska Native, Chinese, Filipino, and Asian Indian. *Pediatrics*. 2008;122:e217-e222.
 25. Rose D, Garwick A. Urban American Indian family caregivers' perceptions of barriers to management of childhood asthma. *J Pediatr Nurs*. 2003;18:2-11.
 26. Van Sickle D, Wright AL. Navajo perceptions of asthma and asthma medications: clinical implications. *Pediatrics*. 2001;108:E11.
 27. Lee AC, Brugge D, Phan L, Woodin M. A comparison of knowledge about asthma between Asians and non-Asians at two pediatric clinics. *J Immigr Minor Heal*. 2007;9:245-254.
 28. Greenfield RO, Lee AC, Tang R, Brugge D. Screening for asthma in Cantonese-speaking immigrant children. *BMC Public Health*. 2005;5:48.
 29. Calmes D, Leake BD, Carlisle DM. Adverse asthma outcomes among children hospitalized with asthma in California. *Pediatrics*. 1998;101:845-850.
 30. Krishnan V, Diette GB, Rand CS, et al. Mortality in patients hospitalized for asthma exacerbations in the United States. *Am J Respir Crit Care Med*. 2006;174(6):633-8.
 31. Davis AM, Kreutzer R, Lipsett M, King G, Shaikh N. Asthma prevalence in Hispanic and Asian American ethnic subgroups: results from the California Healthy Kids Survey. *Pediatrics*. 2006;118:e363-e370.
 32. Wilhelm M, Qian L, Ritz B. Outdoor air pollution, family and neighborhood environment, and asthma in LA FANS children. *Health Place*. 2009;15:25-36.
 33. Lee J-T, Son J-Y, Kim H, Kim S-Y. Effect of air pollution on asthma-related hospital admissions for children by socioeconomic status associated with area of residence. *Arch Environ Occup Health*. 2007;61:123-130.
 34. Self TH, Chrisman CR, Mason DL, Rumbak MJ. Reducing emergency department visits and hospitalizations in African American and Hispanic patients with asthma: a 15-year review. *J Asthma*. 2005;42:807-12.
 35. Coffman JM, Cabana MD, Halpin HA, Yelin EH. Effects of Asthma Education on Children's Use of Acute Care Services: A Meta-analysis. *Pediatrics*. 2008;121:575-586.

Table 1. Patient/hospital characteristics (n=97,379)

Variable	N (%)	Variable	N (%)
Race/Ethnicity		Region	
White	29,496 (30.3)	Northeast	23,378 (24.0)
Black	27,306 (28.0)	Midwest	17,665 (18.1)
Hispanic	17,628 (18.1)	South	39,646 (40.7)
Asian/Pacific Islander	1,787 (1.8)	West	16,690 (17.1)
Native American	726 (0.8)	Location/teaching status	
Other	4,542 (4.7)	Rural	10,705 (11.0)
Missing	15,894 (16.3)	Urban/non-teaching	27,307 (28.0)
Insurance		Urban/teaching	59,367 (61.0)
Public	51,934 (53.3)	Bed size‡	
Private	37,524 (38.5)	Small	13,302 (13.7)
Other	7,748 (8.0)	Medium	22,561 (23.2)
Missing	173 (0.2)	Large	61,516 (63.2)
Median income by zip code		Weekend admission	25,357 (26.0)
\$1 to \$38,999	36,377 (37.4)	Year	
\$39,999 to \$47,999	22,265 (22.9)	2007	20,048 (20.6)
\$48,000 to \$62,999	18,924 (19.4)	2008	18,323 (18.8)
\$63,000+	16,689 (17.1)	2009	22,143 (22.7)
Missing	3,124 (3.2)	2010	19,711 (20.2)
Median Age, years (IQR)	6 (3 to 9)	2011	17,154 (17.6)
Sex		Quarter	
Female	37,081 (38.1)	January to March	22,322 (22.9)
Male	58,973 (60.6)	April to June	18,756 (19.3)
Missing	1,325 (1.4)	July to September	20,246 (20.8)
Secondary Diagnoses		October to December	28,132 (28.9)
Pneumonia	13,314 (13.7)	Missing	7,923 (8.1)
Influenza	1,267 (1.3)	Outcomes	
RSV	1,352 (1.4)	Mortality	26 (0.03)
Obesity/Overweight	2,890 (3.0)	Intubated	290 (0.30)
Obstructive Sleep Apnea	830 (0.9)	Median LOS (IQR), days	2 (1 to 3)
Mood Disorders	750 (0.8)	Median Total Costs (IQR)	\$2950 (\$1990-4610)
Tobacco Addiction	752 (0.8)		
Allergic Rhinitis	1,787 (1.8)		
Eczema	1,050 (1.1)		
Gastroesophageal Reflux	2,632 (2.7)		

IQR = Interquartile Range; LOS = Length of Stay; RSV = Respiratory Syncytial Virus

‡Definition differs depending on hospital location and teaching status¹⁸

Table 2. Independent Predictors of Inpatient Mortality†

Variable	Adjusted OR (95% CI)
Native American (Reference: White)	11.15 (1.09 to 114.49)*
13 to 18 year old age group (Reference: 2 to 5)	7.10 (1.93 to 26.05)**
West region (Reference: Northeast)	14.05 (1.62 to 121.51)*

*p<0.05, **p<0.01; OR = Odds Ratio; CI = Confidence Interval

†The model also adjusted for the following variables that did not reach statistical significance: race/ethnicity (black, Hispanic, and “other”), median income by zip code, age group, sex, diagnosis of pneumonia, region, hospital bed size, and timing of admission (by quarter of year, weekend vs. weekday, and year)

Table 3. Independent Predictors of Inpatient Intubation†

Variable	Adjusted OR (95% CI)
Hispanic (Reference: White)	0.59 (0.36 to 0.94)*
Age, years (Reference: 2 to 5)	
6 to 12	1.64 (1.15 to 2.34)**
13 to 18	4.31 (3.03 to 6.12)***
Secondary Diagnoses	
Pneumonia	2.09 (1.62 to 2.69)***
Influenza	2.96 (1.64 to 5.33)***
RSV	3.26 (1.72 to 6.17)***
Obstructive Sleep Apnea	3.68 (2.00 to 6.77)***
Tobacco Addiction	2.46 (1.28 to 4.72)**
West Region (Reference: Northeast)	1.77 (1.01 to 3.09)*
Urban/teaching hospital (Reference: Rural)	3.69 (1.96 to 6.95)***
Large bed size (Reference: Small) ‡	1.68 (1.13 to 2.48)**

*p<0.05, **p<0.01, ***p<0.001

OR = Odds Ratio; CI = Confidence Interval; RSV = Respiratory Syncytial Virus

† The model also adjusted for the following variables that did not reach statistical significance: race/ethnicity (black, Asian, Native American), and “other” insurance type, median income by zip code, sex, region, hospital bed size and location/teaching status, additional secondary diagnoses (obesity/overweight, mood disorders, allergic rhinitis, eczema, and gastroesophageal reflux), and timing of admission (by quarter of year, weekend vs. weekday, and year)

‡Definition differs depending on hospital location and teaching status¹⁸

Table 4. Independent Predictors of Inpatient Length of Stay (LOS) and Total Costs†

Variable	LOS Increment, days (95% CI)	Costs Increment (95% CI)
Race/Ethnicity (Reference: White)		
Black	NS	+\$310 (+\$145 to +\$531) ***
Hispanic	NS	+\$359 (+\$198 to +\$573) ***
Asian/Pacific Islander	-0.12 (-0.24 to -0.02)*	+\$248 (+\$39 to +\$535) *
Native American	NS	NS
Other	NS	NS
Missing	NS	NS
Insurance (Reference: Public)		
Private	-0.17 (-0.20 to -0.14)***	NS
Other	-0.20 (-0.25 to -0.14)***	-\$122 (-\$198 to -\$18) *
Missing	NS	NS
Median income by zip code (Ref: \$1 to \$38,999)		
\$39,999 to \$47,999	-0.09 (-0.13 to -0.04)**	NS
\$48,000 to \$62,999	-0.17 (-0.23 to -0.11)***	+\$197 (+\$53 to +\$388) **
\$63,000+	-0.30 (-0.36 to -0.22)***	+\$331 (+\$116 to +\$624) **
Missing	NS	-\$569 (-\$701 to -\$364) ***
Age, years (Reference: 2 to 5)		
6 to 12	+0.48 (+0.42 to +0.56)***	+\$275 (+\$189 to +\$384) ***
13 to 18	+0.70 (+0.60 to +0.82)***	+\$500 (+\$357 to +\$676) ***
Sex (Reference: Male)		
Female	+0.14 (+0.10 to +0.18)***	+\$45 (+\$14 to +\$85) **
Missing	NS	+\$767 (+\$280 to +\$1461) ***
Secondary Diagnoses		
Pneumonia	+1.21 (+1.06 to +1.37)***	+\$402 (+\$264 to +\$582)***
Influenza	+0.68 (+0.45 to +0.82)***	NS
RSV	+1.42 (+1.17 to +1.71)***	+\$289 (+\$113 to +\$523) ***
Obesity/Overweight	+0.40 (+0.27 to +0.54)***	+\$170 (+\$50 to +\$331) **
Obstructive Sleep Apnea	+1.22 (+0.90 to +1.59)***	NS
Mood Disorders	NS	NS
Tobacco Addiction	NS	+\$183 (+\$36 to +\$380)*
Allergic Rhinitis	+0.21 (+0.09 to +0.37)**	+\$153 (+\$14 to +\$335) *
Eczema	NS	NS
Gastroesophageal Reflux	+0.88 (+0.67 to +1.11)***	+\$237 (+\$96 to +\$424) ***
Region (Reference: Northeast)		
Midwest	-0.15 (-0.26 to -0.01)*	NS
South	NS	-\$598 (-\$740 to -\$367)***
West	NS	NS
Location/teaching status (Reference: Rural)		
Urban/non-teaching	+0.11 (+0.001 to +0.23)*	-\$200 (-\$327 to -\$15) *
Urban/teaching	NS	NS
Bed size (Reference: Small) ‡		
Medium	+0.20 (+0.003 to +0.44)*	-\$432 (-\$644 to -\$104) *
Large	+0.34 (+0.15 to +0.56)***	-\$443 (-\$638 to -\$136)**
Weekend (Reference: Weekday)	+0.05 (+0.02 to +0.09)**	+\$130 (+\$87 to +\$187) ***
Year (Continuous)	NS	NS
Quarter (Reference: January to March)		
April to June	-0.20 (-0.23 to -0.17)***	NS
July to September	-0.20 (-0.24 to -0.16)***	+\$123 (+\$65 to +\$202) ***
October to December	-0.19 (-0.22 to -0.15)***	+\$136 (+\$73 to +\$218) ***
Missing	NS	NS
LOS	---	+\$863 (+\$631 to +\$1149) ***

*p<0.05, **p<0.01, ***p<0.001; OR = Odds Ratio; CI = Confidence Interval NS=Not Significant; LOS = Length of Stay; RSV = Respiratory Syncytial Virus; †All variables in the table were included in the multivariable analysis

‡Definition differs depending on hospital location and teaching status¹⁸

Appendix 1. Diagnosis Codes

Variable	Code
Pneumonia	CCS 122
Influenza	CCS 123
RSV	ICD-9 079.6, 466.11, 480.1
Obesity/Overweight	ICD-9 278.0, 278.00, 278.01, 278.02
Obstructive Sleep Apnea	ICD-9 327.23
Mood Disorders	CCS 657
Tobacco Addiction	ICD-9 305.1
Allergic Rhinitis	ICD-9 477, 477.0, 477.1, 477.2, 477.8, 477.9
Eczema	ICD-9 691.8
Gastroesophageal Reflux	ICD-9 530.81

CCS = Healthcare Cost and Utilization Project Clinical Classification Software Code

RSV = Respiratory Syncytial Virus

ICD-9 = *International Classification of Diseases, Ninth Revision*

Appendix 2. Predictors of mortality from unadjusted analyses

Variable	OR (95% CI)
Race/ethnicity (Reference: White)	
Black	3.91 (0.95 to 16.16)
Hispanic	2.47 (0.43 to 13.99)
Native American	14.47 (1.34 to 156.12)*
Other	2.19 (0.18 to 25.82)
Missing	3.65 (0.67 to 19.81)
Median income by zip code (Reference: \$1 to \$38,999)	
\$39,999 to \$47,999	0.66 (0.23 to 1.92)
\$48,000 to \$62,999	1.06 (0.42 to 2.66)
\$63,000+	0.17 (0.02 to 1.40)
Missing	0.86 (0.10 to 7.48)
Age, years (Reference: 2 to 5)	
6 to 12	2.94 (0.91 to 9.56)
13 to 18	6.43 (1.60 to 25.78)**
Female (Reference: Male)	0.84 (0.41 to 1.75)
Pneumonia diagnosis	1.46 (0.59 to 3.61)
Region (Reference: Northeast)	
Midwest	10.18 (1.20 to 86.50)*
South	7.62 (0.97 to 59.60)
West	10.05 (1.20 to 83.84)*
Bed size (Reference: Small) ‡	
Medium	4.88 (0.58 to 40.93)
Large	4.52 (0.58 to 35.35)
Quarter (Reference: January to March)	
April to June	0.91 (0.25 to 3.37)
July to September	1.48 (0.42 to 5.18)
October to December	1.03 (0.27 to 3.90)
Missing	1.67 (0.38 to 7.39)
Weekend (Reference: Weekday)	1.55 (0.68 to 3.52)
Year	1.37 (0.99 to 1.97)

*p<0.05, **p<0.01; OR = Odds Ratio; CI = Confidence Interval

‡Definition differs depending on hospital location and teaching status¹⁸

Appendix 3. Predictors of intubation from unadjusted analyses

Variable	OR (95% CI)
Race/Ethnicity (Reference: White)	
Black	1.22 (0.90 to 1.68)
Hispanic	0.69 (0.46 to 1.05)
Asian/Pacific Islander	0.77 (0.27 to 2.16)
Native American	0.48 (0.07 to 3.26)
Other	1.23 (0.71 to 2.14)
Missing	0.91 (0.57 to 1.44)
Insurance (Reference: Public)	
Private	0.94 (0.72 to 1.24)
Other	0.73 (0.45 to 1.20)
Median income by zip code (Reference: \$1 to \$38,999)	
\$39,999 to \$47,999	1.21 (0.89 to 1.64)
\$48,000 to \$62,999	1.10 (0.90 to 1.51)
\$63,000+	1.01 (0.70 to 1.48)
Missing	0.41 (0.19 to 0.95)
Age, years (Reference: 2 to 5)	
6 to 12	1.53 (1.07 to 2.18)
13 to 18	3.97 (2.86 to 5.50)
Sex (Reference: Male)	
Female	0.86 (0.67 to 1.12)
Missing	0.22 (0.03 to 1.63)
Secondary Diagnoses	
Pneumonia	1.86 (1.40 to 2.33)***
Influenza	3.11 (1.69 to 5.73)***
RSV	2.93 (1.60 to 5.35)**
Obesity/Overweight	1.58 (0.91 to 2.74)
Obstructive Sleep Apnea	5.25 (2.98 to 9.27)***
Mood Disorders	1.88 (0.72 to 4.91)
Tobacco Addiction	5.33 (2.92 to 9.70)***
Allergic Rhinitis	1.52 (0.76 to 3.04)
Eczema	1.59 (0.60 to 4.18)
Gastroesophageal Reflux	2.18 (1.31 to 3.64)**
Region (Reference: Northeast)	
Midwest	1.29 (0.78 to 2.14)
South	0.98 (0.62 to 1.55)
West	1.22 (0.72 to 2.07)
Location/teaching status (Reference: Rural)	
Urban/non-teaching	1.02 (0.51 to 2.01)
Urban/teaching	3.30 (1.78 to 6.13)***
Bed size (Reference: Small) ‡	
Medium	1.39 (0.88 to 2.21)
Large	1.51 (1.01 to 2.26)*
Weekend (Reference: Weekday)	1.15 (0.88 to 1.51)
Year (Continuous)	1.09 (0.99 to 1.21)
Quarter (Reference: January to March)	
April to June	0.89 (0.65 to 1.22)
July to September	0.87 (0.63 to 1.18)
October to December	0.71 (0.52 to 0.98)
Missing	0.63 (0.34 to 1.16)

*p<0.05, **p<0.01, ***p<0.001; OR = Odds Ratio; CI = Confidence Interval; RSV = Respiratory Syncytial Virus

‡Definition differs depending on hospital location and teaching status¹⁸

Appendix 4. Predictors of length of stay from unadjusted analyses

Variable	Increment, days (95% CI)
Race/Ethnicity (Reference: White)	
Black	+0.12 (+0.03 to +0.22)**
Hispanic	+0.20 (+0.06 to +0.35)**
Asian/Pacific Islander	-0.31 (-0.46 to -0.15)***
Native American	+0.27 (+0.06 to +0.51)*
Other	+0.11 (-0.07 to +0.30)
Missing	-0.11 (-0.29 to +0.08)
Insurance (Reference: Public)	
Private	-0.30 (-0.35 to -0.23)***
Other	-0.24 (-0.33 to -0.14)***
Missing	-0.27 (-0.66 to +0.17)
Median income by zip code (Reference: \$ 1 to \$38,999)	
\$39,999 to \$47,999	-0.16 (-0.24 to -0.08)***
\$48,000 to \$62,999	-0.28 (-0.37 to -0.18)***
\$63,000+	-0.48 (-0.59 to -0.36)***
Missing	-0.14 (-0.39 to +0.14)
Age, years (Reference: 2 to 5)	
6 to 12	+0.49 (+0.43 to +0.54)***
13 to 18	+0.73 (+0.64 to +0.83)***
Sex (Reference: Male)	
Female	+0.25 (+0.21 to +0.29)***
Missing	-0.19 (-0.38 to +0.03)
Secondary Diagnoses	
Pneumonia	+1.44 (+1.31 to +1.56)***
Influenza	+0.86 (+0.65 to +1.09)***
RSV	+1.68 (1.42 to +1.98)***
Obesity/Overweight	+1.06 (0.90 to +1.24)***
Obstructive Sleep Apnea	+2.25 (1.82 to +2.72)***
Mood Disorders	+0.71 (0.46 to +0.99)***
Tobacco Addiction	+0.50 (0.27 to +0.75)***
Allergic Rhinitis	+0.40 (0.23 to +0.58)***
Eczema	+0.07 (-0.12 to +0.28)
Gastroesophageal Reflux	+1.36 (1.10 to +1.65)***
Region (Reference: Northeast)	
Midwest	-0.25 (-0.44 to -0.04)*
South	+0.07 (-0.11 to +0.26)
West	-0.08 (-0.26 to +0.12)
Location/teaching status (Reference: Rural)	
Urban/non-teaching	-0.009 (-0.14 to +0.13)
Urban/teaching	+0.03 (-0.11 to +0.18)
Bed size (Reference: Small) ‡	
Medium	+0.22 (-0.30 to +0.52)
Large	+0.37 (0.14 to +0.64)**
Weekend (Reference: Weekday)	+0.04 (0.001 to +0.08)*
Year (Continuous)	-0.002 (-0.045 to +0.004)
Quarter (Reference: January to March)	
April to June	-0.33 (-0.38 to -0.27)***
July to September	-0.40 (-0.46 to -0.34)***
October to December	-0.30 (-0.30 to -0.24)***
Missing	-0.01 (-0.22 to +2.76)

*p<0.05, **p<0.01, ***p<0.001; OR = Odds Ratio; CI = Confidence Interval; RSV = Respiratory Syncytial Virus

‡Definition differs depending on hospital location and teaching status¹⁸

Appendix 5. Predictors of total costs from unadjusted analyses

Variable	Increment (95% CI)
Race/Ethnicity (Reference: White)	
Black	+\$910 (+\$320 to +\$1560) **
Hispanic	+\$1820 (+\$1210 to +\$2510) ***
Asian/Pacific Islander	+\$1240 (+\$240 to +\$2400)*
Native American	-\$450 (-\$1250 to +\$500)
Other	+\$430 (-\$380 to +\$1370)
Missing	+\$340 (-\$450 to +\$1260)
Insurance (Reference: Public)	
Private	-\$20 (-\$420 to +\$420)
Other	-\$870 (-\$1270 to -\$430) ***
Missing	-\$1240 (-\$2590 to +\$470)
Median income by zip code (Reference: \$1 to \$38,999)	
\$39,999 to \$47,999	+\$100 (-\$300 to +\$540)
\$48,000 to \$62,999	+\$410 (-\$100 to +\$990)
\$63,000+	+\$670 (-\$80 to +\$1530)
Missing	-\$1430 (-\$1880 to -\$910) ***
Age, years (Reference: 2 to 5)	
6 to 12	+\$1430 (+\$1240 to +\$1620) ***
13 to 18	+\$2540 (+\$2190 to +\$2920) ***
Sex (Reference: Male)	
Female	+\$650 (+\$520 to +\$800)***
Missing	+\$3780 (+\$2020 to +\$5880) ***
Secondary Diagnoses	
Pneumonia	+\$3870 (+\$3350 to +\$4420) ***
Influenza	+\$2580 (+\$1750 to +\$3510) ***
RSV	+\$3520 (+\$2690 to +\$4460) ***
Obesity/Overweight	+\$3360 (+\$2610 to +\$4170) ***
Obstructive Sleep Apnea	+\$7220 (+\$5780 to +\$8850) ***
Mood Disorders	+\$2520 (+\$1610 to +\$3540) ***
Tobacco Addiction	+\$2400 (+\$1570 to +\$3340) ***
Allergic Rhinitis	+\$870 (+\$280 to +\$1540) **
Eczema	+\$540 (-\$260 to +\$1460)
Gastroesophageal Reflux	+\$3880 (+\$3100 to +\$4740) ***
Region (Reference: Northeast)	
Midwest	-\$1700 (-\$2710 to -\$280) *
South	-\$2560 (-\$3260 to -\$1570) ***
West	+\$200 (-\$1140 to +\$2020)
Location/teaching status (Reference: Rural)	
Urban/non-teaching	+\$310 (-\$300 to +\$1020)
Urban/teaching	+\$1950 (+\$1100 to +\$2950) ***
Bed size (Reference: Small) ‡	
Medium	-\$1070 (-\$2020 to +\$210)
Large	-\$1130 (-\$1980 to +\$40)
Weekend (Reference: Weekday)	+\$620 (+\$460 to +\$780) ***
Year (Continuous)	-\$11 (-\$230 to +\$220)
Quarter (Reference: January to March)	
April to June	-\$420 (-\$610 to -\$210) ***
July to September	-\$290 (-\$510 to -\$40) *
October to December	-\$60 (-\$280 to +\$190)
Missing	-\$1390 (-\$2300 to -\$280) ***
LOS	+\$900 (+\$700 to +\$1130) ***

*p<0.05, **p<0.01, ***p<0.001; OR = Odds Ratio; CI = Confidence Interval; LOS = Length of Stay;

RSV = Respiratory Syncytial Virus

‡Definition differs depending on hospital location and teaching status¹⁸