


Journal Pre-proof

Peanut Allergy Diagnosis- a 2020 Practice Parameter Update, Systematic Review, and GRADE Analysis

Matthew Greenhawt, MD, MBA, MSc, Marcus Shaker, MD, MS, Julie Wang, MD, John J. Oppenheimer, MD, Scott Sicherer, MD, Corinne Keet, MD, PhD, Keri Swaggart, MSL, Matthew Rank, MD, Jay M. Portnoy, MD, Jonathan Bernstein, MD, Derek K. Chu, MD, Ph.D, Chitra Dinakar, MD, David Golden, MD, Carolyn Horner, MD, David M. Lang, MD, Eddy S. Lang, MD, David A. Khan, MD, Jay Lieberman, MD, David Stukus, MD, Dana Wallace, MD



PII: S0091-6749(20)31137-4

DOI: <https://doi.org/10.1016/j.jaci.2020.07.031>

Reference: YMAI 14705

To appear in: *Journal of Allergy and Clinical Immunology*

Received Date: 21 April 2020

Revised Date: 14 July 2020

Accepted Date: 17 July 2020

Please cite this article as: Greenhawt M, Shaker M, Wang J, Oppenheimer JJ, Sicherer S, Keet C, Swaggart K, Rank M, Portnoy JM, Bernstein J, Chu DK, Dinakar C, Golden D, Horner C, Lang DM, Lang ES, Khan DA, Lieberman J, Stukus D, Wallace D, Peanut Allergy Diagnosis- a 2020 Practice Parameter Update, Systematic Review, and GRADE Analysis, *Journal of Allergy and Clinical Immunology* (2020), doi: <https://doi.org/10.1016/j.jaci.2020.07.031>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Inc. on behalf of the American Academy of Allergy, Asthma & Immunology.

Peanut Allergy Diagnosis- a 2020 Practice Parameter Update, Systematic Review, and GRADE Analysis

Matthew Greenhawt, MD,MBA,MSc,¹ Marcus Shaker, MD, MS² Julie Wang, MD,³ John J Oppenheimer, MD,⁴ Scott Sicherer, MD,³ Corinne Keet, MD, PhD,⁵ Keri Swaggart, MSL⁶, Matthew Rank, MD,⁷ Jay M Portnoy, MD,⁸ Jonathan Bernstein, MD⁹, Derek K Chu, MD,Ph.D¹⁰, Chitra Dinakar, MD¹¹, David Golden, MD¹², Carolyn Horner, MD¹³, David M. Lang, MD¹⁴, Eddy S. Lang, MD¹⁵, David A. Khan, MD¹⁶, Jay Lieberman,MD¹⁷, David Stukus, MD¹⁸, and Dana Wallace, MD¹⁹.

¹ Section of Allergy and Immunology, Department of Pediatrics, Children's Hospital Colorado, University of Colorado School of Medicine, Aurora, CO

² Dartmouth Geisel School of Medicine, Dartmouth-Hitchcock Medical Center, Section of Allergy and Immunology, Lebanon, NH

³ Division of Pediatric Allergy and Immunology, Department of Pediatrics, Icahn School of Medicine at Mount Sinai and the Jaffe Food Allergy Institute, New York, NY

⁴ Department of Internal Medicine, New Jersey Medical School, Morristown, NJ

⁵Division of Pediatric Allergy and Immunology, Johns Hopkins School of Medicine, Baltimore, MD

⁶Library Services, Children's Mercy Hospital, Kansas City, MO

⁷Division of Allergy, Asthma, and Clinical Immunology, Mayo Clinic, Scottsdale, AZ and Division of Pulmonology Phoenix Children's Hospital Phoenix, AZ

⁸Division of Allergy, Asthma & Immunology, Children's Mercy Hospital, Kansas City, MO

⁹Division of Immunology, Department of Internal Medicine, University of Cincinnati, Cincinnati, Ohio

¹⁰Department of Medicine, McMaster University, Hamilton, Ontario, Canada; Department of Health Research Methods, Evidence & Impact, McMaster University, Hamilton, Ontario, Canada; and The Research Institute of St. Joe's Hamilton, Hamilton, Ontario, Canada

¹¹Division of Allergy and Asthma, Stanford University School of Medicine, Palo Alto, CA

¹²Department of Allergy-Clinical Immunology John Hopkins, Baltimore, MD

¹³Division of Allergy, Immunology, and Pulmonary Medicine, Washington University School of Medicine, St. Louis MO

¹⁴Department of Allergy and Clinical Immunology, Cleveland Clinic, Cleveland, OH

¹⁵ Department of Emergency Medicine, Cumming School of Medicine, University of Calgary, Calgary, Alberta, Canada

¹⁶Division of Allergy & Immunology, Department of Medicine, University of Texas Southwestern Medical Center, Dallas, TX

¹⁷Division of Allergy and Immunology, The University of Tennessee, Memphis, TN

¹⁸Division of Allergy and Immunology, The Ohio State University College of Medicine, Columbus, OH

¹⁹Nova Southeastern University College of Allopathic Medicine, Fort Lauderdale, FL

Corresponding Author:

Matthew Greenhawt, MD, MBA, MSc

Section of Allergy and Immunology

Food Challenge and Research Unit

Children's Hospital Colorado

University of Colorado School of Medicine

13123 E. 16th Ave

Aurora, CO 80045

Phone #: 720-777-2264

Matthew.Greenhawt@childrenscolorado.org

Key words: peanut allergy; Diagnosis; Practice Parameter Update; systematic review; GRADE Analysis; Ara h 2; peanut components; skin prick testing; serologic IgE testing; likelihood ratio; evidence to recommendation; meta-analysis; cost-effectiveness analysis

Abbreviations: Food and Drug Administration (FDA); GRADE (Grading of Recommendations, Assessment, Development, and Evaluation); PICO (Population Intervention Comparator Outcome); skin prick test (SPT); oral food challenge (OFC); food protein induced enterocolitis syndrome (FPIES); National Institutes of Allergy and Infectious Disease (NIAID); quality-adjusted life years (QALY); likelihood ratio (LR); positive predictive value (PPV)

Contact Person: Joint Task Force on Practice Parameters (JTFPP) Liaison: Peris Flagg (American Academy of Allergy, Asthma, and Immunology, 555 E. Wells Street, Suite 1100, Milwaukee, WI 53202, pflagg@aaaai.org); JTFPP.allergy@gmail.com

Previously published Practice Parameters of the Joint Task Force on Practice Parameters for Allergy & Immunology are also available at <http://www.allergyparameters.org>; <http://www.AAAAI.org>, and <http://www.ACAAI.org>.

Disclosures: The JTFPP members and work group members' conflict of interest disclosure forms can be found at www.allergyparameters.org. Matthew Greenhawt has received financial support from Aquestive, Aimmune, Merck, Allergenics, Allergy Therapeutics, Sanofi Genzyme, Genentech, GSK, Merck, Aravax, Protalix, Before Brands, the Institute for Clinical and Economic Review, American College of Allergy, Asthma, and Immunology, DBV, Intromune, and is supported by the Agency of Healthcare Research and Quality; has served on the advisory board of International FPIES Association, the Asthma and Allergy Foundation of America, and the National Peanut Board; is Associate Editor of the Annals of Allergy, Asthma, and Immunology

Marcus Shaker is a member of the Joint Taskforce on Allergy Practice Parameters; has a family member who is CEO of Altrix Medical; serves on the Editorial Board of the Journal of Food Allergy and the Annals of Allergy, Asthma, and Immunology.

Julie Wang has received financial support from ALK Abello, Regeneron, DBV, Aimmune; is an UpToDate author; serves on the Executive Committee of the American Academy of Pediatrics Section on Allergy and Immunology; serves as Vice Chair of the AAAAI Anaphylaxis, Dermatitis, Drug Allergy Interest Section.

John Oppenheimer has received financial support from DBV, TEVA, GlaxoSmithKline adjudication/data safety monitoring board, AstraZeneca, Novartis, and Sanofi; is Associate Editor of the Annals of Allergy, Asthma, and Immunology and AllergyWatch, an American Board of Internal Medicine (ABIM) Council Member and American Board of Allergy and Immunology Liaison to the ABIM, UpToDate Reviewer, American College of Chest Physicians Cough Guideline Committee Member, and WebMD Editor.

Scott Sicherer has received financial support from FARE Clinical Center, John Hopkins UPress, HAL Allergy, AAAAI, UpToDate, and NIAID; is a Medical Advisor at International FPIES, Member of the Executive Committee at the American Academy of Pediatrics.

Corinne Keet received royalties from UpToDate, is an Associate Editor for the Journal of Allergy and Clinical Immunology, and is a Board Member for the American Board of Allergy and Immunology

Keri Swaggart has no conflicts of interest to report.

Matthew Rank has received financial support from the ACAAI, NIH, and Levin Family Foundation; has served as Chair of the AAAAI HEDQ Interest Section, and is Research Director of the Phoenix Children's Hospital Breathmobile.

Jay Portnoy has received financial support from Thermofisher, Kaleo, TEVA, Novartis, Hycor, and Boehringer-Ingelheim.

Jonathan Bernstein has received financial support from Sanofi Regeneron, AstraZeneca, Merck, Optinose, Takeda, CSL Behring, Biocryst, Pharming, the National Institutes of Health (NIH), Taylor Francis, INEOS; is Editor in Chief of the Journal of Asthma, INEOS Medical Immunosurveillance Director, Vice Chair and Lectureship Chair of the American Academy of Allergy, Asthma, and Immunology (AAAAI) Foundation, Chairman of AFI, American College of Asthma, Allergy, and Immunology (ACAAI) Asthma Chair, Scientific Chair, and Young Investigator Award Chair, and serves of the Board of Directors and Scientific Committee of Interasma.

Derek Chu is a CAAIF-CSACI-AllerGen Emerging Clinician-Scientist Research Fellow, supported by the Canadian Allergy, Asthma and Immunology Foundation (CAAIF), the Canadian Society of Allergy and Clinical Immunology (CSACI) and AllerGen NCE Inc. (the Allergy, Genes and Environment Network).

Chitra Dinakar has received financial support from Propeller Health, American College of Asthma, Allergy, and Immunology (stipend for Editorial Board of AllergyWatch), and the American Association of Allergists of Indian Origin; serves on the Board of Directors of the American Academy of Allergy, Asthma, and Immunology and on the Medical Advisory Board of Food Equity Initiative; is Assistant Editor of AllergyWatch.

David Golden has received financial support from Aquestive, Sandoz, ALK-Abello, , Genentech, Stallergenes-Greer, and UpToDate.

Caroline Horner has served as committee chair for the American Academy of Allergy, Asthma, and Asthma Diagnosis and Treatment Interest Section, Interest Section Coordinating Committee, and In-Training Exam Coordinating Committee.

David Lang is on the Editorial Board for Allergy and Asthma Proceedings, Topic Editor for DynaMed, Associate Editor for J Asthma, and Delegate to National Quality Forum representing the American Academy of Allergy, Asthma, and Immunology (AAAAI).

Eddy Lang has no conflicts of interest to report.

David Khan has received financial support from UpToDate and Aimmune; serves on the Board of Directors of the American Academy of Allergy, Asthma, and Immunology, American College of Asthma, Allergy, and Immunology (Chair of Literature Review, Co-Chair of Conjoint Board Review, Texas Allergy, Asthma, and Immunology Society Chair of Meetings Committee, and is Associate Editor of the Journal of Allergy and Clinical Immunology In Practice.

Jay Lieberman has received financial support from the American College of Asthma, Allergy, and Immunology, Aquestive, Aimmune, DBV, Biotest Pharma, and Regeneron; is Associate Editor of the Annals of Allergy, Asthma, and Immunology, Vice Chair for the American College of Asthma, Allergy, and Immunology Food Allergy Committee, and Medical Director for Food Allergy Alliance of the MidSouth.

David Stukus has received financial support from Aimmune, Before Brands, Abbott Nutrition, the American Academy of Pediatrics (AAP), ACAAI; has served as Committee Chair for the AAAAI and ACAAI.

Dana Wallace has received financial support from Mylan, Kaleo, Optinose, ALK, Bryan, and Sanofi

Peanut Allergy Diagnosis- a 2020 Practice Parameter Update, Systematic Review, and GRADE Analysis

Executive Summary

IgE mediated peanut allergy has an estimated prevalence of between 0.2-4.5%, depending on geographic area of the world and the methodology used for assessment. While the prevalence in the US appears to have tripled in a recent 10-year period, in the UK the prevalence seems to have plateaued over a similar period, denoting regional heterogeneity in such trends. Peanut allergy is associated with substantial economic and psychologic burden on families that is associated with poor quality of life and high anxiety related to the potential consequences of their child having a severe allergic reaction. Peanut allergy is often a severe and usually a lifelong allergy that is a leading cause of food-related anaphylaxis. There is one US Food and Drug Administration (FDA) approved peanut allergy treatment, and several other emerging treatments approaching consideration for FDA approval. However, presently peanut allergy is managed through peanut avoidance, and by carrying emergency medication such as auto-injectable epinephrine to treat symptoms that may arise from unintended ingestion.

Given this burden of disease and the consequences of diagnosis, it is important that peanut allergy be accurately diagnosed so that an appropriate treatment plan can be developed. However, a positive peanut test result is not always associated with clinical reactivity. This practice parameter addresses the diagnosis of IgE mediated peanut allergy, both in children and adults as pertaining to three fundamental questions (see text box 1). This parameter exclusively discusses IgE mediated peanut allergy and all references herein pertain to IgE mediated food allergy to peanut only, and not to peanut as a potential trigger in eosinophilic esophagitis or non-IgE mediated food allergy such as food protein induced enterocolitis syndrome. Similarly, emerging technologies such as Basophil Activation Testing were also not included in this analysis.

Diagnostic testing for peanut allergy is used to help make a diagnosis where there is suspicion of a peanut allergy based on the clinical history. Failure to make a correct diagnosis can result in either unnecessary avoidance in a non-allergic person, or erroneous guidance that the patient can safely ingest peanut *ad libitum* when there is in fact an allergy—situations that are both problematic. A correct diagnosis facilitates peanut avoidance and counseling when the patient is at risk of potential life-threatening complications of peanut allergy, and therefore is advised to carry epinephrine for use in case of symptomatic accidental ingestion. Alternatively, exclusion of peanut allergy allows peanut to be incorporated into the diet without concern, eliminating the burden of precautions and fear. Changes in peanut sensitization levels over time, compared to baseline, may be associated with whether the allergic individual is likely to be outgrowing their peanut allergy. Although previous research in patients with established peanut allergy reported clinical diagnostic cut-off points for >95% chance of reaction and for <50% chance of reaction to oral food challenge, these are not necessarily predictive of clinical outcomes in all settings and patients, as they are highly dependent on the baseline prevalence of peanut allergy in the particular population studied, which may vary. This document uses likelihood ratios as the main statistic in this analysis, given this metric is not dependent on a population prevalence of disease, and is more adaptable to individual clinical settings.

The expert panel developed the key (PICO, Population Intervention Comparator Outcome) questions to be addressed, and after systematic review of the literature (>1300 references searched), meta-analysis of the evidence, and GRADE (Grading of Recommendations, Assessment, Development, and Evaluation) analysis of the results (a well-established methodology for developing evidence-based guidelines, see table 1), made recommendations - all of which were conditional in strength, with very low certainty of

evidence. Thresholds for detection of sensitization were at 3mm for skin prick test (SPT), and 0.35 KU_s/L for both whole peanut sIgE and component-specific peanut sIgE, based on the most widely reported levels evaluated in the literature. Additional cut-offs were considered, their use would have posed significant limitation to the analysis, given very limited study numbers reporting these values. Extensive sensitivity analysis was performed to confirm the results.

The expert panel suggested that diagnostic testing for peanut allergy be used in patients with a high pre-test probability of peanut allergy, or prior to an oral food challenge for patients with moderate pre-test probability of peanut allergy, as a preference-sensitive choice, but discourages testing in patients with a low or very low pre-test probability of peanut allergy. If a single diagnostic test is to be used, testing for the Ara h 2 component (CPT code 86003, commercially available in the US under Thermo Fisher ordering reference code f423) would provide the most diagnostic accuracy, as determined by the more optimal positive/negative likelihood ratio among the presently available testing options, provided this becomes more commonly available as a stand-alone test, as opposed to being primarily offered as a panel by laboratories with other peanut components. The literature search did not provide patient-level data to determine the value of testing for peanut components in addition to skin prick test or sIgE to whole peanut to increase diagnostic accuracy, including isolated Ara h 2 in that context. The clinician should not use the results of a SPT, sIgE to whole peanut extract, or sIgE to peanut components to determine an allergy phenotype or to predict the severity of a future reaction (e.g., will the patient have anaphylaxis to peanut). An additional supplemental cost-effectiveness analysis of the potential testing options presented in the e-supplement confirms use of Ara h 2 as the optimal choice compared to peanut SPT and whole peanut sIgE. There remain important knowledge gaps and needs for well-designed studies to address these questions, as well as the need for patient-level data to be made available when reporting test sensitivity/specificity to enhance the ability to perform future meta-analysis that can explore different cut-off levels. These recommendations detailed below are summarized in table 2.

Question 1: Should diagnostic testing for peanut allergy be performed in adults and children with a history of suspected peanut allergy who are requesting evaluation for peanut allergy?

Recommendation 1a: We suggest in favor of diagnostic (skin prick or serum sIgE) testing for peanut allergy in patients with a 1) physician-judged high pre-test probability of peanut allergy, or 2)

prior to an oral food challenge for patients with moderate pre-test probability of peanut allergy, for both of whom shared decision-making has been employed to arrive at the final decision. **Conditional recommendation; Certainty of evidence: very low**

Recommendation 1b: We suggest against diagnostic testing in patients where there is low or very low pre-test probability of peanut allergy. **Conditional recommendation; Certainty of evidence: very low**

Discussion: This question was not searched in a systematic manner as the content experts were unaware of any single research study that addressed this question. The workgroup performed a Pubmed literature search that did not identify any articles that address this question, which by default limits the certainty of evidence. For this reason, the workgroup and JTFPP felt that it would therefore not be an appropriate utilization of valuable resources to conduct a librarian-conducted formal literature search. However, expert evidence was collected both from the content experts and the JTFPP. Expert evidence differs from expert opinion, in that the former does not include a judgment on the evidence and offers a systematic and transparent appraisal of the evidence, which differentiates this as an acceptable alternative to making a recommendation under GRADE. The guideline working group related that when evaluating their collective patient experiences, diagnostic testing could be of value to confirm peanut allergy in high-risk individuals for which an oral challenge might not be advisable or agreed to by patients, but also acknowledged that in a patient presenting with a classical history the diagnosis could be made on the basis of history alone without further testing in some circumstances. The expert panel related that they suggested an oral food challenge when there was a moderate probability of peanut allergy but that a large proportion of their patients may prefer a diagnostic test prior to the oral food challenge. Similarly, the collective personal experience of the expert panel was that diagnostic testing in patients with a low probability of peanut allergy (e.g., sibling has peanut allergy and patient has never ingested peanut) often identified patients who were sensitized but not truly allergic. Unfortunately, many of these patients refused an oral food challenge and likely avoided peanut unnecessarily.

These recommendations are in alignment with previous expert guidelines and practice parameters on food allergy diagnosis and management, which provide similar consensus regarding the indications for testing for the presence of food sensitization, including peanut, in evaluating a possible diagnosis of food allergy. While screening of foods in infants prior to initial food introduction is discouraged, testing to peanut in infants at high-risk for peanut allergy (under the very prescribed context of those infants with

either severe eczema and/or egg allergy) is the one notable exception, which was recommended prior to initial peanut introduction per the 2017 NIAID addendum guidelines.

Question 2a: In the patient presenting for evaluation of suspected peanut allergy, which of the three tests—SPT, sIgE to whole peanut, or Ara h2 would provide the highest diagnostic accuracy as determined by the more optimal positive/negative likelihood ratio?

Question 2b: In a patient presenting for evaluation of suspected peanut allergy, does testing for peanut components in addition to either SPT or sIgE to whole peanut increase the diagnostic accuracy?

Recommendation 2a: We suggest in favor of Ara h2 diagnostic testing in a patient presenting for evaluation of suspected peanut allergy for which a single diagnostic test is to be used, as Ara h2 would provide the best diagnostic accuracy as determined by virtue of more optimal positive/negative likelihood ratios. However, while Ara h 2 has the greatest specificity, it has lower sensitivity than SPT and sIgE, and in a patient with a high prior probability, the clinician may use Ara h 2, SPT, or sIgE to confirm the diagnosis of peanut allergy. **Conditional recommendation. Certainty of evidence: low.**

Recommendation 2b: We suggest against routine use of component testing in addition to either to skin prick test or sIgE to whole peanut to increase diagnostic accuracy. **Conditional recommendation. Certainty of evidence: very low**

Discussion: For GRADE analysis, Ara h 2 was compared to skin prick test and to sIgE to whole peanut for the diagnosis of peanut allergy. The literature search did not provide patient-level data to determine the value of testing for peanut components in addition to or reflexively with skin prick test or sIgE to whole peanut to increase diagnostic accuracy. In addition, expert evidence was not available to assist in answering this question. Thus, the use and value of components, including reflexive use of Ara h 2, remains a knowledge gap. There is an unclear utility for measuring sIgE to any other commercially available peanut components given the limited available data on performance of components beyond Ara h 2. Further research is needed to clarify the value of tandem testing, particular in regards to Ara h 2, Ara h 6, and Ara h 8. While Ara h 2 had the greatest specificity in confirming the diagnosis, it had lower

sensitivity when compared with SPT or sIgE. In evaluating diagnostic accuracy, the summary receiving operating characteristic (ROC) curves demonstrated greatest area under the curve (AUC) for Ara h 2 (0.92) when compared to SPT (0.89) and sIgE (0.81). We caution the clinician that despite undetectable sensitization on SPT, sIgE, or Ara h2 testing, that there is a small possibility the individual could still be peanut allergic, and similarly, that detection of sensitization does not always infer clinical allergy. If clinical suspicion remains elevated, further evaluation through an oral food challenge, is potentially indicated.

Question 3: In the patient presenting for evaluation of suspected peanut allergy, can the results of a diagnostic test be used to predict the severity of a future allergic reaction?

Recommendation 3: We suggest against the clinician using the results of a SPT, sIgE to whole peanut extract, or sIgE to peanut components to determine the severity of an allergy phenotype or to predict the severity of a future reaction. **Conditional recommendation. Certainty of evidence: very low.**

Discussion: There was inadequate patient-level data to formulate a GRADE recommendation on the use of a peanut diagnostic test for predicting the severity of a future allergic reaction across a continuous range of test result values; however, dichotomous cut-off values of 10mm (SPT), 50 KU_A/L, and 2 KU_A/L (Ara h 2) demonstrated low sensitivity and specificity for a future severe reaction.

Abstract: Given the burden of disease and the consequences of a diagnosis of peanut allergy, it is important that peanut allergy be accurately diagnosed so that an appropriate treatment plan can be developed. However, a test that indicates there is peanut sensitization present (e.g., a “positive” test) is not always associated with clinical reactivity. This practice parameter addresses the diagnosis of IgE mediated peanut allergy, both in children and adults as pertaining to three fundamental questions, and based upon the systematic reviews and meta-analysis, makes recommendations for the clinician who is evaluating a patient for peanut allergy. These questions relate to when diagnostic tests should be completed, which diagnostic tests to utilize, and the utility (or lack thereof) of diagnostic testing to predict the severity of a future allergic reaction to peanut.

Introduction:

The following document is a GRADE (Grading of Recommendations, Assessment, Development, and Evaluation) based practice parameter for the use of diagnostic testing in evaluating patients with peanut allergy. This document is divided into several sections: 1) a narrative review primer to provide background and context on peanut allergy and the principles of how to apply diagnostic testing for peanut allergy; 2) a meta-analysis and systematic review of peanut allergy diagnostic testing with GRADE recommendations (including supporting tables and figures); 3) an e-Supplement detailing the literature search terms and features of the included articles; 4) an eSupplement detailing a cost-effectiveness analysis of the use of the diagnostic tests; and 5) an eSupplement containing the GRADE evidence to recommendation framework, detailing the considerations that went into formulating the GRADE recommendation; .

Section 1: A Primer on the Diagnosis of Peanut Allergy

Prevalence of peanut allergy

In the general population, the prevalence of peanut allergy is approximately 1.5% when the diagnosis is based on oral food challenge (OFC) or highly convincing history, and 0.2% to 0.4% when it is based on OFC alone.¹ These values differ based on age, race, ethnicity, and geography, but the evidence is not available to precisely determine what those differences are. Recent Australian data representative of the greater Victoria province in one year olds suggests the rate of peanut allergy could be as high as 3%, with

as many as 23% of these cases resolving by age 4, and 31% by age 6.²⁻⁵ US estimates range between 1.4%-4.5%, based on various indirect methods including phone surveys, internet surveys, and analysis of clinical history and epinephrine prescribing patterns.⁶⁻¹⁰ As well, the prevalence of peanut allergy may change with age. Prevalence estimates also vary depending on how peanut allergy is defined. Many studies use peanut sensitization (at a particular level of detection) to define peanut allergy, while others accept a convincing history of a clinical reaction.^{6,11,12} However, the criterion standard is an oral food challenge in which a clear outcome based on peanut ingestion is determined.⁴ Unsurprisingly, reported prevalence rates are higher in studies that include patients diagnosed based on either peanut sensitization and/or a reported convincing clinical history compared to estimates derived from patients diagnosed objectively through OFC. However, there may be some ethical and practical concern in performing OFC for the purpose of confirming prevalence rates using this criterion standard in such aforementioned individuals who already have a clinical diagnosis.^{6,8} Understanding the prevalence rate of a specific allergy helps to determine the relative likelihood that any patient being evaluated could have the allergy, and sets the basis for interpreting any diagnostic test that may be able to infer likelihood of diagnosis through simple tools like Fagan nomograms.^{13,14} For review, the positive likelihood ratio (in food allergy) represents the percent of individuals with food allergy who have detectable sensitization, divided by the percentage of people without food allergy who also have detectable sensitization (and the negative likelihood ratio the converse). These are dependent on test sensitivity and specificity, and not dependent on disease prevalence, making them ideal for clinical decision-making across different settings. Use of likelihood ratios can be reviewed in more detail elsewhere.¹⁵ In this practice guideline, we emphasize the use of the likelihood ratio as the main tool for clinical decision-making given uncertainties in the prevalence of food allergy in any given clinical population. Therefore, it is essential for a clinician to understand how and when performing specific diagnostic tests would provide the highest (or lowest) utility, to help gauge when such tests would be of value in clinical decision-making.

Making the Diagnosis

Available diagnostic tests for assessing peanut sensitization

Peanut specific IgE can be assessed with either a skin prick test (SPT) or a serologic in-vitro (blood) test. SPT assesses the presence of sIgE through formation of a wheal and erythema following percutaneous introduction of the target allergen. SPTs are based on extracts of whole peanut and therefore do not provide information about sensitization to individual peanut proteins (peanut components), though

extracts of recombinant components have been studied in research situations. Prick-to-prick testing with ingestible peanut products (e.g., peanut butter, powder, or kernels) as an alternative to testing with peanut extracts has been advocated by some, but the reproducibility, validity and reliability of this procedure is not established as a marker of sensitization, and this additional test in combination with the clinical history has uncertain value for clinical decision-making.¹³

A multitude of in-vitro tests for specific IgE are available using a variety of technologies. Modern-day serologic IgE tests rely on allergens that are attached to a solid phase substrate and detect IgE bound to those allergens using anti-human IgE antibodies conjugated to enzymes that create a colored (enzyme-linked immunosorbent assay or ELISA) or fluorescent (fluorescent enzyme immunoassay or FEIA) product. There also are technologies that measure the capture of specific IgE bound to allergen in liquid phase with subsequent detection using an appropriate enzyme-substrate. The amount of sIgE is determined by comparing the dilution curves of the unknown samples with a calibration curve based on samples with known sIgE.¹³ Non-specific IgE binding resulting in false positive results (e.g., falsely indicating sensitization) is a potential risk when samples are assessed from patients that are known to have high total IgE levels, but is accounted for by the manufacturer in how the instruments are calibrated.¹¹ Generally, these tests are considered to be quantitative and to have a relatively low coefficient of variance (e.g., approximately 5%). Most commercially available tests for peanut-specific IgE measure sIgE directed at an extract of whole peanut, similar to what is used in skin testing. However, most allergens contain multiple epitopes, each of which may be associated with the ability to specifically bind IgE, and the potential for resulting distinct symptom patterns.¹⁶ Patients may be sensitized to one or more components, which represent major allergens within peanut that IgE can bind to (such as the major allergens Ara h 1, Ara h 2, or Ara h 3; Ara=arachis hypogaeae, the Latin name for peanut, and major allergens are named based on their Latin names in the order of their discovery). There are now commercially available tests to measure select peanut components. Components are not available for skin testing outside of the research setting.¹⁶

Evaluation of Suspected Peanut Allergy

To properly use any allergy diagnostic test to evaluate for possible peanut allergy, the pre-test probability must be determined, which is accomplished through taking a comprehensive history.^{13,17} Typically, patients present to a clinician for an evaluation of a suspected history of peanut allergy, usually having experienced symptoms (in some form) believed to be attributable to peanut ingestion, which represents a situation in which there is high pre-test probability. However, sometimes tests are performed to evaluate individuals without such a history (possibly as part of a diagnostic testing panel), such as someone who has never eaten peanut before, or even in individuals who eat peanut and do not develop symptoms. As a general rule, persons who can eat peanut without developing symptoms are by definition not allergic and should not be tested for peanut allergy. Furthermore, food allergy testing should be conducted as narrowly as possible, and avoid searching for sensitization to other “common” allergens that may be offered as a several item panel if such allergens were not suspected in the history. Serologic testing can always be ordered “a la carte” as single items, and the allergy clinician should consider taking the time to educate non-specialist referring clinicians as to why use of panel tests is potentially problematic, or avoid use of testing when the history is not clearly indicative of a suspected food allergy.^{1,15} The situation is a bit more nuanced when considering an individual for testing who has never before ingested peanut, or in someone where oropharyngeal symptoms most consistent with pollen food allergy syndrome present distinctly, in the absence of other typical IgE mediated symptoms. In general, the pre-test probability for allergy would be very low, so that even if the test were detecting sensitization, the post-test probability would still remain low. However, there may be certain situations where a patient who has never before ingested peanut has other risk factors, such as moderate or severe eczema poorly responsive to therapy or a history of other food allergy, which may elevate the pre-test probability above that of the general population (but still lower than someone presenting with a history of a suspected reaction). In these scenarios, the clinician may desire to test these patients given the pre-test probability is potentially elevated or for more practical reasons such as if the test result will help the patient to make a decision whether they will introduce peanut. This is an example of preference-sensitive care, and requires delicate handling of the risks and benefits of all available options of how to manage detectable sensitization on testing with lower yet still elevated pre-test probability. With a detectable sensitization obtained in this context, performing an OFC (presuming both clinician and patient are willing) can be very helpful but needs to be balanced by how strongly the clinician and patient believe the positive test result indicates a high probability of clinical allergy and the understanding of the risk and downstream consequences of a conflating sensitization and allergy.^{17,18}

390

391 However, most cases do not present asymptotically. In assessing the clinical history, close
392 attention should be paid to the nature of the presenting symptoms (to make sure these are consistent with
393 mast-cell mediator release characteristic of an IgE mediated reaction), and the timing of when these
394 symptoms developed in association with known or suspected peanut ingestion. Symptoms typically
395 develop within minutes to up to about 2 hours if they are related to the peanut ingestion, and rarely
396 develop outside this time window. Non-classical symptoms or time courses that fall outside this interval
397 should decrease the suspicion of peanut allergy, though the clinician may have to consider the
398 significance of an eruption/exacerbation of atopic dermatitis in a child potentially associated with peanut
399 ingestion several hours after ingestion.^{17,18} Diagnostic testing in the patient with a reasonable pre-test
400 probability, established by eliciting a concerning or likely history of symptom development attributable to
401 peanut ingestion, can then be used to help determine the likelihood of a clinical allergy.^{13,14} This
402 describes a high-utility setting of how such tests can be used. One exception of note is food protein
403 induced enterocolitis syndrome (FPIES) to peanut. This is a non-IgE but immune-mediated reaction,
404 which has a delayed onset presentation (typically 1-4 hours after ingestion), resulting in protracted
405 vomiting to the point that lethargy and color change (e.g., cyanosis, pallor) ensue, and in rare instances,
406 bloody diarrhea may result at 6-12 hours. These symptoms represent this very distinct entity, which is
407 hallmarked by isolated GI involvement. FPIES is a clinical diagnosis, and testing for the presence of IgE
408 for peanut FPIES is not recommended. FPIES diagnosis and management is discussed elsewhere, and this
409 document does not refer to peanut FPIES management.¹⁹

410

411 *Potential Exceptions for Testing*

412 A major possible exception are high-risk infants being considered for early peanut introduction. As
413 specified in the 2017 NIAID Addendum Guidelines for the prevention of peanut allergy, a special case
414 may be made for screening infants who present with severe atopic dermatitis in the first 4-6 months of
415 life that is poorly controlled despite escalating skin care.²⁰ In formulating the Addendum Guidelines for
416 the Prevention of Peanut Allergy, an expert panel appointed by the National Institutes of Allergy and
417 Infectious Disease (NIAID) recommended that this presentation in these infants represents an elevated
418 pre-test probability of some likelihood of “pre-existing” peanut allergy (based on data from the Learning

Early About Peanut Allergy Study which used these particular risk factors). Therefore, in this highly specific subgroup the prior NIAID guideline did recommend strong consideration that either peanut SPT or sIgE testing be obtained and interpreted before early peanut introduction in these infants. However, outside of this very circumscribed group, there are otherwise no formal recommendations that any individual should have peanut SPT or sIgE testing before peanut introduction specifically as a screening measure for risk-assessment.²⁰

Historically, another potential exception involved testing children with moderate to severe atopic dermatitis to the common 8 food allergens (including peanut), even if these foods were never previously consumed. This practice reflected a concern that eczema is a precursor symptom of and a significant risk factor for developing food allergy, and represents a situation where the pre-test probability is potentially raised over that of the baseline general population to some degree. In these children, a diagnosis of allergy was typically made based on research that extrapolated positive predictive values taken from groups of children at referral centers with severe eczema who underwent oral food challenge.²¹ In recent years, this practice has largely fallen out of favor as there has been better understanding of a) the limitations of sensitization as a determinant of clinical allergy, b) the pathogenesis of atopic dermatitis occurring independently and not as a marker pathognomonic for undiagnosed food allergy, c) the risks of prolonged allergen avoidance as a factor that may paradoxically increase the risk of food allergy development, and d) the observation that indiscriminant “screening creep” was occurring in children without risk factors or overt symptoms and the predictive values were being used to establish “diagnosis” out of their very tightly established context.¹ The underlying properties of the diagnostic tests themselves make their use as diagnostic screening measures perilous, given they are poorly specific and of optimal utility in the setting of high pre-test probability. Asymptomatic, clinically irrelevant peanut sensitization is not uncommon.

Interpreting peanut allergy sensitization

Allergy testing only confirms or refutes the presence of sensitization, requires “clinical correlation” not unlike a radiographic image, and does not independently diagnose allergic disease. Pre-test probability can be translated to post-test odds, using the positive or negative likelihood ratios associated

with the sensitivity and specificity of these tests, which can then be used to provide a recommendation regarding diagnosis.^{13,14} Thus the presence/absence of sensitization increases or decreases the estimated likelihood that a patient may experience a reaction following peanut ingestion. The final probability of reaction is dependent both on the pre-test probability and the characteristics of the diagnostic test. While this can be translated using a Fagan nomogram,¹⁴ the process is rather intuitive in clinical practice in many situations. Individuals with a strong history (e.g., high pre-test probability) who are sensitized above a critical threshold can be more confidently diagnosed with peanut allergy, and a person with a non-specific/weak history (e.g. low pre-test probability) and a negative or equivocally positive test indicating the presence of sensitization can be more confidently assessed as not having peanut allergy. In individuals with more questionable histories with a less clear pre-test probability, the test positive or negative likelihood ratio then becomes more crucial in influencing the direction of the decision-making, and ultimately diagnostic confidence may be low enough that an oral food challenge (OFC) still may be necessary to definitively establish diagnosis.^{1,17,18,22}

Clinical Conundrums Related to Testing

As alluded to earlier, there are situations where the clinician may encounter a patient in whom testing was potentially inappropriately obtained, such as in a person with no risk-factors and no history of peanut ingestion leading to symptoms. These individuals may be peanut sensitized, but the sensitization is difficult to interpret given the lack of clinical data to determine context of the test value. Here we see two possible management choices. In clinical practice, many may follow prior data establishing positive predictive values (most representative of small populations of eczematous children undergoing OFC at a referral center)²¹ for large skin tests or elevated peanut sIgE that may result in a potential misdiagnosis of peanut allergy leading to unnecessary avoidance. Alternatively, this could be viewed as a situation where a test was obtained with low pre-test probability, requiring OFC to provide diagnostic clarity.²³ In some scenarios, where it is very likely that sensitization may not be associated with clinical reactivity, there may be benefit to performing a supervised OFC. A less challenging conundrum is the use of so-called “alternative tests” for peanut allergy that are becoming popular, and are frequently utilized by non board-certified allergists or marketed directly to patients to order for use at home without provider involvement. Testing for peanut-specific IgG4 in either the symptomatic or non-symptomatic patient is not indicated, and no role for IgG4 levels in the current diagnostic paradigm exists.^{24,25} The role of IgG4 is not well

understood, but in studies of food oral immunotherapy and pollen/venom immunotherapy, IgG4 levels to the allergen in question have been noted to increase as the patient becomes desensitized. As such, no defined association between allergic reactivity and IgG4 levels exists. In addition, a multitude of other non-validated alternative tests are utilized by alternative medicine practitioners but have no role in the diagnosis of peanut allergy. This includes Mediator Release Testing, ALCAT testing, Nambudripad's Allergy Elimination Technique, muscle-provocation testing, electrodermal analysis, and hair/urine analysis.^{24,25} Providers should be aware of these tests, as well as the lack of evidence supporting use, as patients may either request such testing, or have already been subjected to them. Both the AAAAI and the ACAAI have discouraged use of these alternative tests.

Utility of the Oral Food Challenge (OFC) in Diagnosing Peanut Allergy

The OFC remains the criterion reference standard test to define peanut or any food allergy.^{1,17,18} The modern practice of this procedure was developed by Dr. Charles May in 1976, who described oral food provocation challenge in 38 patients with asthma and suspected immediate hypersensitivity to food.²⁶ The OFC generally provides a definitive diagnosis as the outcome is apparent—under medical supervision to observe the outcome, either the person will tolerate ingestion or react. OFCs are rarely indeterminate, so long as the patient can cooperate and ingest the full challenge dose, or subjective symptoms can be avoided. While the double blind, placebo-controlled food challenge is considered the most objective form of OFC (and decreases the likelihood of subjective symptoms complicating interpreting the outcome), open OFC's are usually sufficient for clinical diagnosis and are more practical to conduct, though this has not been directly studied for comparison and represents expert opinion.¹ Inherent in the label "challenge", this implies the outcome is not known beforehand, and thus any challenge carries a risk of a potential allergic reaction, including anaphylaxis, so the clinician must be prepared to potentially treat, and the patient be made aware of such risks.^{1,22} Detailed guidance on conducting OFCs in patients is provided elsewhere.^{27,28} OFCs are considered both time- and resource-intensive by some, and require dedicated office space and provider expertise, which may make them less appealing to some providers to conduct.²⁹ However, this is a routine office-based procedure with a superb safety record in the hands of experienced providers.^{1,27}

A decision to offer an OFC is complex and individualized, and providers approach this with a variable degree of expertise, comfort, and desire to offer the procedure.²⁹ OFC can be used to rule in as well as rule out a diagnosis. However with very high or very low pre-test probability, the necessity to offer diagnostic OFC may be low (e.g., when either the outcome is very likely to result in a reaction, or very likely to be tolerated).^{1,21,27,30} This procedure becomes of greater importance when the probability of having had a reaction to peanut is poorly determinable based on pre-test probability, and testing does not provide much assistance in formulating post-test odds. In this context the OFC can provide an objective outcome to inform decision-making. However, while in such situations there may be obvious utility to perform an OFC, the decision to ultimately do so may depend on patient-specific and provider-specific factors like anxiety, vulnerability, desire to eat peanut as well as the clinical judgement and willingness of the clinician to perform the procedure.^{17,27,30} Patients and families that are particularly anxious about eating peanut might prefer to avoid peanut, even with a lower probability of reaction, rather than undergo OFC.

Section 2: Systematic Review, Meta-Analysis, and GRADE Recommendationaions on the Use of Peanut Allergy Diagnostic Testing

Overview of guideline development process

This practice parameter was developed using the GRADE approach. GRADE is a well-established methodology for developing evidence-based guidelines, detailed elsewhere.³¹⁻³³ In formulating the replies to our key questions we took into account the quality of evidence for assessing test sensitivity/specificity, combining this with how the recommendations would be implemented and the knowledge translated, including cost-effectiveness of the recommendations. Table 1 details the GRADE recommendations and evidence ratings. For more details of the GRADE process please refer to the Joint Taskforce website primer (www.allergyparameters.org).

In 2017, the Joint Taskforce on Practice Parameters submitted a concept for a peanut allergy clinical practice guideline to the AAAAI/ACAAI parent organizations. The JTFPP identified 4 liaisons to help identify content experts to form a working group. The workgroup conducted periodic calls to develop central questions to be answered through systematic reviews using the GRADE process, develop a search strategy to identify and review the relevant literature. The working group was divided into individual subgroups to evaluate the identified literature and draft the recommendations based upon the GRADE analysis, and following AMSTAR-2 criteria for systematic reviews.³⁴ A working draft was prepared by

the workgroup, which was then reviewed and modified by the JTFPP. Both groups were provided the opportunity to comment, propose changes, and approve or disapprove each statement. Consensus was sought and reached for each recommendation's direction and strength. Actual or potential conflicts of interest were disclosed annually and transparency of discussion was maintained. A final draft was then approved by the JTFPP and sent to AAAAI and ACAAI appointed reviewers who were asked to comment on the statements and the rationale within free text fields. All these comments and suggestions were discussed during an JTFPP teleconference. For each comment or suggestion, the JTF evaluated whether the statement needed to be adapted, again taking into account the balance between desirable and undesirable consequences of the alternative management strategies, the quality of the evidence, and the variability in values and preferences.

Concurrent with the AAAAI and ACAAI review, a working draft of the guideline was then posted on the AAAAI, ACAAI, and JTFPP websites for all members and the public at large to review. For each comment or suggestion, the JTF evaluated whether the statement needed to be adapted, again taking into account the balance between desirable and undesirable consequences of the alternative management strategies, the quality of the evidence, and the variability in values and preferences. The finalized draft was then sent to this journal for additional peer review before publication.

GRADE Methodology

Development of Searchable Questions

Prior to conducting a literature search, 3 pre-specified PICO (Population, Intervention, Comparator, Outcomes) format question were formulated by the workgroup and the JTF as per standard GRADE approach.³⁵ The population for study included published data for patients with known or highly suspected peanut allergy, who underwent oral food challenge (open or blinded) to establish/confirm a clinical outcome of peanut allergy in at least 50% of participants, where both serologic assessment of peanut allergen components (Ara h 1,2,3,6,8) and/or prick skin testing to whole peanut extract or sIgE testing to whole peanut were obtained as markers of peanut sensitization.

The questions developed were the following:

1. In adults and children with a history of suspected peanut allergy and requesting evaluation what are the indications to perform or not perform diagnostic test(s)?

Population: Adults and children presenting for the evaluation of suspected peanut allergy

Intervention: Perform a diagnostic test for peanut allergy based upon history provided

Comparator: Not perform a diagnostic test for peanut allergy based upon history provided

Outcomes: Accuracy of history in determining need for diagnostic testing for peanut allergy

2a. In the patient presenting for evaluation of suspected peanut allergy, should the provider use a skin prick test, a serum-specific IgE test, or both?

Population: Adults and children presenting for the evaluation of peanut allergy

Intervention: Using skin prick testing (SPT), serum specific IgE to whole peanut (sIgE) or both to determine peanut sensitization to assist in the diagnosis of peanut allergy

Comparator: Oral food challenge

Outcomes: Diagnostic accuracy of peanut allergy testing (true/false positive, true/false negative tests)

2b. In the patient presenting for evaluation of suspected peanut allergy, does testing peanut components in addition to SPT or sIgE whole peanut increase diagnostic accuracy?

Population: Adults and children presenting for the evaluation of peanut allergy

Intervention: Using peanut component testing, e.g., Ara h 2, in addition to SPT or sIgE whole peanut to determine peanut sensitization to assist in the diagnosis of peanut allergy

Comparator: Oral food challenge

Outcomes: Diagnostic accuracy of peanut allergy testing (true/false positive, true/false negative tests)

3. In the patient presenting for evaluation of suspected peanut allergy, do the results of diagnostic tests for peanut allergy, in addition to the patient history, help to predict the severity of a future allergic reaction to peanuts?

Population: Adults and children presenting for the evaluation of suspected peanut allergy

Intervention: Performing a diagnostic test(s) for peanut allergy to help predict the severity of a future allergic reaction to peanuts

Comparator: Predicting the severity of a future allergic reaction to peanuts based solely upon the history and without the use of a diagnostic test for peanut allergy

Outcomes: Accurate prediction of the severity of a future allergic reaction to peanuts

Literature Search and Study Eligibility

In conjunction with a medical librarian (KS), a detailed pre-specified search strategy was developed, with input from the working group, as well as based on recently published systematic reviews on peanut allergy diagnostic testing. Study selection was limited to human subjects of any age who were seeking evaluation for the diagnosis of peanut allergy, and English language studies published or in press starting from 1946-2018. The finalized search parameters were then independently run on Medline (PubMed 1946-2018) and Embase (Elsevier 1947-2018) databases, with the results combined and filtered for duplicates. A total of 1,314 potential references were identified and transferred into Covidence for review by 4 workgroup members (MG, MS, JW, JO), where 127 studies were identified for full text review by the same 4 authors, resulting in a final selection of 89 studies for data extraction pertaining to searchable questions under GRADE format. (Figure 1a-d, overall PRISMA diagram and diagrams by individual searchable question; eSupplement 1, literature search strategy and table of included studies). The search results were combined and culled for duplicate entries, then uploaded into Covidence, where a minimum of two study team members independently reviewed each study for eligibility for full-text review, to determine inclusion, with this process repeated to determine the final studies for data extraction. Conflicts regarding inclusion were resolved by a third study team member. Studies where OFC was not performed as part of the assessment accompanying the diagnostic testing were excluded (including cohort and observational studies based on patient-reported or chart-reported history of peanut allergy involving the use of the aforementioned diagnostic tests without OFC confirmation) but was inclusive of either prospective, retrospective, cross-sectional, or case-control methodologies from both pediatric and adult populations. The full-text versions of the final studies meeting inclusion were reviewed for data extraction of the measures of diagnostic accuracy including sensitivity, specificity, positive/negative predictive value, and the number of true positives, false positives, true negatives, and

false negatives. Individual study authors were contacted to provide additional data for the following reasons:

- 1) To clarify information pertaining to number of successful and non-successful challenges relative to a reported cut-off level of the test in question, where such data was not available or calculatable, so that sensitivity and specificity could be calculated (e.g., obtain the cells to inform true/false positive and true/false negative according to our pre-specified thresholds)
- 2) To request data not presented/analyzed in the selected paper according to the cut-off levels chosen as part of this review, to enable re-tallying of the true/false positive and true/false negative cases
- 3) To see if additional data regarding other searchable questions was potentially available, that had not been published

Studies selected for data extraction were excluded if the aforementioned measures of diagnostic testing accuracy were not directly reported in the manuscript; upon final review the population, use/application of the index test, use/application of the reference standard was deemed to not fit the pre-specified inclusion criteria; or the study team could not/did not provide the requested additional details for more tailored data to be reported per our extraction parameters upon being contacted to provide this information.

Outcomes and Data Synthesis

Based on the diagnostic test used, the extracted number of true positives, false positives, true negatives, and false negatives with respect to oral food challenge outcome were recorded into a MS Excel spreadsheet, as classified by a conservative cut-off level of these tests (for diagnosis, >0.35 KU_A/L for sIgE and Ara h 2 sIgE, ≥ 3 mm for SPT; for severity >50 KU_A/L for sIgE, >2 KU_A/L Ara h 2 sIgE, ≥ 10 mm for SPT) relative to the oral food challenge performed in the study. To assess potential influence of Ara h 6 and Ara h 8 on diagnostic accuracy, pre-specified subgroup analyses were planned based on data availability. Meta-analysis of the pooled sensitivity, specificity, positive and negative likelihood ratios (with visual display of these ratios) on a Fagan Nomogram set to a range of potential lower (30%) and higher (70%) situational pre-test probabilities (representing a clinical suspicion of a diagnosis based on history before a test is performed) of a patient having peanut allergy. Data analysis was performed in Stata, version 15 using the MIDAS command (peto method, random effects model).³⁶ Study

heterogeneity was reported by the I^2 statistic. Risk of bias was assessed using the QUADAS-2 tool. Publication bias was assessed using funnel plots when possible.³⁷ GRADEpro software was used to construct the evidence profiles and calculate the absolute effects.³⁸ Pre-specified sensitivity analyses were planned to explore inclusion only of trials with double blinded challenges as opposed to other challenge types, to assess the effects of geographical region of study, and pediatric vs. non-pediatric studies if permissible. Additional post-hoc sensitivity analyses were performed to verify impact of inclusion of any study on the estimates where there was elevated risk of bias based on patient selection and flow/timing, comparison of individual pooled test precision where SPT/sIgE, sIgE/Ara h 2, or all 3 tests were simultaneously performed, (which per the JTFPP was prioritized as the top sensitivity analysis to report despite this being post hoc, given it most directly answers the searched questions). Data were additionally synthesized narratively. The systematic review process followed AMSTAR2 criteria.³⁴ Lastly, cost-effectiveness analysis using simulated cohorts with Markov modeling over a 20-year horizon, from a societal perspective, was performed to assess simulated health and economic benefits of the use of the individual diagnostic tests (see eSupplement 2).

Reaching workgroup consensus on statements and conclusions:

Where GRADE was not appropriate to answer a particular question, the workgroup employed a modified Delphi process for the determination of the “Strength of the recommendation” and the “Statement profile” for each question. The Delphi method is a structured, interactive, decision-making process used by a panel of experts to arrive at a consensus when there are differing views and perspectives.³⁹⁻⁴¹ For any statement or conclusion in which there was a difference of opinion, a modified Delphi method was used. Workgroup members provided anonymous answers via email to the JTFPP administrative director (AD) to the questions being considered. The AD provided via teleconference an anonymous summary of the experts’ answers and reasons they provided for their responses. The workgroup members discussed all the answers and then were encouraged to modify their answers on the next round(s) of email voting and teleconferences until a consensus was reached.

Results

Question 1: Should diagnostic testing for peanut allergy be performed in adults and children with a history of suspected peanut allergy who are requesting evaluation for peanut allergy?

Recommendation 1a: We suggest in favor of diagnostic (skin prick or serum sIgE) testing for peanut allergy in patients with a 1) physician-judged high pre-test probability of peanut allergy, or 2) prior to an oral food challenge for patients with moderate pre-test probability of peanut allergy, with whom shared decision-making has been employed to arrive at the final decision. **Conditional recommendation; Certainty of evidence: very low**

Recommendation 1b: We suggest against diagnostic testing in patients where there is low or very low pre-test probability of peanut allergy. **Conditional recommendation; Certainty of evidence: very low**

Agreement by the workgroup (By Delphi: 1a 9/9 agree; 1b 9/9 agree).

Quality of Evidence: This question was determined to not be searchable under GRADE format.

Evidence Summary

This question was not searched in a systematic manner as the content experts were unaware of any single research study that addressed this question. However, expert evidence was collected both from the content experts, the JTFPP, and the known prior literature most relevant to this topic. Expert evidence differs from expert opinion, in that the former does not include a judgment on the evidence and offers a systematic and transparent appraisal of the evidence.⁴²

Discussion

Testing for peanut allergy is of the highest utility when there is a history of a known or suspected ingestion of peanut leading to symptoms of an IgE mediated reaction. The identification of individuals for whom testing is indicated requires careful consideration of the clinical history and of epidemiologic risk factors which may increase or decrease the odds of having peanut allergy (e.g., severe atopic dermatitis or another food allergy). Persons with no history of peanut ingestion or an unknown history of ingestion

(without other potential risk factors for developing food allergy), or who asymptotically ingest peanut with impunity should generally not be tested for peanut allergy.^{17,18} The estimated pre-test probability of peanut allergy in these situations is very low, and in most circumstances detection of sensitization will not shift the post-test odds of diagnosis appreciably and will require peanut challenge to resolve the diagnosis. Peanut allergy testing itself is not definitively diagnostic of peanut allergy, as asymptomatic sensitization is not uncommon.¹ Therefore, identifying individuals with a strong pre-test probability for peanut allergy is imperative in the optimal use of diagnostic testing and making an accurate diagnosis of peanut allergy.

Apart from the high-risk infant meeting NIAID addendum 1 criteria, there are potential situations where some providers may ascribe a higher pre-test probability of peanut allergy to a child who has never eaten peanut, and feel that testing may be desired. These generally apply to peanut naïve individuals with other potential risk factors for developing food allergy (e.g., moderate to severe eczema and/or other food allergy), where the pre-test probability may be variably elevated but generally perceived as greater than that of the general population, though still lower than someone with a suspected reaction history. For example, consider the cases of the younger sibling of a peanut allergic child whose family is reluctant to introduce peanut; a child with milk, egg, tree nut or other food allergy; or the child with delayed peanut introduction for other reasons. The decision to test in these circumstances represents a preference-sensitive care option, and in the context of shared decision-making and a thorough explanation of the risks and benefits associated with the preference-sensitive care choices, testing for peanut sensitization may be a reasonable choice. This choice is subject to shared decision making with the patient, and consideration of the risks and benefits of the potential use of oral challenge to help confirm the test results, the magnitude of the degree to which the risk is appreciably different than that of the general population, as well as the potential for the likelihood and consequences of overdiagnosis resulting from detection of asymptomatic peanut sensitization if a challenge is not performed. No decision-aid for this has been developed, however, though this would be potentially useful.

To some degree, clinicians should be advised that they should be prepared to offer OFC to patients where the pre-test probability is no higher than moderate, uncertainty remains, and the patient still desires testing. The risks and consequences of a diagnosis of varying potential accuracy or probability related to a potentially false positive detection of sensitization may or may not outweigh the potential benefit

gained through an at-home introduction or an in-office OFC for some families. Table 3 details some considerations for these situations. Testing the younger sibling of a peanut allergic individual (who does not otherwise meet the addendum 1 high-risk criteria) before peanut introduction has not been shown to be cost-effective unless: a) the baseline prevalence of peanut allergy in younger siblings is >11%; b) that every peanut sensitized child undergoes an OFC to determine actual outcome; and c) the health utility detriment from the initial reaction to peanut was only experienced with at-home introduction and not under an OFC in the office. Without OFC being performed, pre-testing was only cost-effective if the baseline prevalence of peanut allergy in younger siblings was >63%.⁴³

More importantly, it is also crucial to consider the patient who presents to the allergist's office with a test indicating detection of peanut sensitization, but has never eaten peanut before. Here, the context (e.g. the presumed pre-test probability) under which the test denoting sensitization was obtained (and its potential interpretation) also requires careful consideration. This as well may represent a situation of a preference-sensitive choice where a role for shared decision-making arises, with consideration for the benefit of performing an OFC to better determine the outcome should be very carefully weighed against the risk of potential misdiagnosis (and recommended avoidance) from a falsely positive test. The presence of the detectable peanut sensitization itself cannot, however, be used as a condition of "elevated" pre-test probability (which is determined solely by the clinical history).

Question 2a: In the patient presenting for evaluation of suspected peanut allergy, which of the three tests—SPT, sIgE to whole peanut, or Ara h 2 would provide the highest diagnostic accuracy as determined by the more optimal positive/negative likelihood ratio?

Question 2b: In a patient presenting for evaluation of suspected peanut allergy, does testing for peanut components in addition to either SPT or sIgE to whole peanut increase the diagnostic accuracy?

759

Recommendation 2a: We suggest in favor of Ara h2 diagnostic testing in a patient presenting for evaluation of suspected peanut allergy for which a single diagnostic test is to be used, as Ara h2 would provide the best diagnostic accuracy as determined by virtue of more optimal positive/negative likelihood ratios. However, while Ara h 2 has the greatest specificity, it has lower sensitivity than SPT and sIgE, and in a patient with a high prior probability, the clinician may use Ara h 2, SPT, or sIgE to confirm the diagnosis of peanut allergy. **Conditional recommendation. Certainty of evidence: low.**

Recommendation 2b: We suggest against routine use of routine use of component testing in addition to either to skin prick test or sIgE to whole peanut to increase diagnostic accuracy. **Conditional recommendation. Certainty of evidence: very low**

760

Clinical Statement: For GRADE analysis, Ara h 2 was compared to skin prick test and sIgE to whole peanut for the diagnosis of peanut allergy. Providers can interchangeably use either SPT or serologic testing for whole peanut extract IgE, taking into account availability of the test, patient preference, safety, cost, and whether there are patient factors that preclude use of one or both tests. Both tests have high sensitivity but poor specificity in identifying oral food challenge reactive patients at cut-off levels of 3mm wheal size SPT or 0.35 KU_A/L peanut-specific IgE. No data were available regarding use of the tests in tandem or reflexively. In sensitivity analyses where both tests were available, there was minimal difference in the overall sensitivity/specificity between these modalities, and these were similar to the precision in the base analyses of each test individually. However, as a single stand-alone test, compared to either SPT or sIgE testing to whole peanut extract, Ara h 2 has the most optimal combination of positive and negative likelihood ratio, and has much higher specificity, likely decreasing the number of false positive cases where sensitization is detected. While Ara h 2 had the greatest specificity in confirming the diagnosis, it had lower sensitivity when compared with SPT or sIgE. In evaluating diagnostic accuracy, the summary receiving operating characteristic (ROC) curves demonstrated greatest area under the curve (AUC) for Ara h 2 (0.92) when compared to SPT (0.89) and sIgE (0.81). Despite the test characteristics, future research is needed to better clarify the role of Ara h 2 as a stand-alone measure of peanut sensitization in the patient seeking evaluation for possible peanut allergy. In studies where Ara h 2 was evaluated with sIgE or where all 3 tests were evaluated, the precision advantage for Ara h 2 did

not change. A potential risk associated with using Ara h 2 as a stand-alone test is that an allergic individual may be sensitized to other components but not to Ara h 2, though this may be balanced by superior test precision of this approach and was accommodated for in design of the meta-analysis which used OFC as the gold standard for diagnosis. We caution the clinician that despite undetectable sensitization on SPT, sIgE, or Ara h2 testing, that there is a small possibility the individual could still be peanut allergic. If clinical suspicion remains elevated, further evaluation through an oral food challenge, is potentially indicated.

The literature search did not provide patient-level data to determine the value of testing for peanut components in addition to skin prick test or sIgE to whole peanut to increase diagnostic accuracy. Thus, the use and value of components, including reflexive use of Ara h 2, remains a knowledge gap. There is an unclear utility for measuring sIgE to any other commercially available peanut components (Ara h 1, Ara h 3, Ara h 6, Ara h 8, Ara h 9) if peanut sIgE is elevated or SPT >3mm (both indicating sensitization), given the limited available data on performance of components beyond Ara h 2.

Evidence Summary (Questions 2a and 2b):

For SPT and sIgE to whole peanut, from the 89 articles selected for final evidence synthesis, 56 directly pertained to this question. Of these, 32 had data available for extraction (5 studies had no data available, 10 authors did not respond to requests for data, and 9 studies had available data but could not be analyzed due to zero-cell interactions in the 2x2 table). A total of 18 studies (n=2124 patients) were pooled for evidence synthesis for SPT⁴⁴⁻⁵⁶ and 30 studies (n=3989 patients) for sIgE.^{44-48,50,51,53,54,56-70} No literature was identified that detailed the simultaneous, tandem, or reflexive use of both SPT and sIgE to whole peanut. Figure 2a details the summary forest plot for the pooled sensitivity, specificity, and both positive and negative likelihood ratios for a prick skin test to whole peanut extract of 3 mm or greater, and Figure 2b for peanut serum-specific IgE of 0.35 KU_A/L or higher. The summary measures for each test are presented in Table 4. Heterogeneity across these studies was high, and this is reflected in a downgrading under inconsistency. Figures 3 and 4 detail Fagan nomograms for a practical general example of how to roughly interpret the utility of these tests, set at a pre-specified pre-test probability of 2% (general population prevalence), 30% (low suspicion) and 70% (high suspicion). These nomograms show that the likelihood ratio for sensitization at 3mm or 0.35 KU_A/L at 2% or 30% pre-test probability

do not translate to post-test odds >50%, but at the 70% pre-test probability this is raised to ~80%. Negative likelihood ratios do largely decrease post-test odds in all three scenarios. Based on these data, both SPT and sIgE to whole peanut can be used interchangeably, and this is a preference-sensitive choice given no discernable advantage in terms of test precision. There were no data noted that indicate using both tests together was disadvantageous. Both SPT and sIgE to whole peanut have similarly high sensitivity but poor specificity, with serologic testing having slightly higher specificity in identifying oral food challenge reactive patients at the assessed cut-off levels. Table 3 additionally includes sensitivity analysis for the individual sensitivity/specificity of SPT and sIgE assessed when both tests were assessed in the same study. The clinician should be advised of the inherent weaknesses of either of these tests having relatively poor specificity, in that this may predispose to a higher rate of falsely positive detection of peanut sensitization.

For Ara h 2 component-specific IgE, from the 89 articles selected for final evidence synthesis, 41 directly pertained to this question. Of these, 24 had data available for extraction (11 authors did not respond to a request for additional data, 6 articles did not have data available). This resulted in a total of 24 studies (n=2289 patients) pooled for evidence synthesis.^{46,47,50,51,57,59,60,64,71-77 53,54,56,65-67,78-80} The summary measures for Ara h 2 are presented in table 4. Figure 5 detail the summary forest plot for the pooled sensitivity and specificity, for Ara h 2 peanut serum-specific IgE of 0.35 KU_A/L or higher. Heterogeneity across these studies was high. Figure 6 details Fagan nomograms for the use of these tests, set at a pre-specified pre-test probability of 2% (population prevalence), 30% (low suspicion) and 70% (high suspicion). These nomograms show that the likelihood ratio for Ara h 2 sensitization at 0.35 KU_A/L at 2% or 30% pre-test probability translate to post-test odds of 10% and 70%, but at the 70% pre-test probability translates to 89% post-test odds. Negative likelihood ratios do largely decrease post-test odds in all three scenarios.

We were unable to find sufficient number of studies to analyze any other individual peanut components or pool the use of component panels. Therefore, we can offer no comment regarding the role or significance of evaluating these other components individually or in aggregate, or what the clinical implications of their use may be. Similarly, there were no studies identified comparing reflexive use of Ara h 2 or any components after SPT or sIgE. There were no studies identified that evaluated the comparative efficacy of Ara h 2 as a stand-alone test compared to any other component or whole peanut PST or sIgE in their use for clinical decision-making. A potential advantage of Ara h 2 relative to SPT

and sIgE to whole peanut is higher specificity, which may reduce the number of falsely positive cases of sensitization identified, though a disadvantage is this could risk a falsely negative case if someone is sensitized to other components but not Ara h 2. However, the high sensitivity and specificity of the test may limit this risk. In studies where Ara h 2 was evaluated with sIgE or where all 3 tests were evaluated, Ara h 2 consistently had slightly lower sensitivity but much higher specificity, and a more optimal positive/negative likelihood ratio, comparatively. This is similar to the difference noted in the base case where the tests were evaluated individually (Table 4).

Quality of Evidence: Tables 5a and 5b details the summary of GRADE evidence for both SPT and sIgE. There is moderate certainty of evidence for use of either test, and the estimate was downgraded one point for risk of bias. Table 6 details the certainty of evidence for the use of Ara h 2. There is low certainty of evidence, and this estimate was downgraded one point for risk of bias and one point for inconsistency.

Discussion

In practice, SPT and sIgE are often used interchangeably and at the preference of the ordering clinician or the family. Many clinicians may use these tests in tandem with one another as well, though no evidence exists to specifically evaluate this practice. However, this meta-analysis highlights that Ara h 2 is the single best choice based on the most optimal combination of positive and negative likelihood ratios, leveraged by a very high specificity that clinically can translate to a much lower change of a falsely positive diagnosis of peanut allergy. The reduction in the false positive cases provides tremendous clinical value. A 2009 systematic review by Chafen et al⁸¹ noted no statistically significant differences between the diagnostic utility of food-specific SPT and sIgE when comparing their summary ROC curves. A 2015 systematic review by Klemans et al noted the sensitivity of peanut SPT was 0.66-1 the specificity 0-0.95, and the positive and negative likelihood ratios between 1-3.91 and 0-0.65 respectively. For peanut sIgE, this had sensitivity between 0.8-1, specificity between 0-0.63, and positive and negative likelihood ratios between 0.95-2.15 and 0-0.56 respectively.⁸² Overall both SPT and sIgE to whole peanut have very similar test precision, with a very slight relative advantage in sensitivity (0.02) and specificity (0.08) for skin testing over sIgE testing. In the setting of the high-risk infant being evaluated for early peanut introduction, the NIAID guideline specifically recommended SPT as the preferred modality when available, though non-allergists can elect to send peanut sIgE and refer patients for further evaluation or recommend at-home introduction in this population.²⁰ This recommendation was based on data from the

LEAP study, suggesting that SPT provided better classification of peanut allergic infants after peanut challenge than serologic testing.⁸³

There is widespread availability of component testing and several publications have concluded that Ara h 2 may have unique diagnostic value, which has led to debate about whether the clinician should routinely test for IgE to peanut components and base diagnostic decisions solely on these results.⁸² In practice, the clinician has the option to request tests for peanut components in combination with whole peanut SPT and/or peanut specific IgE, or request tests for component testing as a stand-alone test (the Ara h 2 reference code is f423 under CPT 86003). While peanut components are run as single individual tests, these are most commonly offered as a panel by commercial labs at present. To date, no practice parameter or clinical practice guideline has advocated selective use of one or a panel of components over whole peanut SPT or sIgE, how components, including just assessment of Ara h 2, could be used in tandem or reflexively with these tests, or specifically recommend how use of components definitively provides a diagnostic advantage.^{1,17,18} There is limited study of other component testing that was found in this literature search. Ara h 6 sensitization is an emerging area of investigation,⁸⁴ and one study of Ara h 8 mono-sensitization suggested a potential role in discriminating asymptomatic peanut sensitization from allergy, more likely to have clinical relevance in geographic areas where birch pollen is endemic.^{85,86} However, we found few studies that reported challenge-proven outcomes meeting our selection criteria for components apart from Ara h 2, and very limited studies that evaluated use of single vs. panels of peanut components. Thus, we are precluded from commenting any further on specific use of components such as Ara h 6 or Ara h 8, and their potential value in assisting the clinician in making a diagnosis of peanut allergy.

No studies were identified evaluating tandem use of SPT and sIgE to whole peanut. Many studies had both SPT and sIgE measured together, and the individual results are incorporated in the respective analyses. However, we offer no recommendation to this tandem approach, perceived to be commonly done in practice. In studies where both SPT and sIgE were reported, the pooled sensitivity/specificity results were very similar to the base analyses, and reflective of those same small differences. Similarly, no studies were identified evaluating reflexive or tandem use of Ara h 2 or any component with SPT and sIgE to whole peanut, and it is unclear how component testing would be optimally positioned in a clinician's arsenal. Future studies are required to further evaluate Ara h 2 as a stand-alone marker, if components should be tested reflexively after sensitization to whole peanut is denoted or even tested at

all. Importantly, in the context of either very strong or very weak pre-test probability, it is debatable if components (including Ara h 2) offer any additional diagnostic leverage over whole peanut testing, or supersedes the OFC if there was any doubt. In such circumstances, even the good positive likelihood ratio associated with Ara h 2 would not likely change the clinical decision-making or provide more value than the OFC.

Ara h 2 may have more value vs. other testing options in the context of a questionable history and whole peanut sensitization given its higher specificity, in particular in areas with high birch (or birch cross-reactive) pollen. However, additional research is needed to more robustly evaluate such use, and we noted insufficient numbers of study specifically for this application. There is no universal cut-off value for any component (including Ara h 2) that can be used to reliably predict peanut allergy--such levels vary considerably by geographic region, population tested, and possibly by age.^{82 87} As was noted in question 1, there may be situations where a clinician may ascribe a higher pre-test probability to a child who has never eaten peanut before (apart from those falling under NIAID Addendum 1 recommendations), and desire to obtain Ara h 2 component testing. Overall, use of Ara h 2 at present is limited in the capacity of a corroborating test, indicated when there is sufficient pre-test probability for peanut allergy, and not in the capacity of a screening test where there is no pre-test probability. This is demonstrated in the Fagan nomograms in figure 6, which may help illustrate practical general examples of how the test may be reasonably interpreted under different hypothetical pre-test probabilities.

There are several other considerations regarding test preference, including safety, cost, patient features that may drive the choice, availability and practice patterns. SPT is associated with an exceptionally rare risk of systemic reactions (0.077%, with 75% of cases attributable to food), though those doing skin testing should be prepared to potentially treat anaphylaxis.⁸⁸ There also are data demonstrating that there are more side effects from sIgE testing vs. SPT based on assessment in the NHANES study. The cost of SPT and sIgE tests varies among different offices and laboratories, but has been reported to be from 2-7 times less expensive per test for SPT (typically ~\$8 per SPT and \$10-20 per allergen for sIgE test, including components, though components are presently available only as a full panel). (<http://health.costhelper.com/allergy-testing.html>) Certain patient-related factors may make SPT difficult to perform, such as inability to stop medications with anti-histamine activity, severe dermatographism, unstable asthma, patients who may be averse to or afraid of the procedure (such as young children) and hard to control eczema with extensive skin involvement.¹³ However, since SPT can be done on the back

or arm or may be possible on other unaffected areas of skin, it is often possible to do the test even with extensive eczema or delay this until the eczema flare has calmed down. The advantage of SPT is that it is a point of care test that can be rapidly performed in clinic, but a trained specialist generally performs this. There are few limitations to sIgE testing, and often multiple allergens can be assessed from 2-5 mL of blood obtained from routine venipuncture. The test is not point of care, however.¹³ As was noted in question 1, there may be situations in which a clinician may ascribe a higher pre-test probability to a child who has never eaten peanut before (apart from those falling under NIAID Addendum 1 recommendations), and desire to obtain peanut PST or sIgE. The Fagan nomograms in figures 3,4, and 6 may help provide guidance for how the test may be reasonably interpreted in such a scenario.

Test thresholds of 3 mm for SPT and 0.35KU_A/L for sIgE and Ara h 2 sIgE were chosen for analysis of this question. These represent sensitization levels at which a patient traditionally would be considered to have a test indicating allergic sensitization. These are the most widely published “cut-off” levels in the literature, though higher levels, including levels indicative of reported positive predictive values have also been reported, and more recently, lower levels of 0.1 KU_A/L are being commonly reported.^{82,89} We considered different levels (both higher and lower) but disfavored such an approach as this would have reduced the number of citations that would have been available, and made the analysis even more dependent on the goodwill of authors sending us data reconfigured to our needs. A problem unique to the newer conventions of reporting to the technical lower limit of detection at 0.1 KU_A/L is that many studies otherwise eligible for inclusion in our search were performed before reporting to this lower standard was available, and would have limited our total numbers. More importantly, we are unaware of any literature indicating that sensitization between 0.1 and 0.34 KU_A/L is of clinical significance, as opposed to ample literature that clearly has defined sensitization >0.35 KU_A/L as significant.¹ Lastly, we did not attempt to provide a positive predictive value (PPV) for these cut-off levels. The PPV is dependent on a population prevalence of disease, which we do not know and did not assess. Instead, we report likelihood ratios and provide example Fagan nomograms for how the test results could be interpreted at a clinic level, which is a more accurate and appropriate analysis.⁹⁰

Question 3: In the patient presenting for evaluation of suspected peanut allergy, can the results of a diagnostic test be used to predict the severity of an allergic reaction?

Recommendation 3: We suggest against the clinician using the results of a SPT, sIgE to whole peanut extract, or sIgE to peanut components to determine an allergy phenotype or to predict the severity of a future reaction. **Conditional recommendation. Certainty of evidence: very low.**

Clinical statement:

There was inadequate patient-level data to formulate a GRADE recommendation on the use of a peanut diagnostic test for predicting the severity of a future allergic reaction across a continuous range of test result values; however, dichotomous cut-off values of 10mm (SPT), 50 KU_A/L, and 2 KU_A/L (Ara h 2) demonstrated low sensitivity and specificity for a future severe reaction.

Evidence Summary:

From the 89 articles selected for final evidence synthesis, 31 directly pertained to this question. Of these, 16 had data available for extraction (12 authors did not respond to a request for additional data, 1 study did not have data available). A total of 18 studies were pooled for evidence synthesis (10 for Ara h 2 at 2 KU_A/L, n=845 patients;^{46,53,54,56,57,60,65,77,91} 13 for whole peanut sIgE at 50 KU_A/L, n=1051 patients;^{46,48,53,54,56,60,70,91, 92} 12 for SPT 10 mm, n=737 patients^{46,53-56,65,70,91,92}). The summary measures for each test are presented in table 4. Figures 7-9 detail the summary forest plot for the pooled sensitivity and specificity for cut off levels for severe reactions for Ara h 2 peanut serum-specific IgE of 2 KU_A/L or higher, whole peanut sIgE at 50 KU_A/L, and for SPT 10 mm. Due to both low sensitivity and specificity, with no individual measure greater than 0.68 for any of these analyses, likelihood ratios and Fagan nomograms were not reported. Heterogeneity across these studies was high. Based on these data, this analysis notes exceptionally poor sensitivity and specificity for these cut-off values, which differs from a similar analysis by Klemans et al in a 2015 systematic review where Ara h 2 as a marker of severity was concluded to have more potential. Klemans et al explored several different cut-off levels than we did in this analysis, though did so with far less studies included per cut-off level investigated.⁸² Therefore, the results of this analysis should be interpreted as a significant caution to clinicians against using the dichotomous sensitization cut-offs studied to whole peanut (skin/blood) or peanut component (blood) as a surrogate to determine if someone will have a future severe reaction or has a “severe” reaction phenotype. This caution is pending further future studies of much higher quality, more consistently defining severity, with less selection bias, and with more patient level data for analysis. There were insufficient numbers

of other studies to comment regarding the role or significance of evaluating these other components individually or in aggregate to determine if there is any test that may infer reaction severity.

Evidence Strength: Tables 7a-c detail the certainty of evidence for the use of Ara h 2, sIgE, and SPT at these stated cut-off levels for the assessment of the severity of a reaction. There is very low certainty of evidence for all three of these measures and this estimate was downgraded one point for risk of bias and two points for inconsistency (based on the high heterogeneity of the sensitivity and specificity of pooled studies, and a different definition of severity among the studies).

Discussion

There is no relationship indicating that the degree of sensitization is predictive of the underlying severity of the reaction to peanut, using either skin or serologic markers, whole allergen or component. This includes any single test, component, or panel of tests. Importantly, the clinician is advised against making the interpretation that sensitization (at the thresholds evaluated) will predict if someone will have a severe reaction or not. Per our meta-analysis, at the cut-off levels reported, there is no clearly predictive relationship with reaction severity from available data (regardless of the criteria used to define severity), at the levels of sensitization evaluated. Severe reactions can still occur with low/lower sensitization levels. Multiple practice parameters, guidelines and systematic reviews have repeatedly emphasized these points.^{1,17,18} A few individual peanut component-based studies have suggested some degree of association between the recognition of discrete levels of Ara h 2 and history of a severe allergic reaction, though a greater number of studies have noted no such association, and many of these have multiple biases.⁸² At our chosen cut-off levels (Ara h 2 2 KU_A/L; PST 10mm, sIgE 50 KU_A/L), we were unable to identify a relationship to severity, though if patient-level data were available for pooling and evaluation of sensitization as continuous variables, it is possible a relationship could exist. We caution that there is very serious risk of bias among even the few numbers of studies we included. In particular, many studies did not assess severity using Ara h 2, and small inclusion numbers may present a misleading estimate due to omission of data.

There is some evidence that singular recognition of Ara h 8 sensitization (in the absence of other component recognition) may be suggestive as a potential discriminator of pollen cross-sensitization in individuals residing in particular geographic areas (e.g., birch-pollen endemic areas such as the northeast US or northern Europe) who are likely to only experience oropharyngeal, transient itching from peanut ingestion (e.g., pollen food allergy syndrome).¹⁶ However, we could not analyze this question due to low study numbers evaluating this relationship that met inclusion criteria (specifically that 50% of the population underwent OFC). Furthermore, while some expert opinions may support that Ara h 8 monosensitization is a potential indicator of pollen-food allergy syndrome and surrogate for low risk of a severe reaction, these findings lack definitive confirmation in this and prior meta-analysis.⁸² Importantly, we found insufficient numbers of studies for components apart from Ara h 2 meeting our criteria to pool for analysis and cannot comment on the clinical utility of these tests without further rigorous study to validate this concept.

Regional geography may influence component sensitization patterns, in particular with the pollen cross-sensitized individuals, which complicate assessing the relationship between sensitization and severity. Two studies have shown differences in component recognition patterns in patients in northern Europe (Sweden), southern Europe (Spain), and the USA (New York City), as well as differing patterns among different regions in the USA (as noted for birch endemic areas such as the northeast) which may complicate the use of any particular component as a phenotypic discriminator.⁸⁵ For instance, in birch endemic areas, Ara h 8 may behave as a cross-sensitizing marker, and has been proposed to help identify such individuals from those recognizing other proteins in peanut. Ara h 9 could have relevance as a component associated with lipid transfer protein syndrome in certain areas of the world (such as the Mediterranean coast), with high potential to cause systemic reaction in sensitized individuals) whereas elsewhere it behaves similarly to Ara h 8 as a marker of tree pollen sensitization.¹⁶ Therefore, it is unclear the degree to which severity of a reaction may be affected by such geographical differences influencing component recognition, and this area of component research remains promising, but at present represents a knowledge gap.

Importantly, there are issues of bias that must strongly be considered regarding the studies noting an association between sensitization levels and severity. Most of these studies suffer from multiple biases, the most concerning of which is patient selection from serum banks within retrospective cohorts, and lack of generalizability of the sample used for analysis. Many of these studies also lack clear comparison to a

reference (“gold”) standard, tended to be conducted only in certain aged samples, and lacked prospective use of an OFC complicating an objective determination of reaction severity. Study of severe reactions is further hampered given a predilection to not challenge strongly sensitized individuals with a supporting clinical history, as well as ethical considerations to promptly treat reactions when individuals are challenged, which preclude determining how severe a reaction could be.

The cut-off levels chosen for this analysis were based on review of the literature, where we could include the maximal number of studies, and represent realistically large sensitization levels. For reasons discussed previously, we do not report to the lower limit of detection, other levels of sensitization, or attempted to derive a PPV for severe reactivity.

Analysis of Health and Economic Benefits of Peanut Diagnostic Strategies

Cost-effectiveness of peanut allergy diagnostic options was evaluated with decision analysis informed by results of the meta-analysis of diagnostic operating characteristics of single ara h 2 sIgE, whole peanut sIgE, and SPT. The results of this, showing Ara h 2 to be the superior choice in terms of overall cost-savings and accumulation in terms of quality-adjusted life years (QALY), are detailed in the eSupplement 2, eTables 1-2, and eFigures 1-4.

Sensitivity Analyses

In our protocol we pre-specified sensitivity analyses based on OFC type, geographical region of where the study was conducted, and patient age. We performed additional post-hoc sensitivity analyses for studies that had high risk of bias where both patient selection and flow/timing were noted to be issues. These results are shown in table 4,8, and supplemental figures 5-11. Additional supplemental figures display summary receiver operating characteristic curves, as an alternative to the sensitivity forrest plots for the skin test, sIgE and Ara h 2 cut offs (supplemental figure 10) and for potential reaction severity (supplemental figure 11).

Finally we performed one additional, more detailed sensitivity analysis of a very limited number of studies to further assess question 3 and explore size of skin prick tests and levels of serum specific IgE to peanut or its components as predictors of severe allergic reactions. Of the available data from the diagnostic dataset, we then meta-analyzed across studies any adjusted measures of association for skin prick tests, serum-specific IgE to either whole peanut or Ara h 2. In some cases, we generated these data using each study’s reported primary data by logistic regression adjusting for at minimum, age and sex.

For skin prick tests of 8mm or greater, random effects meta-analysis of Preece,⁴¹ Van Erp⁵⁴ and Lewis⁹⁰ showed the adjusted odds ratio (aOR) to be 1.05 [95%CI 0.95-1.15] for systemic reactions. Pooling van Erp,⁵⁴ Balmer-Weber,⁵⁵ and Glaumann,⁵⁸ yielded aOR 1.02 (95%CI 1.00-1.05) for Ara h 2 >2 and systemic reactions. While this may be statistically significant, 2%, even 5% (the upper limit of the 95%CI), increase in odds is not clinically significant. Lastly, we pooled Balmer-Weber,⁵⁵ Glaumann,⁵⁸ Lewis,⁹⁰ and Wensing⁶⁸ and found a pooled aOR of 1.78 (95%CI 0.64-4.96) for peanut IgE > 50 kU/L and systemic reactions. These data lend support to the current recommendation and highlight sparsity of data on prognostic performance of these commonly employed biomarkers, leading to imprecision and therefore, less certainty at to which factors to test for and counsel patients around. As well they highlight the need for uniform data collection and reporting in food allergy studies involving tests.

Risk of Bias Assessment

Risk of bias was assessed using the QUADAS-2 assessment tool.³⁷ This noted some instances where high risk was noted pertaining to the studies for either risk of bias or applicability. The results of this are detailed in figure 10. Sensitivity analyses for all 3 searchable questions were completed after removing studies judged to have high risk for bias based on patient selection and flow/timing of the testing and challenge but this did not alter the pooled sensitivity and specificity estimates to an appreciable or significant degree.

General Limitations of this Practice Parameter and GRADE Recommendations

There are multiple limitations to this analysis. Foremost, we were only able to address prespecified 3 questions, including one that was not searchable, in the scope of this analysis. This does not imply that there are other factors or issues within peanut allergy diagnostic testing that are less important. The JTFPP did limit the questions investigated, for pragmatic reasons to ensure we could produce a GRADE based parameter in the timeframe allotted. All parameters going forward give priority to offering focused updates to formerly published documents using GRADE format. Therefore, this document updates the Diagnostic Testing parameter from 2008,¹³ with a focus on the use of diagnostic testing for peanut allergy. GRADE is not the only system for evidence-based reviews, but is the chosen system for the JTFPP. GRADE has multiple noted limitations, including forced downgrading of certainty

and strength of recommendation based on particular study attributes, and a general trend that the overall strength of recommendations are rarely strong.³¹⁻³³ Peanut components were not commercially available before the latter part of the 2000's and thus this may have introduced not-at-random factors about the types of patients studied in those compared to earlier studies when components were not available. Fairly low cut-off levels were chosen in the analysis for reasons detailed in the sub-sections, but this remains a limitation in that the relative precision of the test may perform differently at different levels.

We found a scarcity of available studies in our literature search that met the OFC criteria and explored use of these tests at a general population level. Therefore, most included studies either involved a referral center cohort, or in many cases, a referral center cohort enriched for patients with known sensitization (skin and/or serologic IgE testing) as selection criteria before being offered OFC. In choosing the selection criteria and evaluating studies for final inclusion, it was felt that this was an acceptable approach given that the specialist clinician would generally be dealing with issues surrounding test interpretation in this population, and be less concerned with false negative rates from the general population (which the pooled sensitivity and specificity may inaccurately estimate in this analysis). We have accounted for this by downgrading the risk of bias (on account of risk of bias from patient selection) category in the GRADE certainty of evidence table, which factors into the overall certainty of the recommendations. Additionally, the analyses involve pooling of studies for assessment of severity that did not all use the same severity criteria (they were similar enough to pool but the rankings reflected different criteria that have evolved over time) and most had wide confidence intervals, requiring us to downgrade 2 points for inconsistency.

The limitations of lack of studies evaluating a tandem or reflexive approach, or the robustness of studies pertaining to other components beyond Ara h 2 (necessary to allow for meta-analysis) have already been mentioned, as has the lack of a consistent objective grading criteria, the small number of studies evaluating reaction severity, as well as differences noted in the timing/flow and selection processes of each of these studies. This is accounted for in grading the certainty of evidence and risk of bias. As well, the aforementioned sensitivity analyses were done to further confirm if inclusion of those studies felt to be most at risk would alter the estimates, which they did not. We could not stratify by allergic co-

morbidity (in particular presence of atopic dermatitis) or age with accuracy due to limited available data in the reporting which would allow for such stratifications to be made, though we did perform sensitivity analysis on challenge type, adult vs. pediatric studies, as well as by region of the world (Europe, North America) in which the data were observed. Statistically, the pooling of data are limited by high heterogeneity, with some included studies having high risk of bias.

Knowledge Gaps

Within in the scope of these questions, multiple gaps in the current knowledge base were identified that could not be resolved through our literature search and meta-analysis. These include, but are not limited to:

- a) A lack of identified studies that systematically evaluate when someone should be tested for peanut allergy
- b) A lack of identified studies that evaluate the tandem or reflexive use of whole peanut extract SPT and whole peanut sIgE in combination
- c) A lack of identified studies that evaluate the tandem or reflexive use of whole peanut extract SPT and whole peanut sIgE in combination with peanut components
- d) A lack of identified studies that evaluate the tandem or reflexive use of one or more peanut components
- e) A lack of identified studies that evaluate Ara h 1, Ara h 3, Ara h 6, Ara h 8, and Ara h 9 performance, or if severity or reaction phenotypes are associated with recognition of these components
- f) A lack of identified studies that consistently or systematically study reaction severity using unified criteria or cut-off markers, or evaluate this question at different cut-off levels
- g) A lack of identified studies that study any of the searchable questions at a population level that are less enriched for already sensitized individuals as opposed to within more clustered clinical referral centers
- h) A lack of identified studies that trace longitudinal outcomes and natural history of disease to better understand the full scope of the ramifications of diagnostic testing choices to inform best-practices

- i) A lack of clear understanding and inconsistent use of diagnostic cut-off points for the use of these tests
- j) A lack of consistent reporting at an individual level of allergic co-factors that may influence the performance of these diagnostic tests in relation to the food challenge outcome to assess the influence of such covariates

Text box 2 addresses a number of the key take-home messages and knowledge gaps.

Benefits/Harms of Implementing the Guideline Recommendations

Potential Benefits

The potential benefit of this analysis is the appropriate management of patients with Peanut allergy. See the “Discussion” section for each question in the guideline document for benefits of tests. Cost-effectiveness analysis was undertaken to further explore such health benefits. Please refer to eSupplement 3, which comprehensively details the evidence to recommendation process.

Potential Harms

The potential harms include adverse effects associated with incorrect diagnosis of peanut allergy. See the “Discussion” section for each question in the guideline document for adverse events of specific interventions. Cost-effectiveness analysis was undertaken to further explore such health detriments. Please refer to eSupplement 3, which comprehensively details the evidence to recommendation process.

Qualifying Statements

This clinical practice guideline was designed to facilitate informed decision-making on the diagnosis of children and adults with suspected peanut allergy. It was not intended to define a standard of care, and should not be construed as such. It should not be interpreted as a prescription for an exclusive course of management.

Summary and Conclusions

In making a diagnosis of peanut allergy, it is important to clearly understand the indications for running a diagnostic test. Patients with a history of peanut ingestion leading to symptom development benefit most from peanut allergy diagnostic testing.^{17,18,22} With the exception of patients who are not newborn infants under the age of 4-6 months of life who have either egg allergy or severe eczema,²⁰ there is no clear indication for any form of peanut allergy testing in someone who has not yet eaten peanut and subsequently developed symptoms of an allergic reaction. Testing only determines the presence or absence of peanut sensitization and alone does not allow a definite diagnosis without a history to provide context as to what happens upon peanut ingestion.¹³ Use of the tests in these contexts helps translate the pre-test probability of allergy (e.g. based on the history) into post-test probability of a peanut allergy diagnosis.¹⁴ In some cases, an oral food challenge may be necessary to definitively rule in or rule out a diagnosis, but may be a patient-preference sensitive decision. In terms of choice of tests, when assessing for whole peanut sensitization, there is little practical difference between use of SPT or sIgE—both are highly sensitive but relatively poorly specific, and may be prone to false positive detection of sensitization in certain contexts. Use of testing to the peanut component Ara h 2 has the best profile of high sensitivity, high specificity, and optimal positive/negative likelihood ratio, and is probably the most accurate single test that is available in terms of a test that could be sent with the lowest potential risk of false positive sensitization being detected. However, how this test should be used in the work up of the suspected peanut allergic patient remains unresolved and not prospectively validated in terms of clinical pathways as to how such properties could be leveraged. We do present evidence herein that shows that using Ara h 2 as a sole diagnostic test in the evaluation of peanut allergy could be cost effective, given the cost-savings at a societal level (with downstream costs considered) associated with a significant simulated reduction in the number of false positive cases, as one such possible application of how the test could be used. At the dichotomous cut-offs evaluated, no whole peanut allergen or component test dictates severity of a future reaction, or a reaction phenotype.

1208

1209 **References**

- 1210 1. National Academies of Sciences E, Medicine. Finding a Path to Safety in Food Allergy:
 1211 Assessment of the Global Burden, Causes, Prevention, Management, and Public Policy. Washington, DC:
 1212 The National Academies Press; 2017.
- 1213 2. Peters RL, Allen KJ, Dharmage SC, Tang ML, Koplin JJ, Ponsonby AL, et al. Skin prick test
 1214 responses and allergen-specific IgE levels as predictors of peanut, egg, and sesame allergy in infants. *J*
 1215 *Allergy Clin Immunol* 2013;132:874-80.
- 1216 3. Peters RL, Allen KJ, Dharmage SC, Koplin JJ, Dang T, Tilbrook KP, et al. Natural history of
 1217 peanut allergy and predictors of resolution in the first 4 years of life: A population-based assessment. *J*
 1218 *Allergy Clin Immunol* 2015;135:1257-66.e1-2.
- 1219 4. Osborne NJ, Koplin JJ, Martin PE, Gurrin LC, Lowe AJ, Matheson MC, et al. Prevalence of
 1220 challenge-proven IgE-mediated food allergy using population-based sampling and predetermined
 1221 challenge criteria in infants. *J Allergy Clin Immunol* 2011;127:668-76.e1-2.
- 1222 5. Peters R, Koplin J, Ponsonby A-L, Perret K, Dharmage SC, Allen K. The Natural History of
 1223 Peanut and Egg Allergy and Predictors of Persistence: The Healthnuts Longitudinal Study, 6-Year-Old
 1224 Follow-up. *Journal of Allergy and Clinical Immunology* 2019;143:AB421.
- 1225 6. Gupta RS, Springston EE, Warrier MR, Smith B, Kumar R, Pongracic J, et al. The prevalence,
 1226 severity, and distribution of childhood food allergy in the United States. *Pediatrics* 2011;128:e9-17.
- 1227 7. Sicherer SH, Muñoz-Furlong A, Godbold JH, Sampson HA. US prevalence of self-reported
 1228 peanut, tree nut, and sesame allergy: 11-year follow-up. *Journal of Allergy and Clinical Immunology*
 1229 2010;125:1322-6.
- 1230 8. Bunyavanich S, Rifas-Shiman SL, Platts-Mills TAE, Workman L, Sordillo JE, Gillman MW, et
 1231 al. Peanut allergy prevalence among school-age children in a US cohort not selected for any disease.
 1232 *Journal of Allergy and Clinical Immunology* 2014;134:753-5.
- 1233 9. McGowan EC, Keet CA. Prevalence of self-reported food allergy in the National Health and
 1234 Nutrition Examination Survey (NHANES) 2007-2010. *Journal of Allergy and Clinical Immunology*
 1235 2013;132:1216-9.e5.
- 1236 10. Liu AH, Jaramillo R, Sicherer SH, Wood RA, Bock SA, Burks AW, et al. National prevalence
 1237 and risk factors for food allergy and relationship to asthma: Results from the National Health and
 1238 Nutrition Examination Survey 2005-2006. *Journal of Allergy and Clinical Immunology* 2010;126:798-
 1239 806.e14.
- 1240 11. Bernstein IL, Li JT, Bernstein DI, Hamilton R, Spector SL, Tan R, et al. Allergy Diagnostic
 1241 Testing: An Updated Practice Parameter. *Annals of Allergy, Asthma & Immunology* 2008;100:S1-S148.
- 1242 12. Nelson A. An Interactive Workshop Reviewing Basic Biostatistics and Applying Bayes' Theorem
 1243 to Diagnostic Testing and Clinical Decision-Making. *MedEdPORTAL : the journal of teaching and*
 1244 *learning resources* 2018;14:10771-.
- 1245 13. Grimes DA, Schulz KF. Refining clinical diagnosis with likelihood ratios. *Lancet* 2005;365:1500-
 1246 5.
- 1247 14. Sastre J. Molecular diagnosis in allergy. *Clinical & Experimental Allergy* 2010;40:1442-60.
- 1248 15. Sampson HA, Aceves S, Bock SA, James J, Jones S, Lang D, et al. Food allergy: A practice
 1249 parameter update—2014. *Journal of Allergy and Clinical Immunology* 2014;134:1016-25.e43.
- 1250 16. Guidelines for the Diagnosis and Management of Food Allergy in the United States: Report of the
 1251 NIAID-Sponsored Expert Panel. *Journal of Allergy and Clinical Immunology* 2010;126:S1-S58.

17. Nowak-Węgrzyn A, Chehade M, Groetch ME, Spergel JM, Wood RA, Allen K, et al. International consensus guidelines for the diagnosis and management of food protein-induced enterocolitis syndrome: Executive summary-Workgroup Report of the Adverse Reactions to Foods Committee, American Academy of Allergy, Asthma & Immunology. *J Allergy Clin Immunol* 2017;139:1111-26 e4.
18. Togias A, Cooper SF, Acebal ML, Assa'ad A, Baker JR, Jr., Beck LA, et al. Addendum guidelines for the prevention of peanut allergy in the United States: Report of the National Institute of Allergy and Infectious Diseases-sponsored expert panel. *J Allergy Clin Immunol* 2017;139:29-44.
19. Sampson HA, Ho DG. Relationship between food-specific IgE concentrations and the risk of positive food challenges in children and adolescents. *Journal of allergy and clinical immunology* 1997;100:444-51.
20. Sicherer SH, Sampson HA. Food allergy: A review and update on epidemiology, pathogenesis, diagnosis, prevention, and management. *J Allergy Clin Immunol* 2018;141:41-58.
21. Fleischer DM, Bock SA, Spears GC, Wilson CG, Miyazawa NK, Gleason MC, et al. Oral Food Challenges in Children with a Diagnosis of Food Allergy. *The Journal of Pediatrics* 2011;158:578-83.e1.
22. Hammond C, Lieberman JA. Unproven Diagnostic Tests for Food Allergy. *Immunology and Allergy Clinics of North America* 2018;38:153-63.
23. Kelso JM. Unproven Diagnostic Tests for Adverse Reactions to Foods. *The Journal of Allergy and Clinical Immunology: In Practice* 2018;6:362-5.
24. May CD. Objective clinical and laboratory studies of immediate hypersensitivity reactions to foods in asthmatic children. *J Allergy Clin Immunol* 1976;58:500-15.
25. Nowak-Węgrzyn A, Assa'ad AH, Bahna SL, Bock SA, Sicherer SH, Teuber SS. Work Group report: Oral food challenge testing. *Journal of Allergy and Clinical Immunology* 2009;123:S365-S83.
26. Bird JA, Leonard S, Groetch M, Assa'ad A, Cianferoni A, Clark A, et al. Conducting an Oral Food Challenge: An Update to the 2009 Adverse Reactions to Foods Committee Work Group Report. *J Allergy Clin Immunol Pract* 2020;8:75-90 e17.
27. Pongracic JA, Bock SA, Sicherer SH. Oral food challenge practices among allergists in the United States. *Journal of Allergy and Clinical Immunology* 2012;129:564-6.
28. Stiefel G, Anagnostou K, Boyle RJ, Brathwaite N, Ewan P, Fox AT, et al. BSACI guideline for the diagnosis and management of peanut and tree nut allergy. *Clinical & Experimental Allergy* 2017;47:719-39.
29. Brożek JL, Akl EA, Alonso-Coello P, Lang D, Jaeschke R, Williams JW, et al. Grading quality of evidence and strength of recommendations in clinical practice guidelines. *Allergy* 2009;64:669-77.
30. Brożek JL, Akl EA, Jaeschke R, Lang DM, Bossuyt P, Glasziou P, et al. Grading quality of evidence and strength of recommendations in clinical practice guidelines: Part 2 of 3. The GRADE approach to grading quality of evidence about diagnostic tests and strategies. *Allergy* 2009;64:1109-16.
31. Brożek JL, Akl EA, Compalati E, Kreis J, Terracciano L, Fiocchi A, et al. Grading quality of evidence and strength of recommendations in clinical practice guidelines Part 3 of 3. The GRADE approach to developing recommendations. *Allergy* 2011;66:588-95.
32. Shea BJ, Reeves BC, Wells G, Thuku M, Hamel C, Moran J, et al. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ* 2017;358:j4008.
33. Schünemann H, Brożek J, Guyatt G, Oxman A, editors GRADE handbook for grading quality of evidence and strength of recommendations Updated October 2013 The GRADE Working Group, 2013 Available from guidelinedevelopment.org/handbook

34. Dwamena B, 2007. "MIDAS: Stata module for meta-analytical integration of diagnostic test accuracy studies," Statistical Software Components S456880, Boston College Department of Economics, revised 05 Feb 2009. Downloaded August 2, 2018.
35. Whiting PF, Rutjes AWS, Westwood ME, Mallett S, Deeks JJ, Reitsma JB, et al. QUADAS-2: A Revised Tool for the Quality Assessment of Diagnostic Accuracy Studies. *Annals of Internal Medicine* 2011;155:529-36.
36. GRADEpro Guideline Development Tool [Software] McMaster University, 2015 (developed by Evidence Prime, Inc) Available from gradepro.org Downloaded August 1, 2018.
37. Dalkey N, Helmer O. An Experimental Application of the DELPHI Method to the Use of Experts. *Management Science* 1963;9:458-67.
38. Linstone HA, Turoff M. The Delphi method : techniques and applications. Reading, Mass.: Addison-Wesley Pub. Co., Advanced Book Program; 1975.
39. Brown B. Delphi Process: A methodology used for the elicitation of opinions of experts. In: Corporation TR, ed. Santa Monica, CA1968.
40. Schunemann HJ, Zhang Y, Oxman AD, Expert Evidence in Guidelines G. Distinguishing opinion from evidence in guidelines. *BMJ* 2019;366:l4606.
41. Shaker MS, Iglesia E, Greenhawt M. The Health and Economic Benefits of Approaches for Peanut Introduction in Infants with a Peanut Allergic Sibling. *Allergy* 2019.
42. Abrams EM, Becker AB. Oral food challenge outcomes in a pediatric tertiary care center. *Allergy Asthma Clin Immunol* 2017;13:43.
43. Begin P, Graham F, Killer K, Paradis J, Paradis L, Des Roches A. Introduction of peanuts in younger siblings of children with peanut allergy: a prospective, double-blinded assessment of risk, of diagnostic tests, and an analysis of patient preferences. *Allergy* 2016;71:1762-71.
44. Chinthrajah RS, Purington N, Andorf S, Rosa JS, Mukai K, Hamilton R, et al. Development of a tool predicting severity of allergic reaction during peanut challenge. *Allergy* 2018;121:69-76.e2.
45. Comberiati P, Colavita L, Minniti F, Paiola G, Capristo C, Incorvaia C, et al. Utility of Specific IgE to Ara h 2 in Italian Allergic and Tolerant Children Sensitized to Peanut. *Int J Mol Cell Med* 2016;5:160-6.
46. DunnGalvin A, Daly D, Cullinane C, Stenke E, Keeton D, Erlewyn-Lajeunesse M, et al. Highly accurate prediction of food challenge outcome using routinely available clinical data. *J Allergy Clin Immunol* 2011;127:633-9.e1-3.
47. Johannsen H, Nolan R, Pascoe EM, Cuthbert P, Noble V, Corderoy T, et al. Skin prick testing and peanut-specific IgE can predict peanut challenge outcomes in preschoolchildren with peanut sensitization. *Clin Exp Allergy* 2011;41:994-1000.
48. Klemans RJ, Broekman HC, Knol EF, Bruijnzeel-Koomen CA, Otten HG, Pasmans SG, et al. Ara h 2 is the best predictor for peanut allergy in adults. *J Allergy Clin Immunol Pract* 2013;1:632-8.e1.
49. Klemans RJ, Otte D, Knol M, Knol EF, Meijer Y, Gmelig-Meyling FH, et al. The diagnostic value of specific IgE to Ara h 2 to predict peanut allergy in children is comparable to a validated and updated diagnostic prediction model. *J Allergy Clin Immunol* 2013;131:157-63.
50. Ludman S, Ballabeni P, Eigenmann PA, Wassenberg J. Predicting positive food challenges in children sensitised to peanuts/tree nuts. *Pediatr Allergy Immunol* 2013;24:276-81.
51. Preece K, Bhatia R, Belcher J, Patchett K, McElduff P, Collison A, et al. The fraction of exhaled nitric oxide improves prediction of clinical allergic reaction to peanut challenge in children. *Clin Exp Allergy* 2014;44:371-80.
52. Rajput S, Sharma V, Hughes SM, Ewing CI, Arkwright PD. Allergy testing in predicting outcome of open food challenge to peanut. *J Allergy Clin Immunol* 2018;141:457-8.

53. Song GC, Wang XY, Wang Z, Ruan XL, Yang J, Zhu Z, et al. [Association between serum allergens and asthma in children]. *Zhongguo Dang Dai Er Ke Za Zhi* 2015;17:806-10.
54. van Erp FC, Knulst AC, Kentie PA, Pasmans SG, van der Ent CK, Meijer Y. Can we predict severe reactions during peanut challenges in children? *Pediatr Allergy Immunol* 2013;24:596-602.
55. Ballmer-Weber BK, Lidholm J, Fernandez-Rivas M, Seneviratne S, Hanschmann KM, Vogel L, et al. IgE recognition patterns in peanut allergy are age dependent: perspectives of the EuroPrevall study. *Allergy* 2015;70:391-407.
56. Beigelman A, Strunk RC, Garbutt JM, Schechtman KB, Jaenicke MW, Stein JS, et al. Clinical and laboratory factors associated with negative oral food challenges. *Allergy Asthma Proc* 2012;33:467-73.
57. Ebisawa M, Moverare R, Sato S, Borres MP, Ito K. The predictive relationship between peanut- and Ara h 2-specific serum IgE concentrations and peanut allergy. *J Allergy Clin Immunol Pract* 2015;3:131-2.e1.
58. Glaumann S, Nopp A, Johansson SG, Rudengren M, Borres MP, Nilsson C. Basophil allergen threshold sensitivity, CD-sens, IgE-sensitization and DBPCFC in peanut-sensitized children. *Allergy* 2012;67:242-7.
59. Guilloux L, Morisset M, Codreanu F, Parisot L, Moneret-Vautrin DA. Peanut allergy diagnosis in the context of grass pollen sensitization for 125 patients: roles of peanut and cross-reactive carbohydrate determinants specific IgE. *Int Arch Allergy Immunol* 2009;149:91-7.
60. Gupta RS, Lau CH, Hamilton RG, Donnell A, Newhall KK. Predicting outcomes of oral food challenges by using the allergen-specific IgE-total IgE ratio. *J Allergy Clin Immunol Pract* 2014;2:300-5.
61. Johansson EK, Bergstrom A, Kull I, Lind T, Soderhall C, van Hage M, et al. IgE sensitization in relation to preschool eczema and filaggrin mutation. *J Allergy Clin Immunol* 2017;140:1572-9.e5.
62. Klemans RJ, Liu X, Knulst AC, Knol MJ, Gmelig-Meyling F, Borst E, et al. IgE binding to peanut components by four different techniques: Ara h 2 is the most relevant in peanut allergic children and adults. *Clin Exp Allergy* 2013;43:967-74.
63. Leo SH, Dean JM, Jung B, Kuzeljevic B, Chan ES. Utility of Ara h 2 sIgE levels to predict peanut allergy in Canadian children. *J Allergy Clin Immunol Pract* 2015;3:968-9.
64. Lieberman JA, Glaumann S, Batelson S, Borres MP, Sampson HA, Nilsson C. The utility of peanut components in the diagnosis of IgE-mediated peanut allergy among distinct populations. *J Allergy Clin Immunol Pract* 2013;1:75-82.
65. Nicolaou N, Murray C, Belgrave D, Poorafshar M, Simpson A, Custovic A. Quantification of specific IgE to whole peanut extract and peanut components in prediction of peanut allergy. *J Allergy Clin Immunol* 2011;127:684-5.
66. Perry TT, Matsui EC, Kay Conover-Walker M, Wood RA. The relationship of allergen-specific IgE levels and oral food challenge outcome. *J Allergy Clin Immunol* 2004;114:144-9.
67. Wainstein BK, Yee A, Jelley D, Ziegler M, Ziegler JB. Combining skin prick, immediate skin application and specific-IgE testing in the diagnosis of peanut allergy in children. *Pediatr Allergy Immunol* 2007;18:231-9.
68. Wensing M, Penninks AH, Hefle SL, Koppelman SJ, Bruijnzeel-Koomen CA, Knulst AC. The distribution of individual threshold doses eliciting allergic reactions in a population with peanut allergy. *Journal of allergy and clinical immunology* 2002;110:915-20.
69. Bernard H, Paty E, Mondoulet L, Burks AW, Bannon GA, Wal JM, et al. Serological characteristics of peanut allergy in children. *Allergy* 2003;58:1285-92.

70. Beyer K, Grabenhenrich L, Hartl M, Beder A, Kalb B, Ziegert M, et al. Predictive values of component-specific IgE for the outcome of peanut and hazelnut food challenges in children. *Allergy* 2015;70:90-8.
71. Dang TD, Tang M, Choo S, Licciardi PV, Koplin JJ, Martin PE, et al. Increasing the accuracy of peanut allergy diagnosis by using Ara h 2. *J Allergy Clin Immunol* 2012;129:1056-63.
72. Ebisawa M, Moverare R, Sato S, Maruyama N, Borres MP, Komata T. Measurement of Ara h 1-, 2-, and 3-specific IgE antibodies is useful in diagnosis of peanut allergy in Japanese children. *Pediatr Allergy Immunol* 2012;23:573-81.
73. Eller E, Bindslev-Jensen C. Clinical value of component-resolved diagnostics in peanut-allergic patients. *Allergy* 2013;68:190-4.
74. Keet CA, Johnson K, Savage JH, Hamilton RG, Wood RA. Evaluation of Ara h2 IgE thresholds in the diagnosis of peanut allergy in a clinical population. *J Allergy Clin Immunol Pract* 2013;1:101-3.
75. Kukkonen AK, Pelkonen AS, Makinen-Kiljunen S, Voutilainen H, Makela MJ. Ara h 2 and Ara 6 are the best predictors of severe peanut allergy: a double-blind placebo-controlled study. *Allergy* 2015;70:1239-45.
76. Martinet J, Couderc L, Renosi F, Bobee V, Marguet C, Boyer O. Diagnostic Value of Antigen-Specific Immunoglobulin E Immunoassays against Ara h 2 and Ara h 8 Peanut Components in Child Food Allergy. *Int Arch Allergy Immunol* 2016;169:216-22.
77. Schots M, de Mol AC, Vermeer HJ, Roosen YM, Vriesman AW. Is Ara h 2 indeed the best predictor for peanut allergy in Dutch children? *Diagnosis (Berl)* 2016;3:31-5.
78. Suratannon N, Ngamphaiboon J, Wongpiyabovorn J, Puripokai P, Chatchatee P. Component-resolved diagnostics for the evaluation of peanut allergy in a low-prevalence area. *Pediatr Allergy Immunol* 2013;24:665-70.
79. Chafen JJ, Newberry SJ, Riedl MA, Bravata DM, Maglione M, Suttrop MJ, et al. Diagnosing and managing common food allergies: a systematic review. *JAMA* 2010;303:1848-56.
80. Klemans RJ, van Os-Medendorp H, Blankestijn M, Bruijnzeel-Koomen CA, Knol EF, Knulst AC. Diagnostic accuracy of specific IgE to components in diagnosing peanut allergy: a systematic review. *Clin Exp Allergy* 2015;45:720-30.
81. Du Toit G, Roberts G, Sayre PH, Bahnson HT, Radulovic S, Santos AF, et al. Randomized trial of peanut consumption in infants at risk for peanut allergy. *The New England journal of medicine* 2015;372:803-13.
82. Asarnoj A, Glaumann S, Elfstrom L, Lilja G, Lidholm J, Nilsson C, et al. Anaphylaxis to peanut in a patient predominantly sensitized to Ara h 6. *Int Arch Allergy Immunol* 2012;159:209-12.
83. Vereda A, van Hage M, Ahlstedt S, Ibanez MD, Cuesta-Herranz J, van Odijk J, et al. Peanut allergy: Clinical and immunologic differences among patients from 3 different geographic regions. *J Allergy Clin Immunol* 2011;127:603-7.
84. Asarnoj A, Nilsson C, Lidholm J, Glaumann S, Ostblom E, Hedlin G, et al. Peanut component Ara h 8 sensitization and tolerance to peanut. *J Allergy Clin Immunol* 2012;130:468-72.
85. Valcour A, Jones JE, Lidholm J, Borres MP, Hamilton RG. Sensitization profiles to peanut allergens across the United States. *Annals of allergy, asthma & immunology : official publication of the American College of Allergy, Asthma, & Immunology* 2017;119:262-6 e1.
86. Sellaturay P, Nasser S, Ewan P. The incidence and features of systemic reactions to skin prick tests. *Ann Allergy Asthma Immunol* 2015;115:229-33.
87. Flores Kim J. Diagnostic accuracy, risk assessment, and cost-effectiveness of component-resolved diagnostics for food allergy: A systematic review. *Allergy* 2018;73:1609-21.

88. Brenner H. Measures of differential diagnostic value of diagnostic procedures. *Journal of Clinical Epidemiology* 1996;49:1435-9.
89. Klemans RJ, Blom WM, van Erp FC, Masthoff LJ, Rubingh CM, van der Ent CK, et al. Objective eliciting doses of peanut-allergic adults and children can be combined for risk assessment purposes. *Clin Exp Allergy* 2015;45:1237-44.
90. Lewis SA, Grimshaw KE, Warner JO, Hourihane JO. The promiscuity of immunoglobulin E binding to peanut allergens, as determined by Western blotting, correlates with the severity of clinical symptoms. *Clinical and experimental allergy* 2005;35:767-73.
91. Peeters KA, Koppelman SJ, van Hoffen E, van der Tas CW, den Hartog Jager CF, Penninks AH, et al. Does skin prick test reactivity to purified allergens correlate with clinical severity of peanut allergy? *Clinical and experimental allergy* 2007;37:108-15.
92. Wainstein BK, Studdert J, Ziegler M, Ziegler JB. Prediction of anaphylaxis during peanut food challenge: usefulness of the peanut skin prick test (SPT) and specific IgE level. *Pediatr Allergy Immunol* 2010;21:603-11.

1462

1463

1464

1465

1466

1467

Journal Pre-proof

Table 1: Summary Recommendations in Evaluating the Patient with Suspected Peanut Allergy

Question	Recommendation	Evidence Certainty	Risk of Bias
Should diagnostic testing for peanut allergy be performed in adults and children with a history of suspected peanut allergy who are requesting evaluation for peanut allergy?	We suggest in favor of diagnostic (skin prick or serum sIgE) testing for peanut allergy in patients with a 1) physician-judged high pre-test probability of peanut allergy, or 2) prior to an oral food challenge for patients with moderate pre-test probability of peanut allergy, with whom shared decision-making has been employed to arrive at the final decision. We suggest against diagnostic testing in patients where there is low or very low pre-test probability of peanut allergy.	Very Low	Not Rated
In the patient presenting for evaluation of suspected peanut allergy, which of the three tests—SPT, sIgE to whole peanut, or Ara h2 would provide the highest diagnostic accuracy as determined by the more optimal positive/negative likelihood ratio?	We suggest in favor of Ara h2 diagnostic testing in a patient presenting for evaluation of suspected peanut allergy for which a single diagnostic test is to be used, as Ara h2 would provide the best diagnostic accuracy as determined by virtue of more optimal positive/negative likelihood ratios. However, while Ara h 2 has the greatest specificity, it has lower sensitivity than SPT and sIgE, and in a patient with a high prior probability, the clinician may use Ara h 2, SPT, or sIgE to confirm the diagnosis of peanut allergy.	Low	High
In a patient presenting for evaluation of suspected peanut allergy, does testing for peanut components in addition to either SPT or sIgE to whole peanut increase the diagnostic accuracy?	We suggest against routine use of component testing in addition to either to skin prick test or sIgE to whole peanut to increase diagnostic accuracy.	Very Low	High

In the patient presenting for evaluation of suspected peanut allergy, can the results of a diagnostic test be used to predict the severity of a future allergic reaction?	We suggest against the clinician using the results of a SPT, sIgE to whole peanut extract, or sIgE to peanut components to determine the severity of a previous reaction and/or allergy phenotype or to predict the severity of a future reaction.	Very Low	High
---	--	----------	------

Journal Pre-proof

Table 2: The GRADE System of Recommendations and Evidence Certainty

Strength of Recommendation		
	For the Patient	For the Clinician
Strong	Most individuals in this situation would prefer the recommended course of action and only a small proportion would not.	The attending provider should strongly consider the recommended course of action as a first-line management. Formal decision aids may have less of a role to help individuals make decisions consistent with their values and preferences.
Conditional	The majority of individuals in this situation would prefer the suggested course of action, but many would not.	Different choices may be appropriate for different patients. Decision aids may be useful in helping individuals in making decisions consistent with their values and preferences. Clinicians should expect to spend more time with patients when working towards a decision.
Certainty in estimates of effect / quality rating both for outcome and for an entire evidence base as it pertains to a PICO		
High	There is high confidence that the true effect lies close to that of the estimate of the effect.	
Moderate	There is moderate confidence in the effect estimate. The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.	
Low	There is limited confidence in the effect estimate. The true effect may be substantially different from the estimate of the effect.	
Very Low	There is very little confidence in the effect estimate. The true effect is likely to be substantially different from the estimate of effect	

Additional information regarding GRADE methodology, including how recommendations are formulated and the evidence certainty is rated can be found at Shaker MS, Oppenheimer J, Wallace DV, Golden DBK, Lang DM, Joint Task Force for Allergy Practice P, et al. Making the GRADE in anaphylaxis management: Toward recommendations integrating values, preferences, context, and shared decision making. *Ann Allergy Asthma Immunol* 2020. Mar 19:S1081-1206(20)30153-8. doi:10.1016/j.anai.2020.03.009 and on the Joint Taskforce on Allergy Practice Parameters website at <https://www.allergyparameters.org/resources-for-understanding-grade/>.

Table 3: Situations of Low to Moderate Pre-Test Probability for Peanut Allergy Where Testing May be a Preference-Sensitive Care Option to Offer in the Evaluation of a Patient^a

Situations Where A Clinician Might Be Considering Testing for Peanut Allergy ^b	Pros for Testing	Cons for Testing
A young child >1year but <3 year with multiple asthma hospitalizations, on chronic inhaled steroids, with known milk allergy who has not yet tried peanut	- Possible elevated risk for an additional food allergy in someone who already has one food allergy - Parents may not introduce peanut without a positive test, leading to additional risk from delayed introduction	While the risk could be elevated over baseline, it is unclear if the absolute risk is elevated more than the low probability scenario of a 30% pre-test probability where a positive test was not shown to appreciably shift the post-test odds
A young child >1year but <3 year old without eczema with prior anaphylaxis to one or more foods, but who has not yet tried peanut	- Possible elevated risk for an additional food allergy in someone who already has one food allergy - Parents may not introduce peanut without a positive test, leading to additional risk from delayed introduction	While the risk could be elevated over baseline, it is unclear if the absolute risk is elevated more than the low probability scenario of a 30% pre-test probability where a positive test was not shown to appreciably shift the post-test odds
A child in the first year of life with eczema suspected to be flared by one legume, and anaphylaxis to hummus who has not yet tried peanut	- Possible elevated risk for an additional food allergy in someone who already has one food allergy - Parents may not introduce peanut without a positive test, leading to additional risk from delayed introduction	- While the risk could be elevated over baseline, it is unclear if the absolute risk is elevated more than the low probability scenario of a 30% pre-test probability where a positive test was not shown to appreciably shift the post-test odds - By NIAID addendum criteria, the eczema does not make this child “high-risk”
A 6 month old child with mild eczema tolerating a milk based formula, who has not tried egg or peanut. Their older sibling has milk, egg, and peanut allergy	Parents may be reluctant to introduce peanut without a negative test, based on the experience with the older child, leading to additional risk from delayed introduction	While the risk could be elevated over baseline, it is unclear if the absolute risk is elevated more than the low probability scenario of a 30% pre-test probability where a positive test was not shown to appreciably shift the post-

	Some clinicians ascribe to older literature that has suggested the younger sibling may be at some degree of increased risk of developing peanut allergy, though such literature did not account for the highly important factor of delayed introduction.	test odds - By NIAID addendum criteria, the eczema does not make this child “high-risk” - Recent data has shown that testing the younger sibling is not cost effective until the prevalence of peanut allergy in siblings is shown to be >14% AND all such screened children also undergo an oral food challenge to provide a definitive outcome.
--	--	--

^aSee textbox 3 for explanation of what high, moderate, and low pre-test probability represent in the context of evaluating peanut allergy.

^bThese are hypothetical examples of situations that the workgroup members felt could represent potential scenarios that a clinician may evaluate under the context of a preference-sensitive care option. The choice of specific allergens, ages, and comorbidities are for illustration purposes only. Other allergens, ages, and comorbidities may represent possible presentations for consideration.

Table 4: Summary Statistics with 95% Confidence Intervals for SPT, sIgE, Ara h 2 Peanut Diagnostic Testing and Assessment of Reaction Severity

Diagnostic Test	Outcome	Sensitivity	Specificity	Positive Likelihood Ratio	Negative Likelihood Ratio
SPT 3mm	Diagnosis	0.97 (0.93-0.99)	0.46 (0.29-0.65)	1.82 (1.29-2.57)	0.05 (0.02-0.18)
sIgE 0.35 kU/L	Diagnosis	0.95 (0.91-0.97)	0.38 (0.28-0.48)	1.52 (1.3-1.77)	0.14 (0.08-0.24)
Ara h 2 sIgE 0.35 kU/L	Diagnosis	0.86 (0.81-0.89)	0.84 (0.79-0.89)	5.5 (3.99-7.56)	0.17 (0.13-0.23)
Ara h 2 sIgE 2 kU/L	Severe reaction	0.78 (0.69-0.85)	0.45 (0.28-0.63)	1.4 (1.08-1.83)	0.5 (0.37-0.66)
sIgE 50 kU/L	Severe reaction	0.39 (0.26-0.53)	0.89 (0.75-0.95)	3.4 (1.57-2.03)	0.69 (0.56-0.84)
SPT 10mm	Severe reaction	0.37 (0.22-0.55)	0.62 (0.44-0.77)	0.98 (0.71-1.35)	1 (0.84-1.22)
Sensitivity Analyses					
SPT 3mm ^a	SPT/sIgE Assessed in Same Study	0.98 (0.92-0.99)	0.5 (0.31-0.69)	1.94 (1.32-2.86)	0.04 (0.01-0.15)
sIgE 0.35 kU/L ^a	SPT/sIgE Assessed in Same Study	0.94 (0.9-0.97)	0.46 (0.32-0.6)	1.75 (1.35-2.26)	0.13 (0.07-0.21)
sIgE 0.35 kU/L ^a	sIgE/Ara h 2 Assessed in Same Study	0.95 (0.93-0.97)	0.3 (0.21-0.41)	1.36 (1.19-1.56)	0.47 (0.26-0.87)
Ara h 2 sIgE 0.35 kU/L ^a	sIgE/Ara h 2 Assessed in Same Study	0.85 (0.79-0.9)	0.86 (0.79-0.9)	5.87 (4.02-8.58)	0.18 (0.12-0.25)
SPT 3mm ^a	SPT/sIgE/Ara h 2 Assessed in Same Study	0.98 (0.89-1)	0.39 (0.22-0.6)	1.63 (1.19-2.23)	0.04 (0.01-0.25)
sIgE 0.35 kU/L ^a	SPT/sIgE/Ara h 2 Assessed in Same Study	0.95 (0.91-0.97)	0.4 (0.3-0.5)	1.58 (1.35-1.84)	0.12 (0.07-0.22)
Ara h 2 sIgE 0.35 kU/L ^a	SPT/sIgE/Ara h 2 Assessed in Same Study	0.83 (0.74-0.9)	0.79 (0.73-0.85)	4.03 (3.11-5.21)	0.21 (0.14-0.32)

^aTest sensitivity and specificity are being reported for pooled studies for the particular individual test evaluated in the setting where multiple tests were run simultaneously in patients undergoing oral food challenge. Please refer to table 7 for reporting of additional sensitivity analyses.

Table 5a: GRADE Table of Evidence Certainty, Skin Prick Testing

Question: Should peanut skin prick testing at a threshold of 3mm wheal size be used to diagnose peanut allergy in patients with known or suspected peanut allergy?

Total number of studies/patients entered into the analysis: 18 studies, 2124 patients

Bibliography: Abrahms 2017, Begin 2017, Bernard2003, Chinthrajah 2018, Comberati 2016, Dang 2012, DunnGalvin 2001, Johannsen 2016; Klemans Broekman 2013; Klemans Otte 2013, Leo 2015, Ludman 2013, Preece 2014, Rajput 2018, Rance 2003, Sampson 2017, Song 2015, Van Erp 2013.

Sensitivity	0.97 (95% CI: 0.93 to 0.90)		Prevalences				2%	30%	70%
Specificity	0.46 (95% CI: 0.29 to 0.65)								

Outcome	No of studies (No of patients)	Study design	Factors that may decrease certainty of evidence					Effect per 1,000 patients tested (95% CI)			Test accuracy CoE
			Risk of bias	Indirectness	Inconsistency	Imprecision	Publication bias	pre-test probability of 2%	pre-test probability of 30%	pre-test probability of 70%	
True positives (patients with peanut allergy)	18 studies 961 patients	cross-sectional (cohort type accuracy study)	serious ^a	not serious	serious ^b	not serious	none	19 (18 to 19)	291 (270 to 279)	679 (630 to 651)	⊕⊕○○ LOW
False negatives (patients incorrectly classified as not having peanut allergy)								1 (1 to 2)	9 (21 to 30)	21 (49 to 70)	
True negatives (patients without peanut allergy)	18 studies 1163 patients	cross-sectional (cohort type accuracy study)	serious ^a	not serious	serious ^b	not serious	none	451 (284 to 637)	322 (203 to 455)	138 (87 to 195)	⊕⊕○○ LOW
False positives (patients incorrectly classified as having peanut allergy)								529 (343 to 696)	378 (245 to 497)	162 (105 to 213)	

Explanations

a. Multiple studies had potential for selection bias due to non-consecutive, non-randomized, or otherwise unexplained selective enrollment of the study population within the potentially eligible cohort. Multiple studies with issues relative to the flow/timing of when index diagnostic test performed relative to the reference oral food challenge

b. I² for sensitivity was 90.1% and for specificity was 93%

Table 5b: GRADE Table of Evidence Certainty, Serum IgE Testing

Question: Should peanut serologic IgE testing at a threshold of >0.35 KU/L be used to diagnose peanut allergy in patients with suspected peanut allergy?

Total number of studies/patients entered into the analysis: 30 studies, 3983 patients

Bibliography: Abrahms 2017, Balmer Weber 2015, Begin 2017, Beigelman 2012, Bernard 2003, Beyer 2015, Chinthrajah 2018, Comberiati 2016, Dang 2012, DunnGalvin 2001, Ebisawa 2012, Ebisawa 2015, Eller 2013, Johannsen 2016; Klemans Broekman 2013; Klemans Otte 2013, Leo 2015, Lieberman 2013, Ludman 2013, Martinet 2016, Nicolaou 2011, Preece 2014, Rajput 2018, Rance 2003, Sampson 2017, Song 2015, Van Erp 2013, Wainstein 2007

Sensitivity	0.95 (95% CI: 0.91 to 0.97)							Prevalences			Test accuracy CoE
Specificity	0.38 (95% CI: 0.28 to 0.48)							2%	30%	70%	
Outcome	No of studies (No of patients)	Study design	Factors that may decrease certainty of evidence					Effect per 1,000 patients tested (95% CI)			Test accuracy CoE
			Risk of bias	Indirectness	Inconsistency	Imprecision	Publication bias	pre-test probability of 2%	pre-test probability of 30%	pre-test probability of 70%	
True positives (patients with peanut allergy)	30 studies 2046 patients	cross-sectional (cohort type accuracy study)	serious ^a	not serious	serious ^b	not serious	none	19 (18 to 19)	285 (273 to 291)	665 (637 to 679)	⊕⊕○○ LOW
False negatives (patients incorrectly classified as not having peanut allergy)								1 (1 to 2)	15 (9 to 27)	35 (21 to 63)	
True negatives (patients without peanut allergy)	30 studies 1937 patients	cross-sectional (cohort type accuracy study)	serious ^a	not serious	serious ^b	not serious	none	372 (274 to 470)	266 (196 to 336)	114 (84 to 144)	⊕⊕○○ LOW
False positives (patients incorrectly classified as having peanut allergy)								608 (510 to 706)	434 (364 to 504)	186 (156 to 216)	

Explanations

a. Multiple studies had potential for selection bias due to non-consecutive, non-randomized, or otherwise unexplained selective enrollment of the study population within the potentially eligible cohort. Multiple studies with issues relative to the flow/timing of when index diagnostic test performed relative to the reference oral food challenge

b. I² for sensitivity was 95.9% and for specificity was 92.8%

Table 6: GRADE Table of Evidence Certainty, Ara h 2 sIgE Testing

Question: Should Ara h 2 specific IgE at a threshold of >0.35 KU/L be used to diagnose peanut allergy in patients with suspected peanut allergy?

Total number of studies/patients entered into the analysis: 24 studies, 2289 patients

Bibliography: Balmer Weber 2015, Bernard 2003, Beyer 2015, Chintrajah 2018, Comberiati 2016, Dang 2012, Ebisawa 2012, Ebisawa 2015, Eller 2013, Glaumann 2012, Keet 2013, Klemans Broekman 2013, Klemans Otte 2013, Kukkonen 2015, Leo 2015, Lieberman 2013, Martinet 2016, Nicolaou 2011, Preece 2014, Rajput 2018, Rance 2003, Schots 2016, Suratannon 2013 Van Erp 2013

Sensitivity	0.86 (95% CI: 0.81 to 0.89)					<table><tr><td>Prevalences</td><td>2%</td><td>30%</td><td>70%</td></tr></table>					Prevalences	2%	30%	70%
Prevalences	2%	30%	70%											
Specificity	0.84 (95% CI: 0.79 to 0.89)													

Outcome	№ of studies (№ of patients)	Study design	Factors that may decrease certainty of evidence					Effect per 1,000 patients tested (95% CI)			Test accuracy CoE
			Risk of bias	Indirectness	Inconsistency	Imprecision	Publication bias	pre-test probability of 2%	pre-test probability of 30%	pre-test probability of 70%	
True positives (patients with peanut allergy)	24 studies 1336 patients	cross-sectional (cohort type accuracy study)	serious ^a	not serious	serious ^b	not serious	none	17 (16 to 18)	258 (243 to 267)	602 (567 to 623)	⊕⊕○○ LOW
False negatives (patients incorrectly classified as not having peanut allergy)								3 (2 to 4)	42 (33 to 57)	98 (77 to 133)	
True negatives (patients without peanut allergy)	24 studies 953 patients	cross-sectional (cohort type accuracy study)	serious ^a	not serious	serious ^b	not serious	none	823 (774 to 872)	588 (553 to 623)	252 (237 to 267)	⊕⊕○○ LOW
False positives (patients incorrectly classified as having peanut allergy)								157 (108 to 206)	112 (77 to 147)	48 (33 to 63)	

Explanations

a. Multiple studies had potential for selection bias due to non-consecutive, non-randomized, or otherwise unexplained selective enrollment of the study population within the potentially eligible cohort. Multiple studies with issues relative to the flow/timing of when index diagnostic test performed relative to the reference oral food challenge

b. I² for sensitivity was 81.4 and specificity was 69.7

Table 7a: GRADE Table of Evidence Certainty, Ara h 2 sIgE to Assess Reaction Severity

Question: Should Ara h 2 specific IgE at a threshold of >2 KU/Lbe used to diagnose severe peanut allergy in patients with suspected peanut allergy?

Total number of studies/patients entered into the analysis: 10 studies, 845 patients

Bibliography: Balmer Weber 2015, Chintrajah 2018, Dang 2012, Glaumann 2012, Klemans Broekman 2013; Kukkonen 2015, Leo 2015, Preece 2014, Rajput 2018, Van Erp 2013

Sensitivity	0.78 (95% CI: 0.69 to 0.85)	Prevalences		
Specificity	0.45 (95% CI: 0.28 to 0.63)	2%	30%	70%

Outcome	No of studies (No of patients)	Study design	Factors that may decrease certainty of evidence					Effect per 1,000 patients tested (95% CI)			Test accuracy CoE
			Risk of bias	Indirectness	Inconsistency	Imprecision	Publication bias	pre-test probability of 2%	pre-test probability of 30%	pre-test probability of 70%	
True positives (patients with severe peanut allergy)	10 studies 308 patients	cross-sectional (cohort type accuracy study)	serious ^a	not serious	very serious ^{b,c,d}	not serious	none	16 (14 to 17)	234 (207 to 255)	546 (483 to 595)	⊕○○○ ○ VERY LOW
False negatives (patients incorrectly classified as not having severe peanut allergy)								4 (3 to 6)	66 (45 to 93)	154 (105 to 217)	
True negatives (patients without severe peanut allergy)	10 studies 380 patients	cross-sectional (cohort type accuracy study)	serious ^a	not serious ^b	very serious ^{b,c,d}	not serious	none	441 (274 to 617)	315 (196 to 441)	135 (84 to 189)	⊕○○○ ○ VERY LOW
False positives (patients incorrectly classified as having severe peanut allergy)								539 (363 to 706)	385 (259 to 504)	165 (111 to 216)	

Explanations

a. Multiple studies had potential for selection bias due to non-consecutive, non-randomized, or otherwise unexplained selective enrollment of the study population within the potentially eligible cohort.

Multiple studies with issues relative to the flow/timing of when index diagnostic test performed relative to the reference oral food challenge

b. I² for sensitivity was 68.7% and for specificity was 91.6%

c. The heterogeneity for the estimate was very high

d. The criteria to assess severity was not uniform among all studies included

Table 7b: GRADE Table of Evidence Certainty, Peanut sIgE to Assess Reaction Severity

Question: Should peanut serologic IgE testing at a threshold of >50 KU_A/Lbe used to diagnose severe peanut allergy in patients with suspected peanut allergy?

Total number of studies/patients entered into the analysis: 13 studies, 1051 patients

Bibliography: Chinthrajah 2018, Dang 2012, DunnGalvin 2001, Glaumann 2012, Klemans Broekman 2013; Lewis 2005, Peeters 2007, Preece 2014, Rajput 2018, Song 2015, Van Erp 2013, Wainstein 2007, Wensing 2002

Sensitivity	0.39 (95% CI: 0.26 to 0.53)		Prevalences 2%30%70%			
Specificity	0.89 (95% CI: 0.75 to 0.95)					

Outcome	Ne of studies (Ne of patients)	Study design	Factors that may decrease certainty of evidence					Effect per 1,000 patients tested (95% CI)			Test accuracy CoE
			Risk of bias	Indirectness	Inconsistency	Imprecision	Publication bias	pre-test probability of 2%	pre-test probability of 30%	pre-test probability of 70%	
True positives (patients with severe peanut allergy)	13 studies 256 patients	cross-sectional (cohort type accuracy study)	serious ^a	not serious	very serious ^{b,c,d}	not serious	none	8 (5 to 11)	117 (78 to 159)	273 (182 to 371)	⊕○○○ VERY LOW
False negatives (patients incorrectly classified as not having severe peanut allergy)								12 (9 to 15)	183 (141 to 222)	427 (329 to 518)	
True negatives (patients without severe peanut allergy)	13 studies 795 patients	cross-sectional (cohort type accuracy study)	serious ^a	not serious	very serious ^{b,c,d}	not serious	none	872 (735 to 931)	623 (525 to 665)	267 (225 to 285)	⊕○○○ VERY LOW
False positives (patients incorrectly classified as having severe peanut allergy)								108' (49 to 245)	77 (35 to 175)	33 (15 to 75)	

Explanations

a. Multiple studies had potential for selection bias due to non-consecutive, non-randomized, or otherwise unexplained selective enrollment of the study population within the potentially eligible cohort. Multiple studies with issues relative to the flow/timing of when index diagnostic test performed relative to the reference oral food challenge

b. I² for sensitivity was 75.7% and for specificity was 90.9%

c. The criteria to assess severity was not uniform among all studies

d. The heterogeneity for the estimate was very high

Table 7c: GRADE Table of Evidence Certainty, Peanut sIgE to Assess Reaction Severity

Question: Should peanut skin prick testing at a threshold of 10mm wheal size be used to diagnose severe peanut allergy in patients with suspected peanut allergy?

Total number of studies/patients entered into the analysis: 12 studies, 737 patients

Bibliography: Chinthrajah 2018, Dang 2012, DunnGalvin 2001, Klemans Broekman 2013; Leo 2015, Lewis 2005; Preece 2014, Rajput 2018, Song 2015, Van Erp 2013, Wainstein 2010, Wensing 2002

Sensitivity	0.37 (95% CI: 0.22 to 0.55)	Prevalences				2%	30%	70%
Specificity	0.62 (95% CI: 0.44 to 0.77)							

Outcome	No of studies (No of patients)	Study design	Factors that may decrease certainty of evidence					Effect per 1,000 patients tested (95% CI)			Test accuracy CoE
			Risk of bias	Indirectness	Inconsistency	Imprecision	Publication bias	pre-test probability of 2%	pre-test probability of 30%	pre-test probability of 70%	
True positives (patients with severe peanut allergy)	12 studies 166 patients	cross-sectional (cohort type accuracy study)	serious ^a	not serious	very serious ^{b,c,d}	not serious	none	7 (4 to 11)	111 (66 to 165)	259 (154 to 385)	⊕○○○ VERY LOW
False negatives (patients incorrectly classified as not having severe peanut allergy)								13 (9 to 16)	189 (135 to 234)	441 (315 to 546)	
True negatives (patients without severe peanut allergy)	12 studies 571 patients	cross-sectional (cohort type accuracy study)	serious ^a	not serious	very serious ^{b,c,d}	not serious	none	608 (431 to 755)	434 (308 to 539)	186 (132 to 231)	⊕○○○ VERY LOW
False positives (patients incorrectly classified as having severe peanut allergy)								372 (225 to 549)	266 (161 to 392)	114 (69 to 168)	

Explanations

a. Multiple studies had potential for selection bias due to non-consecutive, non-randomized, or otherwise unexplained selective enrollment of the study population within the potentially eligible cohort.

Multiple studies with issues relative to the flow/timing of when index diagnostic test performed relative to the reference oral food challenge

b. I^2 for sensitivity was 64% and for specificity was 87.9%

c. The criteria to assess severity was not uniform among all studies included

d. The heterogeneity for the estimate was very high

Table 8: Additional Sensitivity Analyses

Test	Outcome	Analyses	Sensitivity	Specificity	Positive LR	Negative LR
SPT 3mm	Diagnosis	Exclusion of studies with high risk of bias	0.96	0.48	1.85	0.08
		Pediatric studies only	0.97	0.52	2.02	0.06
		Open OFC studies only	0.96	0.53	2.04	0.08
		DBPCFC studies only	0.99	0.38	1.60	0.03
		European studies only	0.98	0.56	2.23	0.04
		Non-European studies only	0.97	0.32	1.43	0.09
slgE >0.35	Diagnosis	Exclusion of studies with high risk of bias	0.96	0.44	1.71	0.09
		Pediatric studies only	0.94	0.41	1.59	0.15
		Open OFC studies only	0.94	0.4	1.57	0.15
		DBPCFC studies only	0.97	0.42	1.67	0.07
		European studies only	0.95	0.38	1.53	0.13
		Non-European studies only	0.95	0.37	1.51	0.14
Ara h 2 slgE >0.35	Diagnosis	Exclusion of studies with high risk of bias	0.86	0.81	4.53	0.17
		Pediatric studies only	0.85	0.85	5.67	0.18
		Open OFC studies only	0.85	0.85	5.67	0.18
		DBPCFC studies only	0.87	0.83	5.12	0.16
		European studies only	0.88	0.85	5.87	0.14
		Non-European studies only	0.83	0.84	5.19	0.20
Ara h 2 slgE >2	Severity	Exclusion of studies with high risk of bias	0.75	0.42	1.29	0.60
		Pediatric studies only	0.72	0.49	1.41	0.57
		Open OFC only	0.64	0.43	1.12	0.84
		DBPCFC only	0.8	0.44	1.43	0.45
		European studies only	0.77	0.43	1.35	0.53
		Non-European studies only	0.71	0.44	1.27	0.66
slgE >50	Severity	Exclusion of studies with high risk of bias	0.36	0.88	3.00	0.73
		Pediatric studies only	0.38	0.92	4.75	0.67
		Open OFC only	0.29	0.97	9.67	0.73
		DBPCFC studies only#	0.47	0.71	1.62	0.75
		European studies only	0.38	0.86	2.71	0.72
		Non-European studies only	0.44	0.92	5.50	0.61
SPT 10mm	Severity	Exclusion of studies with high risk of bias	0.41	0.57	0.95	1.04
		Pediatric studies only	0.29	0.71	1.00	1.00
		Open OFC studies only	0.26	0.69	0.84	1.07
		DBPCFC studies only#	0.62	0.41	1.05	0.93
		European studies only	0.39	0.67	1.18	0.91
		Non-European studies only	0.36	0.59	0.88	1.08

eTable 1: Simulation Model Inputs

Variable	Model Reference (sensitivity range)	Source
US Life Table	National Vital Statistics Reports, April 2017	Arias E, Heron M, Xu J. United States Life Tables, 2013. National Vital Statistics Reports 2017; 66(3): 1-64.
Testing characteristics	Skin prick testing: Sn: 0.97 (0.86-0.98); Sp 0.46 (0.17-0.67) Ara h 2: Sn 0.86 (0.72 – 0.90); Sp 0.84 (0.65-0.87) Whole peanut sIgE: 0.95 (0.89-0.97); Sp 0.38 (0.23-0.49)	Meta-analysis
Food allergy fatality	5-19 years: 3.25 per million person years (0.3 – 30) 20 years and older: 1.81 per million person years (1.81-18.1)	Umasunthar T, Leonardi-Bee, Hodes M, et al. Incidence of fatal food anaphylaxis in people with food allergy: a systematic review and meta-analysis. Clinical & Experimental Allergy. 2013; 43: 1333-1341.
Rate of accidental peanut exposure and symptoms in peanut allergic persons	7% per year (5%-45%)	Neuman-Sunshine D, Eckman J, Keet C, Matsui E, Peng R, et al. The natural history of peanut allergy. Ann Allergy Asthma Immunol. 2012; 108: 326-331.
Rate of emergency room visit for severe symptoms in peanut allergic persons	1% per year (0.5%-35%)	Neuman-Sunshine D, Eckman J, Keet C, Matsui E, Peng R, et al. The natural history of peanut allergy. Ann Allergy Asthma Immunol. 2012; 108: 326-331.
Hospitalization following emergency room visit for anaphylaxis	35% (5%-45%)	Robinson M, Greenhawt M, Stukus D. Factors associated with epinephrine administration for anaphylaxis in children before arrival to the emergency department. Ann Allergy Asthma Immunol. 2017; 119: 164-169.
Primary care visits (mean incremental annual cost for food allergy diagnosis)	\$102 (\$94-\$105)	Gupta R, Holdford D, Bilaver L, Dyer A, Holl JL, Meltzer D. The economic impact of childhood food allergy in the United States. JAMA Pediatr 2013;167:1026-31; US Department of Labor, Bureau of Labor Statistics. Available from www.bls.gov .
Allergist visits for food allergy (mean incremental annual cost for food allergy diagnosis)	\$151 (\$140 - \$152)	Gupta R, Holdford D, Bilaver L, Dyer A, Holl JL, Meltzer D. The economic impact of childhood food allergy in the United States. JAMA Pediatr 2013;167:1026-31; US Department of Labor, Bureau of Labor Statistics. Available from www.bls.gov .
Nutritionist visits for food allergy (per year)	\$17 (\$15 - \$18)	Gupta R, Holdford D, Bilaver L, Dyer A, Holl JL, Meltzer D. The economic impact of childhood food allergy in the United States. JAMA Pediatr 2013;167:1026-31; US Department of Labor, Bureau of Labor

		Statistics. Available from www.bls.gov .
Alternative provider visits for food allergy (per year)	\$25 (\$22 - \$27)	Gupta R, Holdford D, Bilaver L, Dyer A, Holl JL, Meltzer D. The economic impact of childhood food allergy in the United States. JAMA Pediatr 2013;167:1026-31; US Department of Labor, Bureau of Labor Statistics. Available from www.bls.gov .
Incremental annual grocery costs (living with food allergy)	\$315 (\$290-330)	Gupta R, Holdford D, Bilaver L, Dyer A, Holl JL, Meltzer D. The economic impact of childhood food allergy in the United States. JAMA Pediatr 2013;167:1026-31; US Department of Labor, Bureau of Labor Statistics. Available from www.bls.gov .
Job-related opportunity costs from food allergy (per year)	\$2,637 (\$0 - \$2,697)	Gupta R, Holdford D, Bilaver L, Dyer A, Holl JL, Meltzer D. The economic impact of childhood food allergy in the United States. JAMA Pediatr 2013;167:1026-31; US Department of Labor, Bureau of Labor Statistics. Available from www.bls.gov .
Personal epinephrine auto-injector	\$726 (\$100-\$800)	Shaker M, Bean K, Verdi M. Economic evaluation of epinephrine autoinjectors for peanut allergy. Ann Allergy Asthma Immunol. 2017; 119(2): 160-163. US Department of Labor, Bureau of Labor Statistics. CPI Inflation calculator. Accessed at https://data.bls.gov on 9/2/18.
sIgE / ara h 2 IgE testing	\$17 per test (\$10-\$117)	Healthcare Bluebook. www.healthcarebluebook.com . Accessed 11/22/18
Skin test cost	\$24 (\$10-\$40)	Physician Fee schedule. Available from http://www.cms.gov/ . Accessed 10/3/17.
Hospitalization	\$5,991 (\$5,732-\$6,066)	Patel DA, Holdford DA, Edwards E, Carroll NV. Estimating the economic burden of food-induced allergic reactions and anaphylaxis in the United States. J Allergy Clin Immunol 2011;128:110-5 e5 US Department of Labor, Bureau of Labor Statistics. CPI Inflation calculator. Accessed at https://data.bls.gov on 9/2/18.
ED visit	\$702 (\$689-\$710)	Patel DA, Holdford DA, Edwards E, Carroll NV. Estimating the economic burden of food-induced allergic reactions and anaphylaxis in the United States. J Allergy Clin Immunol 2011;128:110-5 e5 US Department of Labor, Bureau of Labor Statistics. CPI Inflation calculator. Accessed at https://data.bls.gov on 9/2/18.

Supervised oral food challenge	\$123 (\$110-600)	Doctors office visits billing and charges. Dartmouth-Hitchcock. http://www.dartmouth-hitchcock.org . Accessed March 10, 2017 US Department of Labor, Bureau of Labor Statistics. CPI Inflation calculator. Accessed at https://data.bls.gov on 4/21/19
Start age	0 years (0 years to 8 years)	
Negative health state influence for food allergy and food anaphylaxis	-0.09 (-0.02 - -0.11)	Carroll AE, Downs SM. Improving decision analyses: parent preferences (utility values) for pediatric health outcomes. J Pediatr 2009;155:21-5, 5 e1-5.
Cycle length	1 year	
Time Horizon	20 years (5 years – 80 years)	
Peanut allergy pre-test probability	14% (3% - 90%)	
Annual discount rate	0.03 (0-0.03)	
Probability of spontaneous tolerance	22% (0-22%)	
Probability of identifying false positive test over model horizon	20% (5%-20% over 5-20 years)	

eTable 2: Cost-Effectiveness Comparisons of Use of Peanut SPT, sIgE, and Ara h 2 sIgE Testing


	Cost	QALY	Net Monetary Benefit	Total Rxn	Anaphylaxis	Anaphylaxis Fatality
<u>3% peanut allergy pre-test probability</u>						
Skin Prick Test						
Mean	\$20,734.48	14.43	-\$20,734.48	0.0341	0.0047	0.0000
Std Deviation	\$23,902.82	1.36	\$23,902.82	0.2833	0.0745	0.0000
Ara h 2						
Mean	\$7,669.24	14.79	-\$7,669.24	0.0379	0.0049	0.0000
Std Deviation	\$17,355.08	1.33	\$17,355.08	0.3008	0.0783	0.0000
Whole peanut sIgE						
Mean	\$23,466.54	14.35	-\$23,466.54	0.0345	0.0048	0.0000
Std Deviation	\$24,165.42	1.35	\$24,165.42	0.2852	0.0756	0.0000
<u>14% peanut allergy pre-test probability</u>						
Skin Prick Test						
Mean	\$23,859.49	14.36	-\$23,859.49	0.1555	0.0213	0.0000
Std Deviation	\$25,361.09	1.33	\$25,361.09	0.5784	0.1574	0.0000
Ara h 2						
Mean	\$12,329.23	14.69	-\$12,329.23	0.1725	0.0223	0.0000
Std Deviation	\$22,237.68	1.32	\$22,237.68	0.6169	0.1614	0.0000
Whole peanut sIgE						
Mean	\$26,289.04	14.29	-\$26,289.04	0.1581	0.0212	0.0000
Std Deviation	\$25,304.83	1.32	\$25,304.83	0.5836	0.1574	0.0000
<u>75% peanut allergy pre-test probability</u>						
Skin Prick Test						
Mean	\$41,680.67	13.99	-\$41,680.67	0.8479	0.1200	0.0000
Std Deviation	\$25,973.46	1.24	\$25,973.46	1.1182	0.3571	0.0000
Ara h 2						
Mean	\$38,191.62	14.09	-\$38,191.62	0.9273	0.1206	0.0000
Std Deviation	\$27,947.58	1.28	\$27,947.58	1.1690	0.3588	0.0000

Whole peanut slgE						
Mean	\$42,378.21	13.97	-\$42,378.21	0.8632	0.1205	0.0000
Std Deviation	\$25,494.62	1.24	\$25,494.62	1.1286	0.3579	0.0000


Figure 1: PRISMA Diagrams

A. B.

PRISMA Flow Diagram




PRISMA Flow Diagram



C.

PRISMA Flow Diagram



D.

PRISMA Flow Diagram




Figure 10: Risk of Bias Assessment

Study	Year	Bias				Applicability		
		Patient Selection	Index Test	Reference Standard	Flow and Timing	Patient Selection	Index Test	Reference Standard
Abrams	2017	Red	Green	Red	Red	Yellow	Green	Green
Balmer Weber	2015	Red	Green	Green	Red	Red	Green	Green
Begin	2017	Green	Green	Green	Green	Green	Green	Green
Beigelman	2012	Red	Green	Red	Red	Red	Green	Green
Bernard	2003	Green	Green	Red	Green	Green	Yellow	Green
Beyer	2015	Yellow	Green	Red	Green	Green	Green	Green
Chinthrajah	2018	Red	Green	Yellow	Red	Red	Green	Yellow
Comberiati	2016	Yellow	Green	Red	Green	Yellow	Green	Green
Dang	2012	Red	Green	Red	Green	Yellow	Green	Green
DunnGalvin	2011	Red	Green	Green	Red	Yellow	Green	Green
Ebisawa	2015	Yellow	Green	Green	Red	Green	Green	Green
Ebisawa	2012	Green	Yellow	Green	Green	Green	Yellow	Green
Eller	2013	Red	Yellow	Green	Green	Yellow	Yellow	Green
Glaumann	2012	Red	Yellow	Green	Green	Red	Yellow	Green
Guilloux	2009	Green	Yellow	Green	Red	Yellow	Yellow	Green
Gupta	2014	Red	Yellow	Green	Green	Red	Green	Yellow
Johannsen	2016	Red	Yellow	Green	Green	Red	Yellow	Green
Keet	2013	Red	Yellow	Yellow	Yellow	Red	Yellow	Yellow
Klemans Blom	2015	Red	Yellow	Yellow	Yellow	Yellow	Green	Yellow
Klemans Broekman	2013	Red	Yellow	Green	Red	Green	Yellow	Green
Klemans Liu	2013	Red	Yellow	Red	Red	Red	Yellow	Red
Klemans Otte	2013	Red	Yellow	Green	Green	Green	Green	Green
Kukkonen	2015	Yellow	Yellow	Yellow	Yellow	Yellow	Green	Green
Leo	2015	Red	Yellow	Green	Yellow	Red	Yellow	Green
Lewis	2005	Yellow	Yellow	Yellow	Yellow	Yellow	Green	Green
Lieberman	2013	Red	Yellow	Yellow	Red	Yellow	Green	Yellow
Ludman	2013	Green	Green	Green	Yellow	Yellow	Green	Green
Martinet	2016	Yellow	Yellow	Green	Yellow	Yellow	Yellow	Green
Nicolaou	2011	Red	Yellow	Yellow	Red	Green	Green	Red
Peeters	2007	Red	Yellow	Green	Yellow	Green	Green	Green
Perry	2004	Red	Green	Yellow	Red	Yellow	Green	Green
Preece	2014	Red	Yellow	Yellow	Yellow	Green	Green	Green
Rajput	2018	Red	Green	Green	Yellow	Yellow	Green	Green
Rance	2003	Yellow	Green	Green	Green	Green	Green	Green
Schots	2016	Red	Green	Green	Yellow	Yellow	Green	Green
Song	2015	Red	Green	Green	Green	Red	Green	Green
Suratannon	2013	Red	Green	Yellow	Yellow	Red	Green	Yellow
Van Erp	2013	Red	Green	Green	Yellow	Green	Green	Green
Wainstein	2007	Red	Green	Green	Green	Green	Green	Green
Wainstein	2010	Red	Green	Yellow	Green	Red	Green	Yellow
Wensing	2002	Red	Green	Yellow	Yellow	Red	Green	Yellow

Red: high risk Yellow: Unclear risk Green: low risk

Figure 2: Summary Forest Plots for Sensitivity and Specificity of Skin Prick Testing at 3mm and slgE testing at 0.35KU_A/L






Figure 3: Fagan Nomograms for SPT 3mm Performance at Low, Moderate, and High Pre-Test Probability

a



b



c






Figure 4: Fagan Nomograms for sIgE 0.35KU_A/L Performance at Low, Moderate, and High Pre-Test Probability

a



b



c







Figure 5: Summary Forest Plots for Sensitivity and Specificity of Ara h 2 sIgE testing at 0.35KU_A/L

Figure 6 Fagan Nomograms for Ara h 2 sIgE 0.35KU_A/L Performance at Low, Moderate, and High Pre-Test Probability

a



b



c





Figure 7: Summary Forest Plots for Sensitivity and Specificity of Ara h 2 sIgE testing at 2 KU_A/L Indicating a Severe Reaction









Figure 8: Summary Forest Plots for Sensitivity and Specificity of sIgE testing at 50 KU_A/L Indicating a Severe Reaction


Figure 9: Summary Forest Plots for Sensitivity and Specificity of Skin Prick Testing at 10mm Indicating a Severe Reaction



eFigure 10 : Summary ROC for SPT 3mm, sIgE 0.35KU_A/L, and Ara h 2 0.35KU_A/L


eFigure 11: Summary ROC for Reaction Severity Ara h 2 2KU_A/L, slgE 50 KU_A/L, and SPT 3mm

eFigure 7: Summary Forest Plots for Sensitivity and Specificity of Skin Prick Testing at 3mm, sIgE testing at 0.35KU_A/L and Ara h 2 sIgE .35KU_A/L When All Tests Run Simultaneously





eFigure : Summary Forest Plots for Risk of Bias Removed Sensitivity and Specificity of Skin Prick Testing at 3mm and sIgE testing at 0.35KU_A/L

a




b




eFigure 9: Summary Forest Plots for Risk of Bias Removed Sensitivity and Specificity of Ara h 2 sIgE testing at 0.35KU_A/L


eFigure 1: Outcomes of Using Diagnostic Testing for Peanut Allergy




eFigure 2: Decision Model for Assessing the Cost Effectiveness of the Use of Diagnostic Testing




eFigure 3: Deterministic Sensitivity Analysis of the Threshold of Ara h 2 Specificity Where Stand-Alone Use Is Cost-Effective




eFigure 4: Probabilistic Sensitivity Analysis of Stand Alone Ara h 2 Use




eFigure 5: SPT 3mm, slgE 0.35 KU_A/L, and Ara h 2 slgE 0.35KU_A/L Performance at a 50% Pre-Test Probability

SPT




slgE




Ara h 2

eFigure 6 : Summary Forest Plots for Sensitivity and Specificity of Skin Prick Testing at 3mm and sIgE testing at 0.35KU_A/L When Both Tests Run Simultaneously

A



B



QUESTION

In patients presenting for evaluation of suspected peanut allergy, which of the three tests—Skin prick test, sIgE to whole peanut, or Ara h2 would provide the most diagnostic accuracy as determined by the more optimal positive/negative likelihood ratio?

POPULATION:	Adults and children presenting for evaluation of peanut allergy
INTERVENTION:	Using peanut skin prick testing (SPT), serum specific IgE to whole peanut (sIgE), or Ara h 2 serum specific IgE (Ara h 2 sIgE) to determine peanut sensitization to assist in the diagnosis peanut allergy
COMPARATOR	Oral Food challenge
OUTCOMES	Diagnostic accuracy of peanut allergy testing as determined by the more optimal positive/negative likelihood ratio.
PURPOSE OF THE TEST:	TO DETECT SENSITIZATION TO PEANUT PROTEIN
ROLE OF THE TEST:	DETECTABLE OR NON-DETECTABLE SENSITIZATION CAN BE USED TO HELP INCREASE OR DECREASE THE LIKELIHOOD OF PEANUT ALLERGY BASED ON THE PRESENTING PATIENT HISTORY
LINKED-RECOMMENDATIONS	AD LIBITUM PEANUT INGESTION, SUPERVISED ORAL FOOD CHALLENGE TO PEANUT, PEANUT AVOIDANCE WITH/WITHOUT TREATMENT
ANTICIPATED OUTCOMES:	Appropriate selection of the test to improve the likelihood of correct diagnosis
SETTING:	Patients presenting a to an allergist or a primary care provider for evaluation of suspected peanut allergy
PERSPECTIVE:	Patients and clinicians want to know the best diagnostic test to perform to help confirm the patients' history of suspected peanut allergy. Clinicians want to know when an oral food challenge should be performed, when it is safe to advise a patient to eat peanut, and when peanut should be avoided due to risk of an allergic reaction and consider seeking treatments
BACKGROUND:	Peanut allergy affects between 1.4% to 4.5% of the US population. This can be a potentially severe and life-long condition associated with reduced health and economic outcomes. Soon to be approved treatments can offer limited protection to a small amount of peanut but no therapy can cure the condition, but being on treatment still implies the patient is peanut allergic and must otherwise avoid intended peanut ingestion and carry emergency medication. Approximately 20%-34% will outgrow their peanut allergy. With the advent of available treatment options, it is imperative to understand how to use available diagnostic tests and interpret their results to aid in making an accurate diagnosis of peanut allergy.
SUBGROUPS:	Persons with a severe allergic reaction occurring during an observed oral food challenge; Persons with low, medium, and high pre-test probability of a suspected peanut allergy
CONFLICT OF INTERESTS:	See main document

ASSESSMENT

Problem Is the problem a priority?		
JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
☐ No ☐ Probably no ☐ Probably yes ☒ Yes ☐ Varies ☐ Don't know	The following studies support that peanut allergy, among other food allergies, is a major public health issue for children and adults in westernized countries. National Academies of Sciences E, Medicine. Finding a Path to Safety in Food Allergy: Assessment of the Global Burden, Causes, Prevention, Management, and Public Policy. Washington, DC: The National Academies Press; 2017. Sicherer SH, Sampson HA. Food allergy: A review and update on epidemiology, pathogenesis, diagnosis, prevention, and	The precise prevalence of peanut allergy is uncertain, given variation in the methods used to determine prevalence, and practice variation where detectable sensitization may be considered as clinical allergy without a history of symptoms arising from peanut ingestion in some circumstances. This may complicate using peanut allergy prevalence as an estimation of

	management. The Journal of allergy and clinical immunology 2018;141:41-58. Sampson HA, Aceves S, Bock SA, James J, Jones S, Lang D, et al. Food allergy: A practice parameter update—2014. Journal of Allergy and Clinical Immunology 2014;134:1016-25.e43. Osborne NJ, Koplin JJ, Martin PE, Gurrin LC, Lowe AJ, Matheson MC, et al. Prevalence of challenge-proven IgE-mediated food allergy using population-based sampling and predetermined challenge criteria in infants. The Journal of allergy and clinical immunology 2011;127:668-76.e1-2. Gupta RS, Springston EE, Warrier MR, Smith B, Kumar R, Pongracic J, et al. The prevalence, severity, and distribution of childhood food allergy in the United States. Pediatrics 2011;128:e9-17. Chafen JJ, Newberry SJ, Riedl MA, Bravata DM, Maglione M, Suttrop MJ, et al. Diagnosing and managing common food allergies: a systematic review. JAMA 2010;303:1848-56 Klemans RJ, van Os-Medendorp H, Blankestijn M, Bruijnzeel-Koomen CA, Knol EF, Knulst AC. Diagnostic accuracy of specific IgE to components in diagnosing peanut allergy: a systematic review. Clinical and experimental allergy : journal of the British Society for Allergy and Clinical Immunology 2015;45:720-30. Gupta RS, Warren CM, Smith BM, et al. The public health impact of parent-reported childhood food allergies in the United States. Pediatrics. 2018;142:e20181235.	pre-test probability difficult in certain contexts. The FDA has fast-tracked the development of commercial therapies to address the growing prevalence of peanut allergy. Overdiagnosis and unwarranted practice variation can create a significant healthcare burden and family quality of life. All stakeholders desire a fast, reliable diagnostic tool.
--	---	--

Test accuracy

How accurate is the test?

JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS																				
○ Very inaccurate ○ Inaccurate ○ Accurate ○ Very accurate ● Varies ○ Don't know	Pooled Sensitivity/Specificity positive/negative likelihood ratio (with 95% CI): <table><tr><th>Test</th><th>Sensitivity</th><th>Specificity</th><th>Positive Likelihood</th><th>Negative Likelihood</th></tr><tr><td>SPT 3mm</td><td>0.97 (0.93-0.99)</td><td>0.46 (0.29-0.65)</td><td>1.82 (1.29-2.57)</td><td>0.05 (0.02-0.18)</td></tr><tr><td>slgE 0.35 kU/L</td><td>0.95 (0.91-0.97)</td><td>0.38 (0.28-0.48)</td><td>1.52 (1.3-1.77)</td><td>0.14 (0.08-0.24)</td></tr><tr><td>Ara h 2 slgE 0.35 kU/L</td><td>0.86 (0.81-0.89)</td><td>0.84 (0.79-0.89)</td><td>5.5 (3.99-7.56)</td><td>0.17 (0.13-0.23)</td></tr></table> Both SPT and whole peanut slgE have high pooled sensitivity but relatively poor specificity for the diagnosis of peanut allergy proven by oral food challenge. Ara h 2 slgE has somewhat reduced sensitivity to SPT or slgE, but has enhanced specificity relative to these tests, and the most optimal positive/negative likelihood ration combination. Despite the individual test precision, the interpretation of the test of choice is highly dependent on an adequate suspicion of significant pre-test probability, reflected by a reasonable history that the patient had ingested peanut and demonstrated symptoms characteristic of an IgE mediated reaction. Using thresholds evaluated in the present meta-analysis, all three tests are suboptimal screening measures due to poor specificity and a high likelihood of detecting asymptomatic sensitization, potentially resulting in a false positive diagnosis. Irrespective of the test used, there are limited situations where a positive result alone relays adequate post-test odds of a peanut allergy diagnosis without the need to do a follow up oral food challenge. Based on the current analysis, in situations of low to moderate pre-test probability, detectable Ara h 2 slgE translates to higher post-test odds of peanut allergy, compared to SPT and whole peanut slgE. <u>Where there is moderate to high pre-test probability, choice of test is less likely to influence the post-test odds, as best illustrated by the Faqan nomograms in figures 3, 4 and 6.</u>	Test	Sensitivity	Specificity	Positive Likelihood	Negative Likelihood	SPT 3mm	0.97 (0.93-0.99)	0.46 (0.29-0.65)	1.82 (1.29-2.57)	0.05 (0.02-0.18)	slgE 0.35 kU/L	0.95 (0.91-0.97)	0.38 (0.28-0.48)	1.52 (1.3-1.77)	0.14 (0.08-0.24)	Ara h 2 slgE 0.35 kU/L	0.86 (0.81-0.89)	0.84 (0.79-0.89)	5.5 (3.99-7.56)	0.17 (0.13-0.23)	Skin prick testing is the traditional test of choice of the board-certified allergist and otolaryngologist with focused allergy sub-training. It is a point-of-care test that is easy to perform, exceptionally safe, inexpensive, and reliable under contexts where there is a reasonable suspicion for allergy. The advantage of this test is that the clinician can detect if sensitization is present or absent during the visit, though ambient dermatographism can affect interpretation. Serologic tests are usually performed outside of the clinical encounter. In the patient where there is strong clinical suspicion for peanut allergy, detecting sensitization through the SPT at the time of the encounter can help make the diagnosis in real-time, and allow for the patient to be counseled on avoidance and anaphylaxis management. There may be consideration that given the enhanced likelihood ratio combination that Ara h 2 is the most optimal confirmatory test to be sent after detection of sensitization on skin prick testing.
Test	Sensitivity	Specificity	Positive Likelihood	Negative Likelihood																		
SPT 3mm	0.97 (0.93-0.99)	0.46 (0.29-0.65)	1.82 (1.29-2.57)	0.05 (0.02-0.18)																		
slgE 0.35 kU/L	0.95 (0.91-0.97)	0.38 (0.28-0.48)	1.52 (1.3-1.77)	0.14 (0.08-0.24)																		
Ara h 2 slgE 0.35 kU/L	0.86 (0.81-0.89)	0.84 (0.79-0.89)	5.5 (3.99-7.56)	0.17 (0.13-0.23)																		

Desirable Effects

How substantial are the desirable anticipated effects?

JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
○ Trivial ○ Small ● Moderate ○ Large	In setting of high pre-test probability, detection of peanut sensitization with any of the 3 tests can significantly increase the post-test odds of a peanut allergy diagnosis as shown in Figures 4 and 5.	The main advantage to the SPT over serologic IgE tests is that this is a point-of-care test that can help facilitate a diagnosis being made during the encounter. No test is a substitute or surrogate for taking a good history.

○ Varies ○ Don't know		
Undesirable Effects How substantial are the undesirable anticipated effects?		
JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
○ Large ○ Moderate ○ Small ○ Trivial ● Varies ○ Don't know	In settings of low to moderate pre-test probability, detection of sensitization with any of the three tests still translates to a similar low to moderate post-test odds (figures 3, 4, and 6), and considering these results indicative of peanut allergy may significantly risk a false positive diagnosis.	Clinician Vantage: The level of sensitization above the positive threshold cannot be used to predict the risks of a future reaction. Likewise, test sensitization below the positive threshold in the setting of a history suggestive of high risk, cannot exclude peanut allergy. Test results, whether positive or negative, may still require an oral food challenge be performed to clarify the diagnosis Patient vantage: Patients may have variable preferences regarding having a false positive diagnosis than a false negative diagnosis, and therefore patients may prefer an oral food challenge after the test results are known, in particular when considering entering into possible treatment for peanut allergy. The clinician should be aware of the role for shared decision making and the need for decision-aids to help patients consider their options and to make the most appropriate decisions.
Certainty of the evidence of test accuracy What is the overall certainty of the evidence of test accuracy?		
JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
○ Very low ○ Low ● Moderate ○ High ○ No included studies	18 studies were pooled for evidence synthesis for use of SPT at a threshold of 3mm, with sensitivity of 97% and specificity of 46%. 30 studies were pooled for evidence synthesis for peanut sIgE at a threshold of >0.35 KU _A /L, with sensitivity of 95% and specificity of 38%. 24 studies were pooled for Ara h 2 sIgE >0.35 KU _A /L, with sensitivity of 0.86 and specificity of 0.84. There was high heterogeneity among the pooled studies, and serious risk of bias, but no serious risk of indirectness, imprecision, or inconsistency. Sensitivity analysis where studies with high risk for both patient selection and flow/timing were removed had similar pooled sensitivity and specificity for all three tests. Overall there is moderate certainty in the evidence for each of the 3 tests. (Please see tables 4 and 5).	Where there is high pre-test probability, detection of peanut sensitization using any of the 3 tests can greatly increase the post-test odds of a peanut allergy diagnosis. Absence of sensitization in such patients can be helpful in lowering the odds that peanut allergy is present. The choice of which test to use is also not crucial in this setting. The Fagan nomograms in figures 4 and 5 demonstrate how the likelihood ratios translate to post-test odds in these situations, and based on these post-test odds some clinicians may feel an oral food challenge is still necessary to confirm the diagnosis. Ara h 2 may perform better than SPT or sIgE where the pre-test probability is low to moderate, but is unlikely to allow the clinician and patient to be provided with the degree of certainty to where an oral food challenge would be unnecessary to confirm a diagnosis.
Certainty of the evidence of test's effects What is the overall certainty of the evidence for any critical or important direct benefits, adverse effects or burden of the test?		
JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
○ Very low ○ Low	There are no included studies that detail the overall certainty or importance of direct benefits, adverse effects or burden of the tests.	Please refer to the explanation in the above box. While one may question why patients with low suspicion for peanut allergy

○ Moderate ○ High ● No included studies		require testing, there may be a role for shared decision making where the risks and benefits of potential overdiagnosis vs. misdiagnosis are clearly explained, given some patients may clearly prefer a test be run, notwithstanding the pre-test probability.
---	--	---

Certainty of the evidence of management's effects

What is the overall certainty of the evidence of effects of the management that is guided by the test results?

JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
○ Very low ● Low ○ Moderate ○ High ○ No included studies	Identification of the trigger of a previous episode of anaphylaxis can lead to a reduction in the risk of future anaphylactic events. Treatment options based on a positive diagnosis of peanut allergy include avoidance and carriage of epinephrine. Additionally, for some patients there may be an opportunity for treatments that desensitize the patient to the point of being able to tolerate a low threshold dose of peanut. However, we have very low certainty in the evidence that by making a diagnosis of peanut allergy that the above described options provide an unequivocal benefit for the patient. National Academies of Sciences E, Medicine. Finding a Path to Safety in Food Allergy: Assessment of the Global Burden, Causes, Prevention, Management, and Public Policy. Washington, DC: The National Academies Press; 2017. Sicherer SH, Sampson HA. Food allergy: A review and update on epidemiology, pathogenesis, diagnosis, prevention, and management. The Journal of allergy and clinical immunology 2018;141:41-58. Sampson HA, Aceves S, Bock SA, James J, Jones S, Lang D, et al. Food allergy: A practice parameter update—2014. Journal of Allergy and Clinical Immunology 2014;134:1016-25.e43. Chu DK, Wood RA, French S, et al. Oral immunotherapy for peanut allergy (PACE): a systematic review and meta-analysis of efficacy and safety. Lancet. 2019;393:2222-2232. Robinson M, Greenhawt M, Stukus D. Factors associated with epinephrine administration for anaphylaxis in children before arrival to the emergency department. Ann Allergy Asthma Immunol. 2017;119:164-169. https://icer-review.org/wp-content/uploads/2018/12/ICER_PeanutAllergy_Final_Report_071019.pdf Shaker M, Greenhawt M. Estimation of health and economic benefits of commercial peanut immunotherapy products: a cost-effectiveness analysis. JAMA Netw Open. 2019;2:e193242.	

Certainty of the evidence of test result/management

How certain is the link between test results and management decisions?

JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
○ Very low ● Low ○ Moderate ○ High ○ No included studies	As reflected in the Fagan nomograms in figures 3, 4, and 6, an oral food challenge may often still be necessary to provide a definitive diagnosis and management strategy despite a positive test result, given that the systematic review suggests that even with very high pre-test probability, the post-test odds do not eclipse 90% (coming closest for the use of Ara h 2). Moreover, even with no detectable sensitization, the post-test odds are still 2-3%.	

Certainty of effects

What is the overall certainty of the evidence of effects of the test?

JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
-----------	-------------------	---------------------------

○ Very low ● Low ○ Moderate ○ High ○ No included studies	All three of the diagnostic tests for peanut allergy have high sensitivity though detection of peanut specific IgE does not always translate in to post-test odds sufficient enough to support a diagnosis of peanut allergy without a confirmatory oral food challenge. Conversely, the absence of detectable sensitization using any of these three tests should translate to very low post-test odds of a diagnosis of peanut allergy. Therefore, despite which test is used, in many cases an oral food challenge will still be indicated, and thus there is low certainty in the effects of the test in providing benefit for the patient from this perspective. Once a diagnosis is made through either testing or oral food challenge, there is low certainty in the benefits of the available treatment options. Chu DK, Wood RA, French S, et al. Oral immunotherapy for peanut allergy (PACE): a systematic review and meta-analysis of efficacy and safety. <i>Lancet</i>. 2019;393:2222-2232. https://icer-review.org/wp-content/uploads/2018/12/ICER_PeanutAllergy_Final_Report_071019.pdf Shaker M, Greenhawt M. Estimation of health and economic benefits of commercial peanut immunotherapy products: a cost-effectiveness analysis. <i>JAMA Netw Open</i>. 2019;2:e193242. National Academies of Sciences E, Medicine. Finding a Path to Safety in Food Allergy: Assessment of the Global Burden, Causes, Prevention, Management, and Public Policy. Washington, DC: The National Academies Press; 2017.	The use of these diagnostic tests at the stated thresholds (SPT 3 mm or greater, sIgE >0.35 KU_A/L, Ara h 2 > 2 KU_A/L) is most helpful in situations of high (>70%) pre-test probability in shifting the post-test odds appreciably, and can provide moderate to high certainty of a diagnosis. Testing should be undertaken with extreme caution in patients with low pre-test probability.
--	---	---

Values

Is there important uncertainty about or variability in how much people value the main outcomes?

JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
○ Important uncertainty or variability ○ Possibly important uncertainty or variability ● Probably no important uncertainty or variability ○ No important uncertainty or variability	Quality of life and qualitative research has indicated a negative effect based upon a poor understanding of the implications of diagnostic testing in terms of the severity and prognosis of the patient's allergy. Kao LM, Greenhawt MJ, Warren CM, et al. Parental and parent-perceived child interest in clinical trials for food allergen immunotherapy. <i>Ann Allergy Asthma Immunol</i>. 2018;120:331-333.e1. Lewis MO, Brown-Whitehorn TF, Cianferoni A, Rooney C, Spergel JM. Peanut-allergic patient experiences after epicutaneous immunotherapy: peanut consumption and impact on QoL. <i>Ann Allergy Asthma Immunol</i>. 2019; 123:101-103. Ward C, Greenhawt M. Differences in caregiver food allergy quality of life between tertiary care, specialty clinic, and caregiver-reported food allergic populations. <i>J Allergy Clin Immunol Pract</i>. 2016;4:257-264. Waggoner MR. Parsing the peanut panic: the social life of a contested food allergy epidemic. <i>Soc Sci Med</i>. 2013;90:49-55. Greenhawt M, Marsh R, Gilbert H, Sicherer S, DunnGalvin A, Matlock D. Understanding caregiver goals, benefits, and acceptable risks of peanut allergy therapies. <i>Ann Allergy Asthma Immunol</i>. 2018;121:575-579. Greenhawt M, DunnGalvin A. Preliminary psychometric analyses and clinical performance of a caregiver self-efficacy scale for food allergy self-management. <i>Ann Allergy Asthma Immunol</i>. 2018;120:73-79. Greenhawt M, Dunn Galvin A, Chalil JM, Prinz M, Rogers M, Green TD. Patient and Caregiver Burden of Peanut Allergy: An Ethnographic Study. Presented at the 2019 EAACI Pediatric Allergy and Asthma Management Conference, Florence, Italy, October 18, 2019.	Both patients and clinicians highly value an accurate diagnosis, but may be concerned about the undesirable effects highlighted above. There is emerging evidence that uncertainty of what diagnostic test results imply at the time of diagnosis may have detrimental effects on patients and their families. There are no published data on the values and preferences of patients and families regarding performing diagnostic testing for food allergy. Specifically there are no data regarding the potential harms of a false-positive test result as compared with the potential harms of a missed diagnosis (false-negative test result), or how the future implications of the erroneous diagnosis may be handled. This could encompass a scenario where a false negative test results in no diagnosis being given, but the individual later eats a peanut containing item and has a reaction, or alternatively (and more likely), the scenario of someone diagnosed as peanut allergic based on positive testing (without a history of ingestion), who later "outgrows" the allergy and may be resentful of the possibility of a false positive diagnosis.

Balance of effects

Does the balance between desirable and undesirable effects favor the intervention or the comparison?

JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
-----------	-------------------	---------------------------

○ Favors the comparison ○ Probably favors the comparison ○ Does not favor either the intervention or the comparison ○ Probably favors the intervention ○ Favors the intervention ● Varies ○ Don't know	The oral food challenge is the most accurate, definitive assessment of peanut allergy. However, in situations where there is high pre-test probability for peanut allergy, the three diagnostic tests can greatly assist in increasing (sensitization detected) or significantly decreasing (no sensitization detected) the post-test odds of having peanut allergy, and confirmatory oral food challenge may not always be required. Outside of a strong stated preference where there is low pre-test probability, the comparator test (oral food challenge) has more desirable effects than the intervention diagnostic tests, and can be used to avoid diagnostic misclassification. Ward C, Greenhawt M. Differences in caregiver food allergy quality of life between tertiary care, specialty clinic, and caregiver-reported food allergic populations. <i>J Allergy Clin Immunol Pract.</i> 2016;4:257-264. National Academies of Sciences E, Medicine. Finding a Path to Safety in Food Allergy: Assessment of the Global Burden, Causes, Prevention, Management, and Public Policy. Washington, DC: The National Academies Press; 2017. Sampson HA, Aceves S, Bock SA, James J, Jones S, Lang D, et al. Food allergy: A practice parameter update—2014. <i>Journal of Allergy and Clinical Immunology</i> 2014;134:1016-25.e43. Franxman T, Howe L, Teich E, Greenhawt M. Oral food challenge improves food allergy quality of life. <i>J Allergy Clin Immunol Pract</i>: 2015; 3: 50-56 Kansen, HM, Le, T-M, Meijer, Y, et al. The impact of oral food challenges for food allergy on quality of life: A systematic review. <i>Pediatr Allergy Immunol.</i> 2018; 29: 527– 537. https://doi.org/10.1111/pai.12905	The risks of a false positive test are significant and may lead to prolonged unnecessary avoidance and costs, as well as potential stigma related to being classified as being peanut allergic. Particularly at young ages, over-diagnosis by isolated positive tests of sensitization may also lead to a lost opportunity to establish peanut tolerance.
Resources required How large are the resource requirements (costs)?		
JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
○ Large costs ○ Moderate costs ○ Negligible costs and savings ○ Moderate savings ○ Large savings ● Varies ○ Don't know	There are no studies that directly investigated the resources requirements	All the possible interventions and the comparator tests do require resources in terms of both direct and indirect costs. These costs and cost burdens may vary depending on the healthcare system in question but are likely already nested into the cost of normal practice operation. Newer management options based on test results may have additional costs that have not been studied. Costs may vary based on the particular healthcare system and geography but these largely fall into overlapping ranges across the US. Skin testing (CPT code 95004) may have more variability in terms of cost and reimbursement than serologic IgE testing (CPT code 86003) based on a selected sample of US cities in different parts of the country, detailed below: Lebanon, NH: 95004 code \$9-28; 86003 code \$15-\$98 New York City: 95004 code \$9-28; 86003 code \$15-\$98 Winston-Salem, NC: 95004 code \$8-23; 86003 code \$15-\$98 Miami, FL: 95004 code \$8-25; 86003 code \$15-\$98 Kansas City, MO: 95004 code \$8-25; 86003 code \$15-\$98 Duluth, MN: 95004 code \$9-25; 86003 code \$15-\$98 Denver, CO: 95004 code \$8-23; 86003 code \$15-\$98 Eugene, OR: 95004 code \$9-28; 86003 code \$15-\$98 Los Angeles, CA: 95004 code \$8-25; 86003 code \$15-\$98

Certainty of evidence of required resources

What is the certainty of the evidence of resource requirements (costs)?

JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
○ Very low ○ Low ○ Moderate ○ High ● No included studies	There are no studies that directly assessed the certainty of evidence of resource requirement.	There would not be any anticipated new resources needed to support the use of any of these tests that are not already established and in use in clinical practice. There may be additional resources required to offer Ara h 2 as a stand-alone test, as opposed to a full component panel. Initially more expensive but then cheaper later. Operating costs vary from region to region and depend on practice location, personnel experience, and practice volume. While 95% of practicing allergists offer oral food challenges, only 17% perform more than 10 per month, which could complicate access to confirm diagnostic test results.

Cost effectiveness

Does the cost-effectiveness of the intervention favor the intervention or the comparison?

JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
○ Favors the comparison ○ Probably favors the comparison ○ Does not favor either the intervention or the comparison ○ Probably favors the intervention ○ Favors the intervention ● Varies ○ No included studies	The cost-effectiveness of diagnostic testing varies based on which test is chosen. A cost-benefit analysis as part of this document shows that use of skin prick testing as opposed to use of Ara h 2 testing is not cost-effective and is associated with higher societal costs related to the risk of false positive results, leading to a patient who is not truly peanut allergic being managed as such. Skin prick testing remains associated with higher costs and lower benefits as a choice of test (e.g. "dominated" in economic terms) in the analysis until the specificity of Ara h 2 decreases significantly from the values identified in the meta-analysis. Deterministic sensitivity analysis did not reveal other factors related to assessing a patient for peanut allergy with diagnostic testing, that, if changed, could make this test more cost-effective than Ara h 2.	Skin prick testing to peanut has lower specificity than Ara h 2 testing, and will result in more falsely positive diagnoses identified, resulting in lower QALY accumulation. However, with the marginal increase in sensitivity, SPT would result in a slightly lower rate of peanut allergic reactions.

Equity

What would be the impact on health equity?

JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
○ Reduced ○ Probably reduced ○ Probably no impact ○ Probably increased ○ Increased ○ Varies ● Don't know	There are no studies that directly assessed the impact on equity.	Serologic testing is more widely available and less dependent on allergy specialists which may improve equity potentially, whereas skin testing is the opposite. Certain states have different reimbursement rules/rates for skin vs. serologic testing, which could reduce equity if certain of these tests are not available, based on location or insurance.

Acceptability

Is the intervention acceptable to key stakeholders?

JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
○ No ○ Probably no ○ Probably yes ○ Yes ○ Varies ● Don't know	There are not studies that directly assess if the intervention is acceptable to key stakeholders	Clinician vantage: Multiple prior practice parameters have echoed these findings; however, there is well-known practice variation with respect to indication for testing, and interpretation of tests in certain contexts. The clinician may not accept or follow guidelines that advise against their current practices, their training, or their comfort level with decision-making. Patient/Advocate Vantage: Patients may have variable preferences regarding how diagnostic testing is used, and as stated above may differentially value having a false positive diagnosis than a false negative diagnosis, or having to undergo an oral challenge to confirm a diagnosis. There could be a role for shared decision making and a decision-aid to help patients consider their options.

Feasibility

Is the intervention feasible to implement?

JUDGEMENT	RESEARCH EVIDENCE	ADDITIONAL CONSIDERATIONS
○ No ○ Probably no ○ Probably yes ○ Yes ○ Varies ● Don't know	There are no studies that evaluated the feasibility of implementing these findings.	This should be feasible to implement but implementation could be limited by lack of availability of Ara h 2 as a stand-alone test. Variable reimbursement of allergy testing services may also limit access to care and implementation. An even more problematic implementation would be if there are sufficient resources at all allergy practices to support an increased need for subsequent oral food challenge to confirm diagnosis when indicated. Not all allergy practices offer oral food challenges and most primary care providers would not be conducting oral food challenges.

SUMMARY OF JUDGEMENTS

	JUDGEMENT						
PROBLEM	No	Probably no	Probably yes	Yes		Varies	Don't know
TEST ACCURACY	Very inaccurate	Inaccurate	Accurate	Very accurate		Varies	Don't know
DESIRABLE EFFECTS	Trivial	Small	Moderate	Large		Varies	Don't know
UNDESIRABLE EFFECTS	Large	Moderate	Small	Trivial		Varies	Don't know
CERTAINTY OF THE EVIDENCE OF TEST ACCURACY	Very low	Low	Moderate	High			No included studies
CERTAINTY OF THE EVIDENCE OF TEST'S EFFECTS	Very low	Low	Moderate	High			No included studies
CERTAINTY OF THE EVIDENCE OF MANAGEMENT'S EFFECTS	Very low	Low	Moderate	High			No included studies
CERTAINTY OF THE EVIDENCE OF TEST RESULT/MANAGEMENT	Very low	Low	Moderate	High			No included studies
CERTAINTY OF EFFECTS	Very low	Low	Moderate	High			No included studies
VALUES	Important uncertainty or variability	Possibly important uncertainty or variability	Probably no important uncertainty or variability	No important uncertainty or variability			
BALANCE OF EFFECTS	Favors the comparison	Probably favors the comparison	Does not favor either the intervention or the comparison	Probably favors the intervention	Favors the intervention	Varies	Don't know
RESOURCES REQUIRED	Large costs	Moderate costs	Negligible costs and savings	Moderate savings	Large savings	Varies	Don't know
CERTAINTY OF EVIDENCE OF REQUIRED RESOURCES	Very low	Low	Moderate	High			No included studies
COST EFFECTIVENESS	Favors the comparison	Probably favors the comparison	Does not favor either the intervention or the comparison	Probably favors the intervention	Favors the intervention	Varies	No included studies
EQUITY	Reduced	Probably reduced	Probably no impact	Probably increased	Increased	Varies	Don't know
ACCEPTABILITY	No	Probably no	Probably yes	Yes		Varies	Don't know
FEASIBILITY	No	Probably no	Probably yes	Yes		Varies	Don't know

TYPE OF RECOMMENDATION

Strong recommendation against the intervention ○	Conditional recommendation against the intervention ○	Conditional recommendation for either the intervention or the comparison ○	Conditional recommendation for the intervention ●	Strong recommendation for the intervention ○
---	--	---	--	---

CONCLUSIONS

Recommendation

Recommendations 1a: We suggest in favor of diagnostic (skin prick or serum sIgE) testing for peanut allergy in patients with a 1) physician-judged high pre-test probability of peanut allergy, or 2) prior to an oral food challenge for patients with moderate pre-test probability of peanut allergy, with whom shared decision-making has been employed to arrive at the final decision. Conditional recommendation; Certainty of evidence: very low

Recommendation 1b: We suggest against diagnostic testing in patients with a low or very low pre-test probability of peanut allergy. Conditional recommendation; Certainty of evidence: very low

Recommendation 2a: We suggest in favor of Ara h2 diagnostic testing in a patient presenting for evaluation of suspected peanut allergy for which a single diagnostic test is to be used, as Ara h2 would provide the best diagnostic accuracy as determined by the more optimal positive/negative likelihood ratio. Conditional recommendation. Certainty of evidence: moderate

Recommendation 2b: We suggest against component testing in addition to either skin prick test or sIgE to whole peanut to increase diagnostic accuracy. Conditional recommendation. Certainty of evidence: moderate.

Recommendation 3: We suggest against the clinician using the results of a SPT, sIgE to whole peanut extract, or sIgE to peanut components to determine the severity of a previous reaction and/or allergy phenotype or to predict the severity of a future reaction, Conditional recommendation. Certainty of evidence: very low.

Technical remarks:

It is critical to consider diagnostic test performance in the context of the pre-test probability of peanut allergy. The clinician should recognize the circumstances where one or more of the peanut diagnostic tests may not translate to a clinically meaningful improved post-test odds of peanut allergy. Except in cases of high pre-test probability, it is likely that an oral food challenge will be needed to establish the diagnosis of peanut allergy, regardless of the results of the selected diagnosis test(s).

Certain tests may be more appropriate than others in particular situations. We suggest that the choice of SPT, sIgE, or Ara h 2 sIgE is not critical in circumstances where there is high pre-test probability of peanut allergy.

While testing of patients with low pre-test probability is not generally recommended, if the decision is made to test in these circumstances, from a test precision standpoint, use of Ara h 2 rather than SPT or sIgE can help decrease misclassification of patients as peanut allergic, leading to less harm through falsely positive diagnosis. When testing individuals with low pre-test probability, it is recommended that an oral food challenge still be performed to validate the clinical significance of the detection of sensitization, given that the low pre-test probability in the setting of detectable sensitization translates to only moderate post-test odds of a diagnosis.

Justification

Overall justification

In patients with a high pre-test probability for peanut allergy, SPT, sIgE, and Ara h 2 sIgE are highly sensitive and reliable tests that can be considered for routine use in the diagnosis of peanut allergy.

Detailed justification

Test accuracy

These are tests with high sensitivity

Desirable Effects

Detection of sensitization in an individual with likely or suspected peanut allergy will aid considerably in confirming the diagnosis. Choice of test in circumstances where there is high pre-test probability is not critical. The absence of sensitization is helpful in ruling out the diagnosis (although in many cases, oral food challenge will still be necessary).

Subgroup considerations

Severity of reaction was investigated as a potential subgroup. These tests do not perform well to identify individuals for potentially severe reactions at the dichotomous thresholds investigated. Data are limited that may better inform if these tests have higher or lower value within other particular subgroups. In infants meeting high-risk criteria for early peanut introduction, SPT is often used; however, incorporation of Ara h 2 might result in a lower rate of over-diagnosis. Unfortunately, evidence comparing these tests in this particular population is lacking.

Implementation considerations

These are tests that are already in routine use or routinely available for use; however, testing for Ara h 2 as a single component would be needed to implement routine use of this test. In many instances, the clinician is already using these tests in tandem, potentially. If the clinician starts with the SPT in the office setting, using the Fagan nomogram in figure 3, the post-test odds could then reasonably be used as the pre-test odds for choice of a confirmatory test, represented by figure 3 for sIgE or figure 6 for Ara h 2. In this setting, given higher specificity and higher positive likelihood ratio, Ara h 2 may be the better choice of a confirmatory test if it could be obtained as a stand-alone measure.

Monitoring and evaluation

Additional meta-analysis at different cut-off points may help inform decision making, in particular for the severity subgroup or use of tests in sequence/tandem. We would recommend to journal editors that there be a requirement for future reporting of studies investigating diagnostic test precision in relation to food challenge outcome that raw data be included as a supplemental text denoting the challenge outcome, the numeric quantity of the test, the sequence of testing run if multiple tests were assessed simultaneously, and any data on severity of the reaction. This would allow for a repository to be created that would greatly assist with updating practice guidelines. For study authors to make such deidentified data available, it would enable more direct assessment of test performance as a continuous variable, which would allow for different diagnostic thresholds could be directly assessed and compared for the purposes of meta-analysis and systematic review, as opposed to having to rely on dichotomous assessment of pre-selected thresholds and potential back calculation of sensitivity/specificity. These factors serve as distinct limitations with regard to this particular document.

Research priorities

Additional studies in more unselected populations, and at a population level are needed. Future research studies reporting diagnostic sensitivity and specificity should report the true/false positive and true/false negative patient level results to assist in future meta-analyses where cut off levels would be easier to assess. If these data were available, it would have permitted analysis of the sensitization levels as a continuous variable rather than a dichotomous variable and potentially allowed better comparison of tests used sequentially or in tandem. Better data are needed to help inform what defines low, moderate, or high pre-test probability in a patient being assessed for possible peanut allergy, as well as to understand how clinicians and patients may perceive risk.

Text Box 1: GRADE Questions Evaluated in this Practice Parameter

PICO Questions: GRADE Analysis of Diagnostic Testing in the Diagnosis of Peanut Allergy

1. Should diagnostic testing for peanut allergy be performed in adults and children with a history of suspected peanut allergy who are requesting evaluation for peanut allergy?
- 2a. In the patient presenting for evaluation of suspected peanut allergy, which of the three tests—SPT, sIgE to whole peanut, or Ara h2 would provide the highest diagnostic accuracy as determined by the more optimal positive/negative likelihood ratio?
- 2b. In a patient presenting for evaluation of suspected peanut allergy, does testing for peanut components in addition to either SPT or sIgE to whole peanut increase the diagnostic accuracy?
3. In the patient presenting for evaluation of suspected peanut allergy, can the results of a diagnostic test be used to predict the severity of a future allergic reaction?

Text Box 2: Defining Allergic Sensitization and a “Positive Test”

Allergic sensitization is denoted by the presence of detectable allergen-specific IgE, either through a serologic assay or through skin prick testing. All tests for sensitization have a threshold where the test is considered to be positive, as well as either a detection limit or a reporting limit. For skin prick testing, the most commonly reported convention for where a test is considered “positive” for the presence of allergen specific IgE is when the allergen-specific test is 3mm of wheal diameter greater than that of a simultaneously placed glycerinated saline control. As discussed in the 2008 Diagnostic Testing Practice Parameter (www.allergyparameters.org) different testing devices produce some degree of variation in the size range of negative controls, as does variation related to the tester. Wheal size is recommended to be measured as the average length of the two longest bisecting planes, though many clinics may elect to measure the longest single plane.

For serum-specific IgE tests using fluorescent enzymatic immunoassay (FEIA) detection, the instruments have both detection limits and reporting limits that have influenced test results. However, each instrument has particular reporting and detection ranges, and these differ between commercial tests. The technical detection limit for these machines is typically 0.1 KU_A/L, and antibody levels above this threshold are reported as they are detected, to an upper reporting limit of 100 KU_A/L. Quantification of levels >100 KU_A/L is possible through sample dilution. For many years, the reporting limit was conventionally set at < 0.35 KU_A/L, though in recent years, this has been replaced by the detection limit of 0.1 KU_A/L. Using the older convention of the 0.35 KU_A/L reporting limit, “positive” sensitization was considered to be 0.36 KU_A/L or higher. With the newer convention of using the 0.1 KU_A/L detection limit as the reporting limit, “positive” sensitization would therefore be 0.11 KU_A/L. This creates a conundrum of how to interpret sensitization between 0.11 KU_A/L and 0.35 KU_A/L, which prior to the change in reporting convention would have fallen into the “negative” range. It is debatable that such sensitization is clinically relevant, or that many clinicians would only consider sensitization above 0.7 KU_A/L as clinically relevant. Nonetheless, studies may report positive sensitization at 0.11 KU_A/L in a binary fashion. One additional classification that is seen are classes representing sextiles of IgE quantity detected between the upper and lower reporting limits. These are arbitrary conventions that date back to the quartiles originally described for Radioallorbsorbent Testing, adjusted for the FEIA method. Class 0 represents levels below the reporting limit, and class 1 typically starts at the reporting lower limit, ranging to class 6 representing the highest levels detectable which are reported. These class designations have no clinical relevance in and of themselves, and no reference to class designations is made in this document.

In this document, if the term positive is used, in relation to either form of test it is in this sense that this refers to positive detection of sensitization (e.g. a positive test). Unequivocally, positive detection of sensitization is unrelated to a positive clinical diagnosis of allergy. A positive diagnosis is predicated on both a demonstrated clinical history of allergy and the presence of detectable sensitization, or in very circumscribed instances, very high levels of sensitization in infants with very particular pre-existing risk factors who have never ingested peanut previously.

Text Box 3: Examples of Pre-test Probability in Determining if Diagnostic Testing is Indicted

1. **High pre-test probability** should be considered as a situation where there was ingestion of peanut and typical IgE mediated symptoms of an allergic reaction resulted, either directly observed or reported; or an infant meeting NIAID early peanut introduction high-risk criteria prior to peanut introduction. Testing is of the highest utility in these scenarios and peanut sensitization above a certain threshold is of high likelihood to be associated with the highest post-test odds of a diagnosis of peanut allergy.
2. **Moderate pre-test probability** should be considered as a situation where there is less clarity that peanut was ingested and resulted in IgE mediated symptoms, but some consideration for this in explaining an allergic reaction under evaluation. In some instances it may represent situations where the patient has not previously consumed peanut but could be considered at a risk greater than the general population for peanut allergy based on the presence of certain types of other food allergies, certain atopic comorbidities (e.g., severe eczema), or certain children outside the first year of life with delayed peanut introduction. Testing is of unclear utility in these situations, and not necessarily associated with post-test odds that clarify clinical decision making. An oral food challenge may be required to definitively establish a diagnosis when there is peanut sensitization above a certain threshold.
3. **Low pre-test probability** should be considered a situation where there is very little uncertainty that the person is peanut tolerant (e.g. eats peanut without becoming symptomatic), that peanut was unrelated to the allergic reaction being evaluated (e.g. it is clear that a single allergen other than peanut likely caused the aforementioned reaction and the product was peanut-free, or peanut is being tested solely because it is part of a multi-allergen panel and there is no specific independent concern for peanut allergy itself), family history of peanut allergy or allergic disease, general curiosity about what someone could speculatively be “allergic to”, or for an infant meeting addendum 2 or 3 criteria for NIAID early peanut introduction guidelines prior to peanut introduction. In some instances it may represent situations where the patient has not previously consumed peanut but the clinician may have concern that the patient is at a risk greater than the general population for peanut allergy based on the presence of certain types of other food allergies or concern for cross-reactivity, certain atopic comorbidities (e.g., mild or moderate eczema), or certain children outside the first year of life with delayed peanut introduction but who have no baseline risk factors. Testing in these situations is of exceptionally limited to no utility whatsoever, is not associated with any shift of post-test odds over baseline, and is not indicated. An oral food challenge is likely required to establish that the peanut sensitization detected is clinically irrelevant.

Text Box 4: Key Questions in Peanut Allergy Diagnostic Testing

- **Are there any clinical indications to obtain peanut allergy testing for a patient who is eating peanut without immediate onset or reproducible symptoms?**

In general, no. However, rare exceptions to this include part of the evaluation of patients with eosinophilic esophagitis where dietary elimination is considered as a treatment option, which is a highly specific context with very particular (non-IgE mediated) symptoms, which is beyond the scope of this document.

- **Which test should be ordered in the evaluation of patients who have never ingested peanut, i.e. prior to early introduction for at risk infants?**

Peanut skin prick and serum IgE testing is poorly specific and in general should not be used as a screening tool for someone who has never eaten peanut before and developed symptoms. When used as part of the early introduction guidelines for infants less than 6 months of age who have severe eczema and/or egg allergy, both skin prick and serum peanut IgE tests can be utilized. There is no current role for component testing in this context. Even in patients with very high peanut specific IgE levels, without a history denoting that they have eaten peanut and become symptomatic, these are very difficult to interpret, and OFC should be considered to definitively diagnose such patients. A shared decision-making approach can be employed here, given that some parents and clinicians may strongly feel that such test values represent a high likelihood of clinical allergy, despite the absent history.

- **Are there cut-off levels for peanut skin prick or serum IgE testing that diagnoses peanut allergy?**

A universal cut-off level does not exist. These are technically difficult to generate, given that these are based on accurately knowing the population prevalence of peanut allergy. Cut-off levels are only relative probabilities that are imperfect and have an error rate that will potentially misclassify individuals. When prevalence of disease is not known, the likelihood ratio is a more applicable test. This tells the likelihood of a positive test in someone with the disease compared to the likelihood of a positive test in someone without disease, and can help convert the pre-test probability that someone has the disease to a post-test odds using a Fagan nomogram. Thus, as stand-alone measures, neither skin prick nor serum IgE test results can be interpreted as diagnostic for peanut allergy.

- **Should peanut allergy testing be considered in children with moderate-to-severe atopic dermatitis?**

Atopic dermatitis is caused by changes in the epidermal skin barrier and is generally not due to food allergy, though children with persistent and refractory moderate-to-severe atopic dermatitis may be at higher risk of developing food allergy. Peanut allergy testing should not be a standard part of the evaluation for any patient with atopic dermatitis. However, in a very small subset of infants and young children with severe, treatment- refractory atopic dermatitis may benefit from select food allergy, including peanut allergy testing if the clinical history suggests peanut has not yet been introduced, or there is suspicion that peanut ingestion is temporally associated with flares.

- **Should children with a family history of peanut allergy in another sibling be evaluated for peanut allergy prior to this being introduced?**

Screening of younger siblings for peanut allergy should not be routinely performed, and there is no evidence that such individuals are at higher risk for developing peanut allergy based just on the sibling history alone. To facilitate timely introduction and prevent delay, there could be consideration for a role for testing when parents are overly anxious about introducing peanut and will not introduce peanut to their child through any other means. However, such testing must be

interpreted properly and a positive result not be considered diagnostic for peanut allergy. In these situations, either skin prick or serum IgE testing may be utilized. Data exists to show that this practice is not cost-effective until there is a much higher baseline prevalence of peanut allergy in the population, and then only cost-effective if sensitized children undergo challenge rather than avoid peanut based on strong sensitization. There is no indication to utilize component testing in this context.

- **Are all patients with detectable Ara h 2 clinically allergic to peanuts?**

No. Detectable isolated sensitization to Ara h 2 is not diagnostic for peanut allergy, and a diagnosis can only be made where the individual is sensitized in the context of a known or suspected reaction after eating peanut. There are no well-established cut off levels for Ara h 2 at this time that indicate the presence of allergy versus sensitization. However, when compared to whole peanut skin prick and sIgE tests, Ara h 2 testing has vastly increased specificity, though this is still largely dependent on the context in which any testing is indicated. Patients may have detectable Ara h 2 but exhibit no clinical reactivity upon ingestion of peanut.

- **Does component testing predict the severity of future reactions?**

No test, including components, has good sensitivity or specificity to indicate the severity of symptoms of a future reaction. Component testing may have a potential role to help identify sensitization patterns that indicate recognition of cross sensitization with pollen allergens as opposed to more primary allergens unique to peanut, though the clinical significance of this is still to be defined.

- **When should component testing be ordered as the initial diagnostic test?**

The role of component testing is evolving, and it is unclear how and when these tests should be used. Comparatively, testing for Ara h 2 compared to whole peanut skin prick and sIgE testing does have significantly higher specificity, which may translate to a lower likelihood of a false positive diagnosis if testing is run the right context. Moreover, in this context, use of Ara h 2 as a stand-alone test is highly cost-effective. However, there is a present knowledge gap if Ara h 2 should be the initial test ordered.