

ORIGINAL ARTICLE

EXPERIMENTAL ALLERGY AND IMMUNOLOGY

Randomized controlled trial of fish oil supplementation in pregnancy on childhood allergies

D. J. Palmer^{1,2}, T. Sullivan³, M. S. Gold⁴, S. L. Prescott², R. Heddle^{5,6}, R. A. Gibson⁷ & M. Makrides^{1,4}

¹Women's & Children's Health Research Institute, North Adelaide, SA; ²School of Paediatrics and Child Health, University of Western Australia, Subiaco, WA; ³Data Management & Analysis Centre, University of Adelaide, Adelaide, SA; ⁴School of Paediatrics and Reproductive Health, University of Adelaide, Adelaide, SA; ⁵SA Pathology, Royal Adelaide Hospital, Adelaide, SA; ⁶Flinders University of South Australia, Bedford Park, SA; ⁷School of Agriculture, Food and Wine, University of Adelaide, Glen Osmond, SA, Australia

To cite this article: Palmer DJ, Sullivan T, Gold MS, Prescott SL, Heddle R, Gibson RA, Makrides M. Randomized controlled trial of fish oil supplementation in pregnancy on childhood allergies. *Allergy* 2013; DOI: 10.1111/all.12233.

Keywords

allergy prevention; eczema; fatty acids; pregnancy; randomized controlled trial.

Correspondence

Prof. Maria Makrides, Women's & Children's Health Research Institute, 72 King William Road, North Adelaide, SA 5006, Australia.
Tel.: +61-(0)8-8161-6067
Fax: +61-(0)8-8239-0267
E-mail: maria.makrides@health.sa.gov.au

Accepted for publication 1 July 2013

DOI:10.1111/all.12233

Edited by: Bodo Niggemann

Abstract

Background: Diets high in n-3 long chain polyunsaturated fatty acids (LCPUFA) may modulate the development of IgE-mediated allergic disease and have been proposed as a possible allergy prevention strategy. The aim of this study was to determine whether n-3 LCPUFA supplementation of pregnant women reduces IgE-mediated allergic disease in their children.

Methods: Follow-up of children ($n = 706$) at hereditary risk of allergic disease in the Docosahexaenoic Acid to Optimise Mother Infant Outcome randomized controlled trial. The intervention group ($n = 368$) was randomly allocated to receive fish oil capsules (providing 900 mg of n-3 LCPUFA daily) from 21 weeks' gestation until birth; the control group ($n = 338$) received matched vegetable oil capsules without n-3 LCPUFA. The diagnosis of allergic disease was made during medical assessments at 1 and 3 years of age.

Results: No differences were seen in the overall percentage of children with IgE-mediated allergic disease in the first 3 years of life between the n-3 LCPUFA and control groups (64/368 (17.3%) vs 76/338 (22.6%); adjusted relative risk 0.78; 95% CI 0.58–1.06; $P = 0.11$). Eczema was the most common allergic disease; 13.8% of children in the n-3 LCPUFA group had eczema with sensitization compared with 19.0% in the control group (adjusted relative risk 0.75; 95% CI 0.53–1.05; $P = 0.10$).

Conclusions: Overall, n-3 LCPUFA supplementation during pregnancy did not significantly reduce IgE-associated allergic disease in the first 3 years of life. Further studies should examine whether the nonsignificant reductions in IgE-associated allergies are of clinical and public health significance.

Changing dietary patterns have favoured increased intake of n-6 polyunsaturated fatty acids (PUFA) from linoleic acid (18:2, n-6)-rich vegetable oils, especially as margarine and vegetable oils have become common dietary staples over the past 40 years. This, together with an increase in

consumption of meat-based products, has led to an increase of arachidonic acid (AA, 20:4, n-6) in tissues. AA gives rise to eicosanoids such as prostaglandin E_2 that can enhance the synthesis of T helper type 2 cytokines and immunoglobulin E (IgE) antibodies potentially leading to sensitization to allergens. However, when diets are high in n-3 long chain (LC) PUFA (from fatty fish and fish oils), they are readily incorporated into cellular phospholipids and thereby displace AA and alter membrane composition and fluidity. This leads to a range of immunological effects, including reduction in prostaglandin E_2 synthesis (1, 2) providing a plausible mechanism by which diets high in n-3 LCPUFA

Abbreviations

AA, arachidonic acid; DHA, docosahexaenoic acid; DOMInO, Docosahexaenoic acid to Optimise Mother Infant Outcome; EPA, eicosapentaenoic acid; IgE, immunoglobulin E; PUFA, polyunsaturated fatty acids; LCPUFA, long chain polyunsaturated fatty acids; RCT, randomized controlled trial.

may modulate the development of IgE-mediated allergic disease.

In support of this biological mechanism, epidemiological studies have reported that increased maternal fish intake during pregnancy is associated with reduced atopic or allergic outcomes in children (3–7). Furthermore, evidence from randomized controlled trials (RCTs) (8–11) involving fish oil supplementation of pregnant women has found beneficial effects on reduced allergen sensitization and allergic disease outcomes in the offspring. However, previous RCTs have only reported allergic disease outcomes at 1 year (8, 9), 2 years (12) and 16 years of age (10), and none have reported outcomes between 2 and 16 years of age.

Our study was specifically designed to assess the effect of n-3 LCPUFA supplementation, predominantly as DHA, in pregnancy on the cumulative incidence of IgE-mediated allergic disease in the first 3 years of life. Outcome data focussing on eczema and food allergy over the first year of life have been previously published (11).

Methods

Subjects and study design

The present study is an allergy follow-up of a subset of children whose mothers were participants in the Docosahexaenoic Acid (DHA) to Optimise Mother Infant Outcome (DOMInO) trial (13). This subset of children all had a mother, father or sibling with a history of medically diagnosed allergic diseases (asthma, allergic rhinitis, eczema), and they were enrolled at either of the two study centres in Adelaide, Australia. Full details of entry into this allergy follow-up study have been previously published (11). Written informed consent was sought before birth and included consent for their offspring to participate in both the 1 and 3 year of age allergy follow-up assessments. Approval for this study was granted by the Human Research Ethics Committees of each centre, Women's and Children's Hospital, Adelaide and Flinders Medical Centre, Adelaide. This allergy follow-up study was registered at www.anzctr.org.au as ACTRN12610000735055. The Docosahexaenoic Acid to Optimise Mother Infant Outcome (DOMInO) trial was registered at www.anzctr.org.au as ACTRN12605000569606.

The dietary treatments for the DOMInO trial have been previously described (13). Briefly, women allocated to the n-3 LCPUFA group were asked to consume three 500 mg capsules of fish oil concentrate, providing 800 mg of DHA and 100 mg of eicosapentaenoic acid (EPA); women in the control group were asked to take three 500 mg vegetable oil capsules without n-3 LCPUFA daily. This was a double-blinded study; all capsules were similar in size, shape and colour. Women took capsules from 21 weeks' gestation until delivery.

Early childhood allergic disease outcome assessments and definitions

The primary outcome was diagnosis of IgE-associated allergic diseases (eczema, asthma, allergic rhinitis or food allergy) at 1

or 3 years of age. Participating children attended a medical review appointment at 1 and 3 years of age, conducted by one of six medical practitioners (who made the allergic disease diagnosis for a period covering the previous 12 months by taking a structured history and doing a standardized clinical examination) and one of six experienced research nurses (who performed skin prick testing). All were blinded to treatment group allocation and had quality assurance reviews every 6 months with one of the investigators (DJP). Data were also collected on possible confounding variables, including details of number of other children in the home, use of house dust mite covers for mattresses and pillows, presence of a cat as a pet and household smoking exposure.

Sensitization was defined as a positive skin prick test (wheal ≥ 3 mm above negative control) to at least one of the allergens assessed. At 1 year of age, the food allergens tested were whole hens' egg, cows' milk, wheat, tuna and peanut, and the aeroallergens tested were ryegrass pollen, olive tree pollen, *Alternaria tenuis*, cat hair and house dust mite (*Dermatophagoides pteronyssinus*). At 3 years of age, the same allergens were tested with the addition of two foods (cashew nut and sesame seed) and one aeroallergen (house dust mite, *Dermatophagoides farinae*). The cows' milk allergen extract became unavailable from the supplier for an extended period during the 3-year assessments and consequently was excluded from the definition of sensitization at 3 years.

Eczema was defined as the presence of eczema (criteria according to (14)) on medical review or a history of an itchy rash distributed to the facial, flexural or extensor surface of the skin that had followed a fluctuating or chronic course. We defined IgE-associated eczema or atopic eczema as eczema with sensitization to at least one of the allergens assessed. IgE-associated food allergy was defined as a history of immediate (within 60 min) skin rash (hives, rash or swelling) with or without respiratory symptoms (cough, wheeze, stridor), gastrointestinal symptoms (abdominal pain, vomiting, loose stools) or cardiovascular symptoms (collapse) following ingestion of a food and sensitization to the implicated food. Asthma was defined a history of 3 or more episodes of wheeze with the episodes less than 6 weeks apart and/or daily use of asthma medication. Allergic rhinitis was defined as a history of sneezing or a runny or blocked nose accompanied by itchy-watery eyes when there have not been symptoms to suggest an upper respiratory tract infection. IgE-associated asthma/allergic rhinitis was defined as asthma/allergic rhinitis along with sensitization to at least one of the aeroallergens tested.

Statistical methods

With >328 children per treatment group, we would be able to detect an absolute reduction of 10% (relative reduction of 33%) in the cumulative incidence of IgE-mediated allergic disease from 30% to 20% with >80% power ($\alpha = 0.05$) over the first 3 years of life. Such a reduction was realistic based on the pilot data from Dunstan et al. (8).

All analyses were performed according to the intention to treat principle. Multiple imputation was used to deal

with missing data, with 50 complete data sets imputed for analysis. Overall, the missing at random assumption of the multiple imputation approach appeared reasonable for these data. Binary outcomes were analysed using log binomial regression models, with treatment effects expressed as relative risks. Rare binary outcomes were compared between groups using Fisher's exact tests on the original (unimputed) data. Negative binomial regression models were used to analyse count outcomes, with the effect of treatment expressed as a ratio of means. Both unadjusted and adjusted analyses were performed, with adjustment for the stratification variables of centre and parity as well as the prespecified baseline variables of infant sex and maternal history of allergic disease. The adjusted analyses were considered to be the primary analyses. Potential confounding variables (environmental characteristics measured after

randomization) that may influence allergic disease outcomes were compared between groups using Mann–Whitney and chi-square tests. Statistical significance was assessed at the two-sided $P < 0.05$ level. All analyses were performed using SAS version 9.3.

Results

The allergy follow-up trial profile is shown in Fig. 1. Enrolment began on 20th March 2006 and ended on 8th May 2008, with a total of 706 infants recruited into the study. Data collection for the medical assessments at 3 years of age was completed on 1st September 2011. 638/706 (90.4%) children attended a medical review at 3 years of age with 587/706 (83.1%) of children having skin prick testing to determine their sensitization status.




Figure 1 Trial participant outcomes flow diagram.

Demographic and family characteristics

The participants consisted of 337/706 (47.7%) males, and 281/706 (39.8%) were the first born (parity zero). Mean maternal age at trial entry was 29.6 years (SD 5.7 years) and 92/706 (13.0%) of the participating mothers smoked during pregnancy. All participants had at least one first-degree relative with a history of medically diagnosed allergic disease; 656/706 (92.9%) had at least one parent and 206/706 (29.2%) had both parents with a history of allergic disease. Other baseline demographic characteristics of the trial participants and their families have been previously reported (11). Early childhood (0–3 years) home environmental characteristics are shown in Table 1; there were no statistically significant differences between the groups.

Early childhood allergic disease outcomes

In the n-3 LCPUFA intervention group, 17.3% of children were diagnosed with IgE-mediated allergic diseases (asthma, allergic rhinitis, eczema and/or food allergy) in the first 3 years of life compared with 22.6% in the control group (adjusted relative risk 0.78; 95% CI 0.58–1.06; $P = 0.11$; Table 2). As expected, the most common IgE-mediated allergic disease in the first 3 years of life was eczema with sensitization, affecting a total of 115/706 (16.3%) of the participating children. There was a lower, but not statistically significant, incidence of eczema with sensitization in the n-3 LCPUFA group (13.8% vs 19.0% control group, adjusted relative risk 0.75; 95% CI 0.53–1.05; $P = 0.10$; Table 2). Overall, 4.6% of the children were diagnosed with at least one IgE-mediated food allergy through to 3 years of age, and egg allergy was the most common, affecting 2.8% of the children. There were no differences between the n-3 LCPUFA

and control groups in the percentage of children diagnosed with food allergy or respiratory allergic diseases (asthma, allergic rhinitis) with sensitization through to 3 years of age (Table 2). The percentage of children diagnosed with allergic disease without sensitization (non-IgE-mediated) did not differ between groups at either time point (Tables 2 and 3).

Table 3 reports the allergic disease outcomes at 3 years of age and demonstrates that there were no differences between the n-3 LCPUFA and control groups. The percentage of children diagnosed with eczema with sensitization at 3 years of age was 13.0% and higher than that reported at 1 year of age (9.2% (11)). Of the 92 children diagnosed with IgE-associated eczema at 3 years of age, 54% were new cases (25 in each group) who did not have IgE-associated eczema at 1 year of age.

There was no difference between the groups in sensitization to at least one allergen at 1 or 3 years of age, with 29.4% of children in the n-3 LCPUFA group sensitized compared with 35.2% in the control group (adjusted relative risk 0.85; 95% CI 0.68–1.06; $P = 0.14$; Table 2). At 3 years of age, 24.6% of children in the n-3 LCPUFA group compared to 26.1% in the control group were sensitized to at least one allergen (adjusted relative risk 0.96; 95% CI 0.74–1.25; $P = 0.76$; Table 4). *Alternaria tenuis* (grass mould) (total of 7.8%), ryegrass pollen (total of 7.1%) and *D. Pteronyssinus* (total of 6.8%) were the most common allergens that children were sensitized to at 3 years (Table 4). Despite no differences in cat ownership between the groups in the first 3 years of life (Table 1), 6.8% of children in the n-3 LCPUFA group compared to 3.4% in the control group were sensitized to cat at 3 years of age; however, this difference did not reach statistical significance (adjusted relative risk 1.95; 95% CI 0.98–3.89; $P = 0.06$; Table 4). Figure 2 illustrates the changing profile of sensitization status from predominately food allergen sensitization at 1 year of age to increased aeroallergen sensitization at 3 years of age. Egg sensitization was found in a total of 86/706 (12.2%) of the children at 1 year of age, but only 27/706 (3.8%) at 3 years of age.

Discussion

This study is the largest randomized controlled trial of fish oil supplementation during pregnancy and was designed to resolve uncertainties surrounding the use of n-3 LCPUFA supplementation in pregnancy as an allergic disease preventative strategy for children with hereditary risk. Overall fish oil supplementation in pregnancy did not significantly reduce the cumulative incidence of IgE-associated allergic disease over the first 3 years of life. Our study was powered to detect a 33% relative reduction in allergic disease, and it is therefore not surprising that the differences noted did not reach statistical significance. The nonsignificant risk reductions of up to 22% may still be of public health significance as the burden and cost of allergic disease on affected families are high, and fish oil intervention is safe and relatively cheap.

The lower incidence of IgE-associated eczema observed at 1 year of age in the n-3 LCPUFA group did not persist at

Table 1 Early childhood home environmental characteristics. Values are numbers (Percentages) unless stated otherwise

Characteristic	n-3 LCPUFA	Control	P-value
Number of other children in home at 3 years of age: median (IQR)*	1 (0–2)	1 (0–2)	0.53
House dust mite covers for mattresses in first 3 years of life†	200/353 (56.7)	179/330 (54.2)	0.53
House dust mite covers for pillows in first 3 years of life†	93/342 (27.2)	105/322 (32.6)	0.13
Cat as pet in first 3 years of life†	161/354 (45.5)	135/326 (41.4)	0.29
Smoker in household in first 3 years of life†	172/352 (48.9)	148/330 (44.8)	0.29

*Groups compared using Mann–Whitney test on original data.

†Groups compared using chi-square test on original data.

IQR, inter quartile range; n-3 LCPUFA, long chain polyunsaturated fatty acid.

Table 2 Allergic disease in the first 3 years of life (cumulative incidence based on 1 and 3 year assessments)

	n-3 LCPUFA (<i>n</i> = 368) (%) [*]	Control (<i>n</i> = 338) (%) [*]	Unadjusted RR (95% CI)	Unadjusted <i>P</i> -value	Adjusted [†] RR (95% CI)	Adjusted [†] <i>P</i> -value
Allergic disease	64 (17.3)	76 (22.6)	0.77 (0.56–1.04)	0.09	0.78 (0.58–1.06)	0.11
with sensitization						
Eczema with	51 (13.8)	64 (19.0)	0.73 (0.51–1.03)	0.07	0.75 (0.53–1.05)	0.10
sensitization						
Food allergy with	18 (5.0)	14 (4.2)	1.20 (0.60–2.41)	0.60	1.25 (0.63–2.49)	0.53
sensitization						
Asthma with	6 (1.8)	5 (1.7)		1.00		
sensitization [‡]						
Allergic rhinitis	18 (4.8)	20 (5.9)	0.82 (0.44–1.54)	0.54	0.82 (0.43–1.53)	0.53
with sensitization						
Sensitization	108 (29.4)	119 (35.2)	0.83 (0.67–1.04)	0.11	0.85 (0.68–1.06)	0.14
Allergic disease	124 (33.7)	117 (34.6)	0.97 (0.79–1.21)	0.81	0.95 (0.77–1.18)	0.67
without sensitization						

^{*}All data are based on analysis of 50 imputed datasets. For n-3 LCPUFA and control groups, data are number of subjects (percentage).

[†]Adjusted for centre, parity, maternal history and sex.

[‡]Fisher's exact test using original data, adjusted/imputed analyses not done due to rarity of outcomes.

CI, confidence interval; n-3 LCPUFA, long chain polyunsaturated fatty acid; RR, relative risk.

Table 3 Allergic disease at 3 years of age

	n-3 LCPUFA (<i>n</i> = 368) (%) [*]	Control (<i>n</i> = 338) (%) [*]	Unadjusted RR (95% CI)	Unadjusted <i>P</i> -value	Adjusted [†] RR (95% CI)	Adjusted [†] <i>P</i> -value
Allergic disease	55 (14.8)	59 (17.6)	0.84 (0.59–1.19)	0.33	0.86 (0.61–1.21)	0.38
with sensitization						
Eczema with	44 (11.8)	48 (14.1)	0.84 (0.57–1.25)	0.39	0.86 (0.58–1.27)	0.45
sensitization						
Food allergy	14 (3.8)	9 (2.8)	1.36 (0.59–3.11)	0.47	1.48 (0.67–3.30)	0.34
with sensitization						
Asthma with	6 (1.8)	5 (1.6)		1.00		
sensitization [‡]						
Allergic rhinitis	18 (4.8)	20 (5.9)	0.82 (0.44–1.54)	0.54	0.82 (0.43–1.53)	0.53
with sensitization						
Allergic disease	91 (24.7)	86 (25.3)	0.98 (0.74–1.28)	0.87	0.95 (0.72–1.25)	0.72
without sensitization						

^{*}All data are based on analysis of 50 imputed datasets. For n-3 LCPUFA and control groups, data are number of subjects (percentage).

[†]Adjusted for centre, parity, maternal history and sex.

[‡]Fisher's exact test using original data, adjusted/imputed analyses not done due to rarity of outcomes.

CI, confidence interval; n-3 LCPUFA, long chain polyunsaturated fatty acid; RR, relative risk.

3 years of age. As more than half of the children diagnosed with eczema with sensitization at 3 years of age did not have this diagnosis at 1 year of age, it is interesting to question whether these new cases could have been reduced if the n-3 LCPUFA supplementation had continued beyond pregnancy during early childhood. A recent postnatal (0–6 months of age), n-3 LCPUFA supplementation trial (*n* = 420) (15) found no overall effect on allergic disease outcomes at 1 year of age, but there was a significant reduction in eczema diagnosis in those infants who had higher n-3 LCPUFA levels at 6 months of age. This may be an important influencing factor as Furuholm et al. (12) (n-3 LCPUFA supplementation during pregnancy and first 3.5 months of lactation) also

found higher maternal, and infant proportions of DHA and EPA in plasma phospholipids were associated with lower prevalence of IgE-associated allergic disease in a dose-dependent manner. Interestingly compared with our study, other studies have used higher doses of n-3 LCPUFA that have ranged from 2700 mg/day (10, 12) to 3700 mg/day (8), and it may be that higher n-3 LCPUFA doses are needed to result in reduced allergic disease outcomes. Dose, timing and duration of n-3 LCPUFA supplementation are important considerations and worthy of further investigation. A limitation of our study was that we have not taken blood samples from the children and hence cannot report on their n-3 LCPUFA plasma phospholipids levels at 1 or 3 years of age.

Table 4 Sensitization to allergen skin prick test extracts at 3 years of age


	n-3 LCPUFA (n = 368) (%)*	Control (n = 338) (%)*	Unadjusted RR (95% CI)	Unadjusted P-value	Adjusted [†] RR (95% CI)	Adjusted [†] P-value
Any sensitization	91 (24.6)	88 (26.1)	0.94 (0.72–1.23)	0.67	0.96 (0.74–1.25)	0.76
Egg sensitization	11 (3.1)	16 (4.7)	0.65 (0.30–1.41)	0.28	0.68 (0.32–1.45)	0.32
Peanut sensitization	13 (3.4)	20 (5.8)	0.59 (0.29–1.20)	0.14	0.64 (0.31–1.29)	0.21
Cashew nut sensitization	5 (1.4)	12 (3.5)	0.41 (0.14–1.18)	0.10	0.44 (0.15–1.26)	0.13
Sesame seed sensitization [‡]	2 (0.7)	3 (1.1)		0.67		
Wheat sensitization [‡]	6 (2.0)	2 (0.7)		0.29		
Fish sensitization [‡]	2 (0.7)	1 (0.4)		1.00		
Ryegrass pollen sensitization	22 (6.0)	27 (8.3)	0.72 (0.41–1.26)	0.25	0.76 (0.44–1.32)	0.33
Olive tree sensitization	14 (3.7)	9 (2.7)	1.38 (0.58–3.24)	0.46	1.45 (0.62–3.40)	0.39
<i>Dermatophagoides</i> <i>pteronysinus</i> sensitization	23 (6.2)	25 (7.5)	0.83 (0.48–1.46)	0.52	0.82 (0.47–1.43)	0.49
<i>Dermatophagoides</i> <i>farinae</i> sensitization	16 (4.5)	17 (4.9)	0.92 (0.46–1.82)	0.80	0.93 (0.47–1.84)	0.84
Cat sensitization	25 (6.8)	12 (3.4)	1.97 (0.98–3.95)	0.06	1.95 (0.98–3.89)	0.06
<i>Alternaria tenuis</i> sensitization	32 (8.8)	23 (6.8)	1.29 (0.76–2.19)	0.34	1.31 (0.78–2.22)	0.31

*All data are based on analysis of 50 imputed datasets. For n-3 LCPUFA and control groups, data are number of subjects (percentage).

[†]Adjusted for centre, parity, maternal history and sex.

[‡]Fisher's exact test using original data, adjusted/imputed analyses not done due to rarity of outcomes.

CI, confidence interval; n-3 LCPUFA, long chain polyunsaturated fatty acid; RR, relative risk.

**Figure 2** Allergen sensitization profile at 1 and 3 years of age.

The pattern of allergic disease is known to differ with age, with the greatest incidence of food allergy and atopic dermatitis/eczema being in the first few years of life, while asthma and allergic rhinitis continue to rise until adulthood (16). Similarly, the changing sensitization pattern was as expected; the incidence of food allergen sensitization decreased and the aeroallergen sensitization increased between 1 to 3 years of age (17, 18). Specifically, the frequency of egg sensitization significantly reduced, and the significant reduction in egg sensitization at 1 year of age with n-3 LCPUFA treatment (9.3% vs 15.4%; adjusted relative risk 0.62; 95% CI 0.41–0.93; $P = 0.02$) (11) disappeared by 3 years of age. One could

question whether the timing of various allergen exposures during the first few years of life and the corresponding timing of the n-3 LCPUFA supplementation may influence the pattern of sensitization.

Conclusion

Overall, n-3 LCPUFA supplementation during pregnancy did not result in a significant reduction in IgE-associated allergic disease in the first 3 years of life to the magnitude originally predicted. Collectively, results from this and other n-3 LCPUFA supplementation RCTs suggest that the dose,

timing and duration of n-3 LCPUFA supplementation may influence sensitization and allergic disease outcomes. Clearly, further follow-up studies are required to definitively determine whether there is benefit in n-3 LCPUFA supplementation as an allergy prevention strategy.

Acknowledgments

We thank the families who participated and the following investigators and research team who supported the data collection: Anderson A, Griffith E, Warcup S, Tan F, Tye D, Quinn P, Di Giovine P, Shahin T, O'Grady M, Oates C, Scott A, Helbers W, Calderisi D, Garreffa H, Naccarella N, Loudis H and Crabb M. We also acknowledge the support of the Docosahexaenoic Acid to Optimise Mother Infant Outcome (DOMInO) trial steering committee: Makrides M, Gibson RA, McPhee AJ, Yelland L, Quinlivan J and Ryan P.

Funding declaration

The trial was supported by a grant from the Australian National Health and Medical Research Council (ID 399389) and a grant from the Australian Egg Corporation Limited.

References

- Calder PC, Yaqoob P, Thies F, Wallace FA, Miles EA. Fatty acids and lymphocyte functions. *Br J Nutr* 2002;**87**(Suppl 1):S31–S48.
- Gotttrand F. Long-chain polyunsaturated fatty acids influence the immune system of infants. *J Nutr* 2008;**138**:1807S–1812S.
- Calvani M, Alessandri C, Sopo SM, Panetta V, Pingitore G, Tripodi S et al. Consumption of fish, butter and margarine during pregnancy and development of allergic sensitizations in the offspring: role of maternal atopy. *Pediatr Allergy Immunol* 2006;**17**:94–102.
- Romieu I, Torrent M, Garcia-Esteban R, Ferrer C, Ribas-Fito N, Anto JM et al. Maternal fish intake during pregnancy and atopy and asthma in infancy. *Clin Exp Allergy* 2007;**37**:518–525.
- Salam MT, Li YF, Langholz B, Gilliland FD. Maternal fish consumption during pregnancy and risk of early childhood asthma. *J Asthma* 2005;**42**:513–518.
- Sausenthaler S, Koletzko S, Schaaf B, Lehmann I, Borte M, Herbarth O et al. Maternal diet during pregnancy in relation to eczema and allergic sensitization in the offspring at 2 y of age. *Am J Clin Nutr* 2007;**85**:530–537.
- Willers SM, Devereux G, Craig LC, McNeill G, Wijga AH, Abou El-Magd W et al. Maternal food consumption during pregnancy and asthma, respiratory and atopic symptoms in 5-year-old children. *Thorax* 2007;**62**:773–779.
- Dunstan JA, Mori TA, Barden A, Beilin LJ, Taylor AL, Holt PG et al. Fish oil supplementation in pregnancy modifies neonatal allergen-specific immune responses and clinical outcomes in infants at high risk of atopy: a randomized, controlled trial. *J Allergy Clin Immunol* 2003;**112**:1178–1184.
- Furuhjelm C, Warstedt K, Larsson J, Fredriksson M, Bottcher MF, Falth-Magnusson K et al. Fish oil supplementation in pregnancy and lactation may decrease the risk of infant allergy. *Acta Paediatr* 2009;**98**:1461–1467.
- Olsen SF, Osterdal ML, Salvig JD, Mortensen LM, Rytter D, Secher NJ et al. Fish oil intake compared with olive oil intake in late pregnancy and asthma in the offspring: 16 year of registry-based follow-up from a randomized controlled trial. *Am J Clin Nutr* 2008;**88**:167–175.
- Palmer DJ, Sullivan T, Gold MS, Prescott SL, Heddle R, Gibson RA et al. Effect of n-3 long chain polyunsaturated fatty acid supplementation in pregnancy on infants' allergies in first year of life: randomised controlled trial. *BMJ* 2012;**344**:e184.
- Furuhjelm C, Warstedt K, Fageras M, Falth-Magnusson K, Larsson J, Fredriksson M et al. Allergic disease in infants up to 2 years of age in relation to plasma omega-3 fatty acids and maternal fish oil supplementation in pregnancy and lactation. *Pediatr Allergy Immunol* 2011;**22**:505–514.
- Makrides M, Gibson RA, McPhee AJ, Yelland L, Quinlivan J, Ryan P. Effect of DHA supplementation during pregnancy on maternal depression and neurodevelopment of young children: a randomized controlled trial. *JAMA* 2010;**304**:1675–1683.
- Hanifin JM, Rijken G. Diagnostic features of atopic dermatitis. *Acta Dermatovener* 1980;**92**:44–47.
- D'Vaz N, Meldrum SJ, Dunstan JA, Martino D, McCarthy S, Metcalfe J et al. Postnatal fish oil supplementation in high-risk infants to prevent allergy: randomized controlled trial. *Pediatrics* 2012;**130**:674–682.
- Spergel JM. From atopic dermatitis to asthma: the atopic march. *Ann Allergy Asthma Immunol* 2010;**105**:99–106; quiz 107–9, 117.
- Dean T, Venter C, Pereira B, Arshad SH, Grundy J, Clayton CB et al. Patterns of sensitization to food and aeroallergens in the first 3 years of life. *J Allergy Clin Immunol* 2007;**120**:1166–1171.
- Kulig M, Bergmann R, Klettke U, Wahn V, Tacke U, Wahn U. Natural course of sensitization to food and inhalant allergens during the first 6 years of life. *J Allergy Clin Immunol* 1999;**103**:1173–1179.

The treatment and placebo capsules were donated by Efamol, UK.

Author contributions

Conception and design (MM, MG, SP and RG), acquisition of data (DP, MM, TS, MG, SP and RH), analysis and interpretation of data (DP, MM, TS, MG, SP, RH and RG), drafting of the manuscript (DP, MM and TS), critical revision of the manuscript (all), statistical analysis (DP, MM and TS), obtained funding (MM, MG, SP, RG and DP) and study supervision (DP, MM, MG and RH).

Conflicts of interest

RAG serves on a scientific advisory board for Fonterra, SLP serves on scientific advisory boards for the Nestle Nutrition Institute and Danone, and MM serves on scientific advisory boards for the Nestle Nutrition Institute, Danone and Fonterra. All associated honoraria for MM and RAG are paid to their institutions for professional development of students and early career researchers. No other authors declare a conflict of interest.