

Maternal prenatal psychological distress associates with offspring early-life wheezing – FinnBrain Birth Cohort

Emma Puosi , Laura S. Korhonen, Linnea Karlsson, Eeva-Leena Kataja, Heikki Lukkarinen, Hasse Karlsson, Minna Lukkarinen

First published: 30 November 2021

<https://doi-org.bibliosan.clas.cineca.it/10.1111/pai.13706>

Funding information:

This work was supported by the Academy of Finland (#134950, # 308176, #325292) (HK, LKa, LKa), Aivosäätiö (ELK), Allergy Research Foundation (EP), Alli Paasikivi Foundation (ELK), the Finnish Medical Foundation (LKa, LKo, ML), the Finnish State Grants for Clinical Research (LKa, LKo, ML), the Instrumentarium Science Foundation (LKo), the Yrjö Jahnsson Foundation (LKo), Jane and Aatos Erkko Foundation (HK), Jenny and Antti Wihuri Foundation (ELK), Juho Vainio Foundation (EP, LKo, ML), Päivikki and Sakari Sohlberg Foundation (EP, LKo, ML), Signe and Ane Gyllenberg Foundation (EP, LKa, LKo, ML), the Southwestern Finland Foundation of Allergy Research (EP), and the TYKS Foundation (EP)—all in Finland. None of the funding sources had a role in study design, data collection, analyses, interpretation of data, writing of the report, or decision to submit this manuscript for publication.

Abstract

Background

Exposure to prenatal maternal psychological distress may contribute to the development of childhood atopic disorders. Little is known about the importance of distress severity and its duration for the risk. Our aim was to investigate how chronic maternal depressive and anxiety symptoms across gestation influence the risk of wheezing and eczema at child age 24 months.

Methods

The study population was drawn from the FinnBrain Birth Cohort Study, including 1305 mother-infant dyads followed across gestation until the child age of 24 months when the outcomes were mother-reported wheezing ever and doctor-diagnosed eczema. To investigate the risk of wheezing phenotypes, wheezing with and without eczema was separated. Maternal distress was assessed with the Edinburgh Postnatal Depression

Scale for depressive and the Symptom Checklist-90 for anxiety symptoms three times during pregnancy, and the chronicity was demonstrated using symptom trajectories composed by latent growth mixture modeling.

Results

Of the children, 219/1305 (17%) had wheezing ever and 285/1276 (22%) had eczema. Risk of wheezing ever was elevated with maternal consistently high depressive symptoms (adjusted odds ratio 2.74; 95% confidence interval 1.37–5.50) or moderate and increasing anxiety symptoms (1.94; 1.06–3.54, respectively). Similarly, wheezing without eczema was associated with consistently high depressive (3.60; 1.63–7.94, respectively) and moderate and increasing anxiety symptoms (2.43; 1.21–4.91, respectively).

Conclusions

Maternal chronic psychological distress across gestation was associated with toddler wheezing and especially wheezing without other atopic features (eczema). This finding supports the theory of intrauterine programming effect by maternal psychological distress on offspring immune system and respiratory morbidity.

Abbreviations

CI

confidence interval

EPDS

Edinburgh Postnatal Depression Scale

gwk

gestational week

IQR

interquartile range

ISAAC

The International Study of Asthma and Allergies in Childhood

OR

odds ratio

PD

psychological distress

SCL-90

Symptom Checklist-90, anxiety subscale

SD

standard deviation

Key message

Maternal prenatal chronically elevated psychological distress symptoms were

associated with offspring wheezing and eczema at child age 24 months, suggesting altered immunology and increased risk of atopic diseases.

1 INTRODUCTION

Wheezing respiratory illness in early childhood is an expiratory breathing difficulty with multifactorial background, and it may indicate the development of asthma when occurring recurrently.¹ The incidence of childhood asthma has increased during the last decades, and the subsequent socioeconomic burden is considerable.¹ While multiple contributors, for example, hygiene hypothesis have been discovered, much of the increased incidence remains unexplained.¹ Current asthma research in children has mainly focused on postnatal exposures even though factors during gestation such as maternal psychological distress (PD) symptoms during gestation have recently been associated with elevated risk of offspring atopic diseases.²⁻¹² This association could result from the programming effect of maternal PD predisposing the offspring to aberrant immune responses and altered lung development by interfering with immune functions alongside other integrated systems.¹³ Animal studies have shown that chronic prenatal PD exposes the offspring to immune system changes,⁵ while only little is known how chronic maternal PD symptoms affect the immune system in human fetuses/infants.¹⁴ Majority of previous prospective birth cohort studies on maternal PD during gestation and later risk of offspring wheezing and asthma included only one timepoint during gestation, and only two investigated stress separately in two trimesters.^{6-8, 10, 12, 15-19} It is suggested that exposure to maternal PD during end-gestation, rather than mid-gestation, predisposes older children to atopic diseases, which could be the effect of a sensitive window but also the cumulating effect of chronic PD symptoms.² However, our recently published study indicated that chronic PD symptoms across gestation increase the risk for food allergy at the age of 6 months suggesting an early programming effect and altered immune responses¹⁴. Other previous studies have not used standardized analyses on severity and duration of maternal PD symptoms covering the entire gestation.¹⁴ Based on this study, we hypothesized that chronic maternal PD symptoms (depression or anxiety) also influence the offspring risk for later wheezing and eczema.^{2, 6} We aimed to investigate how chronic PD symptoms associate with (1) wheezing ever or doctor-diagnosed eczema at the age of 24 months and further with (2) with atopic and non-atopic wheezing subtypes, which may serve as indicators for later childhood asthma phenotypes.

2 METHODS

2.1 Study subjects and design

The FinnBrain Birth Cohort Study is population-based, prospective pregnancy cohort investigating the long-term effects of prenatal and early-life exposures on child health

outcomes.²⁰ Originally, the cohort consists of 3808 mothers, 2623 fathers, and 3837 children. The families were recruited by research nurses when attending the free-of-charge ultrasounds at gestational weeks (gwks) 12 in three maternity welfare clinics in the Southwestern Hospital District and Åland Islands in Finland 12/2011–04/2015. To this study, 1368 mother-infant dyads with available primary outcome data at the child age of 24 months were eligible (Figure S1). Of these, the children born preterm (gwks <37) were excluded since the preterm delivery is a risk factor for wheezing (Figure S1).²¹ The study conforms to the Declaration of Helsinki, was approved by the Ethics Committee of the Hospital District of Southwest Finland (ETMK:57/180/2011), and commenced only after obtaining written informed consent from the guardians.

2.2 Exposures

At gwks 14, 24, 34, and at child age 24 months, the maternal PD symptom scores were assessed using the Edinburgh Postnatal Depression Scale (EPDS, range 0–30 points) for the depressive and the Symptom Checklist-90, anxiety scale (SCL-90, range 0–40 points) for the anxiety symptoms.^{20, 22} Prenatal symptom chronicity and severity assessment was based on the EPDS and SCL-90 scores during entire gestation and modeled as symptom trajectories using latent growth mixture modeling with imputed missing data in Mplus 6.0 (StatModel).^{23, 24} Five latent curves of depressive symptoms and three of anxiety symptoms were chosen for analyses based on the inspection of Akaike's information criterion and Bayesian's information criterion and entropy value (Figures 1 and 2, Table S6). Data for maternal body mass index, smoking, and delivery were derived from the Finnish Medical Birth Register and other data from the parental questionnaires (Table 1). Parental education level was used as the indicator of socioeconomical status. It was divided into three groups (low: ≤ 12 , mid: 12–15, and high: >15 years).²⁰ Atopic diseases were identified from the question “Have you or at least one of your children had allergies, allergic rhinitis and/or eczema?” (yes/no).





FIGURE 1[Open in figure viewer](#) | [PowerPoint](#)

Five latent growth curve trajectories of depressive symptoms illustrate the chronicity and severity of the maternal psychological distress symptoms during entire gestation. The depressive symptoms were assessed using the Edinburgh Postnatal Depression Scale (EPDS, range 0–30 points)

**FIGURE 2**[Open in figure viewer](#) | [PowerPoint](#)

Three latent curves of anxiety symptoms illustrate the chronicity and severity of the maternal psychological distress symptoms during entire gestation. The anxiety symptoms with the Symptom Checklist-90, anxiety scale (SCL-90, range 0–40 points)

TABLE 1. Population characteristics and attrition analysis

Exposures	Cohort <i>N</i> = 3837	Included <i>N</i> = 1305	Excluded <i>N</i> = 2532	<i>p</i> ^a
Maternal characteristics during gestation				
Maternal age at birth, years (SD)	31 (4.5)	31 (4.4)	30 (4.8)	<.001
	<i>n</i> = 3684	<i>n</i> = 1305	<i>n</i> = 2532	
Maternal pre-pregnancy body mass index, kg/m ² median (IQR)	24 (22:27)	23 (21:27)	24 (21:27)	.16

Exposures	Cohort	Included	Excluded	<i>p</i> ^a
	<i>N</i> = 3837	<i>N</i> = 1305	<i>N</i> = 2532	
	<i>n</i> = 3684	<i>n</i> = 1287	<i>n</i> = 2297	
Maternal smoking, nr (%)				
Quit in early pregnancy	274/3748 (7.1)	65/1285 (5.1)	209/2396 (8.7)	<.001
Through pregnancy	207/3748 (5.4)	37/1285 (2.9)	170/2396 (7.1)	

Abbreviations: BMI, body mass index; CI, confidence interval; EPDS, The Edinburgh Postnatal Depression Scale; gwks, gestational weeks; IQR, interquartile range; N/A, not applicable; OR, odds ratio; SCL-90, The Symptom Checklist-90, anxiety scale; SD, standard deviation.

^a Indicates the comparisons with “excluded” as a reference group.

2.3 Outcomes

The primary outcomes were the maternal report of wheezing ever and doctor-diagnosed eczema at the child age of 24 months according to the standardized questionnaire of The International Study of Asthma and Allergies in Childhood (ISAAC).²⁵ To demonstrate the susceptibility to diverged immunology after prenatal PD exposure, we categorized the wheezing ever into atopic and non-atopic wheezing subtypes by adding the information of existing doctor-diagnosed eczema. For this purpose, we formed the four-class combination outcome: “wheezing with eczema,” “wheezing without eczema,” “eczema without wheezing,” and “neither wheezing nor eczema.”

2.4 Statistics

Characteristics and subgroup comparisons were done using the *t* test, Mann-Whitney *U* test, or the Pearson chi-square (Tables 1-3). In the primary analyses, the risk for wheezing ever or doctor-diagnosed eczema ever was assessed with the binary logistic regression, first in univariable and then in multivariable analyses. The multivariable analyses were adjusted for known risk factors; for wheezing, prenatal maternal smoking, parental education level, child's sex, and maternal history of asthma¹; whereas for eczema, child's sex, maternal history of atopic diseases, and parental education level.²⁶ The impact of chronic depressive and anxiety symptoms on the outcomes was analyzed using symptom trajectories with multinomial logistic regression, where the trajectory of consistently low symptoms was the reference group (Table 4, Figures 1 and 2). In the secondary analyses, the risk for the four-

class combination outcome was assessed using the multinomial logistic regression, where the “neither wheezing nor eczema” was the reference group (Table [S3](#)). These analyses were adjusted for parental education level, child's sex, maternal prenatal smoking, and history of asthma and/or atopic diseases as a combined variable. In the supplementary analyses, distress symptom scores were studied as continuous variables at each timepoint to illustrate the role of distress exposure separately in each trimester (Table [S2](#)). Correlation analysis of perinatal maternal PD symptom scores was performed (Table [S4](#)) following the inclusion of maternal PD symptoms at the child age of 24 months as a potential confounder of reporting bias for the pediatric outcomes (Table [S5](#)).²⁷ The *p* value <.05 was regarded as significant, and 95% confidence interval (CI) was used. Analyses were performed using the IBM SPSS 27.0 software (SPSS Inc.).

TABLE 2. Population characteristics of the included at 24 months

Exposures	No wheezing (<i>n</i> = 1086)	Wheezing ever (<i>n</i> = 219)	<i>p</i> ^a	No eczema (<i>n</i> = 991)	Doctor- diagnosed eczema (<i>n</i> = 285)	<i>p</i> ^a
Maternal characteristics during gestation						
Maternal age at birth, years (SD)	31 (4.4) <i>n</i> = 912	31 (4.2) <i>n</i> = 183	.42	31 (4.3) <i>n</i> = 991	31 (4.4) <i>n</i> = 285	.10
Maternal pre- pregnancy body mass index, kg/m ² median (IQR)	23 (21:26) <i>n</i> = 1070	24 (21:28) <i>n</i> = 217	.21	23 (21:26) <i>n</i> = 977	24 (21:27) <i>n</i> = 283	.86
Maternal smoking, nr (%)						
Quit in early pregnancy	52/1068 (4.9)	13/217 (6.0)	.79	50/975 (5.1)	15/283 (5.3)	.93
Through pregnancy	31/1068 (2.9)	6/217 (2.8)		27/975 (2.8)	9/283 (3.2)	
Maternal ^b	78/1042 (7.5)	30/207 (15)	.001	76/950 (8.0)	10/272 (9.2)	.53

Abbreviations: BMI, body mass index; CI, confidence interval; EPDS, The Edinburgh Postnatal Depression Scale; gwks, gestational weeks; IQR, interquartile range; N/A, not applicable; OR, odds ratio; SCL-90, The Symptom Checklist-90,

anxiety scale; SD, standard deviation. Bold values indicate significance of $P < .05$.

^a Indicates the group comparisons.

^b The correlation between maternal and sibling asthma in the cohort, $p = .003$

^c The correlation between maternal and sibling atopic diseases in the cohort, $p < .001$.

TABLE 3. Population characteristics of the included at 24 months

Exposures	Wheezing with eczema (<i>n</i> = 53)	Wheezing without eczema (<i>n</i> = 154)	Eczema without wheezing (<i>n</i> = 231)	Neither wheezing nor eczema (<i>n</i> = 829)	<i>p</i> ^a
Maternal characteristics during gestation					
Maternal age at birth, years (SD)	31 (3.4) <i>n</i> = 53	32 (4.4) <i>n</i> = 154	32 (4.4) <i>n</i> = 231	32 (4.4) <i>n</i> = 829	.27
Maternal pre- pregnancy body mass index, kg/m ² median (IQR)	24 (21:29) <i>n</i> = 53	24 (21:27) <i>n</i> = 152	24 (21:27) <i>n</i> = 229	23 (21:26) <i>n</i> = 817	.63
Maternal smoking, nr (%)					
Quit in early pregnancy	4/53 (7.5)	8/152 (5.3)	11/229 (4.8)	41/815 (5.0)	.71
Through pregnancy	3/53 (5.7)	2/152 (1.3)	6/229 (2.6)	25/815 (3.1)	
Maternal asthma ^b	7/48 (15)	18/147 (12)	18/223 (8.1)	58/795 (7.3)	.090

Abbreviations: BMI, body mass index; CI, confidence interval; EPDS, The Edinburgh Postnatal Depression Scale; gwks, gestational weeks; IQR, interquartile range; N/A, not applicable; OR, odds ratio; SCL-90, The Symptom Checklist-90, anxiety scale; SD, standard deviation. Bold values indicate significance of $P < .05$.

^a Indicates the comparisons with “Neither wheezing nor eczema” as the reference group.

^b The correlation between maternal and sibling asthma in the cohort, $p = .003$

^c The correlation between maternal and sibling atopic diseases in the cohort, $p < .001$

TABLE 4. Risk of wheezing or doctor-diagnosed atopic eczema at 24 months

Maternal reported exposures during gestation	Wheezing ever			Doctor-diagnosed eczema		
	OR	95% CI	<i>p</i> ^a	OR	95% CI	<i>p</i> ^a
Unadjusted analyses						
Maternal atopic diseases	1.4	1.0–1.8	.046	1.8	1.4–2.4	<.001
Maternal asthma	2.1	1.3–3.3	.001	1.2	0.73–1.9	.53
Maternal smoking during pregnancy						
Quit in early pregnancy ^a	1.2	0.67–2.3	.50	1.0	0.57–1.9	.90
Through pregnancy ^a	0.96	0.40–2.3	.93	1.2	0.54–2.49	.71
Older children with mother-reported atopic diseases	1.3	0.84–1.9	.26	2.0	1.4–2.8	<.001
Older children with mother-reported asthma	2.6	1.3–5.3	.008	0.87	0.38–2.0	.75
Male sex vs. female sex	1.9	1.4–2.6	<.001	1.1	0.87–1.5	.34
Parental education level						
Up to 12 years ^a	2.6	1.4–4.7	.002	1.4	0.95–1.9	.090
Between 13 and 15 years ^a	1.8	1.1–3.0	.032	1.3	0.95–1.8	.11
Over 15 years ^a	1.0			1.0		

Abbreviations: BMI, body mass index; CI, confidence interval; EPDS, The Edinburgh Postnatal Depression Scale; gwks, gestational weeks; IQR, interquartile range; N/A, not applicable; OR, odds ratio; SD, standard deviation; SCL-90, The Symptom Checklist-90, anxiety scale. Bold values indicate significance of $P < .05$.

^a Indicates the comparison with "No wheezing" or "No eczema" as the reference groups.

3 RESULTS

3.1 Study population and characteristics

At the child age of 24 months, maternal questionnaires including data for wheezing were available from 1368/3837 (36%) and for eczema from 1335/3837 (35%) of the cohort children (Figure S1). Children whose mothers did not return the 24-month questionnaire were excluded. The non-responding mothers were younger, had higher total sum scores in the EPDS and SCL-90 questionnaires, and smoked more during gestation (Table 1, Table S1, and

Figure S1). Additionally, children born preterm (gwks <37) were excluded, and from them, 63/1368 (5%) had available data on wheezing ever and 59/1335 (4%) on doctor-diagnosed eczema. Finally, 1305/3837 (34%) full-term children with available outcome data were included (Figure S1). From these 1305 children, the number of children whose mothers had reported wheezing ever was 219 (17%) and doctor-diagnosed eczema 285 (22%) (Table 2 and Figure S1). Based on the four-class combination outcome, the number of children with wheezing with eczema was 53 (4%), wheezing without eczema 154 (12%), eczema without wheezing 231 (18%), or neither wheezing nor eczema 829 (66%) (Table 3).

The children with wheezing ever had more family atopy than non-wheezing children: maternal asthma (15% vs. 8%; $p = .001$) and atopic diseases (49% vs. 42%; $p = .046$), and older siblings with mother-reported asthma (6% vs. 2%; $p = .006$), which correlated with maternal asthma ($p < .001$) (Table 2). The children with eczema had more maternal atopic diseases than children without eczema (54% vs. 40%; $p < .001$) (Table 2). Male sex dominated among children with wheezing (66%; $p < .001$). No differences were found regarding maternal smoking during gestation or delivery type (Table 2).

3.2 The primary analysis

3.2.1 Risk of wheezing at age 24 months

In the unadjusted analyses, wheezing ever associated with maternal asthma and atopic diseases, gestational weeks at delivery, male sex, mother-reported asthma in older siblings, and parental education level (Tables 2 and 3). In the EPDS trajectory model, the risk was elevated with the “Consistently High” trajectory (odds ratio [OR] 2.74; 95% CI 1.37–5.50; $p = .004$) (Table 3, Figures 1 and 2). Likewise, SCL-90 trajectory of “Moderate and Increasing” symptoms was associated with increased risk of wheezing (Table 4, Figures 1 and 2). In the adjusted analyses, the results remained unchanged for both depressive and anxiety symptom trajectories (Table 4).

3.2.2 Risk of eczema at 24 months

In the unadjusted analysis, doctor-diagnosed eczema at 24 months associated with atopic diseases of mothers and older siblings (Table 3). Prenatal maternal depressive or anxiety symptom trajectories did not associate with doctor-diagnosed eczema (Table 3, Figures 1 and 2). After adjustments, the “High and Decreasing” SCL trajectory was associated with increased risk for eczema (OR 1.89; 95% CI 1.00–3.55; $p = .049$) (Table 4).

3.3 The secondary analysis

3.3.1 Risk in four-class combination outcome

In the unadjusted analysis, the risk for wheezing without eczema was elevated with the EPDS trajectory of “Consistently High” (OR 3.40; 95% CI 1.75–7.81; $p = .001$) and SCL-90 trajectory of “Moderate and Increasing” (2.08; 1.05–4.14; $p = .036$, respectively). The risk for eczema

without wheezing was elevated with the SCL-90 trajectory of “High and Decreasing” (2.19; 1.14–4.21; $p = .019$) (Table [S3](#)). Adjustments did not change the results (Table [S3](#)). In all trajectory analyses, the “neither wheezing nor eczema” was the reference group.

3.4 The supplementary analyses

The PD symptom scores were studied as continuous variables at each timepoint to illustrate the role of distress exposure separately in each trimester. The risk of wheezing ever was increased in the first and last trimesters when using the continuous EPDS scores and in the second and last trimesters when using the continuous SCL-90 scores (Table [S2](#)).

The chronic nature of maternal PD symptoms is illustrated with correlation analysis between PD symptoms during gestation and at 24 months (Table [S4](#)). The associations between EPDS trajectories during gestation and wheezing remained unchanged after adjusting for maternal PD symptoms at the child age of 24 months, but not between SCL-90 trajectories (Table [S5](#)). However, the associations remained unchanged when the risk of wheezing and eczema was examined in the four-class combination outcome of “Wheezing without eczema” and “Eczema without wheezing” (Table [S5](#)).

4 DISCUSSION

In keeping with earlier studies, we demonstrate that maternal PD during gestation associates with atopic disorders in offspring.^{3, 7, 8, 11, 16, 18, 19} Furthermore, we found that mother-experienced PD symptoms that are chronically elevated across gestation, assessed in three timepoints, associated with offspring wheezing and eczema at the age of 24 months. This is the first study on early childhood asthma which observed maternal PD across gestation and applied latent growth curve modeling in composing the trajectories for chronicity/severity of the PD symptoms.¹⁴ Our finding is in line with our previous study where maternal chronic PD associated with offspring food allergy at the infant age of 6 months.¹⁴ Our study also sheds new light into the effects of chronic maternal PD resulting in varied risk for different wheezing subtypes.²

The novelty of our study was the creation of PD symptom trajectories enabling the investigation of symptom severity and chronicity on the outcomes. In previous studies on prenatal PD and atopic diseases, the questionnaires and distress types and timing have been heterogenous.² The studies to date have used different validated questionnaires of depressive and anxiety symptoms^{3, 7, 9} but also self-reported negative life events or other stressful situations.^{8, 12, 18, 19} The timing of these exposures during gestation has varied between studies,² but also inside studies, a single time point at any time during gestation,⁸ at one trimester,^{7, 10–12, 18, 19} at two trimesters (2nd and 3rd),⁴ or before and after gwks 18.³ Derived from previous studies the third trimester may be the most vulnerable time regarding PD exposure.² However, our study shows that chronic depressive and anxiety symptoms may be risk factor for wheezing and eczema and this finding is endorsed by the

aligned results of PD symptoms per the time point assessment. We adjusted analyses for maternal atopic diseases or asthma since the correlation between mother/sibling atopic diseases and asthma was significant. Male sex associated with wheezing, as expected.²¹ Possible reporting bias because of maternal distress symptoms was acknowledged, and therefore, we also adjusted for the PD symptoms at 24 months.²⁷ Also, we acknowledge that the attrition could affect the generalizability of the results particularly in the population with accumulated risk factors as maternal PD and lower socioeconomic status were related to attrition. The potential bias resulting from these characteristics in the attrition may dilute the observed associations as the subjects with higher PD symptoms were underrepresented.

To provide new insights into underlying aberrant immune system of different early childhood wheezing subtypes, a four-class combination outcome was applied. For this, we combined wheezing ever and doctor-diagnosed eczema to model the simplified outcomes of “wheezing with eczema,” “wheezing without eczema,” “eczema without wheezing,” and “neither wheezing nor eczema.” They were intended to anticipate the atopic/non-atopic wheezing subtypes, and further, the early phenotypes of atopic/non-atopic asthma. Noteworthy is that exposure to prenatal chronic PD was associated with wheezing without eczema, but not with wheezing with eczema or eczema without wheezing, suggesting that prenatal chronic PD predisposes the offspring toward non-atopic wheezing subtype. However, we acknowledge the statistical power issue caused by relatively small cell sizes, and therefore, conclusions should be made with caution. We hope that these preliminary findings inspire future research as they support the previous studies showing that the susceptibility to diverged immune responses leading to school-age atopic or non-atopic asthma can be seen during the first year of life implying that asthma programming occurs prenatally because of genetic predisposition and environmental exposures.^{2, 5, 28, 29} In addition, previously has been shown that school-age atopic and non-atopic asthma have different early-life risk factors; that is, atopic asthma has concomitant sensitization and/or eczema in infancy whereas maternal smoking during gestation has associated with non-atopic asthma.^{28, 29} In current study, maternal prenatal smoking did not associate with wheezing ever or wheezing without eczema, and importantly, maternal depressive and anxiety symptoms did not associate with smoking either. This underlines the notion that prenatal PD may be an independent risk factor for offspring non-atopic wheezing subtype. Prenatal exposures may partly explain the non-genetic variability seen in childhood asthma phenotypes interfering with the immune system and inducing the shift toward Th2-dominant predisposition. The allergen-specific immunoglobulin E was not tested here since sensitization to aeroallergens develops later in general population and its value indicating asthma risk in early life is decreased.³⁰ Possible simultaneously mediating mechanisms might include the elevated levels of maternal glucocorticoids and/or cytokines resulting in placental dysregulation, impaired fetal maturation of HPA axis, and altered glucocorticoid levels.^{2, 5}

The strengths include the large and non-selected sample of 1305 mother-infant dyads from prospective, population-based pregnancy cohort without major exclusion criteria. It can be perceived as an unselected sample of the general population and therefore represents the maternal and paternal prenatal period of depressive and anxiety symptoms adequately. In general population, the prevalence of wheezing ever before the child age of three years is approximately 20% and for childhood eczema approximately 20%.^{26, 31} In current study, the prevalence of wheezing ever and eczema was within expected. The outcomes were based on the structured and widely used ISAAC questionnaire which makes comparison between studies feasible.²⁵ In earlier studies on prenatal PD exposure, the outcomes have varied greatly including mother-reported wheezing,¹⁰ mother-reported doctor-diagnosed wheeze and asthma,^{3, 7} mother-reported doctor-diagnosed eczema,^{8, 9, 11} only mother-reported,^{4, 12} and ISAAC-based eczema,¹¹ ranging between 1 and 14 years.² The use of mother-reported questionnaires poses a possible reporting bias, also in our study. The review of medical charts would have provided more accurate outcomes, but as our study population originates from large birth cohort with multiple simultaneous multi-disciplined sub-studies, the number of questions were restricted. We would have benefitted from the wheezing severity or number of wheezing episodes, but it was unachievable to examine children during every wheezing episode.

In conclusion, mother-experienced chronic PD symptoms during gestation were associated with elevated risk of offspring wheezing and non-atopic type of wheezing during the first two years of life. The findings highlight the influence of prenatal PD exposure on immune system and support the theory of intrauterine programming effect of maternal PD leading to atopic diseases and respiratory morbidity.²

ACKNOWLEDGMENT

We thank the families participating in the study and the study nurses and staff at the FinnBrain Birth Cohort Study for their assistance.

CONFLICT OF INTEREST

None.

AUTHOR CONTRIBUTIONS

Emma Puosi: Conceptualization (lead); Data curation (lead); Formal analysis (lead); Funding acquisition (supporting); Investigation (lead); Methodology (lead); Visualization (lead); Writing – original draft (lead); Writing – review & editing (equal). **Laura S. Korhonen:** Data curation (supporting); Investigation (supporting); Methodology (supporting); Resources (supporting); Writing – original draft (supporting); Writing – review & editing (equal). **Linnea Karlsson:** Conceptualization (supporting); Funding acquisition (supporting); Project administration (lead); Resources (supporting); Writing – review & editing (equal). **Eeva-Leena Kataja:**

Conceptualization (supporting); Data curation (supporting); Formal analysis (supporting); Methodology (supporting); Visualization (supporting); Writing – original draft (supporting); Writing – review & editing (equal). **Heikki Lukkarinen:** Conceptualization (lead); Investigation (supporting); Methodology (equal); Supervision (equal); Writing – review & editing (equal). **Hasse Karlsson:** Conceptualization (supporting); Funding acquisition (supporting); Project administration (lead); Resources (supporting); Writing – review & editing (equal). **Minna Lukkarinen:** Conceptualization (lead); Data curation (equal); Funding acquisition (supporting); Methodology (lead); Supervision (lead); Writing – original draft (supporting); Writing – review & editing (lead).

PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1111/pai.13706>.

Open Research



PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1111/pai.13706>.

Supporting Information



Filename	Description
pai13706-sup-0001-FigureS1.pdf PDF document, 102.6 KB	Figure S1
pai13706-sup-0002-Supinfo.docx Word 2007 document , 77 KB	Supplementary Material

Please note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries (other than missing content) should be directed to the corresponding author for the article.

REFERENCES



- 1 Beasley R, Semprini A, Mitchell EA. Risk factors for asthma: is prevention possible? *Lancet*. 2015;

386: 1075- 1085. doi:[10.1016/S0140-6736\(15\)00156-7](https://doi.org/10.1016/S0140-6736(15)00156-7)

[Crossref](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

2 Flanigan C, Sheikh A, DunnGalvin A, Brew BK, Almquist C, Nwaru BI. Prenatal maternal psychosocial stress and offspring's asthma and allergic disease: a systematic review and meta-analysis. *Clin Exp Allergy*. 2018; **48**(4): 403- 414. [10.1111/cea.13091](https://doi.org/10.1111/cea.13091)

[Wiley Online Library](#) | [CAS](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

3 Cookson H, Granell R, Joinson C, Ben-Shlomo Y, Henderson AJ. Mothers' anxiety during pregnancy is associated with asthma in their children. *J Allergy Clin Immunol*. 2009; **123**(4): 847-853.e11. doi:[10.1016/j.jaci.2009.01.042](https://doi.org/10.1016/j.jaci.2009.01.042) .

<http://www.ncbi.nlm.nih.gov/bibliosan.clas.cineca.it/pubmed/19348924>

[Crossref](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

4 Hartwig IRV, Sly PD, Schmidt LA, et al. Prenatal adverse life events increase the risk for atopic diseases in children, which is enhanced in the absence of a maternal atopic predisposition. *J Allergy Clin Immunol*. 2014; **134**: 160- 169. doi:[10.1016/j.jaci.2014.01.033](https://doi.org/10.1016/j.jaci.2014.01.033)

[Crossref](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

5 Veru F, Laplante DP, Luheshi G, King S. Prenatal maternal stress exposure and immune function in the offspring. *Stress*. 2014; **17**(2): 133- 148. doi:[10.3109/10253890.2013.876404](https://doi.org/10.3109/10253890.2013.876404)

[Crossref](#) | [CAS](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

6 Andersson NW, Hansen MV, Larsen AD, Hougaard KS, Kolstad HA, Schlünssen V. Prenatal maternal stress and atopic diseases in the child: a systematic review of observational human studies. *Allergy*. 2016; **71**(1): 15- 26. doi:[10.1111/all.12762](https://doi.org/10.1111/all.12762) .

<http://www.ncbi.nlm.nih.gov/bibliosan.clas.cineca.it/pubmed/26395995>

[Wiley Online Library](#) | [CAS](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

7 Guxens M, van der Sonnenschein Voort AMM, Tiemeier H, et al. Parental psychological distress during pregnancy and wheezing in preschool children: the Generation R Study. *J Allergy Clin Immunol*. 2014; **133**(1): 59- 67.e12. [10.1016/j.jaci.2013.04.044](https://doi.org/10.1016/j.jaci.2013.04.044) .

<http://www.ncbi.nlm.nih.gov/bibliosan.clas.cineca.it/pubmed/23777854>

[Crossref](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

8 Wang IJ, Wen HJ, Chiang TL, Lin SJ, Chen PC, Guo YL. Maternal employment and atopic dermatitis in children: a prospective cohort study. *Br J Dermatol*. 2013; **168**: 794- 801. doi: [10.1111/bjd.12195](https://doi.org/10.1111/bjd.12195)

[Wiley Online Library](#) | [CAS](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

9 Cheng TS, Chen H, Lee T, et al. An independent association of prenatal depression with wheezing and anxiety with rhinitis in infancy. *Pediatr Allergy Immunol*. 2015; **26**(8): 765- 771. doi: [10.1111/pai.12453](https://doi.org/10.1111/pai.12453) . <http://www.ncbi.nlm.nih.gov/bibliosan.clas.cineca.it/pubmed/26235785>

[Wiley Online Library](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

10 Chiu CY, Huang YL, Tsai MH, et al. Sensitization to food and inhalant allergens in relation to

atopic diseases in early childhood: a birth cohort study. *PLoS One*. 2014; **9**(7):e102809. doi: [10.1371/journal.pone.0102809](https://doi.org/10.1371/journal.pone.0102809)

[Crossref](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

11 Chang HY, Suh DI, Yang SI, et al. Prenatal maternal distress affects atopic dermatitis in offspring mediated by oxidative stress. *J Allergy Clin Immunol*. 2016; **138**: 468- 475.e5. doi: [10.1016/j.jaci.2016.01.020](https://doi.org/10.1016/j.jaci.2016.01.020)

[Crossref](#) | [CAS](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

12 Wen HJ, Wang YJ, Lin YC, et al. Prediction of atopic dermatitis in 2-yr-old children by cord blood IgE, genetic polymorphisms in cytokine genes, and maternal mentality during pregnancy. *Pediatr Allergy Immunol*. 2011; **22**: 695- 703. doi:[10.1111/j.1399-3038.2011.01177.x](https://doi.org/10.1111/j.1399-3038.2011.01177.x)

[Wiley Online Library](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

13 Rosa MJ, Lee AG, Wright RJ. Evidence establishing a link between prenatal and early-life stress and asthma development. *Curr Opin Allergy Clin Immunol*. 2018; **18**(2): 148- 158. doi: [10.1097/ACI.0000000000000421](https://doi.org/10.1097/ACI.0000000000000421) .

<https://pubmed-ncbi-nlm-nih.gov.bibliosan.clas.cineca.it/29369067/>

[Crossref](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

14 Lukkarinen M, Puosi E, Kataja E, et al. Maternal psychological distress during gestation is associated with infant food allergy. *Pediatr Allergy Immunol*. 2021; **32**(4): 787- 792. doi: [10.1111/pai.13449](https://doi.org/10.1111/pai.13449)

[Wiley Online Library](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

15 de Marco R, Pesce G, Girardi P, et al. Foetal exposure to maternal stressful events increases the risk of having asthma and atopic diseases in childhood. *Pediatr Allergy Immunol*. 2012; **23**(8): 724- 729. doi:[10.1111/j.1399-3038.2012.01346.x](https://doi.org/10.1111/j.1399-3038.2012.01346.x) .

<http://www.ncbi.nlm.nih.gov.bibliosan.clas.cineca.it/pubmed/22957808>

[Wiley Online Library](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

16 Reyes M, Perzanowski MS, Whyatt RM, et al. Relationship between maternal demoralization, wheeze, and immunoglobulin E among inner-city children. *Ann Allergy Asthma Immunol*. 2011; **107**(1): 42- 49.e1. doi:[10.1016/j.anai.2011.03.004](https://doi.org/10.1016/j.anai.2011.03.004) .

<http://www.ncbi.nlm.nih.gov.bibliosan.clas.cineca.it/pubmed/21704884>

[Crossref](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

17 Pacheco-Gonzalez RM, Mallol J, Solé D, et al. Factors associated with the time to the first wheezing episode in infants: a cross-sectional study from the International Study of Wheezing in Infants (EISL). *NPJ Prim Care Respir Med*. 2016; **26**(1): 15077. doi:[10.1038/npjpcrm.2015.77](https://doi.org/10.1038/npjpcrm.2015.77) .

<http://www.nature.com.bibliosan.clas.cineca.it/articles/npjpcrm201577>

[Crossref](#) | [PubMed](#) | [Google Scholar](#)

18 Chiu Y-HM, Coull BA, Cohen S, Wooley A, Wright RJ. Prenatal and postnatal maternal stress and wheeze in urban children: effect of maternal sensitization. *Am J Respir Crit Care Med*. 2012; **186**(2): 147- 154. doi:[10.1164/rccm.201201-0162OC](https://doi.org/10.1164/rccm.201201-0162OC)

[Crossref](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

19 Rosa MJ, Just AC, Ortiz TY, et al. Prenatal and postnatal stress and wheeze in Mexican children: sex-specific differences. *Ann Allergy Asthma Immunol*. 2016; **116**: 306- 312.e1. doi: [10.1016/j.anai.2015.12.025](https://doi.org/10.1016/j.anai.2015.12.025)

[Crossref](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

20 Karlsson L, Tolvanen M, Scheinin NM, et al. Cohort profile: the FinnBrain Birth Cohort Study (FinnBrain). *Int J Epidemiol*. 2018; **47**(1): 15- 16j. doi:[10.1093/ije/dyx173](https://doi.org/10.1093/ije/dyx173) . <http://www.ncbi.nlm.nih.gov/bibliosan.clas.cineca.it/pubmed/29025073>

[Crossref](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

21 Jaakkola JJK, Ahmed P, Ieromnimon A, et al. Preterm delivery and asthma: a systematic review and meta-analysis. *J Allergy Clin Immunol*. 2006; **118**: 823- 830. doi:[10.1016/j.jaci.2006.06.043](https://doi.org/10.1016/j.jaci.2006.06.043)

[Crossref](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

22 Bergink V, Kooistra L, van den Lambregtse Berg MP, et al. Validation of the Edinburgh Depression Scale during pregnancy. *J Psychosom Res*. 2011; **70**(4): 385- 389. doi: [10.1016/j.jpsychores.2010.07.008](https://doi.org/10.1016/j.jpsychores.2010.07.008) .

<https://pubmed-ncbi-nlm-nih.gov.bibliosan.clas.cineca.it/21414460/>

[Crossref](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

23 Korja R, Nolvi S, Kataja EL, et al. The courses of maternal and paternal depressive and anxiety symptoms during the prenatal period in the Finnbrain birth cohort study. *PLoS One*. 2018; **13**(12):e0207856. doi:[10.1371/journal.pone.0207856](https://doi.org/10.1371/journal.pone.0207856)

[Crossref](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

24 Muthén L, Muthén B. *Mplus User's Guide*. 6th ed. Muthén & Muthén; 2012. doi: [10.1111/j.1600-0447.2011.01711.x](https://doi.org/10.1111/j.1600-0447.2011.01711.x)

[Google Scholar](#)

25 Asher MI, Keil U, Anderson HR, et al. International study of asthma and allergies in childhood (ISAAC): rationale and methods. *Eur Respir J*. 1995; **8**(3): 483- 491. doi:

[10.1183/09031936.95.08030483](https://doi.org/10.1183/09031936.95.08030483) . <http://erj.ersjournals.com/content/8/3/483>

[Crossref](#) | [CAS](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

26 Abuabara K, Yu AM, Okhovat J-P, Allen IE, Langan SM. The prevalence of atopic dermatitis beyond childhood: a systematic review and meta-analysis of longitudinal studies. *Allergy*. 2018; **73**(3): 696- 704. doi:[10.1111/all.13320](https://doi.org/10.1111/all.13320) .

<http://www.ncbi.nlm.nih.gov/bibliosan.clas.cineca.it/pubmed/28960336>

[Wiley Online Library](#) | [CAS](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

27 Korhonen LS, Karlsson L, Scheinin NM, et al. Prenatal maternal psychological distress and offspring risk for recurrent respiratory infections. *J Pediatr*. 2019; **208**: 229- 235.e1. doi:

[10.1016/j.jpeds.2018.12.050](https://doi.org/10.1016/j.jpeds.2018.12.050) .

<https://pubmed-ncbi-nlm-nih.gov.bibliosan.clas.cineca.it/30723014/>

[Crossref](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

28 Lukkarinen M, Koistinen A, Turunen R, Lehtinen P, Vuorinen T, Jartti T. Rhinovirus-induced first wheezing episode predicts atopic but not nonatopic asthma at school age. *J Allergy Clin Immunol*. 2017; **140**(4): 988- 995. doi:[10.1016/j.jaci.2016.12.991](https://doi.org/10.1016/j.jaci.2016.12.991) .

<http://www.ncbi.nlm.nih.gov/bibliosan.clas.cineca.it/pubmed/28347734>

[Crossref](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

29 Goksör E, Alm B, Pettersson R, et al. Early fish introduction and neonatal antibiotics affect the risk of asthma into school age. *Pediatr Allergy Immunol*. 2013; **24**(4): 339- 344. doi:

[10.1111/pai.12078](https://doi.org/10.1111/pai.12078)

[Wiley Online Library](#) | [CAS](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

30 Nissen SP, Kjaer HF, Høst A, Nielsen J, Halken S. The natural course of sensitization and allergic diseases from childhood to adulthood. *Pediatr Allergy Immunol*. 2013; **24**: 549- 555. doi:

[10.1111/pai.12108](https://doi.org/10.1111/pai.12108)

[Wiley Online Library](#) | [PubMed](#) | [Web of Science®](#) | [Google Scholar](#)

31 Uphoff EP, Bird PK, Antó JM, et al. Variations in the prevalence of childhood asthma and wheeze in MeDALL cohorts in Europe. *ERJ Open Res*. 2017; **3**(3):e00150-2016. doi:

[10.1183/23120541.00150-2016](https://doi.org/10.1183/23120541.00150-2016) . <https://pubmed-ncbi-nlm-nih-gov.ezproxy.utu.fi/28845428/>

[Crossref](#) | [Google Scholar](#)

[Download PDF](#)

About Wiley Online Library

Privacy Policy

Terms of Use

Cookies

Accessibility

Publishing Policies

Help & Support

Contact Us

Training and Support

DMCA & Reporting Piracy

Opportunities


Subscription Agents

Advertisers & Corporate Partners

Connect with Wiley

The Wiley Network
Wiley Press Room

Copyright © 1999-2021 John Wiley & Sons, Inc. All rights reserved



1 **Maternal prenatal psychological distress associates with offspring early-life wheezing –**
2 **FinnBrain Birth Cohort**

3

4 Emma Puosi, MD^{1,2,3} Laura S. Korhonen, MD, PhD^{1,2,3} Linnea Karlsson, MD, PhD^{1,3,4} Eeva-Leena Kataja,
5 MSc, PhD^{1,3,5} Heikki Lukkarinen, MD, PhD^{1,2} Hasse Karlsson, MD, PhD^{1,3,4} Minna Lukkarinen, MD,
6 PhD^{1,2,3}

7

8 From ¹FinnBrain Birth Cohort Study, Turku Brain and Mind Center, Department of Clinical Medicine,
9 University of Turku;

10 ²the Department of Pediatrics and Adolescent Medicine, University of Turku, and Turku University
11 Hospital;

12 ³Centre for Population Health Research, University of Turku, and Turku University Hospital;

13 ⁴the Department of Psychiatry, University of Turku, and Turku University Hospital;

14 ⁵the Department of Psychology, University of Turku; all in Turku, Finland.

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

METHODS

The following criteria were used to evaluate the fit including information-theoretic models (*i.e.*, Akaike's Information Criterion, AIC; Bayesians Information Criterion, BIC) where lower AIC and BIC values represent a better fit¹, and the entropy index measure (the entropy value ranges between 0 and 1) where values > 0.80 represent a good fit.² Table S5 presents the resultant fit indices.

In addition, a visual inspection of the plots produced by different models were used to evaluate the meaningfulness and interpretability of the models. Finally, latent groups sizes were also considered when choosing the final models. The numbers of missing values (*i.e.*, thus imputed values) per questionnaire (EPDS, Edinburgh postnatal depression scale and SCL-90, Symptom Checklist 90, anxiety subscale) and per measurement (gestational weeks 14/24/34) point were: EPDS; 81/50/57 and SCL- 90; 81/51/61.

REFERENCES

1. Nylund KL, Asparouhov T, Muthén BO. Deciding on the Number of Classes in Latent Class Analysis and Growth Mixture Modeling: A Monte Carlo Simulation Study. <https://doi.org/10.1080/10705510701575396> [Internet]. 2007 [cited 2021 Sep 15];14:535–69. Available from: <https://www.tandfonline.com/doi/abs/10.1080/10705510701575396>
<https://doi.org/10.1080/10705510701575396>
2. NAGIN D. Group-Based Modeling of Development [Internet]. Cambridge, MA and London, England: Harvard University Press; 2005. Available from: <http://www.degruyter.com/view/books/9780674041318/9780674041318/9780674041318.xml>
<https://doi.org/10.4159/9780674041318>
3. Lubke G, Muthén BO. Performance of Factor Mixture Models as a Function of Model Size, Covariate Effects, and Class-Specific Parameters. <https://doi.org/10.1080/10705510709336735> [Internet]. 2007 [cited 2021 Sep 15];14:26–47. Available from: <https://www.tandfonline.com/doi/abs/10.1080/10705510709336735>
<https://doi.org/10.1080/10705510709336735>
4. Asparouhov T, Muthén B. Using Mplus TECH11 and TECH14 to test the number of latent classes. 2012;

TABLE S1. Population characteristics and attrition with continuous exposures.

Maternal characteristics during	Cohort N = 3837	Included N = 1305	Excluded N = 2532	<i>p</i> *
EPDS point medians (range 0-30)				
gwks 14 (IQR)	4.0 (2.0:7.0) n = 3050	4.0 (2.0:7.0) n = 1234	5.0 (2.0:8.0) n = 1816	0.001
gwks 24 (IQR)	4.0 (2.0:7.0) n = 2770	4.0 (2.0:7.0) n = 1259	4.0 (2.0:8.0) n = 1511	<0.001
gwks 34 (IQR)	4.0 (2.0:7.0) n = 2602	4.0 (1.1:7.0) n = 1260	4.0 (2.0:7.2) n = 1342	0.001
SCL point medians (range 0-40)				
gwks 14 (IQR)	2.0 (0.0:5.0) n = 3052	2.0 (0.0:4.0) n = 1234	2.0 (0.0:5.0) n = 1818	0.16
gwks 24 (IQR)	3.0 (1.0:5.6) n = 2767	2.0 (1.0:5.0) n = 1258	3.0 (1.0:6.0) n = 1509	0.030
gwks 34 (IQR)	2.0 (0.0:5.0) n = 2598	2.0 (0.0:4.0) n = 1256	2.0 (0.0:5.0) n = 1342	0.33

Abbreviations: EPDS, The Edinburgh Postnatal Depression Scale; IQR, Interquartile range; SCL-90, The Symptom Checklist-90, anxiety scale.

*Indicates the group comparisons.

174
175
176

TABLE S2. The risk of wheezing and/or doctor-diagnosed atopic eczema at 24 months by exposure to maternal PD symptoms per the time point of assessment.

Maternal reported exposures during pregnancy	Wheezing ever			Doctor-diagnosed eczema			Wheezing with eczema			Wheezing without eczema			Eczema without wheezing		
<EPDS points as continuous variable, risk increase/point	OR	95% CI	<i>p</i> *	OR	95% CI	<i>p</i> *	OR	95% CI	<i>p</i> **	OR	95% CI	<i>p</i> **	OR	95% CI	<i>p</i> **
gwks 14	1.04	1.0-1.08	0.033	1.01	0.98-1.05	0.52	1.03	0.96-1.11	0.46	1.05	1.01-1.10	0.020	1.02	0.98-1.06	0.33
gwks 24	1.02	0.98-1.06	0.29	1.02	0.99-1.06	0.22	1.01	0.94-1.09	0.30	1.03	0.98-1.07	0.23	1.03	0.99-1.07	0.12
gwks 34	1.04	1.01-1.08	0.020	1.01	0.97-1.04	0.73	1.03	0.96-1.10	0.37	1.05	1.01-1.09	0.019	1.01	0.97-1.05	0.57
<SCL-90 points as continuous variable, risk increase/point															
gwks 14	1.03	1.00-1.07	0.082	1.01	0.97-1.04	0.80	0.98	0.90-1.07	0.69	1.05	1.01-1.09	0.028	1.02	0.98-1.06	0.31
gwks 24	1.05	1.01-1.08	0.006	0.99	0.96-1.03	0.74	1.03	0.96-1.09	0.44	1.05	1.01-1.09	0.011	1.01	0.97-1.04	0.72
gwks 34	1.04	1.00-1.08	0.031	1.02	0.99-1.05	0.25	1.04	0.98-1.11	0.20	1.05	1.01-1.09	0.018	1.02	0.99-1.06	0.21
Adjusted analysis															
<EPDS points as continuous variable, risk increase/point															
gwks 14	1.04	0.98-1.11	0.19	1.01	0.98-1.05	0.52	1.03	0.95-1.10	0.50	1.05	1.01-1.11	0.018	1.02	0.98-1.06	0.29
Maternal asthma	1.3	0.6-2.7	0.46	N/A			N/A								
Maternal atopic diseases	N/A			1.0	1.4-2.4	<0.001									
Maternal asthma and/or atopic diseases	N/A						1.8	0.99-3.3	.055	1.4	0.99-2.0	0.054	1.9	1.4-2.6	<0.001
Male sex vs. female sex	1.2	0.8-2.0	0.38	1.1	0.86-1.5	0.36	3.5	1.7-7.1	<0.001	1.5	1.02-2.1	0.04	1.0	0.74-1.4	0.98
Maternal smoking during pregnancy	5.0	0.59-43	0.14	N/A			0.69	0.19-2.6	0.59	2.2	0.64-7.5	0.22	1.7	0.67-4.2	0.27
Parental education level															
Up to 12 years	2.2	1.2-4.3	0.015	1.4	0.94-2.0	0.10	1.4	0.61-3.0	0.46	1.1	0.64-7.5	0.77	1.4	0.94-2.1	0.10
Between 13-15 years	1.8	1.1-3.1	0.030	1.4	1.0-1.9	0.039	1.3	0.65-2.7	0.45	1.1	0.71-1.6	0.74	1.5	1.0-2.1	0.031
Over 15 years	1.0			1.0			1.0			1.0			1.0		
gwks 24	1.01	0.97-1.05	0.69	1.00	0.97-1.04	0.88	1.02	0.94-1.10	0.67	1.02	0.97-1.07	0.46	1.03	1.00-1.07	0.094
Maternal asthma	2.1	1.3-3.4	0.002	N/A			N/A								
Maternal atopic diseases	N/A			1.9	1.4-2.5	<0.001									
Maternal asthma and/or atopic diseases	N/A						1.8	0.95-3.3	0.073	1.4	1.00-2.2	0.051	1.9	1.4-2.6	<0.001
Male sex vs. female sex	1.9	1.4-2.6	<0.001	1.1	0.83-1.5	0.54	3.6	1.7-7.7	<.001	1.5	1.04-2.2	0.032	0.96	0.71-1.3	0.82
Maternal smoking during pregnancy	0.87	0.38-2.0	0.75	N/A			0.60	0.16-2.3	0.45	1.9	0.55-6.5	0.31	1.6	0.64-4.1	0.31
Parental education level															
Up to 12 years	1.1	0.75-1.7	0.58	1.4	0.94-1.9	0.1	1.5	0.60-3.1	0.45	1.1	0.71-1.8	0.59	1.4	0.93-2.1	0.11
Between 13-15 years	1.1	0.80-1.6	0.48	1.4	0.96-1.9	0.054	1.4	0.66-2.8	0.40	1.2	0.76-1.7	0.5	1.4	1.01-2.1	0.047
Over 15 years	1.0			1.0			1.0			1.0			1.0		
gwks 34	1.05	1.01-1.09	0.016	1.02	0.98-1.10	0.29	1.04	0.97-1.12	0.27	1.06	1.01-1.10	0.014	1.0	0.98-1.1	0.44
Maternal asthma	2.1	1.3-3.3	0.003	N/A			N/A								
Maternal atopic diseases	N/A			1.8	1.4-2.4	<0.001									
Maternal asthma and/or atopic diseases	N/A						1.9	1.0-3.4	0.049	1.3	0.87-1.8	0.22	1.8	1.4-2.5	<0.001
Male sex vs. female sex	1.9	1.4-2.6	<0.001	1.1	0.83-1.5	0.52	3.4	1.7-7.1	<0.001	1.5	1.06-2.2	0.02	0.95	0.70-1.3	0.74
Maternal smoking during pregnancy	0.90	0.39-2.1	0.81	N/A			0.61	0.16-2.3	0.46	1.9	0.55-6.5	0.32	1.5	0.58-3.7	0.41
Parental education level															
Up to 12 years	1.1	0.71-1.6	0.73	1.3	0.87-1.8	0.23	1.2	0.55-2.8	0.61	1.1	0.68-1.8	0.69	1.4	0.89-2.0	0.15

Between 13-15 years	1.1	0.77-1.6	0.61	1.4	0.99-1.9	0.055	1.3	0.65-2.7	0.43	1.1	0.74-1.7	0.61	1.4	1.0-2.1	0.041
Over 15 years	1.0			1.0			1.0			1.0			1.0		

177

178

SCL-90 points as continuous variable, risk increase/point															
gwks 14	1.03	0.99-1.07	0.11	1.0	0.97-1.0	0.92	0.98	0.90-1.07	0.71	1.05	1.01-1.09	0.02	1.02	0.98-1.06	0.34
Maternal asthma	2.2	1.4-3.5	<0.001	N/A			N/A								
Maternal atopic diseases	N/A			1.8	1.4-2.4	<0.001									
Maternal asthma and/or atopic diseases	N/A			N/A			1.8	0.99-3.1	0.054	1.4	0.99-2.0	0.056	1.9	1.4-2.6	<0.0001
Male sex vs. female sex	1.8	1.3-2.5	<0.001	1.1	0.86-1.5	0.35	3.5	1.7-7.1	<0.001	1.5	1.04-2.2	0.3	1.0	0.74-1.4	0.97
Maternal smoking during pregnancy	0.98	0.43-2.2	0.96	N/A			0.64	0.17-2.4	0.52	2.1	0.62-7.3	0.23	1.7	0.66-4.2	0.28
Parental education level															
Up to 12 years	1.1	0.72-1.6	0.74	1.3	0.94-1.9	0.11	1.4	0.62-3.0	0.45	1.1	0.67-1.7	0.79	1.4	0.94-2.1	1.0
Between 13-15 years	1.1	0.74-1.5	0.77	1.4	1.0-1.9	0.042	1.3	0.64-2.7	0.47	1.1	0.71-1.6	0.76	1.5	1.0-2.1	0.032
Over 15 years	1.0			1.0			1.0			1.0			1.0		
gwks 24	1.04	1.00-1.07	0.028	1.00	0.97-1.04	0.89	1.03	0.96-1.1	0.40	1.05	1.01-1.09	0.015	1.01	0.97-1.05	0.59
Maternal asthma	2.1	1.3-3.4	0.002	N/A			N/A								
Maternal atopic diseases	N/A			1.9	1.4-2.5	<0.001									
Maternal asthma and/or atopic diseases	N/A						1.8	0.94-3.3	0.075	1.5	1.05-2.2	0.027	1.9	1.4-2.6	<0.001
Male sex vs. female sex	1.9	1.4-2.6	<0.001	1.1	0.83-1.5	0.54	3.6	1.7-7.8	<0.001	1.5	1.05-2.2	0.027	0.96	0.70-1.3	0.78
Maternal smoking during pregnancy	0.93	0.41-2.1	0.86	N/A			0.62	0.16-2.3	0.47	2.0	0.58-6.9	0.27	1.6	0.62-4.0	0.35
Parental education level															
Up to 12 years	1.1	0.73-1.7	0.64	1.4	0.94-1.9	0.10	1.36	0.60-3.11	0.46	1.1	0.69-1.8	0.64	1.4	0.95-2.1	0.09
Between 13-15 years	1.1	0.80-1.6	0.48	1.4	1.0-1.9	0.054	1.4	0.67-2.9	0.38	1.2	0.77-1.8	0.49	1.5	1.0-2.1	0.04
Over 15 years	1.0			1.0			1.0			1.0			1.0		
gwks 34	1.04	1.01-1.08	0.023	1.02	0.98-1.05	0.61	1.05	0.98-1.13	0.18	1.06	1.05-2.22	0.027	1.0	1.7	0.18
Maternal asthma	2.1	1.3-3.4	0.002	N/A			N/A								
Maternal atopic diseases	N/A			1.8	1.4-2.4	<0.001									
Maternal asthma and/or atopic diseases	N/A						1.9	1.01-3.4	0.046	1.9	0.55-6.5	0.31	1.8	1.3-2.5	<0.001
Male sex vs. female sex	1.9	1.4-2.6	<0.001	1.1	0.83-1.5	0.52	3.4	1.7-7.1	<0.001	1.5	1.05-2.2	0.026	0.97	0.71-1.3	0.82
Maternal smoking during pregnancy	0.89	0.39-2.1	0.79	N/A			0.60	0.16-2.2	0.45	1.9	0.55-6.5	0.31	1.5	0.59-3.8	0.41
Parental education level															
Up to 12 years	1.1	0.73-1.7	0.65	1.3	0.87-1.8	0.23	1.2	0.55-2.8	0.60	1.1	0.69-1.8	0.66	1.3	0.87-2.0	0.19
Between 13-15 years	1.1	0.77-1.6	0.60	1.4	0.99-1.9	0.055	1.3	0.66-2.7	0.43	1.1	0.74-1.7	0.60	1.4	1.0-2.0	0.045
Over 15 years	1.0			1.0			1.0			1.0			1.0		

Abbreviations: BMI, Body mass index; CI, Confidence interval; EPDS, The Edinburgh PostnatalDepression Scale; gwks, Gestational weeks; IQR, Interquartile range; N/A, Not applicable; OR,Odds ratio; P, P- value; SD, Standard deviation; SCL-90, The Symptom Checklist-90, anxiety scale.

*Indicates the group comparisons.

**Indicates the group comparisons with "Neither wheezing nor eczema" as the reference group

179

180

181

TABLE S3. Risk of wheezing and/or doctor-diagnosed atopic eczema at 24 months.

Maternal reported exposures during gestation	Wheezing with eczema			Wheezing without eczema			Eczema without wheezing		
	OR	95% CI	<i>p</i> **	OR	95% CI	<i>p</i> **	OR	95% CI	<i>p</i> **
Unadjusted analyses									
Maternal atopic diseases	1.7	0.97-3.1	0.064	1.4	0.97-2.0	0.078	1.9	1.4-2.6	<0.001
Maternal asthma	2.2	0.93-5.1	0.072	1.8	1.0-3.1	0.045	1.1	0.64-1.9	0.70
Maternal smoking during pregnancy									
* Quit in early pregnancy	1.6	0.55-4.6	0.40	1.0	0.47-2.2	0.94	0.95	0.48-1.9	0.88
* Through pregnancy	2.0	0.57-6.7	0.29	0.42	0.10-1.8	0.23	0.85	0.034-2.1	0.72
Older children with mother-reported atopic diseases	2.0	0.95-4.1	0.070	1.5	0.89-2.4	0.13	2.1	1.4-3.1	<0.001
Older children with mother-reported asthma	1.8	0.40-7.9	0.45	2.7	1.2-6.0	0.018	0.94	0.35-2.5	0.90
Male sex vs. female sex	3.7	1.9-7.2	<0.001	1.5	1.1-2.1	0.023	0.97	0.72-1.3	0.83
Parental education level									
* Up to 12 years	1.5	0.72-3.1	0.28	1.0	0.65-1.6	0.91	1.4	0.92-2.0	0.12
* Between 13-15 years	1.2	0.61-2.4	0.59	1.0	0.68-1.5	0.94	1.4	0.96-1.9	0.09
*Over 15 years	1.0			1.0			1.0		
EPDS trajectories with "Consistently Low" as reference									
"Moderate and Stable"	1.21	0.63-2.34	0.57	1.26	0.83-1.91	0.28	1.29	0.92-1.82	0.14
"Low and Increasing"	1.77	0.40-7.93	0.45	0.97	0.28-3.35	0.97	1.03	0.38-2.79	0.96
"Consistently High"	0.84	0.11-6.47	0.87	3.40	1.75-7.81	0.001	1.36	0.57-3.29	0.49
"High and Decreasing"	0.60	0.08-4.56	0.62	1.54	0.65-3.63	0.32	1.67	0.83-3.36	0.15
SCL-90 trajectories with "Consistently Low" as reference									
"High and Decreasing"	0.65	0.09-4.91	0.67	1.54	0.66-3.63	0.32	2.19	1.14-4.21	0.019
"Moderate and Increasing"	2.57	0.96-6.89	0.061	2.08	1.05-4.14	0.036	1.72	0.92-3.24	0.090
Adjusted analyses									
EPDS trajectories with "Consistently Low" as the reference									
"Moderate and Stable"	1.21	0.60-2.45	0.59	1.17	0.76-1.81	0.49	1.31	0.92-1.88	0.13
"Low and Increasing"	2.72	0.56-13.18	0.21	1.27	0.36-4.48	0.71	0.96	0.31-2.95	0.96
"Consistently High"	0.86	0.11-6.80	0.89	3.60	1.63-7.94	0.002	1.43	0.58-3.52	0.44
"High and Decreasing"	0.64	0.08-4.98	0.67	1.68	0.70-4.00	0.25	1.98	0.97-4.08	0.062
Maternal atopic diseases	N/A								
Maternal asthma									
Maternal asthma and/or atopic diseases	1.8	0.97-3.3	0.063	1.4	0.98-2.0	0.069	1.9	1.4-2.6	<0.001
Male sex vs. female sex	3.5	1.7-7.3	<0.001	1.6	1.0-2.1	0.056	0.97	0.71-1.3	0.83
Maternal smoking during pregnancy	0.70	0.18-2.6	0.60	2.2	0.62-7.6	0.22	1.8	0.71-4.6	0.21

Parental education level									
* Up to 12 years	1.4	0.62-3.1	0.43	1.0	0.65-1.7	0.83	1.4	0.90-2.0	0.14
* Between 13-15 years	1.3	0.65-2.7	0.44	1.1	0.71-1.6	0.76	1.4	1.0-2.1	0.040
*Over 15 years	1.0			1.0			1.0		
SCL-90 trajectories with "Consistently Low" as the reference									
"High and Decreasing"	0.71	0.09-5.46	0.74	1.85	0.77-4.44	0.17	2.5	1.28-5.02	0.008
"Moderate and Increasing"	2.44	0.80-7.34	0.12	2.43	1.21-4.91	0.013	1.7	0.89-3.34	0.11
Maternal atopic diseases	N/A								
Maternal asthma									
Maternal asthma and/or atopic diseases	1.8	0.99-3.1	<0.001	1.4	0.98-2.0	0.063	1.9	1.4-2.5	<0.001
Male vs. female sex	3.5	1.7-7.3	<0.001	1.5	1.0-2.2	0.031	0.98	0.73-1.3	0.98
Maternal smoking during pregnancy	0.70	0.19-2.6	0.60	2.3	0.66-7.8	0.72	1.9	0.73-4.7	0.20
Parental education level									
* Up to 12 years	1.4	0.63-3.1	0.42	1.1	0.68-1.7	0.72	1.4	0.91-2.0	0.14
* Between 13-15 years	1.3	0.64-2.6	0.47	1.1	0.69-1.6	0.83	1.4	1.00-2.0	0.050
*Over 15 years	1.0			1.0			1.0		

BMI, Body mass index; CI, Confidence interval; EPDS, The Edinburgh Postnatal Depression Scale; gwks, Gestational weeks; IQR, Interquartile range; N/A, Not applicable; OR, Odds ratio; SD, Standard deviation; SCL-90, The Symptom Checklist-90, anxiety scale.

*Indicates the comparison with "No wheezing" or "No eczema" as the reference groups.

**Indicates the comparison with "Neither wheezing nor eczema" as the reference group.

198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230

TABLE S4. The correlation of maternal psychological distress sympoms during gestation and at child age of 24 months.

	Maternal distress symptoms at child age 24 months			
Maternal distress symptoms during gestation	EPDS		SCL-90	
EPDS	R	<i>p</i>	<i>R</i>	<i>p</i>
gwks 14	0.43	<0.001	0.36	<0.001
gwks 24	0.43	<0.001	0.37	<0.001
gwks 34	0.46	<0.001	0.39	<0.001
SCL-90				
gwks 14	0.32	<0.001	0.41	<0.001
gwks 24	0.35	<0.001	0.46	<0.001
gwks 34	0.38	<0.001	0.51	<0.001

EPDS, The Edinburgh Postnatal Depression Scale; gwks, Gestational weeks; R, Pearson correlation coefficient; SCL-90, The Symptom Checklist-90, anxiety scale.

The EPDS and SCL-90 were measured and interpreted here as a continuous variables.

TABLE S5. Risk of wheezing and/or doctor-diagnosed eczema at 24 months.

	Wheezing ever			Doctor-diagnosed eczema			Wheezing with eczema			Wheezing without eczema			Eczema without wheezing		
Adjusted analyses	OR	95% CI	<i>p</i> *	OR	95% CI	<i>p</i> *	OR	95% CI	<i>p</i> **	OR	95% CI	<i>p</i> **	OR	95% CI	<i>p</i> **
EPDS trajectories during gestation with "Consistently Low" as the reference															
"Moderate and Stable"	1.0	0.68-1.5	0.98	1.2	0.82-1.6	0.42	0.96	0.46-2.0	0.92	1.1	0.69-1.7	0.71	1.2	0.83-1.8	0.31
"Low and Increasing	1.4	0.49-3.8	0.55	1.0	0.38-2.6	1.0	1.9	0.38-9.7	0.43	1.1	0.31-4.0	0.86	0.83	0.26-2.6	0.75
"Consistently High"	2.3	1.1-4.7	0.030	0.75	0.32-1.8	0.51	0.49	0.058-4.1	0.51	3.0	1.3-7.0	0.011	1.2	0.46-3.0	0.73
"High and Decreasing"	1.0	0.45-2.2	0.99	1.4	0.70-2.7	0.35	0.51	0.065-4.1	0.53	1.6	0.65-3.8	0.32	1.8	0.88-3.8	0.10
Maternal asthma and/or atopic diseases	1.4	1.0-1.8	0.53	1.7	1.3-2.3	<0.001	1.6	0.89-3.0	0.11	1.4	0.97-2.0	0.072	1.9	1.4-2.5	<0.001
Male vs. female sex	1.8	1.3-2.5	<0.001	1.1	0.84-1.5	0.48	3.9	1.9-8.3	<0.001	1.5	1.0-2.1	0.047	0.95	0.70-1.3	0.75
Maternal smoking during pregnancy	1.2	0.51-2.9	0.67	N/A			0.71	0.19-2.7	0.61	2.2	0.63-7.7	0.21	1.8	0.72-4.7	0.20
Parental education level															
Up to 12 years	1.1	0.73-1.6	0.65	1.3	0.91-1.9	0.16	1.5	0.65-3.3	0.36	1.1	0.66-1.7	0.81	1.3	0.89-2.0	0.17
Between 13-15 years	1.1	0.76-1.6	0.65	1.4	1.0-1.9	0.051	1.4	0.67-2.9	0.37	1.1	0.72-1.6	0.67	1.4	0.99-2.0	0.055
Over 15 years	1.0			1.0			1.0			1.0			1.0		
Maternal EPDS score at child age 24 months	1.0	0.99-1.07	0.11	1.04	1.0-1.1	0.043	1.1	1.0-1.2	0.015	1.0	0.98-1.1	0.24	1.0	0.99-1.07	0.21
SCL-90 trajectories during gestation with "Consistently Low" as the reference															
"High and Decreasing"	1.2	0.56-2.4	0.68	1.7	0.92-3.3	0.091	0.61	0.079-4.8	0.64		2.4	1.2-4.8	2.4	1.2-4.8	0.016
"Moderate and Increasing"	1.6	0.86-3.1	0.13	1.3	0.70-2.4	0.43	1.8	0.57-5.7	0.32	2.2	1.0-4.7	0.040	1.5	0.77-3.0	0.23
Maternal asthma and/or atopic diseases	1.4	1.0-1.9	0.044	1.7	1.3-2.3	<0.001	1.7	0.91-3.1	0.098	1.4	0.98-2.0	0.068	1.8	1.3-2.5	<0.001
Male vs. female sex	1.9	1.4-2.6	<0.001	1.1	0.85-1.5	0.43	3.9	1.8-8.2	<0.001	1.5	1.0-2.2	0.032	0.97	0.71-1.3	0.83
Maternal smoking during pregnancy	1.2	0.52-2.9	0.64	N/A			0.72	0.19-2.7	0.63	2.3	0.67-7.9	0.19	1.9	0.74-4.7	0.19
Parental education level															
Up to 12 years	1.1	0.76-1.7	0.55	1.3	0.92-1.9	0.14	1.4	0.64-3.2	0.38	1.1	0.68-1.7	0.74	1.3	0.89-2.0	0.17
Between 13-15 years	1.1	0.76-1.9	0.67	1.4	1.0-1.9	0.052	1.4	0.66-2.8	0.41	1.1	0.71-1.6	0.76	1.4	0.98-2.0	0.067
Over 15 years	1.0			1.0			1.0			1.0			1.0		
Maternal EPDS score at child age 24 months	1.0	1.0-1.07	0.083	1.0	1.0-1.1	0.087	1.1	1.0-1.2	0.027	1.0	0.99-1.1	0.12	1.0	0.99-1.1	0.11

Abbreviations: BMI, Body mass index; CI, Confidence interval; EPDS, The Edinburgh Postnatal Depression Scale; gwks, Gestational weeks; IQR, Interquartile range; N/A, Not applicable; OR, Odds ratio; SD, Standard deviation; SCL-90, The Symptom Checklist-90, anxiety scale.

*Indicates the comparisons with "Wheezing ever" or "Eczema".

**Indicates the comparison with "Neither wheezing nor eczema" as the reference group

TABLE S6. The resultant fit indices of the distress trajectory classes.

	Log L.	AIC	BIC	Entropy	Class Proportions	Average Latent Class Posterior Probabilities
EPDS						
1 Class	-10174.013	20364.026	20405.647	1.000	1.000	1.000
2 Class	-10070.258	20162.516	20219.745	0.789	.82/.18	.96/.85
3 Class	-10013.646	20055.291	20128.129	.84	.03/.72/.26	.88/.95/.87
4 Class	-9972.127	19978.253	20066.698	.84	.04/.02/.71/.23	.80/.89/.94/.82
5 Class	-9925.952	19891.905	19995.958	.84	.23/.68/.03/.04/.02	.83/.93/.83/.75/.91
SCL-90						
1 Class	-9973.333	19962.666	20004.287	1.000	1.000	1.000
2 Class	-9734.885	19491.771	19549.000	.96	.07/.93	.94/.99
3 Class	-9570.181	19168.362	19241.199	.96	.04/.06/.90	.91/.93/.99
4 Class	-9470.686	18975.371	19063.817	.95	.87/.005/.06/.07	.98/.97/.92/.91

AIC, Akaike's Information Criterion; BIC, Bayesian information criteria; EPDS, the Edinburgh Postnatal Depression Scale; SCL-90, the Symptom Checklist-90, anxiety scale.

275
276

277 **FIGURE LEGENDS**

278

279 **FIGURE S1:** Study flow chart. *gwks*, Gestational weeks.