

The Journal of Immunology

This information is current as
of November 6, 2010

**A Cell-Impermeable Cyclosporine A
Derivative Reduces Pathology in a
Mouse Model of Allergic Lung
Inflammation**

Molly A. Balsley, Miroslav Malesevic, Erik J. Stemmy,
Jason Gigley, Rosalyn A. Jurjus, Dallen Herzog, Michael
I. Bukrinsky, Gunter Fischer and Stephanie L. Constant

J. Immunol. published online Nov 5, 2010;
doi:10.4049/jimmunol.1001707

**Supplementary
Data**

[http://www.jimmunol.org/cgi/content/full/jimmunol.1001707/D
C1](http://www.jimmunol.org/cgi/content/full/jimmunol.1001707/D
C1)

Subscriptions

Information about subscribing to *The Journal of Immunology* is
online at <http://www.jimmunol.org/subscriptions/>

Permissions

Submit copyright permission requests at
<http://www.aai.org/ji/copyright.html>

Email Alerts

Receive free email alerts when new articles cite this article. Sign
up at <http://www.jimmunol.org/subscriptions/etoc.shtml>

A Cell-Impermeable Cyclosporine A Derivative Reduces Pathology in a Mouse Model of Allergic Lung Inflammation

Molly A. Balsley,* Miroslav Malesevic,[†] Erik J. Stemmy,* Jason Gigley,*
Rosalyn A. Jurjus,* Dallen Herzog,* Michael I. Bukrinsky,* Gunter Fischer,^{†,1}
and Stephanie L. Constant*,¹

Although the main regulators of leukocyte trafficking are chemokines, another family of chemotactic agents is cyclophilins. Intracellular cyclophilins function as peptidyl-prolyl *cis-trans* isomerases and are targets of the immunosuppressive drug cyclosporine A (CsA). Cyclophilins can also be secreted in response to stress factors, with elevated levels of extracellular cyclophilins detected in several inflammatory diseases. Extracellular cyclophilins are known to have potent chemotactic properties, suggesting that they might contribute to inflammatory responses by recruiting leukocytes into tissues. The objective of the present study was to determine the impact of blocking cyclophilin activity using a cell-impermeable derivative of CsA to specifically target extracellular pools of cyclophilins. In this study, we show that treatment with this compound in a mouse model of allergic lung inflammation demonstrates up to 80% reduction in inflammation, directly inhibits the recruitment of Ag-specific CD4⁺ T cells, and works equally well when delivered at 100-fold lower doses directly to the airways. Our findings suggest that cell-impermeable analogs of CsA can effectively reduce inflammatory responses by targeting leukocyte recruitment mediated by extracellular cyclophilins. Specifically blocking the extracellular functions of cyclophilins may provide an approach for inhibiting the recruitment of one of the principal immune regulators of allergic lung inflammation, Ag-specific CD4⁺ T cells, into inflamed airways and lungs. *The Journal of Immunology*, 2010, 185: 000–000.

Chemokines are critical factors responsible for the recruitment and migration of leukocytes from the circulation into inflamed tissues. Several different members of the CC chemokine family have been described as playing critical roles in allergic lung inflammation and pathology by initiating and promoting the recruitment of eosinophils and T lymphocytes to lung tissues and airways following allergen challenge (1). In an effort to reduce allergic disease pathology, many studies have focused on intervention techniques that specifically target chemokines or chemokine receptors on the cell surface of proinflammatory leukocyte subsets. This approach is made difficult due to the multiple contributing factors released, the redundancy of chemokines, and the promiscuous nature of associations between chemokines and chemokine receptors.

Cyclophilins are ubiquitously distributed intracellular proteins found in all eukaryotic cells. This family of proteins is best known to function in protein folding as peptidyl-prolyl *cis-trans* isomer-

ases (PPIases) (2). However, there is also evidence that cyclophilins function intracellularly in several other capacities, including signal transduction, protein trafficking and chaperoning, apoptosis, and HIV infection (3–5). Cyclophilins can also be secreted (6, 7) and have been demonstrated to function in an extracellular capacity as potent chemoattractants for various leukocyte subsets (8–10), via their PPIase active site (9). High levels of extracellular cyclophilins, notably cyclophilin (Cyp)A and CypB, have been observed in patients with ongoing inflammatory diseases. For example, elevated levels of extracellular cyclophilins have been detected in the synovial fluid of patients with rheumatoid arthritis (11), in tissue from individuals with vascular smooth muscle cell disease (12), and in the serum of patients with severe sepsis (13). Following from these observations, we have previously proposed that the presence of extracellular cyclophilins in inflamed tissues may contribute to inflammatory processes via their capacity to induce leukocyte chemoattraction (14).

CypA is well established as the primary intracellular protein target for the immunosuppressive drug cyclosporine A (CsA) (15), which is widely administered to patients to prevent allograft rejection. When bound to CypA, the CsA/CypA complex acts to inhibit the TCR-dependent activation pathway and production of IL-2 in T lymphocytes. Importantly, CsA has been shown to inhibit the enzymatic activity of many mammalian cyclophilins in the nanomolar range of K_i values (16–18). Therefore, any function of cyclophilins requiring the catalytic site, including chemotaxis, is blocked by CsA. In the present studies we exploited the CsA/CypA interaction and used analogs of CsA to target cyclophilin activity and investigate the potential to reduce inflammatory responses in vivo in a mouse model of allergic lung inflammation. Importantly, we made use of CsA analogs that were nonimmunosuppressive so as not to affect T cell activation. The nonimmunosuppressive cell-permeable analog NIM811 (Melle-4-cyclosporine) differs from CsA by a group replacement at position 4 that prevents NIM811/CypA complexes from binding to and inhibiting calcineurin, although a high-affinity

*Department of Microbiology, Immunology, and Tropical Medicine, The George Washington University Medical Center, Washington, DC 20037; and [†]Max Planck Research Unit for Enzymology and Protein Folding, Halle, Germany

¹S.L.C. and G.F. contributed equally to this work.

Received for publication May 24, 2010. Accepted for publication October 9, 2010.

This work was supported by National Institutes of Health Grant R01-AI67254 (to S.L.C.) and by Deutsche Forschungsgemeinschaft Grant SFB 610 (to G.F.).

Address correspondence and reprint requests to Dr. Stephanie L. Constant, The George Washington University Medical Center, Ross Hall Room 738, 2300 I Street NW, Washington, DC 20037. E-mail address: mtmslc@gwumc.edu

The online version of this article contains supplemental material.

Abbreviations used in this paper: AHR, airway hyperresponsiveness; BAL, bronchoalveolar lavage; C_{dyn} , dynamic lung compliance; CsA, cyclosporine A; Cyp, cyclophilin; GBD, gel band density; i.n., intranasal(ly); PAS, periodic acid-Schiff; PPIase, peptidyl-prolyl *cis-trans* isomerase; R_L , lung resistance; Tg, transgenic.

Copyright © 2010 by The American Association of Immunologists, Inc. 0022-1767/10/\$16.00

interaction with active residues of cyclophilin is maintained (19). Our laboratory has previously made use of NIM811 to investigate the contribution of cyclophilins to leukocyte recruitment in a mouse model of acute lung injury. We demonstrated that intervention with NIM811 significantly reduced the number of neutrophils accumulating in lungs following intranasal delivery of LPS (10). In recent studies we reported the generation of a new cell-impermeable analog of CsA, MM218 (20). This compound contains a highly charged moiety that prohibits its passage through the plasma membrane, thereby preventing it from interacting with intracellular cyclophilins and mediating intracellular immunosuppressive activity. The latter was confirmed by *in vitro* T cell proliferation experiments (20). Compared to NIM811, this compound enabled us to specifically target the extracellular pool of cyclophilins and confirm the contribution of extracellular cyclophilins alone to leukocyte recruitment during inflammatory responses. The present studies demonstrate that this cell-impermeable analog of CsA, MM218, is indeed highly effective at inhibiting leukocyte recruitment, along with other inflammatory parameters, in an experimental model of acute allergic lung inflammation. Such findings suggest these types of analogs might be considered as therapeutic agents for reducing inflammatory diseases associated with leukocyte recruitment into tissues.

Materials and Methods

Mouse model of allergic lung inflammation

All experiments were conducted under the approval of The George Washington University Institutional Animal Care and Use Committee. A previously described (21) mouse model of acute allergic lung inflammation was used for most studies. Briefly, BALB/c mice (purchased from the National Cancer Institute, Bethesda, MD) were primed *i.p.* with 100 μ g OVA protein mixed 1:1 with Injectal alum (Pierce, Rockford, IL) and then challenged with 50 μ g OVA in PBS by intranasal (*i.n.*) delivery for 4 consecutive days (days 7–10). Most analyses were conducted 2 d later (day 12). For experiments in which airway chemokines were measured, the mice were sacrificed 6 h after the third OVA challenge (day 9). For the kinetics studies, groups of mice were sacrificed every other day from day 6 to day 16 after priming.

Inducing allergic lung inflammation by adoptive transfer of OVA-specific CD4⁺ T cells

DO11.10 mice carrying a transgenic (Tg) TCR specific for pOVA_{323–339} were purchased from The Jackson Laboratory (Bar Harbor, ME) and were maintained on BALB/c mice. Female progeny were used as the source of OVA-specific CD4⁺ T cells for adoptive transfer into BALB/c recipients. Populations of OVA-specific Th2 CD4⁺ T cells were generated *in vitro* as described by previous investigators (22) using 5 μ g/ml pOVA_{323–339} (GenScript, Piscataway, NJ), 15 ng/ml IL-4 (PeproTech, Rocky Hill, NJ), 25 U/ml IL-2 (Roche Diagnostics, Indianapolis, IN), and 2.5 μ g/ml anti-IFN- γ (eBioscience, San Diego, CA). After 3 d of culture, 3×10^6 Th2-differentiated CD4⁺ T cells were delivered via tail vein injection to female BALB/c mice, and 24 h after transfer (day 1) the mice were given *i.n.* challenges with 50 μ g OVA for 4 consecutive days. Analyses were conducted 48 h after the final challenge.

In vivo drug intervention

NIM811 CsA (19) was provided by Novartis Institute for Biomedical Research (Cambridge, MA), and MM218 CsA (20) was synthesized at the Max Planck Research Unit for Enzymology of Protein Folding (Halle, Germany). Based on preliminary dosing studies, OVA-primed/challenged mice were given 200- μ g doses of MM218 (0.67 mM in PBS), NIM811 (1.66 mM in diluent), or diluent (16% EtOH in 15% polyoxyethylated castor oil [Cremophor EL; BASF, Mount Olive, NJ]) by *i.p.* injection on days 7, 9, and 11 of the model regimen, and on days 1, 3, and 5 of the transfer regimen.

Tissue isolation and processing

Immediately following sacrifice, blood was collected by cardiac puncture, allowed to clot for several hours at room temperature, and the serum was separated by centrifugation. For the analysis of circulating leukocytes,

blood was collected in heparin and the RBCs were lysed using ammonium chloride buffer. Bronchoalveolar lavage (BAL) was performed after blood collection to isolate airway cells and fluid. The fluid and cells were then separated by centrifugation and used for different analyses. Intact lungs were perfused with 20 ml cold PBS, finely chopped, and crushed through a metal strainer to generate single-cell suspensions. Spleen and lymph node cell suspensions were generated by crushing the individual organs between glass slides followed by filtration through nylon mesh.

Flow cytometric analysis and tissue histology

For flow cytometric analysis, BAL and lung tissue cells were treated with ammonium chloride lysis buffer, counted, and then stained with various combinations of allophycocyanin-anti-GR-1 (neutrophils), FITC-anti-CD8 (CD8⁺ T cells), PE-anti-CD11c (alveolar monocytes), and PE-Cy5-anti-mouse CD4 combined with FITC-anti-mouse CD62L to identify effector/memory CD4⁺ T cells (CD4⁺/CD62L⁺). Cells from the transfer studies were costained with PE-Cy5-anti-mouse CD4 and biotinylated-KJ1-26 followed by FITC-avidin. All Abs were purchased from eBioscience. Eosinophils were identified by forward light scatter/side light scatter distribution and confirmed by cyto-spin analysis. For lung histology, whole lungs were first perfused with PBS and then infused with 1 ml cold periodate-lysine-paraformaldehyde fixative as previously described (21). Fixed lungs were embedded in paraffin and 6- μ m tissue sections were stained with H&E or periodic acid-Schiff (PAS) by Histoserv (Gaithersburg, MD).

Western blot analysis

Equivalent volumes (20 μ l) of BAL fluid from individual mice were separated under reducing conditions using 4–12% Bis-Tris SDS-PAGE gels, followed by blotting and detection using rabbit anti-human CypA (U.S. Biological, Swampscott, MA) and anti-human CypB (ABR, Rockford, IL) Abs as previously described (21). Densitometric analysis was conducted using ImageQuant software.

Measuring cytokines, chemokines, and OVA-specific IgE

To measure airway cytokines, BAL fluids were pooled within treatment groups and then concentrated 8-fold using 3-kDa cut-off Centricon columns (Millipore, Billerica, MA). To test cytokine production by OVA-specific CD4⁺ T cells, suspensions of mononuclear cells were generated from spleens isolated from individual mice. The cells were then cultured at 4×10^6 cells/ml for 4 d with different doses of OVA protein (0–100 μ g/ml) to induce cytokine production. Levels of cytokines in BAL fluid and culture supernatants were measured using mouse IL-5 and IL-13 ELISA kits (eBioscience). Levels of eotaxin/CCL11, TARC/CCL17, and MDC/CCL22 were measured in individual BAL samples using ELISA kits (R&D Systems, Minneapolis, MN). Serum from individual animals was analyzed for the presence of OVA-specific IgE using an ELISA kit purchased from MD Bioproducts (St. Paul, MN).

Measuring airway hyperresponsiveness

On day 12 of the regimen, mice were anesthetized by 300 μ l *i.p.* injection of ketamine (10 mg/ml) and xylazine (1 mg/ml) and a tracheostomy tube was inserted. Measurements of airway resistance and compliance were conducted using the FinePointe RC system (Buxco Research Systems, Wilmington, NC). Mice were sequentially challenged with aerosolized PBS (baseline) followed by increasing doses of methacholine ranging from 1.56 to 25 mg/ml. Maximum resistance and average compliance values were recorded during a 3 min period following each challenge.

Chemotaxis assays

Activated CD4⁺ T cells were generated from total spleen cells isolated from C57BL/6 mice (purchased from National Cancer Institute) by overnight stimulation with 10 μ g/ml Con A. Purified CD4⁺ T cells were set up in 48-well Boyden chemotaxis chambers (Neuroprobe, Gaithersburg, MD) as previously described (21). Positive and negative controls included 1 ng/ml RANTES (PeproTech) and medium alone (RPMI 1640 plus 1% BSA fraction V), respectively. Recombinant cyclophilins were optimized for use at 100 ng/ml for CypA (Calbiochem, San Diego, CA) and 200 ng/ml for CypB (purified in-house). For blocking studies, 2 μ M MM218 was added to the upper and lower wells of the chamber. A chemotactic index was generated by dividing the number of migrated cells in each well by the average number of cells that had migrated to medium alone.

Statistical analysis

Statistically significant differences between groups were determined using the Student *t* test for all studies.

Results

Extracellular cyclophilins are produced throughout effector cell recruitment phase

To determine the optimal timing for targeting cyclophilin activity in our model of allergic lung inflammation, we initially established the kinetics of production of extracellular CypA and CypB, the two most abundant members of the cyclophilin family, relative to the kinetics of leukocyte recruitment within lung airways and tissues. Mice were sacrificed every other day from day 6 (1 d prior to first i.n. OVA challenge) to day 16 (6 d after final i.n. OVA challenge) of our regimen, and airway and lung tissue cells were analyzed by flow cytometry. In previous (21) and present (Supplemental Fig. 1) studies we have observed that this OVA priming/challenge regimen induces an inflammatory response characterized by an influx of monocytes, eosinophils, and CD4⁺ T cells, with minimal recruitment of CD8⁺ T cells and neutrophils. Due to the dominant pathological contribution of eosinophils and CD4⁺ T cells in this model of allergic inflammation (21), our subsequent leukocyte recruitment analyses were focused on monitoring changes in these two effector cell subsets. Fig. 1A shows significant recruitment of both eosinophils and effector/memory CD4⁺ T cells following serial challenge with OVA protein, with peak recruitment between days 12 and 14. When analyzing the BAL fluid from these mice, we observed that levels of both extracellular CypA and CypB increased over the course of leukocyte recruitment (Fig. 1B). While the kinetics of production varied between the two cyclophilins, elevated levels were present throughout the time course of recruitment.

Nonimmunosuppressive CsA analog (NIM811) significantly reduces leukocyte numbers in lung tissue and airways of allergic mice

We next investigated the contribution of cyclophilins in this model by administering nonimmunosuppressive analogs of CsA to inhibit cyclophilin activity. Because extracellular cyclophilins were present throughout the time course of leukocyte recruitment (Fig.

1B), we spanned our intervention regimens throughout the airway challenge phase, with drug treatment given on days 7, 9, and 11 of the regimen. We initially tested a cell-permeable non-immunosuppressive analog of CsA, NIM811 (19), which we had previously shown was highly effective at reducing the influx of neutrophils in a mouse model of LPS-induced acute lung inflammation (10). In the present model of allergic lung inflammation, intervention with NIM811 significantly reduced (29–50%) the number of effector/memory CD4⁺ T cells and eosinophils both in lung tissues and airways (Fig. 2). Importantly, no changes in cell numbers or cell distribution were observed in the blood, spleen, or peripheral lymph nodes of NIM811-treated mice (Supplemental Fig. 2A), ruling out the likelihood of systemic cytotoxic effects. Taken together, these data suggest that leukocyte recruitment to lung tissues and airways can be significantly reduced in experimental acute allergic lung inflammation by blocking the activity of cyclophilins with a nonimmunosuppressive analog of CsA.

Cell-impermeable CsA analog (MM218) also significantly reduces leukocyte numbers in lung tissue and airways of allergic mice

To focus on the contribution of extracellular cyclophilins during allergic lung inflammation, we next tested the capacity of a recently described cell-impermeable analog of CsA, MM218 (20), to reduce leukocyte recruitment in our mouse model. MM218 was found to reduce the number of effector/memory CD4⁺ T cells and eosinophils in lung tissues and airways of allergic mice by a highly significant 75–88% (Fig. 3). This major reduction in inflammatory infiltrates was supported by lung tissue sections stained with H&E to detect inflammatory infiltrates from untreated versus MM218-treated mice (Fig. 4). As with NIM811, analysis of leukocyte numbers in the spleen, peripheral lymph nodes, and blood demonstrated no evidence that treatment with MM218 had induced systemic cytotoxicity (Supplemental Fig. 2B). Further evidence that MM218 does not mediate any significant toxic effects is provided by our recently published studies showing no impact of




FIGURE 1. Extracellular cyclophilins are generated throughout leukocyte recruitment phase of allergic lung inflammation. Groups of OVA-primed/challenged mice were sacrificed every other day starting at day 6 through day 16 of the inflammatory response. *A*, Numbers of effector/memory CD4⁺ T cells and eosinophils were determined in BAL fluid of individual mice by flow cytometric analysis. Data show mean $\pm$ SE cell numbers, with $n = 10$ mice per time point. *B*, Western blot analysis was conducted on cleared fluid from individual BAL samples to determine levels of extracellular CypA and CypB. Data show mean $\pm$ SE gel band density (GBD) for each time point as determined by densitometric analysis. These data are representative of three independent experiments.




FIGURE 2. NIM811 intervention reduces leukocyte numbers in lung tissue and airways of allergic mice. Mice were primed/challenged with OVA, with some groups treated i.p. with NIM811, diluent alone, or untreated, on days 7, 9, and 11 of the regimen. *A*, NIM811 reduces the number of effector/memory CD4⁺ T cells in airway BAL (50%) and lung tissue (38%). *B*, NIM811 reduces the number of eosinophils in airway BAL (48%) and lung tissue (29%). Data show mean $\pm$ SE cell numbers for each group ($n = 12$ mice/group) with a Student *t* test analysis used to determine statistical significance. *** $p < 0.001$; ** $p < 0.01$; * $p < 0.05$, levels of statistical difference comparing OVA plus NIM811 to OVA plus diluent groups. These data are representative of three independent experiments.




FIGURE 3. MM218 intervention significantly reduces leukocyte numbers in lung tissue and airways of allergic mice. Mice were primed/challenged with OVA, with groups treated i.p. with MM218 or PBS on days 7, 9, and 11 of the regimen. Cells were also collected from naive BALB/c mice to serve as a baseline. **A**, MM218 significantly reduces the number of effector/memory CD4⁺ T cells in airway BAL (88%) and lung tissue (75%). **B**, MM218 reduces the number of eosinophils in airway BAL (88%) and lung tissue (79%). Data show mean $\pm$ SE cell numbers for each group ($n = 6$) with a Student *t* test analysis used to determine statistical significance. *** $p < 0.001$; ** $p < 0.01$, levels of statistical difference comparing OVA plus MM218 to OVA plus PBS groups. These data are representative of six independent experiments.

the drug on in vitro T cell activation and proliferation (20). To rule out the possibility that MM218 treatment might be acting by inhibiting the production of other known eosinophil and T cell-attracting chemokines, levels of three airway chemokines associated with leukocyte recruitment in allergic airway inflammation (eotaxin/CCL11, MDC/CCL22, and TARC/CCL17) were compared in BAL fluid from MM218-treated versus untreated mice. Because these chemokines are secreted acutely following airway challenge, measurements were conducted on BAL fluid collected 6 h after OVA challenge. While numbers of airway eosinophils and effector CD4⁺ T cells were reduced by 56–65%,

levels of all three chemokines were comparable between treatment groups, with no significant reductions observed in the MM218-treated mice (Supplemental Fig. 3). The possibility remains that chemokines other than those measured in the present study may have been impacted; however, eotaxin, MDC, and TARC are considered to be major contributors to leukocyte recruitment in allergic responses (1).

Cell-impermeable CsA intervention significantly reduces airway mucus and Th2 cytokine levels

We next investigated whether MM218 would affect other immune parameters of inflammation associated with allergic lung inflammation. Indeed, lung tissue sections stained with PAS to detect airway mucus hypersecretion demonstrated a marked reduction in the frequency and abundance of staining of PAS-positive airways in MM218-treated versus untreated mice (Fig. 4). BAL fluid from the two groups of mice was also assessed for the presence of airway Th2 cytokines, specifically IL-13 and IL-5. Fig. 5A shows that levels of both IL-5 and IL-13 were significantly reduced in the airways of mice treated with MM218 (reductions of 76 and 83%, respectively). These data fit well with the observed reduction in effector/memory CD4⁺ T cells in lung tissues and airways (Fig. 3), since CD4⁺ T cells are known to be a major source of the two cytokines. To rule out the possibility that MM218 treatment might be impacting CD4⁺ T cell priming, thereby reducing the total number of Th2 CD4⁺ T cells generated in vivo, spleen cells from OVA-primed/challenged mice treated, or not treated, with MM218 were isolated and restimulated in vitro with OVA protein to induce cytokine production. As shown in Fig. 5B, culture supernatants from the MM218-treated mice showed no reduction in levels of Th2 cytokines, suggesting that the treatment with MM218 did not adversely affect in vivo T cell priming. Moreover, levels of serum OVA-specific IgE were found to be equivalent between MM218-treated and untreated mice (Fig. 5C), providing additional evidence that Th2 T cell priming was not affected by the drug.

Airway hyperresponsiveness is significantly reduced in allergic mice treated with cell-impermeable CsA

We next investigated whether the observed reductions in leukocyte infiltrates and Th2 cytokines levels mediated by MM218 treatment would also be associated with improved lung function. For these studies airway function was assessed by measuring changes in lung resistance and compliance in response to increasing doses of inhaled methacholine to induce bronchoconstriction. Measurements were conducted on individual mice from groups that were OVA primed/challenged, with and without MM218 treatment, versus naive mice. Measurements of airway resistance and compliance for the three groups are shown in Fig. 6. As expected, airway resistance was significantly increased and lung compliance was decreased in OVA primed/challenged mice, compared with naive mice. Strikingly, mice treated with MM218 showed levels of airway resistance and lung compliance that were almost equivalent to those of naive mice. Collectively, these data demonstrate the capacity of cell-impermeable CsA to reduce several parameters of allergic lung inflammation associated with leukocyte recruitment, including airway hyperresponsiveness (AHR).

Cell-impermeable CsA inhibits migration of activated CD4⁺ T cells in vitro and in vivo

The above findings suggest that extracellular cyclophilins likely contribute to allergic lung inflammation by promoting the recruitment of proinflammatory leukocytes into the respiratory tract and that blocking the function of these cyclophilins using cell-





FIGURE 4. MM218 intervention reduces inflammation in lung tissue of allergic mice. Mice were primed/challenged with OVA, with groups treated i.p. with MM218 or PBS on days 7, 9, and 11 of the regimen. Whole lungs isolated from naive, OVA plus PBS, and OVA plus MM218 mice were paraffin-embedded and 6- μ m sections cut and stained with H&E or PAS. Images (original magnification $\times 10$) show tissue areas surrounding bronchioles. Arrows on H&E sections denote inflammatory foci. Scale bar, 10 μ m. These data are representative of three independent experiments.

FIGURE 5. MM218 intervention reduces Th2 cytokine levels in airways of allergic mice but does not affect Th2 T cell priming. Mice were primed/challenged with OVA, with groups treated i.p. with MM218 or PBS on days 7, 9, and 11 of the regimen. **A**, BAL fluid was collected and pooled for each group, followed by 8-fold concentration. Levels of IL-5 and IL-13 cytokines were measured in the resulting fluid by ELISA. Data show the mean $\pm$ SE for replicate wells ($n = 4$). *** $p < 0.001$, Student t test for both cytokines comparing OVA plus MM218 to OVA plus PBS groups. **B**, Spleens were collected from the same groups of mice and cell suspensions were generated and stimulated in vitro with different doses of OVA protein to induce cytokine production. After 4 d of culture, supernatants were collected and tested for levels of IL-5 and IL-13. Data show the mean $\pm$ SE for replicate wells ($n = 4$). **C**, Serum was collected from individual mice and levels of OVA-specific IgE measured by ELISA. Data shown mean $\pm$ SE titers, with $n = 6$ mice/group. These data are representative of two to three independent experiments.



impermeable CsA impacts this recruitment. In previous studies, we showed that both CypA and CypB have the capacity to induce potent migration of activated CD4⁺ T cells in vitro (21). However, we (21) and others (23) have found that the same cyclophilins are unable to initiate signaling events or to induce migration of eosinophils. These findings have led us to previously suggest that, in vivo, extracellular cyclophilins likely mediate their chemotac-

tic effect directly on effector/memory T cells and indirectly on eosinophils via T cell-derived cytokines (21). To provide supporting evidence that cell-impermeable CsA has the capacity to directly inhibit the migration of CD4⁺ T cells, in vitro chemotaxis assays were conducted in which activated CD4⁺ T cells were stimulated either with CypA or CypB in the presence or absence of MM218. Fig. 7A shows that MM218 was indeed able to completely inhibit the migration of activated CD4⁺ T cells in response to both CypA and CypB. This effect was shown to be cyclophilin-specific, as the presence of MM218 did not alter the response of the same cells to another T cell-attracting chemokine, RANTES (Fig. 7B).

Further supporting evidence that cell-impermeable CsA can impact directly on the recruitment of activated CD4⁺ T cells was provided by in vivo studies using a well-established T cell transfer model of allergic lung inflammation. In this model, OVA-specific activated Th2 CD4⁺ T cells are generated in vitro from DO11.10 TCR Tg mice and then transferred into naive wild-type recipients, followed by intranasal OVA challenge to induce inflammatory responses in the lung (22). This approach enables populations of Ag-specific T cells to be tracked as they migrate into lung tissues and airways and to observe any changes in cell recruitment during drug intervention. Fig. 8 shows a robust recruitment of OVA-specific (KJ1-26⁺) CD4⁺ T cells in response to OVA challenge in recipient mice that received OVA-primed (activated) DO11.10 TCR Tg CD4⁺ T cells, but not in recipient mice that received unprimed (naive) DO11.10 CD4⁺ T cells. Mice treated with cell-impermeable CsA showed a significant reduction in the number of OVA-specific (KJ1-26⁺) CD4⁺ T cells recruited both into lung tissues and airways. Taken together, the results from these studies provide further compelling evidence that cell-impermeable CsA has the capacity to directly inhibit the recruitment of activated CD4⁺ T cells in response to extracellular cyclophilins both in vitro and in vivo.

Cell-impermeable CsA is effective at reducing leukocyte recruitment when administered locally to the airways

There is presently much interest to develop drug therapies and small molecule antagonists that have specific targets and can be administered locally to asthma patients (24). Having demonstrated the efficacy of our cell-impermeable CsA analog to reduce allergic lung inflammation when delivered systemically, we next tested




FIGURE 6. MM218 intervention reduces AHR in allergic mice. Mice were primed/challenged with OVA, with groups treated i.p. with MM218 or PBS on days 7, 9, and 11 of the regimen. On day 12, mice were individually anesthetized with an i.p. injection of ketamine/xylazine, a tracheostomy tube was inserted, and mice were attached to a respirator. The mice were then sequentially challenged for 3 min with increasing doses of methacholine (1.5625–25 mg/ml). Measurements of airway resistance (R_L) and compliance (C_{dyn}) were determined using the Buxco FinePointe RC and associated software. Data show mean $\pm$ SE R_L and C_{dyn} measurements at individual doses of methacholine for each group of mice ($n = 10$ /group). *** $p < 0.001$; ** $p < 0.01$; * $p < 0.05$, Student t test analysis levels of statistical difference at various doses of methacholine between OVA plus MM218 and OVA plus PBS groups. These data are representative of three independent experiments.




FIGURE 7. MM218 inhibits in vitro $CD4^+$ T cell migration in response to extracellular cyclophilins. *A*, Activated $CD4^+$ T cells were generated in vitro and set up in Boyden chambers using recombinant CypA and CypB with or without MM218. RANTES was used as a positive control for T cell migration. Data show mean $\pm$ SE chemotactic indices for each group ($n = 6$ wells/group). $**p < 0.01$; $***p < 0.001$, Student *t* test levels of statistical differences between with MM218 and without MM218 for CypA and CypB stimulation, respectively. *B*, The same assay was used with activated $CD4^+$ T cells set up either with RANTES or CypA with or without MM218. Data show mean $\pm$ SE chemotactic indices for each group ($n = 6$ wells/group). $***p < 0.001$, Student *t* test levels of statistical difference between with MM218 and without MM218 for CypA stimulation only. These data are representative of three independent experiments.

whether it would also be effective when administered directly to the airways. For these studies, OVA-primed/challenged mice were given three doses of either 10 μ g or 2 μ g MM218 by i.n. delivery. As shown in Fig. 9, i.n. administration of MM218 was able to induce a very marked and significant reduction in the number of effector/memory $CD4^+$ T cells and eosinophils detected in the airways of allergic mice. Similar reductions of both leukocyte subsets were observed in lung tissues (data not shown). It is particularly striking that a dose as low as 2 μ g was found to be effective because this dose is 100-fold lower than the optimal dose used for systemic administration. These findings provide promising evidence that cell-impermeable CsA analogs, such as MM218, may have potential for therapeutic use.

Discussion

Cyclophilins, traditionally known as intracellular PPIases, have been shown to be potent chemoattractants for leukocytes. Furthermore, elevated levels of extracellular cyclophilins have been observed in several inflammatory diseases, implicating a role for these proteins in promoting inflammation. In the present study, we used a mouse model of acute allergic lung inflammation characterized by a major influx of eosinophils and activated Th2 $CD4^+$ T cells to investigate the contribution of extracellular cyclophilins to the pathology of allergic lung inflammation. We made use of two different types of nonimmunosuppressive analogs of CsA to block cyclophilin activity.




FIGURE 8. MM218 intervention inhibits in vivo migration of OVA-specific $CD4^+$ T cells to lung tissues and airways of allergic mice. Activated OVA-specific Th2 $CD4^+$ T cells were transferred into BALB/c mice. A separate group of mice received naive OVA-specific $CD4^+$ T cells. All recipient mice were i.n. challenged with OVA for 4 d. Groups of mice ($n = 6$) were treated i.p. with MM218 or PBS during OVA challenge. Data show significant reductions in the number of OVA-specific (KJ1-26 $^+$) effector/memory $CD4^+$ T cells in the airways and lung tissues of MM218-treated mice given activated $CD4^+$ T cells (filled bars). No response was induced in mice given naive $CD4^+$ T cells (open bars). $*p < 0.05$; $***p < 0.001$, Student *t* test levels of statistical difference in the airways and lung tissue, respectively when comparing OVA plus MM218 to OVA plus PBS groups. These data are representative of two independent experiments.

Our findings demonstrate that elevated levels of extracellular CypA and CypB are present throughout the challenge phase of the inflammatory response. Interestingly, the release of CypA and CypB appeared to be differentially regulated since both their initial




FIGURE 9. MM218 is effective when administered directly to the airways. Mice were primed/challenged with OVA, with groups of mice ($n = 6$) given different doses of MM218 by i.n. delivery during challenge. Data show a significant reduction in effector/memory $CD4^+$ T cells and eosinophils in MM218-treated mice. $***p < 0.001$; $**p < 0.01$, Student *t* test levels of statistical difference comparing OVA plus MM218 to OVA plus PBS groups. These data are representative of two independent experiments.

secretion and peak production followed different kinetics profiles. For example, CypA levels showed a marked increase after only a single OVA challenge, whereas little increase was observed in CypB levels until after four challenges. Such findings suggest that the two cyclophilins may regulate different aspects of leukocyte recruitment, with CypA contributing more to initial recruitment and CypB being more important for sustaining leukocyte influx. We were unable to provide direct evidence for this possibility due to the absence of reagents with the capacity to inhibit the function of individual members of the cyclophilin family. However, a new small molecule compound designed to specifically target CypA, and not CypB, was recently described (25). Although this compound is not currently compatible for *in vivo* use due to its poor solubility in aqueous diluents, studies are currently underway to design more soluble versions of such compounds (G. Fischer, unpublished observations). Additionally, we have also initiated studies to develop an immunohistochemical approach that should enable us to investigate any differential kinetics of expression between CypA and CypB, as well as their cell source. Thus, in the present studies our approach was to inhibit the function of all cyclophilins present throughout the leukocyte recruitment phase.

Our results show that targeting the function of cyclophilins, using either a nonimmunosuppressive cell-permeable analog of CsA (NIM811) or a cell-impermeable analog of CsA (MM218), significantly reduces the number of effector/memory CD4⁺ T cells and eosinophils in the airways and lungs of allergic mice. Although the present studies were focused on investigating changes in populations of proinflammatory effector cell subsets, we noted that MM218 treatment also reduced the influx of monocytic cells into airways (Supplemental Fig. 1). This finding is not surprising due to the fact that murine monocytes express CD147, the cell surface receptor for extracellular cyclophilins (9), and they readily migrate in response to cyclophilins (26). Interestingly, the reduction in monocyte numbers was not as marked as that seen in eosinophils and T cells (64% for monocytes versus >80% for the other two subsets). This likely reflects the fact that most resident lung leukocytes are monocytes (macrophages and dendritic cells), and thus fewer are proportionally recruited relative to other subsets of leukocytes. These findings further support the idea that MM218 inhibits active leukocyte recruitment and has little impact on resident leukocytes.

As part of our studies, we also demonstrated that cell-impermeable MM218 CsA has the capacity to inhibit the recruitment of activated CD4⁺ T cells both *in vitro* (using cyclophilin-specific chemotaxis assays) and *in vivo* (using an adoptive cell transfer approach). As previously discussed (21), we think that extracellular cyclophilins likely mediate a chemotactic effect directly on CD4⁺ T cells and indirectly on eosinophils via alterations in Th2 cytokine production. The net result is that inhibiting the function of extracellular cyclophilins affects the recruitment of both leukocyte subsets. The capacity to reduce the influx of effector/memory CD4⁺ T cells into the pulmonary microenvironment is of particular importance in the context of immune-mediated lung inflammation since this population regulates most parameters of allergic lung pathology (27). Indeed, several different studies have shown that inhibition of the cytokines IL-5 and IL-13 alone can lead to reduction in AHR, mucus hypersecretion, IgE generation, as well as eosinophil influx (28–31).

Of particular interest is the finding that treatment with the cell-impermeable MM218 analog was markedly more effective at reducing leukocyte influx (75–88% reduction) compared with the nonimmunosuppressive cell-permeable NIM811 analog (29–50% reduction). Furthermore, the augmented effectiveness of MM218 was achieved using smaller amounts of compound per injection

(0.67 versus 1.66 mM of MM218 and NIM811, respectively). We think the likely explanation is that the cell-impermeable nature of MM218 prevents it from being trapped at intracellular sinks that are known to sequester cell-permeable cyclosporines, such as unmodified CsA and NIM811 (32, 33). This enables the entire amount of MM218 CsA to be available for the inhibition of extracellular cyclophilins. Indeed, studies using fluorescent-tagged CsA have demonstrated that cell-permeable analogs accumulate at very high concentrations within the cell cytosol (20), which would reduce their availability to interact with extracellular ligands. We were particularly pleased to observe that a localized delivery of MM218 was as effective at reducing leukocyte recruitment into allergic lungs as systemic delivery. Moreover, this effect was mediated using 100-fold lower doses of the drug. These findings provide strong supporting evidence that the primary target of our cell-impermeable CsA analog is extracellular cyclophilins generated in the respiratory tract, which has significant positive implications from a therapeutic perspective.

Collectively, our results demonstrate the potent capacity of a cell-impermeable analog of CsA, MM218, to reduce several parameters of acute allergic lung inflammation and provide preliminary evidence that targeting extracellular cyclophilins could have therapeutic potential for reducing tissue inflammation. A CsA derivative that is highly effective at reducing allergic lung inflammation might be questioned in the light of human data reporting only modest improvements following CsA treatment (34). However, several alternative studies have demonstrated CsA to have a potent inhibitory effect on allergic lung inflammation both in animal models (35–37) and in humans (38–40), particularly in cases of severe refractory asthma (41). Regardless of these contradictory findings, it is clear that the side effects associated with high doses and the need for long-term usage reduce the appeal of CsA as a drug of choice (34). We propose that the use of cell-impermeable derivatives of CsA, as described in the present studies, offer a safer and more efficacious approach than unmodified CsA due to their remaining in extracellular spaces, rather than being sequestered within cytoplasmic compartments. By remaining extracellular, cell-impermeable CsA derivatives have no impact on the function of intracellular proteins and will require lower doses for optimal inhibition of extracellular cyclophilin function. An additional advantage of cell-impermeable derivatives is their high solubility in aqueous solutions, making them more manageable for local delivery, for example via inhalation. Interestingly, a study in which CsA was complexed to cyclodextrin to improve its solubility demonstrated an equivalent reduction in murine allergen-induced eosinophilia at 9-fold lower doses than noncomplexed CsA (36), suggesting that CsA functions more efficiently in water-soluble forms. In conclusion, we propose that using drugs and other compounds that remain outside of cells, such as the current cell-impermeable CsA, will enable us to more specifically target extracellular molecules and reduce side effects resulting from uptake by nontarget cells.

Acknowledgments

We thank Dr. Fabrice Allain and Dr. Christophe Vanpouille for providing the necessary reagents to generate recombinant CypB.

Disclosures

The authors have no financial conflicts of interest.

References

1. Velazquez, J. R., and L. M. Teran. 2010. Chemokines and their receptors in the allergic airway inflammatory process. *Clinic. Rev. Allerg. Immunol.* DOI: 10.1007/s12016-010-8202-6.

2. Galat, A. 1993. Peptidylproline *cis-trans*-isomerases: immunophilins. *Eur. J. Biochem.* 216: 689–707.
3. Wang, P., and J. Heitman. 2005. The cyclophilins. *Genome Biol.* 6: 226.
4. Schiene, C., and G. Fischer. 2000. Enzymes that catalyse the restructuring of proteins. *Curr. Opin. Struct. Biol.* 10: 40–45.
5. Bukrinsky, M. I. 2002. Cyclophilins: unexpected messengers in intercellular communications. *Trends Immunol.* 23: 323–325.
6. Sherry, B., N. Yarlett, A. Strupp, and A. Cerami. 1992. Identification of cyclophilin as a proinflammatory secretory product of lipopolysaccharide-activated macrophages. *Proc. Natl. Acad. Sci. USA* 89: 3511–3515.
7. Jin, Z. G., M. G. Melaragno, D. F. Liao, C. Yan, J. Haendeler, Y. A. Suh, J. D. Lambeth, and B. C. Berk. 2000. Cyclophilin A is a secreted growth factor induced by oxidative stress. *Circ. Res.* 87: 789–796.
8. Allain, F., C. Vanpouille, M. Carpentier, M. C. Slomianny, S. Durieux, and G. Spik. 2002. Interaction with glycosaminoglycans is required for cyclophilin B to trigger integrin-mediated adhesion of peripheral blood T lymphocytes to extracellular matrix. *Proc. Natl. Acad. Sci. USA* 99: 2714–2719.
9. Yurchenko, V., G. Zybarth, M. O'Connor, W. W. Dai, G. Franchin, T. Hao, H. Guo, H. C. Hung, B. Toole, P. Gallay, et al. 2002. Active site residues of cyclophilin A are crucial for its signaling activity via CD147. *J. Biol. Chem.* 277: 22959–22965.
10. Arora, K., W. M. Gwinn, M. A. Bower, A. Watson, I. Okwumabua, H. R. MacDonald, M. I. Bukrinsky, and S. L. Constant. 2005. Extracellular cyclophilins contribute to the regulation of inflammatory responses. *J. Immunol.* 175: 517–522.
11. Billich, A., G. Winkler, H. Aschauer, A. Rot, and P. Peichl. 1997. Presence of cyclophilin A in synovial fluids of patients with rheumatoid arthritis. *J. Exp. Med.* 185: 975–980.
12. Suzuki, J., Z.-G. Jin, D. F. Meoli, T. Matoba, and B. C. Berk. 2006. Cyclophilin A is secreted by a vesicular pathway in vascular smooth muscle cells. *Circ. Res.* 98: 811–817.
13. Tegeder, I., A. Schumacher, S. John, H. Geiger, G. Geisslinger, H. Bang, and K. Brune. 1997. Elevated serum cyclophilin levels in patients with severe sepsis. *J. Clin. Immunol.* 17: 380–386.
14. Yurchenko, V., S. Constant, E. Eisenmesser, and M. Bukrinsky. 2010. Cyclophilin-CD147 interactions: a new target for anti-inflammatory therapeutics. *Clin. Exp. Immunol.* 160: 305–317.
15. Handschumacher, R. E., M. W. Harding, J. Rice, R. J. Drugge, and D. W. Speicher. 1984. Cyclophilin: a specific cytosolic binding protein for cyclosporin A. *Science* 226: 544–547.
16. Kallen, J., C. Spitzfaden, M. G. Zurini, G. Wider, H. Widmer, K. Wüthrich, and M. D. Walkinshaw. 1991. Structure of human cyclophilin and its binding site for cyclosporin A determined by X-ray crystallography and NMR spectroscopy. *Nature* 353: 276–279.
17. Ke, H., Y. Zhao, F. Luo, I. Weissman, and J. Friedman. 1993. Crystal structure of murine cyclophilin C complexed with immunosuppressive drug cyclosporin A. *Proc. Natl. Acad. Sci. USA* 90: 11850–11854.
18. Mikol, V., J. Kallen, and M. D. Walkinshaw. 1994. X-ray structure of a cyclophilin B/cyclosporin complex: comparison with cyclophilin A and delineation of its calcineurin-binding domain. *Proc. Natl. Acad. Sci. USA* 91: 5183–5186.
19. Billich, A., F. Hammerschmid, P. Peichl, R. Wenger, G. Zenke, V. Quesniaux, and B. Rosenwirth. 1995. Mode of action of SDZ NIM 811, a non-immunosuppressive cyclosporin A analog with activity against human immunodeficiency virus (HIV) type 1: interference with HIV protein-cyclophilin A interactions. *J. Virol.* 69: 2451–2461.
20. Malešević, M., J. Kühling, F. Erdmann, M. A. Balsley, M. I. Bukrinsky, S. L. Constant, and G. Fischer. 2010. A cyclosporin derivative discriminates between extracellular and intracellular cyclophilins. *Angew. Chem. Int. Ed. Engl.* 49: 213–215.
21. Gwinn, W. M., J. M. Damsker, R. Falahati, I. Okwumabua, A. Kelly-Welch, A. D. Keegan, C. Vanpouille, J. J. Lee, L. A. Dent, D. Leitenberg, et al. 2006. Novel approach to inhibit asthma-mediated lung inflammation using anti-CD147 intervention. *J. Immunol.* 177: 4870–4879.
22. Cohn, L., R. J. Homer, A. Marinov, J. Rankin, and K. Bottomly. 1997. Induction of airway mucus production by T helper 2 (Th2) cells: a critical role for interleukin 4 in cell recruitment but not mucus production. *J. Exp. Med.* 186: 1737–1747.
23. Cormier, S. A., A. G. Taranova, C. Bedient, T. Nguyen, C. Protheroe, R. Pero, D. Dimina, S. I. Ochkur, K. O'Neill, D. Colbert, et al. 2006. Pivotal advance: eosinophil infiltration of solid tumors is an early and persistent inflammatory host response. *J. Leukoc. Biol.* 79: 1131–1139.
24. Holgate, S. T., and R. Polosa. 2008. Treatment strategies for allergy and asthma. *Nat. Rev. Immunol.* 8: 218–230.
25. Daum, S., M. Schumann, S. Mathea, T. Aumüller, M. A. Balsley, S. L. Constant, B. F. de Lacroix, F. Kruska, M. Braun, and C. Schiene-Fischer. 2009. Isoform-specific inhibition of cyclophilins. *Biochemistry* 48: 6268–6277.
26. Damsker, J., I. Okwumabua, T. Pushkarsky, K. Arora, M. I. Bukrinsky, and S. L. Constant. 2009. Targeting the chemotactic function of CD147 reduces collagen-induced arthritis. *Immunology* 126: 55–62.
27. Cohn, L., J. A. Elias, and G. L. Chupp. 2004. Asthma: mechanisms of disease persistence and progression. *Annu. Rev. Immunol.* 22: 789–815.
28. Leckie, M. J. 2003. Anti-interleukin-5 monoclonal antibodies: preclinical and clinical evidence in asthma models. *Am. J. Respir. Med.* 2: 245–259.
29. Kumar, R. K., C. Herbert, D. C. Webb, L. Li, and P. S. Foster. 2004. Effects of anticytokine therapy in a mouse model of chronic asthma. *Am. J. Respir. Crit. Care Med.* 170: 1043–1048.
30. Wills-Karp, M., J. Luyimbazi, X. Xu, B. Schofield, T. Y. Neben, C. L. Karp, and D. D. Donaldson. 1998. Interleukin-13: central mediator of allergic asthma. *Science* 282: 2258–2261.
31. Walter, D. M., J. J. McIntire, G. Berry, A. N. J. McKenzie, D. D. Donaldson, R. H. DeKruyff, and D. T. Umetsu. 2001. Critical role for IL-13 in the development of allergen-induced airway hyperreactivity. *J. Immunol.* 167: 4668–4675.
32. Shibata, N., H. Shimakawa, T. Minouchi, and A. Yamaji. 1993. Pharmacokinetics of cyclosporin A after intravenous administration to rats in various disease states. *Biol. Pharm. Bull.* 16: 1130–1135.
33. Ryffel, B., B. M. Foxwell, A. Gee, B. Greiner, G. Woerly, and M. J. Mihatsch. 1988. Cyclosporine: relationship of side effects to mode of action. *Transplantation* 46(Suppl.): 90S–96S.
34. Evans, D. J., P. Cullinan, and D. M. Geddes. 2001. Cyclosporin as an oral corticosteroid sparing agent in stable asthma. *Cochrane Database Syst. Rev.* 2: CD002993.
35. Yeum, H. S., Y. C. Lee, S. H. Kim, S. S. Roh, J. C. Lee, and Y. B. Seo. 2007. *Fritillaria cirrhosa*, *Anemarrhena asphodeloides*, Lee-Mo-Tang and cyclosporine A inhibit ovalbumin-induced eosinophil accumulation and Th2-mediated bronchial hyperresponsiveness in a murine model of asthma. *Basic Clin. Pharmacol. Toxicol.* 100: 205–213.
36. Fukaya, H., A. Iimura, K. Hoshiko, T. Fuyumuro, S. Noji, and T. Nabeshima. 2003. A cyclosporin A/maltosyl- α -cyclodextrin complex for inhalation therapy of asthma. *Eur. Respir. J.* 22: 213–219.
37. Roh, S. S., S. H. Kim, Y. C. Lee, and Y. B. Seo. 2008. Effects of Radix Adenophorae and cyclosporine A on an OVA-induced murine model of asthma by suppressing T cells activity, eosinophilia, and bronchial hyperresponsiveness. *Mediators Inflamm.* 2008: 781425.
38. Khan, L. N., O. M. Kon, A. J. Macfarlane, Q. Meng, S. Ying, N. C. Barnes, and A. B. Kay. 2000. Attenuation of the allergen-induced late asthmatic reaction by cyclosporin A is associated with inhibition of bronchial eosinophils, interleukin-5, granulocyte macrophage colony-stimulating factor, and eotaxin. *Am. J. Respir. Crit. Care Med.* 162: 1377–1382.
39. Sihra, B. S., O. M. Kon, S. R. Durham, S. Walker, N. C. Barnes, and A. B. Kay. 1997. Effect of cyclosporin A on the allergen-induced late asthmatic reaction. *Thorax* 52: 447–452.
40. Fukuda, T., J. Asakawa, S. Motojima, and S. Makino. 1995. Cyclosporine A reduces T lymphocyte activity and improves airway hyperresponsiveness in corticosteroid-dependent chronic severe asthma. *Ann. Allergy Asthma Immunol.* 75: 65–72.
41. Polosa, R., and J. Morjaria. 2008. Immunomodulatory and biologic therapies for severe refractory asthma. *Respir. Med.* 102: 1499–1510.




Fig. S1. Distribution of leukocytes in airways of allergic mice after MM218 treatment. BAL fluid was collected from individual OVA primed/challenged mice treated i.p. with MM218 or PBS on days 7, 9 and 11 of the regimen. Leukocytes were analyzed by flow cytometry to establish the relative distribution of neutrophils, CD8⁺ T cells, monocytes, eosinophils and CD4⁺ T cells for the two groups, compared to airway cells from naïve mice. Data show mean cell numbers for each group (n=6 per group) $\pm$ SE with a Student's t-test analysis used to determine statistically significant differences between OVA+MM218 and OVA+PBS groups. *** = $P < 0.001$; * = $P < 0.05$.




Fig. S2. Mice treated with either NIM811 or MM218 do not exhibit systemic leukopenia. Blood, spleen and mediastinal LN were collected from individual OVA primed/challenged mice treated i.p. (A) with NIM811 or PBS, or (B) with MM218 or PBS, on days 7, 9 and 11 of the regimen. Total leukocyte numbers were counted by hemacytometry. Data show mean $\pm$ SE cell numbers for each group (n=6 mice per group).




Fig. S3. Levels of airway eotaxin, MDC and TARC chemokines are not reduced by MM218 treatment. Mice were primed i.p. with OVA/Alum (day 0) and then given i.n. OVA challenges on days 7, 8 and 9. Groups of mice were treated i.p. with MM218 or PBS on days 7 and 9. BAL fluid was collected from individual animals 6 h after the third OVA challenge (day 9) and levels of the chemokines eotaxin, MDC and TARC were measured by ELISA. BAL fluid from naïve mice was included for comparison. Data show mean $\pm$ SE levels of each chemokine for the three groups (n=6-8 mice per group). These data were generated from a combination of two independent experiments.