

A substantial neutrophilic inflammation as regular part of severe type 2 chronic rhinosinusitis with nasal polyps

Tim Delemarre, MSc,^a Gabriele Holtappels, BSc,^a Natalie De Ruyck, MSc,^a Nan Zhang, MD, PhD,^a Hans Nauwynck, DVM, PhD,^b Claus Bachert, MD, PhD,^{a,c,*} and Elien Gevaert, PhD^{a,*} *Ghent, Belgium; and Stockholm, Sweden*

Background: Chronic rhinosinusitis with nasal polyps (CRSwNP) is generally associated with severe type 2 immune reactions in the white population. However, recent findings suggest an additional role for neutrophils in severe type 2 inflammation.

Objective: This study aimed to characterize the neutrophilic inflammation in CRSwNP and its relation to eosinophilic inflammation in severe type 2 immune reactions.

Methods: The presence and activation of neutrophils and eosinophils was analyzed in CRS without NP and CRSwNP by measuring cell and activation markers via immunohistochemistry, immunofluorescence, Luminex assay, ELISA, UniCAP, fluorescence-activated cell sorting, and PCR. Differential neutrophil migration was assessed via Boyden-chamber assay and neutrophil survival was analyzed via flow cytometry.

Results: Both CRS without NP and CRSwNP displayed variable degrees of eosinophilic and neutrophilic inflammation, with a profound neutrophilic infiltration and activation in type 2 CRSwNP, associated with eosinophil extracellular traps cell death and Charcot-Leyden crystals, but independent of IL-17. Neutrophil extracellular traps cell death in CRSwNP was associated with bacterial colonization, however, neutrophils were less prone to undergo neutrophil extracellular traps cell death in the tissue of patients with severe type 2 CRSwNP. Neutrophils did not show increased migration nor survival in the CRSwNP environment *in vitro*.

Conclusions: This study demonstrated a severe neutrophilic inflammation associated with severe eosinophilic type 2 inflammatory CRSwNP, the role of which needs further study. (J Allergy Clin Immunol 2020;■■■:■■■-■■■.)

Key words: CRSsNP, CRSwNP, type 2 inflammation, neutrophils, NETosis, IL-17, Charcot-Leyden crystals

Abbreviations used

CitH3: Citrullinated histone 3
CLC: Charcot-Leyden crystal
CRS: Chronic rhinosinusitis
CRSsNP: Chronic rhinosinusitis without nasal polyps
CRSwNP: Chronic rhinosinusitis with nasal polyps
EETosis: Eosinophil extracellular traps cell death
Gal10: Galectin-10
MBP: Major basic protein
NETosis: Neutrophil extracellular traps cell death

Chronic rhinosinusitis (CRS) is a chronic inflammation of the sinonasal mucosa and paranasal sinuses that has a substantial effect on quality of life and daily functioning of the patients. It is an increasing upper airway health problem that currently affects between 6% and 15% of the white population.^{1,2} CRS is clinically subdivided in CRS with nasal polyps (CRSwNP) and CRS without nasal polyps (CRSsNP). In the literature, CRSwNP is generally the more severe phenotype and is typically characterized by a type 2 eosinophilic inflammation, while CRSsNP is considered a type 1 neutrophilic inflammation.³ While this traditional type 2–type 1 classification in CRS is still valid in general, recent reports in asthma suggest that the role of neutrophils in type 2 airway inflammation might be far more important than was initially thought.^{4–6}

Therapies for CRSwNP nowadays mainly focus on targeting type 2 inflammation and considerable progress has been made in endotyping and treatment.^{7–10} Typically, patients with CRSwNP frequently suffer from recurrence after surgery or need repeated oral corticosteroid courses with potential long-term risks.¹¹ Non-responsiveness to oral corticosteroids may partially be caused by a coexisting neutrophilic inflammation, as described in both asthma and CRSwNP.^{12,13}

In asthma, the presentation of a mixed eosinophilic-neutrophilic inflammation, associated with a mixed type 2–type 17 inflammation, is linked to a more severe and harder to control phenotype.⁴ A recent cluster analysis from our group showed that clusters with the most severe clinical phenotype were characterized by a severe type 2 inflammation, as well as by elevated levels of neutrophilic marker proteins.¹⁴ On the other hand, the cluster analysis showed a group of patients with CRSwNP with only neutrophil markers elevated.¹⁴ During neutrophil extracellular traps cell death (NETosis) and eosinophil extracellular traps cell death (EETosis), cytosol and granule proteins are mixed with condensed DNA and released in the extracellular space, where they can affect the local inflammation.¹⁵ Recent studies described

From ^athe Upper Airways Research Laboratory, Faculty of Medicine, and ^bthe Laboratory of Virology, Faculty of Veterinary Medicine, Ghent University; Ghent University, Ghent; and ^cthe Division of Ear, Nose and Throat Diseases, Department of Clinical Science, Intervention and Technology, Karolinska Institute, Stockholm.

*These authors shared project supervision.

E.G. was supported by a post-doctoral research fellowship from the Fonds Wetenschappelijk Onderzoek (FWO) (FWO/DPO/108). C.B. and E.G. were supported by grants from FWO Flanders (1515516N, EOS GOG2318N), the Interuniversity Attraction Poles Grant P7/30, and Sanofi (A17/TT/1942, A19/TT/0828).

Disclosure of potential conflict of interest: The authors declare that they have no relevant conflicts of interest.

Received for publication June 16, 2020; revised August 11, 2020; accepted for publication August 25, 2020.

Corresponding author: Claus Bachert, MD, PhD, Upper Airways Research Laboratory, Department Head and Skin, Ghent University, UZ Ghent, C. Heymanslaan 10, 9000 Ghent, Belgium. E-mail: Claus.Bachert@ugent.be.

0091-6749/\$36.00

© 2020 American Academy of Allergy, Asthma & Immunology

<https://doi.org/10.1016/j.jaci.2020.08.036>

EETosis in CRSwNP tissue, associated with Charcot-Leyden crystal (CLC) deposition; CLCs were found to be responsible for the induction and maintenance of neutrophilic inflammation in CRSwNP.¹⁵⁻¹⁸ We therefore sought to further characterize the driving forces of neutrophilic inflammation and its relationship to eosinophilic inflammation in severe type 2 immune reactions.

METHODS

Sample collection

Tissue and blood samples were collected from patients of the Department of Otorhinolaryngology at Ghent University Hospital (Belgium) after receiving written informed consent for inclusion. Nasal polyps and/or sinus mucosal tissue samples from patients undergoing functional endoscopic sinus surgery for CRSwNP ($n = 56$) or CRSsNP ($n = 35$) and inferior turbinate from healthy patients (undergoing surgery for anatomical obstruction) ($n = 27$) were collected at the Ghent University Hospital between February 2012 and June 2018. Tissue samples were either snap-frozen or fixed in paraformaldehyde and embedded in paraffin. For the isolation of neutrophils from peripheral blood, 100 mL blood was obtained from healthy volunteers and processed within 15 minutes after collection. None of the patients took oral or intranasal glucocorticoids, nor antibiotics within 4 weeks before surgery. Clinical data and symptom surveys were collected from all patients that were enrolled in the study (Table I). The study was approved by the local Ethics Committee of Ghent University Hospital.

Staining

Immunohistochemistry. To prepare tissue slides for immunohistochemistry and immunofluorescence, nasal polyp and control tissues were fixed in 4% paraformaldehyde and embedded in paraffin. These 4 μ m thick slides were first deparaffinized in xylene and rehydrated in decreasing concentrations of ethanol. Slides were blocked with 7.5% BSA in PBS and incubated with primary antibodies (monoclonal mouse anti-human major basic protein, BMK13 [MONOSAN, Sanbio, Uden, The Netherlands]; monoclonal mouse anti-human neutrophilic elastase, NP57 [Dako, Thermo Fisher Scientific, Waltham, Mass]; polyclonal rabbit anti-human histone H3 [Abcam, Cambridge, United Kingdom]) for 16 hours at 4°C. The primary antibodies were visualized via an alkaline phosphatase-linked secondary antibody kit according to manufacturer's instructions (Dako). Finally, the slides were counterstained with hematoxylin for 1 minute and mounted with Aqueatex mounting medium (Merck, Kenilworth, NJ). Isotype controls were included for all samples. Total number of positive cells were counted and expressed as number of cells/mm² tissue.

Immunofluorescence. Paraffin slides were prepared and deparaffinized as mentioned before. After antigen retrieval (EET-, citrullinated histone 3 [CitH3]-, and galectin-10 [Gal10]-staining), slides were blocked with 7.5% BSA in PBS and incubated with primary antibodies (monoclonal mouse anti-human major basic protein, BMK13 [MONOSAN]; monoclonal mouse anti-human neutrophilic elastase, NP57 [Dako]; polyclonal rabbit anti-human histone H3 [Abcam]; monoclonal mouse anti-human galectin-10, 561603 [R&D Systems, Minneapolis, Minn]) for 16 hours at 4°C. The primary antibodies were visualized via fluorescein isothiocyanate-conjugated polyclonal goat anti-mouse or anti-rabbit IgG secondary antibody (Thermo Fisher Scientific). Slides were mounted with 4'-6-diamidino-2-phenylindole, dihydrochloride-containing mounting medium for nuclear counterstain. The slides were analyzed with a confocal laser-scanning microscope (Leica Microsystems, Wetzlar, Germany). For each patient, NETs, EETs and CLCs were counted in 10 random selected fields throughout the tissue. The number of cells involved in NET or EET formation were counted and normalized to the number of neutrophils or eosinophils, respectively, and expressed as percentage of NET- or EET-generating neutrophils or eosinophils.

Cytokine and protein measurements

Snap frozen tissues were weighed, homogenized, and centrifuged as described previously.¹⁹ The samples were assayed for IL-5, IL-6, IL-8, IL-17, myeloperoxidase, and TNF- α using commercially available Luminex kits from R&D, and IFN- γ using a commercially available Quantikine ELISA from R&D. Levels were measured on a Bio-Plex 200 Array Reader (Bio-Rad, Hercules, Calif). Eosinophilic cationic protein, IgE, and *Staphylococcus aureus* enterotoxin-specific IgE (staphylococcal enterotoxin A, staphylococcal enterotoxin C, and toxic shock syndrome toxin-1) were measured using the UniCAP method (Thermo Fisher Scientific, Phadia AB, Uppsala, Sweden) according to the manufacturer's instructions. Concentrations below detection limit were considered negative and were given a value one-half of the detection limit.

Quantitative PCR

RNA was extracted from snap-frozen tissue samples as described previously.¹⁵ In summary, RNA was isolated with the RNeasy Mini Kit (QIAGEN, Hilden, Germany), and subsequently cDNA was synthesized using the iScript Advanced cDNA Synthesis Kit (Bio-Rad). Quantitative real-time PCR was used to quantify mRNA levels of Gal10. The calibrated normalized relative quantities were calculated with the qBase+ software (Biogazelle, Zwijnaarde, Belgium). The primers were commercially purchased from Bio-Rad.

Flow cytometry analysis

The fraction of activated neutrophils in tissue of patients with CRSsNP and those with CRSwNP was analyzed via flow cytometry, based on reduced expression of CD62L. Single-cell suspension was prepared as described before.²⁰ Cells were labeled with near-infrared live-dead dye (Invitrogen, Thermo Fisher Scientific), PE-Cy7 anti-CD45 (BioLegend, San Diego, Calif), PerCP-Cy5.5 anti-CD11b (BioLegend), Pacific blue anti-CD14 (BioLegend), PE anti-CD15 (BioLegend), fluorescein isothiocyanate anti-CD16 (BD Pharmingen, BD, Franklin Lakes, NJ), and antigen presenting cell anti-CD62L (BioLegend). Neutrophils were gated by subsequent gating for viable cells (live-dead staining), single cells (forward scatter area/forward scatter height), leukocytes (CD45⁺), myeloid cells (CD11b⁺), granulocytes (CD14^{dim}), and neutrophils (CD16⁺/CD15⁺), based on fluorescence minus one controls, included for each marker. Activated neutrophils are expressed as percentage of CD62L⁻ neutrophils.

Neutrophil protease activity assays

Neutrophil elastase and cathepsin G activity were analyzed in tissues of controls and patients with CRSsNP and those with CRSwNP. Homogenates were obtained from frozen tissue as described before¹⁹ and dissolved in PBS + Pen-Strep. Fluorometric neutrophil elastase activity assay kit (Sigma-Aldrich, St Louis, Mo) and colorimetric cathepsin G activity assay kit (Abcam) were used as instructed by the manufacturers. Duplicates and background controls were included for all samples in both assays.

Boyden chamber assay

Peripheral blood neutrophils were collected from whole blood of healthy volunteers as described before.¹⁶ Tissue homogenates of controls and patients with CRSsNP and those with CRSwNP were obtained as described above and dissolved in RPMI + Pen-Strep. Lower compartments were filled with RPMI + 2% FCS + Pen-Strep (tissue culture medium) as negative control or 100 μ g/mL tissue homogenate of healthy controls or patients with CRSsNP or CRSwNP. Subsequently, for each condition, 10⁵ neutrophils were primed for 20 minutes with 100 ng/mL GM-CSF (PeproTech, Cranbury, NJ) and then allowed to migrate to the lower compartment through 5- μ m pore size filters (VWR International, Secaucus, NJ) for 90 minutes at 37°C. The migrated neutrophils were collected, stained with a May Grunwald Giemsa stain and quantified for each condition. Two independent experiments were performed

TABLE I. Patient characterization

	Controls	CRSsNP	CRSwNP
Total cases	27	35	56
Sex			
Female	7	12	19
Male	20	23	37
Age (y)	30 (20-58)	44 (18-67)	46 (19-92)
Ethnicity	White	White	White
Previous surgeries			
Yes	3	12	33
No	27	23	23
No. of surgeries	0-1	0-3	0-4
Allergy			
Present	12	18	33
Not present	14	17	22
Missing cases	1	—	1
Asthma			
Present	4	4	28
Not present	23	31	27
Missing cases	—	—	1
Tissue concentrations			
IgE (U/g)	66.56 ± 76.81	578.46 ± 1688.78	932.26 ± 1329.99
SE-IgE (UA/g)	1.07 ± 2.72	0.18 ± 2.27	4.16 ± 5.43
IL-5 (pg/g)	4.84 ± 10.21	89.72 ± 122.83	537.32 ± 680.13
ECP (μg/g)	0.46 ± 0.67	5.67 ± 5.41	41.29 ± 123.88
IL-17 (pg/g)	52.27 ± 81.37	55.83 ± 89.83	42.59 ± 89.58
TNF-α (pg/g)	7.99 ± 15.51	18.65 ± 23.94	22.99 ± 23.88
MPO (μg/g)	103.43 ± 54.16	344.85 ± 270.22	559.48 ± 379.88

ECP, Eosinophilic cationic protein; SE, *Staphylococcus aureus* enterotoxin-specific. Allergy was defined as present when the patient had a positive skin prick test for ≥1 of the inhalant allergens commonly tested in our region.

Values are n, median (range), or mean ± SD.

and results were expressed as migration index compared with tissue culture medium.

Neutrophil survival assay

Peripheral blood neutrophils were isolated and tissue homogenates were prepared as described before. We incubated 10^6 neutrophils in 100 μg/mL tissue homogenate of healthy controls or patients with CRSsNP or CRSwNP for 8 hours at 37°C. After incubation, apoptotic neutrophils were stained using an Annexin V apoptosis detection kit (BD Pharmingen) and analyzed using FACS Canto II with FACS Diva software (BD Bioscience, San Jose, Calif).

Statistical analysis

Statistical analysis was performed using the Prism GraphPad version 8 software program (GraphPad Software, San Diego, Calif). Mann-Whitney *U* test was used to evaluate statistical differences between 2 groups. Kruskal-Wallis test followed by a multiple comparison test was used to evaluate statistical differences among multiple groups. Correlations were analyzed with a nonparametric Spearman correlation test. Significances were expressed as **P* < .05, ***P* < .01, ****P* < .001, and *****P* < .0001. *P* values ≤ .05 were considered as statistically significant.

RESULTS

Severe neutrophilic infiltration and inflammation in CRSwNP versus CRSsNP

In line with previous reports, we found that CRSwNP is clearly associated with type 2 inflammation, while CRSsNP is generally more related with type 1 inflammation. Tissue levels of IL-5 were significantly increased in patients with CRSwNP compared with in those with CRSsNP (*P* < .0001) and in healthy controls (*P*

<.001) (see Fig E1, A, in this article's Online Repository at www.jacionline.org), and tissue levels of IFN-γ were significantly (*P* < .05) increased in patients with CRSsNP compared with in those with CRSwNP and in healthy controls (Fig E1, B). The characterization of the studied patients is presented in Table I.

However, characterization of the tissue in terms of eosinophilic or neutrophilic infiltration showed a more versatile image than was anticipated based on cytokine profiles in both patients with CRSwNP and those with CRSsNP. Both patient groups contained cases with low granulocyte infiltration, predominantly neutrophilic, predominantly eosinophilic, and mixed neutrophilic-eosinophilic infiltration (Fig 1, A). Interestingly, most of the patients with severe type 2 CRSwNP had a mixed neutrophilic-eosinophilic inflammation.

Based on these observations in the tissue, we quantified the number of eosinophils and neutrophils in both patient groups. Numbers of eosinophils were significantly elevated in patients with CRSwNP compared with in those with CRSsNP (*P* < .0001) and in healthy controls (*P* < .001) (see Fig E2, A, in this article's Online Repository at www.jacionline.org). In line with this, we found that tissue eosinophilic cationic protein levels were significantly increased in patients with CRSwNP compared with in those with CRSsNP (*P* < .01) and in healthy controls (*P* < .0001) (Fig E2, B). Numbers of neutrophils were significantly elevated in patients with CRSwNP compared with in those with CRSsNP (*P* < .05) and in healthy controls (*P* < .01) (Fig 1, B). In line with this, we found tissue levels of myeloperoxidase were significantly (*P* < .01) elevated in patients with CRSwNP compared with in controls (Fig 1, C), and IL-8 and IL-6 were




FIG 1. Differential neutrophilic and eosinophilic infiltration in CRSsNP and CRSwNP, with increased neutrophilic inflammation in CRSwNP. **A**, Inflammatory subgroups of CRSwNP and CRSsNP based on the mean neutrophil counts (respectively, 65 and 28 cells/mm²) and the mean eosinophil counts (respectively, 811 and 16 cells/mm²). **B**, Quantification of neutrophils per mm² in tissue of healthy controls and patients with CRSsNP and those with CRSwNP. **C-F**, Myeloperoxidase (MPO), IL-8, IL-6, and IL-17 protein levels in the tissue of healthy controls and patients with CRSsNP and those with CRSwNP, measured via UniCAP and ELISA. **G**, Ratio of neutrophils in tissue of patients with CRSsNP and CRSwNP with high and low protein levels of IL-17, based on the median tissue concentration of IL-17 in both groups (respectively, 27.28 and 15.02 pg/mL). Levels of statistical significance are expressed as * $P < .05$ and ** $P < .01$.

significantly increased in patients with CRSwNP compared with in those with CRSsNP (respectively, $P < .05$ and $P < .01$) and in healthy controls ($P < .01$ and $P < .05$) (Fig 1, D and E). These findings showed a profound neutrophilic inflammation, coexisting with eosinophilic inflammation in patients with severe type 2 CRSwNP.

Interestingly, no intergroup differences were observed in tissue levels of IL-17, a major driving force for the recruitment, survival, and activation of neutrophils (Fig 1, F). To investigate the possible involvement of a type 17 inflammation, IL-17 levels were related to the number of neutrophils in the tissue. Interestingly, while the number of neutrophils was significantly ($P < .05$) elevated in the patients with CRSsNP with high IL-17 levels compared with the group with low IL-17, we found no significant increase in ratio of neutrophils in patients with CRSwNP with high IL-17 compared with in the patients with CRSwNP with low IL-17 levels (Fig 1, G).

Neutrophil infiltration associated with type 2 inflammation in CRSwNP

As was recently reported, the presence of CLCs—the crystallized form of Gal10 protein—results from the process of EETosis

and is a hallmark of type 2 inflammation in airway immunity.^{17,18} Therefore, Gal10 mRNA expression levels were measured and found to be significantly upregulated in patients with CRSwNP compared with in healthy controls ($P < .0001$) and in patients with CRSsNP ($P < .05$). In addition, Gal10 mRNA expression was also significantly ($P < .01$) upregulated in patients with CRSsNP compared with in healthy controls (Fig 2, A). No significant influence of comorbid asthma or allergy on Gal10 mRNA levels was found in the patient groups (data not shown). The numbers of CLCs were quantified and found to be significantly ($P < .01$) upregulated in patients with CRSwNP compared with in healthy controls (Fig 2, B). Interestingly, the number of neutrophils correlated positively with the relative fractions of EETs ($P < .001$; $r = 0.8047$) (Fig 2, C) and the numbers of CLCs ($P < .01$; $r = 0.6504$) (Fig 2, D) in patients with type 2 CRSwNP, but not in patients with CRSsNP and in controls.

Increased neutrophil activation in type 2 CRSwNP

Because both neutrophils and its activating cytokines are increased in CRSwNP, we analyzed and compared neutrophil activity in both CRSsNP and CRSwNP tissue. Neutrophils are known to shed CD62L on activation. Therefore, we studied the




FIG 2. Elevated CLC deposition in CRSwNP associated with neutrophilic infiltration. **A**, Relative Gal10 gene expression in the tissue of healthy controls and patients with CRSsNP and those with CRSwNP, expressed as calibrated normalized relative quantity (CNRQ) values. **B**, Quantification of CLCs per high power field (hpf) in the tissue of healthy controls and patients with CRSsNP and those with CRSwNP. Scatter plots of numbers of neutrophils per mm² with ratios of EETs (**C**) and numbers of CLCs per mm² in tissue of patients with CRSwNP (**D**), showing significant correlations ($P < .001$; $r = 0.8047$; and $P < .01$; $r = 0.6504$), determined via Spearman correlation. Levels of statistical significance are expressed as * $P < .05$, ** $P < .01$, and **** $P < .0001$.

neutrophil activity directly by analyzing tissue-derived neutrophils on reduced CD62L expression via fluorescence-activated cell sorting. The percentages of activated (CD62L⁺) neutrophils were significantly ($P < .05$) increased in tissue of patients with type 2 CRSwNP compared with in those with CRSsNP (Fig 3, A). In addition, the proteolytic activity of cathepsin G was significantly ($P < .01$) increased in the tissue of patients with CRSwNP compared with in controls, and the activity of neutrophil elastase was significantly ($P < .05$) increased in patients with CRSwNP compared with in controls and patients with CRSsNP (Fig 3, C and D).

Neutrophils are more prone to go into NETosis in CRSsNP tissue

Similar to eosinophils, neutrophils have the ability to generate extracellular traps in a process called NETosis. We quantified NETs in the tissue and found that NETosis was present in 67% of all patients with CRSsNP and in 64% of the patients with CRSwNP. The rate of NETosis was significantly ($P < .05$) increased in patients with CRSsNP compared with in healthy controls (Fig 4, A). Tissues were also stained for CitH3, an early marker of NETosis, and the presence of CitH3⁺ neutrophils was quantified and found significantly ($P < .01$) increased in patients with CRSsNP compared with in controls (Fig 4, B). Elastase

was stained positive on the same spots as CitH3, which confirmed the specificity of the CitH3 staining in neutrophils.

In CRSwNP tissue, NETosis was mainly found at the edges of (denuded) epithelium (Fig 4, C) and was colocalized with signs of bacterial colonization (Fig 4, D). In CRSsNP tissue, NETosis was mainly found in the stroma and underneath a clear thickened basement membrane associated with denuded epithelium (Fig 4, E).

Decreased neutrophil survival in CRSwNP cytokine environment

Differential neutrophil migration among the different endotypical environments was studied via Boyden-chamber assays, but did not reveal significant differences in neutrophil migration between the different patient groups (Fig 5, A). To study the influence of the cytokine environment on neutrophil survival, peripheral blood neutrophils were cultured in tissue homogenates of the different patient groups and the rate of apoptosis was analyzed after 24 hours of incubation via fluorescence-activated cell sorting analysis. Neutrophils cultured in CRSsNP homogenates showed significantly ($P < .05$) lower rates of apoptosis compared with neutrophils cultured in homogenates of healthy controls and patients with CRSwNP (Fig 5, B).

To investigate the possible involvement of IL-17 in neutrophil survival, the percentages of apoptotic neutrophils after incubation with homogenates were related to levels of IL-17 in these




FIG 3. Increased neutrophilic activation in CRSwNP. **A**, Fluorescence-activated cell sorting analysis showed significantly ($P < .05$) increased ratios of activated (CD62L⁻) neutrophils in mucosal tissue of patients with CRSwNP compared with those with CRSsNP. Significantly increased activity of cathepsin G ($P < .01$) and neutrophil elastase ($P < .05$) was observed in mucosal tissue of patients with CRSwNP compared with the tissue of controls and patients with CRSsNP. Levels of statistical significance are expressed as * $P < .05$ and ** $P < .01$.

homogenates. Interestingly, no correlation was found in CRSwNP homogenates, while neutrophil apoptosis was significantly ($P < .05$; $r = 0.5912$) inversely correlated with levels of IL-17 in the CRSsNP homogenates (Fig 5, C and D).

DISCUSSION

While we confirmed the general consideration that CRSsNP is mainly mediated by non-type 2 and CRSwNP by a type 2 inflammatory pattern, we here demonstrate a very heterogeneous image of neutrophilic and eosinophilic inflammation among all patients with CRS.^{3,14,21} Both neutrophil tissue counts and neutrophil-related protein levels demonstrated an increased neutrophilic inflammation in the CRSwNP group of patients. This is in line with a previous cluster analysis showing a markedly increased presence of neutrophilic marker proteins in the high type 2 clusters containing the most severe and difficult to treat patients with CRSwNP.¹⁴ Some reports claim that the number of neutrophils is elevated in CRSsNP compared with in CRSwNP, while other studies find that the numbers are comparable or elevated in CRSwNP.²²⁻²⁵ These contradicting results are more than likely attributed to the existence of different CRS endotypes of which some have a clear neutrophilic inflammation coexisting with eosinophilia.

Interestingly, neutrophils are also more activated, and neutrophilic proteases elastase and cathepsin G have increased activity

in type 2 CRSwNP. This indicates that neutrophils are not only more frequent in a type 2 environment, but they also have a higher potential to affect local inflammation via increased proteolytic activity. Elastase and cathepsin G are known to enhance secretion and activation of IL-1 family cytokines as IL-1 β and IL-33.²⁶ These cytokines are key players in the induction of type 2 responses as they function as chemoattractant for T_H2 cells and stimulate the production of type 2 cytokines in eosinophilic nasal polyps.²⁷⁻²⁹ In addition, neutrophil elastase itself has a major impact on airway inflammation as it enhances goblet cell hyperplasia and mucus production.³⁰

Despite the elevated number of neutrophils and increased neutrophil activity in patients with CRSwNP, no elevated neutrophil migration nor survival in CRSwNP tissue homogenates could be found *in vitro*. We speculate that neutrophil clearance is affected in CRSwNP tissue or that the active recruitment of neutrophils into CRSwNP tissue requires intact tissue. In severe asthma, a mixed neutrophilic-eosinophilic inflammation is associated with a mixed type 2/17 inflammation. Despite of the observed similarities in mixed airway inflammation, no remarkable elevation of IL-17 was observed in patients with CRSwNP, nor was there any correlation between IL-17 levels and number of neutrophils in these patients. This is in line with a recent cluster analysis by Tomassen et al¹⁴ showing increased levels of neutrophil related proteins, but no elevated levels of IL-17 in patients with high type 2 CRSwNP. We recently showed that CLCs,




FIG 4. Increased NETosis in CRSsNP. **A**, Quantification of NETs in the tissue of healthy controls and patients with CRSsNP and those with CRSwNP, expressed as percentage of neutrophils generating NETs. **B**, Quantification of CitH3-positive neutrophils in the tissue of healthy controls and patients with CRSsNP and those with CRSwNP, based on immunohistochemistry staining. **C-E**, Immunofluorescent staining of CitH3 and elastase, showing the presence of NETosis in CRSwNP and CRSsNP. **C**, NETosis at the edges of denuded epithelium in CRSwNP, and **(D)** colocalized with signs of bacterial colonization (arrowhead). **E**, NETosis underneath a clear thickened basement membrane in CRSsNP; white line represents the putative epithelial layer. Levels of statistical significance are expressed as $*P < .05$.

prominently present in patients with severe CRSwNP, can evoke a neutrophilic inflammation.¹⁶ In addition, increased neutrophilic influx was observed in murine lungs on injection with CLCs.¹⁷ In line with these findings, we showed here that the number of neutrophils in the tissue was correlated with the extent of EETosis and CLC deposition in CRSwNP. Despite the presence of CLCs in patients with CRSsNP, however, no correlation with neutrophil infiltration was found.³¹ This could possibly be attributed to the differences in CLC-size we observed in both groups, because the size of extracellular pathogens is also known to affect mechanism of clearance by neutrophils.³² *Staphylococcus aureus* colonization—another hallmark of CRSwNP—is also linked to increased neutrophil migration in CRSwNP and could therefore have a prominent role *in vivo* triggering neutrophilia in CRSwNP.³³⁻³⁵ Because the general bacterial load does not differ between CRSsNP and CRSwNP, we believe that the increased neutrophil inflammation observed in CRSwNP is not only caused by bacterial colonization.³⁶ However, variations in the composition of sinus bacterial microbiota were previously shown to be linked to CRS heterogeneity and may therefore also contribute to differences in neutrophilic activation and NETosis.^{36,37} In this study, differences in local bacterial composition in relation

to neutrophilic inflammation were not analyzed, but it would be interesting to do so for future studies.

A considerable fraction of neutrophils was found to undergo NETosis in both the CRSsNP and CRSwNP groups of patients. Both relative and absolute numbers of neutrophils undergoing NETosis were increased in CRSsNP. This was in line with our finding of increased neutrophil survival in CRSsNP. NETosis has been proposed as a type of slow programmed cell death that is prone to occur in neutrophils with increased survival.³⁸ While CitH3 is an early and consistent marker of NETosis, NETs themselves can be quite fragile. Extracellular traps can be disintegrated in a relatively fast manner, especially in a severe type 2 environment with increased numbers of macrophages and bacterial DNases.³⁹⁻⁴¹ Multiple stimulants can induce NET formation and even the type of NETosis—vital or suicidal—is dependent on its environmental incentives.⁴² It has recently been reported that CLCs could induce NETosis and neutrophilic inflammation in CRSwNP, and that NETosis could induce a type 2 response via release of extracellular double-stranded DNA.^{5,16} In CRSwNP, NETosis was mainly located at sites of sometimes denuded epithelium. Interestingly, neutrophilic mediators such as cathepsin G and elastase are extensively released during




FIG 5. Decreased neutrophil apoptosis in CRSsNP. **A**, Quantification of neutrophils migrated to tissue homogenates of healthy controls and patients with CRSsNP and those with CRSwNP. **B**, Ratios of apoptotic neutrophils after 24-hour culturing in tissue homogenates of healthy controls and patients with CRSsNP and those with CRSwNP. **C**, **D**, Scatter plots of neutrophil apoptosis (%) and levels of IL-17 in homogenates, showing a significant ($P < .05$; $r = -0.5912$) inversely correlation in CRSsNP (**C**) and no correlation in CRSwNP (**D**), determined via Spearman correlation. Levels of statistical significance are expressed as * $P < .05$ and ** $P < .01$.

NETosis and stimulate type 2 immune responses as described above.^{26,43}

While NETosis in CRSwNP was mainly found subepithelially and closely collocated with signs of bacterial infection, whereas NETosis in CRSsNP was mainly found deeper in the tissue. It remains to be clarified whether the NETosis is activated via similar pathways in the different endotypes. NETosis in CRSwNP may be driven by type 2-related mediators or may be related to the increased bacterial colonization and biofilm formation at sites of denuded epithelium. In CRSsNP, NETosis is mainly found in the stroma and underneath a clearly thickened membrane, without a clear association with bacterial colonization. This might indicate that other mediators such as IFN- γ or IL-12, both known inducers of NETosis and specifically elevated in those patients, might initiate the NETosis in the CRSsNP tissue.^{44,45}

Due to predominant T_H2 inflammatory pattern, therapies treating chronic airway diseases nowadays—as anti-IL5(R α), anti-IL4R/IL13R, or glucocorticoids—mainly focus on targeting the eosinophilic T_H2 inflammation.^{7-10,13} With the establishment of these therapies, considerable advances have been made, but still some patients do not respond to the treatment.^{11,46,47} We here showed that the contribution of neutrophils—by themselves via increased activity, and via increased influx and NETosis or in combination with eosinophilic inflammation—could be far more important than initially thought, especially in a T_H2 context. Interestingly, as CLCs are highly stable and remain present at

sites of inflammation for months, they could still affect neutrophilic inflammation long after the start of treatment.^{16,17} In addition, recent studies showed reduced responsiveness to corticosteroids in neutrophilic asthma and CRSwNP.^{12,13} All this indicates that neutrophilic inflammation is currently overlooked in assignments of therapies targeting the eosinophilic T_H2 inflammation and may need more attention to enable further improvement of treatments of the patients. Therefore, we speculate that the identification of increased neutrophil activation markers in patients with CRS may evolve as one of the critical parameters in determining treatment response in patients with CRS. The development and coadministration of novel drugs targeting CLC-activated neutrophils may complete the response to the type 2 cytokine antagonists in patients with a mixed eosinophilic-neutrophilic inflammation.

Overall, the behavior of neutrophils observed in CRSwNP is comparable to neutrophilic behavior in infections or other chronic type 2-mediated airway diseases. At sites of infection, neutrophils also induce NET formation and secrete proteases to promote bacterial clearance.⁴⁸ In asthma, neutrophils display a life span that is 5 days longer through decreased apoptosis.³⁰ This increased life span may set the basis for increased activation and development of NETosis, which could contribute to the severity of the disease. Also in nasal turbinate samples of patients with allergic rhinitis, the percentage of activated neutrophils was significantly higher than the percentage of non-activated

neutrophils, while this was not the case in the tissue of patients who were not allergic.⁴⁹ The difference between regular infections and infections in chronic type 2 diseases seems to be the triggers of the local neutrophilic inflammation. Migration of neutrophil to sites of infections are mainly mediated by damage-associated molecular pattern, IL-8, and leukotrienes, while in type 2 inflammations, this traditional way of neutrophilic migration seems to be overruled by mediators of eosinophilic type 2 inflammation such as CLCs.^{16,17,48} Also at sites of infection, neutrophils release proteins and form NETs in direct response to bacterial invasion, while in CRSwNP, these functions are not exclusively triggered by bacterial infection, but also by type 2 mediators such as eosinophils and CLCs.

In conclusion, we here demonstrated a profound involvement of neutrophils in severe type 2 CRSwNP, induced by products of eosinophilic inflammation, and independent of IL-17. Neutrophils have a high potential to affect local inflammation in CRSwNP through an increased activation status, increased proteolytic activity of elastase and cathepsin G, the formation of NETs, and finally to aggravate the type 2 inflammation. These mechanisms could result in a circle of aggravation of inflammation in patients with the most severe CRSwNP with mixed eosinophilic-neutrophilic inflammation, leading to the insensitivity to glucocorticosteroids and type 2 biologics; this hypothesis needs to be further supported.

Key messages

- Type 2 CRSwNP displays a severe neutrophilic inflammation, characterized by increased neutrophil infiltration and activation.
- The presence of CLCs is associated with neutrophilic infiltration in CRSwNP.
- Neutrophils are less prone to generate NETs in CRSwNP tissue and NETosis in CRSwNP is associated with bacterial colonization.

REFERENCES

- Khan A, Vandeplas G, Huynh TMT, Joish VN, Mannent L, Tomassen P, et al. The Global Allergy and Asthma European Network (GALEN) rhinosinusitis cohort: a large European cross-sectional study of chronic rhinosinusitis patients with and without nasal polyps. *Rhinology* 2019;57:32-42.
- Dietz de Loos D, Lourijsen ES, Wildeman MAM, Freling NJM, Wolvers MDJ, Reitsma S, et al. Prevalence of chronic rhinosinusitis in the general population based on sinus radiology and symptomatology. *J Allergy Clin Immunol* 2019;143:1207-14.
- Van Zele T, Claeys S, Gevaert P, Van Maele G, Holtappels G, Van Cauwenberge P, et al. Differentiation of chronic sinus diseases by measurement of inflammatory mediators. *Allergy* 2006;61:1280-9.
- Ray A, Kolls JK. Neutrophilic inflammation in asthma and association with disease severity. *Trends Immunol* 2017;38:942-54.
- Toussaint M, Jackson DJ, Swieboda D, Guedan A, Tsourouktsoglou TD, Ching YM, et al. Host DNA released by NETosis promotes rhinovirus-induced type-2 allergic asthma exacerbation. *Nat Med* 2017;23:681-91.
- Moore WC, Hastie AT, Li X, Li H, Busse WW, Jarjour NN, et al. Sputum neutrophil counts are associated with more severe asthma phenotypes using cluster analysis. *J Allergy Clin Immunol* 2014;133:1557-63.e5.
- Bachert C, Han JK, Desrosiers M, Hellings PW, Amin N, Lee SE, et al. Efficacy and safety of dupilumab in patients with severe chronic rhinosinusitis with nasal polyps (LIBERTY NP SINUS-24 and LIBERTY NP SINUS-52): results from two multicentre, randomised, double-blind, placebo-controlled, parallel-group phase 3 trials. *Lancet* 2019;394:1638-50.
- Bachert C, Zhang N, Hellings PW, Bousquet J. Endotype-driven care pathways in patients with chronic rhinosinusitis. *J Allergy Clin Immunol* 2018;141:1543-51.
- Bachert C, Sousa AR, Lund VJ, Scadding GK, Gevaert P, Nasser S, et al. Reduced need for surgery in severe nasal polyposis with mepolizumab: randomized trial. *J Allergy Clin Immunol* 2017;140:1024-31.e14.
- Pauwels B, Jonstam K, Bachert C. Emerging biologics for the treatment of chronic rhinosinusitis. *Expert Rev Clin Immunol* 2015;11:349-61.
- Van Zele T, Holtappels G, Gevaert P, Bachert C. Differences in initial immunoprofiles between recurrent and nonrecurrent chronic rhinosinusitis with nasal polyps. *Am J Rhinol Allergy* 2014;28:192-8.
- Green RH, Brightling CE, Woltmann G, Parker D, Wardlaw AJ, Pavord ID. Analysis of induced sputum in adults with asthma: identification of subgroup with isolated sputum neutrophilia and poor response to inhaled corticosteroids. *Thorax* 2002;57:875-9.
- Wen W, Liu W, Zhang L, Bai J, Fan Y, Xia W, et al. Increased neutrophilia in nasal polyps reduces the response to oral corticosteroid therapy. *J Allergy Clin Immunol* 2012;129:1522-8.e5.
- Tomassen P, Vandeplas G, Van Zele T, Cardell LO, Arebro J, Olze H, et al. Inflammatory endotypes of chronic rhinosinusitis based on cluster analysis of biomarkers. *J Allergy Clin Immunol* 2016;137:1449-56.e4.
- Gevaert E, Zhang N, Krysko O, Lan F, Holtappels G, De Ruyck N, et al. Extracellular eosinophilic traps in association with *Staphylococcus aureus* at the site of epithelial barrier defects in patients with severe airway inflammation. *J Allergy Clin Immunol* 2017;139:1849-60.e6.
- Gevaert E, Delemarre T, De Volder J, Zhang N, Holtappels G, De Ruyck N, et al. Charcot-Leyden crystals promote neutrophilic inflammation in patients with nasal polyposis. *J Allergy Clin Immunol* 2020;145:427-30.e4.
- Persson EK, Verstraete K, Heyndrickx I, Gevaert E, Aegerter H, Percier J-M, et al. Protein crystallization promotes type 2 immunity and is reversible by antibody treatment. *Science* 2019;364:eaaw4295.
- Ueki S, Tokunaga T, Melo RCN, Saito H, Honda K, Fukuchi M, et al. Charcot-Leyden crystal formation is closely associated with eosinophil extracellular trap cell death. *Blood* 2018;132:2183-7.
- Zhang N, Van Crombruggen K, Holtappels G, Lan F, Katotomichelakis M, Zhang L, et al. Suppression of cytokine release by fluticasone furoate vs. mometasone furoate in human nasal tissue ex-vivo. *PLoS One* 2014;9:e93754.
- Derycke L, Zhang N, Holtappels G, Dutre T, Bachert C. IL-17A as a regulator of neutrophil survival in nasal polyp disease of patients with and without cystic fibrosis. *J Cyst Fibros* 2012;11:193-200.
- Ahern S, Cervin A. Inflammation and endotyping in chronic rhinosinusitis-a paradigm shift. *Medicina (Kaunas)* 2019;55:95.
- Polzehl D, Moeller P, Riechelmann H, Perner S. Distinct features of chronic rhinosinusitis with and without nasal polyps. *Allergy* 2006;61:1275-9.
- Shi LL, Xiong P, Zhang L, Cao PP, Liao B, Lu X, et al. Features of airway remodeling in different types of Chinese chronic rhinosinusitis are associated with inflammation patterns. *Allergy* 2013;68:101-9.
- Soler ZM, Sauer D, Mace J, Smith TL. Impact of mucosal eosinophilia and nasal polyposis on quality-of-life outcomes after sinus surgery. *Otolaryngol Head Neck Surg* 2010;142:64-71.
- Tokunaga T, Sakashita M, Haruna T, Asaka D, Takeno S, Ikeda H, et al. Novel scoring system and algorithm for classifying chronic rhinosinusitis: the JESREC Study. *Allergy* 2015;70:995-1003.
- Clancy DM, Henry CM, Sullivan GP, Martin SJ. Neutrophil extracellular traps can serve as platforms for processing and activation of IL-1 family cytokines. *FEBS J* 2017;284:1712-25.
- Schmitz J, Owyang A, Oldham E, Song Y, Murphy E, McClanahan TK, et al. IL-33, an interleukin-1-like cytokine that signals via the IL-1 receptor-related protein ST2 and induces T helper type 2-associated cytokines. *Immunity* 2005;23:479-90.
- Teufelberger AR, Nordengrun M, Braun H, Maes T, De Grove K, Holtappels G, et al. The IL-33/ST2 axis is crucial in type 2 airway responses induced by *Staphylococcus aureus*-derived serine protease-like protein D. *J Allergy Clin Immunol* 2018;141:549-59.e7.
- Kato A. Immunopathology of chronic rhinosinusitis. *Allergol Int* 2015;64:121-30.
- Silvestre-Roig C, Hidalgo A, Soehnlein O. Neutrophil heterogeneity: implications for homeostasis and pathogenesis. *Blood* 2016;127:2173-81.
- Delemarre T, Holtappels G, De Ruyck N, Zhang N, Nauwynck H, Bachert C, et al. Type 2 inflammation in chronic rhinosinusitis without nasal polyps: another relevant endotype. *J Allergy Clin Immunol* 2020;146:337-43.e6.
- Warnatsch A, Tsourouktsoglou TD, Branzk N, Wang Q, Reincke S, Herbst S, et al. Reactive oxygen species localization programs inflammation to clear microbes of different size. *Immunity* 2017;46:421-32.

33. Wang X, Du J, Zhao C. Bacterial biofilms are associated with inflammatory cells infiltration and the innate immunity in chronic rhinosinusitis with or without nasal polyps. *Inflammation* 2014;37:871-9.
34. Van Zele T, Gevaert P, Watelet JB, Claeys G, Holtappels G, Claeys C, et al. *Staphylococcus aureus* colonization and IgE antibody formation to enterotoxins is increased in nasal polyposis. *J Allergy Clin Immunol* 2004;114:981-3.
35. Huvenne W, Callebaut I, Reekmans K, Hens G, Bobic S, Jorissen M, et al. *Staphylococcus aureus* enterotoxin B augments granulocyte migration and survival via airway epithelial cell activation. *Allergy* 2010;65:1013-20.
36. Chalermwatanachai T, Zhang N, Holtappels G, Bachert C. Association of mucosal organisms with patterns of inflammation in chronic rhinosinusitis. *PLoS One* 2015;10:e0136068.
37. Cope EK, Goldberg AN, Pletcher SD, Lynch SV. Compositionally and functionally distinct sinus microbiota in chronic rhinosinusitis patients have immunological and clinically divergent consequences. *Microbiome* 2017;5:53.
38. Gray RD, Hardisty G, Regan KH, Smith M, Robb CT, Duffin R, et al. Delayed neutrophil apoptosis enhances NET formation in cystic fibrosis. *Thorax* 2018;73:134-44.
39. Farrera C, Fadeel B. Macrophage clearance of neutrophil extracellular traps is a silent process. *J Immunol* 2013;191:2647-56.
40. Thammavongsa V, Missiakas DM, Schneewind O. *Staphylococcus aureus* degrades neutrophil extracellular traps to promote immune cell death. *Science* 2013;342:863-6.
41. Storisteanu DML, Pocock JM, Cowburn AS, Juss JK, Nadesalingam A, Nizet V, et al. Evasion of neutrophil extracellular traps by respiratory pathogens. *Am J Respir Cell Mol Biol* 2017;56:423-31.
42. Yipp BG, Kubes P. NETosis: how vital is it? *Blood* 2013;122:2784-94.
43. Lefrancais E, Roga S, Gautier V, Gonzalez-de-Peredo A, Monsarrat B, Girard JP, et al. IL-33 is processed into mature bioactive forms by neutrophil elastase and cathepsin G. *Proc Natl Acad Sci U S A* 2012;109:1673-8.
44. Martinelli S, Urošević M, Daryadel A, Oberholzer PA, Baumann C, Fey MF, et al. Induction of genes mediating interferon-dependent extracellular trap formation during neutrophil differentiation. *J Biol Chem* 2004;279:44123-32.
45. Keeter WC, Moriarty A, Butcher MJ, Ma KW, Nadler JL, Galkina E. IL-12 induced STAT4 activation plays a role in pro-inflammatory neutrophil functions. *J Immunol* 2018;200(suppl 1):166.59.
46. Calus L, Van Bruaene N, Bosteels C, Dejonckheere S, Van Zele T, Holtappels G, et al. Twelve-year follow-up study after endoscopic sinus surgery in patients with chronic rhinosinusitis with nasal polyposis. *Clin Transl Allergy* 2019;9:30.
47. DeConde AS, Mace JC, Levy JM, Rudmik L, Alt JA, Smith TL. Prevalence of polyp recurrence after endoscopic sinus surgery for chronic rhinosinusitis with nasal polyposis. *Laryngoscope* 2017;127:550-5.
48. De Oliveira S, Rosowski EE, Huttenlocher A. Neutrophil migration in infection and wound repair: going forward in reverse. *Nat Rev Immunol* 2016;16:378-91.
49. Arebro J, Ekstedt S, Hjalmarsson E, Winqvist O, Kumlien Georen S, Cardell LO. A possible role for neutrophils in allergic rhinitis revealed after cellular subclassification. *Sci Rep* 2017;7:43568.




FIG E1. Cytokine levels in different patient groups. **A**, IL-5 protein levels in the tissue of healthy controls and patients with CRSsNP and those with CRSwNP, measured via Luminex assay. **B**, IFN-γ protein levels in the tissue of healthy controls and patients with CRSsNP and those with CRSwNP, measured via ELISA. Levels of statistical significance are expressed as $*P < .05$, $***P < .001$, and $****P < .0001$.




FIG E2. Increased eosinophilia in CRSwNP. **A**, Quantification of eosinophils per mm² in tissue of healthy controls and patients with CRSsNP and those with CRSwNP. **B**, ECP levels in the tissue of healthy controls and patients with CRSsNP and those with CRSwNP, measured via UniCAP. Levels of statistical significance are expressed as ** $P < .01$, *** $P < .001$, and **** $P < .0001$.