

Dupilumab for Atopic Dermatitis during Pregnancy and Breastfeeding: clinical experience in 13 patients

Dear editor,

Atopic dermatitis (AD) affects patients of reproductive age and can worsen or start during pregnancy.¹ The high prevalence of AD in childbearing age women highlights the need to determinate the safety of biologics during pregnancy and lactation. Currently, data is limited regarding the use of dupilumab in pregnant/breastfeeding women.² The aim of this report is to describe our experience using dupilumab during pregnancy/breastfeeding in 13 women with AD.

We conducted a multicentre retrospective case series study with eleven pregnant and two breastfeeding women included. All suffered from moderate-to-severe AD with mean EASI score before initiating dupilumab of 34.9 (0-72). The mean time of exposure to dupilumab was of 6.8 ± 2.9 and 11.5 ± 3.2 months during pregnancy and breastfeeding, respectively. Patients' characteristics, treatments received before dupilumab, reasons that led to initiate this treatment as well as the cause of treatment discontinuation are mentioned in Table 1. The 83.3% of the patients reached EASI-75 at week 16 and this percentage was even better at weeks 24 (90.9%) and 52 (90.0%). At week 16, pruritus NRS and DLQI scores were considerably reduced, maintaining these good results at weeks 24 and 52 (Fig. 1). Regarding materno-fetal outcomes, eleven pregnant women gave birth to twelve babies. Eight were eutocic, vaginal deliveries and three were dystocic deliveries (one vacuum-assisted and two caesarean-deliveries). Ten (90.9%) were full-term pregnancies, without complications during delivery and giving birth to normal weight babies. One was a twin pregnancy that required a caesarean-section, giving birth two premature babies (week 35) with low

weigh (1.5 and 2.0kg respectively) but with a proper development and weight gain. None of the women suffered a miscarriage neither complications during pregnancy or puerperium. Moreover, none of the babies have suffered from AD to date.

Management of AD during both pregnancy and breastfeeding is challenging, especially in severe forms, due to the lack of clinical studies investigating effects of novel treatments on fertility, pregnancy and lactation. Dupilumab, as other IgG antibodies, is expected to cross the placental barrier, especially in the second and third trimesters of pregnancy.³ However, studies using animal models have shown no direct or indirect harmful effects in terms of reproductive toxicity.⁴ The European Medicines Agency reported that the spontaneous abortion rate in patients treated with dupilumab did not exceed that of the general population.⁴ Additionally, a study that assessed the safety profile of dupilumab during pregnancy using pharmacovigilance data did not find any adverse effects.⁵ Moreover, seven case reports published of pregnant women treated with dupilumab resulted in livebirths, with only one premature birth and another infant with low birth weight reported.^{1,2,6-10} Regarding breastfeeding, secretion and absorption of dupilumab is thought to be minimal due to its high molecular weight and its degradation by acid and proteolytic enzymes in the infant's gut, respectively. Notwithstanding, during the first three days postpartum there are wide gaps between the breast alveolar cells which may allow the passage of immunoglobulins through them.¹ For these reasons it is recommended to avoid breastfeeding or the use of dupilumab during a short postpartum period and in preterm infants who are likely to have an immature gastrointestinal tract. There is only one case report published of a woman treated with dupilumab whom breastfeed her infant for at least 4 months in which no complications were reported.²

In conclusion, data is limited regarding the use of dupilumab during pregnancy/breastfeeding. We report the largest case series yet described of patients with moderate-to-severe AD treated with this drug during pregnancy/lactation. In our series maternal-fetal outcomes were excellent, highlighting that dupilumab use in this subgroup of patients has its place but should be preceded by careful assessment.

References

1. Lobo Y, Lee RC, Spelman L. Atopic Dermatitis Treated Safely with Dupilumab during Pregnancy: A Case Report and Review of the Literature. *Case Rep Dermatol*. 2021;**13**:248-256.
2. Kage P, Simon J.C, Treudler R. A case of atopic eczema treated safely with dupilumab during pregnancy and lactation. *J Eur Acad Dermatol Venereol* 2020; **34**:e256-e257.
3. Vestergaard C, Wollenberg A, Barbarot S, et al. European task force on atopic dermatitis position paper: treatment of parental atopic dermatitis during preconception, pregnancy and lactation period. *J Eur Acad Dermatol Venereol*. 2019;**33**:1644–59.
4. European Medicines Agency (EMA) Committee for Medicinal Products for Human Use. (CHMP). Dupixent CHMP assessment report on extension of marketing authorisation and an extension of indication variation February 2019. Available from: https://www.ema.europa.eu/en/documents/variation-report/dupixent-h-c-4390-x-0004-g-e-par-assessment-report-extension_en.pdf (last accessed 1 December 2022).
5. Khamisy-Farah R, Damiani G, Kong JD, et al. Safety profile of Dupilumab during pregnancy: a data mining and disproportionality analysis of over 37,000 reports from the WHO individual case safety reporting database (VigiBase™). *Eur Rev*

6. Mian M, Dunlap R, Simpson E. Dupilumab for the treatment of severe atopic dermatitis in a pregnant patient: a case report. *JAAD Case Rep* 2020;**6**:1051-1052.
7. Gracia-Darder I, Pons De Ves J, Reyero Cortina M, Martín-Santiago A. Patient with atopic dermatitis, hyper IgE syndrome and ulcerative colitis, treated successfully with dupilumab during pregnancy. *Dermatol Ther* 2022; **35**: e15237.
8. Costley M, Murphy B. Severe atopic dermatitis treated successfully with dupilumab throughout pregnancy. *Clin Exp Dermatol* 2022; **47**:960-961.
9. Akhtar N.H, Khosravi-Hafshejani T, Akhtar D, et al. The use of dupilumab in severe atopic dermatitis during pregnancy: a case report. *Allergy Asthma Clin Immunol* 2022;**18**:9.
10. Riquelme-McLoughlin C, Mascaró J.M. Treatment of pemphigoid gestationis with dupilumab. *Clin Exp Dermatol* 2021;**46**:1578-1579.

Tables and figures:

Figure 1. Evolution of EASI (A), itch (B) and DLQI (C) scores in weeks 0,16, 24 and 52 in 12 patients (P) during pregnancy and/or breastfeeding.

Table 1. Patient population characteristics