

MISS JESSY ELST (Orcid ID : 0000-0003-3506-8200)

DR ATHINA VAN GASSE (Orcid ID : 0000-0003-1657-5135)

MR MICHEL VAN HOUDT (Orcid ID : 0000-0002-9510-6961)

DR VITO SABATO (Orcid ID : 0000-0002-1321-314X)

Article type : Review Article

Mast cell activation tests by flow cytometry: a new diagnostic asset?

Elst Jessy, MSc¹, van der Poorten Marie-Line M, MD^{1, 2}, Van Gasse Athina L, MD, PhD^{1,2}, De Puyssseleir Leander, MD¹, Hagendorens Margo M., MD, PhD^{1,2}, Faber Margaretha A, MD, PhD¹, Van Houdt Michel, MLT¹, Passante Egle, PhD³, Bahri Rajia⁴, PhD, Walschot Mark, MD¹, Mertens Christel, MLT¹, Bridts Chris H, MLT¹, Sabato Vito, MD, PhD^{1, 5}, Ebo Didier G, MD, PhD^{1,5*}

¹ Faculty of Medicine and Health Science, Department of Immunology – Allergology – Rheumatology , Antwerp University Hospital and the Infla-Med Centre of Excellence, University of Antwerp, Antwerp University Hospital, Antwerp, Belgium.

² Faculty of Medicine and Health Science, Department of Paediatrics, University of Antwerp, Antwerp University Hospital, Antwerp, Belgium.

³ GlaxoSmithKline, GSK House, 80 Great West Road, Brentford, Middlesex, TW89GS

⁴ Lydia Becker Institute of Immunology and Inflammation, Division of Musculoskeletal &

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/CEA.13984](#)

This article is protected by copyright. All rights reserved

Dermatological Sciences, School of Biological Sciences, Faculty of Biology, Medicine and Health, University of Manchester, Manchester Academic Health Science Centre, Manchester UK

⁵ Department of Immunology and Allergology, AZ Jan Palfijn Gent, Ghent, Belgium

* Correspondence to:

Ebo Didier, MD, PhD

Laboratory of Immunology University of Antwerp

CDE T5.95 Universiteitsplein 1

2610 Antwerpen

Belgium

Tel: ++ 32 (0) 3 2652595

Fax: ++ 32 (0) 3 2652655

immuno@uantwerpen.be

Short title: Mast cell activation test: potentials and limitations

Acknowledgments: Vito Sabato is a Senior Clinical Researcher of the Research Foundation Flanders/Fonds Wetenschappelijk Onderzoek (FWO: 1804518N). Didier Ebo is a Senior Clinical Researcher of the Research Foundation Flanders/Fonds Wetenschappelijk Onderzoek (FWO: 1800614N). The Department of Immunology – Allergology – Rheumatology is a center of excellence of the World Allergy Organization.

Authorship: JE, DE, VS initiated the concept and contributed made the first draft. RB and EP participated to the development of the document. All authors participated in proofreading and revising the final manuscript.

Conflict of Interest statement: The authors declare no conflict of interest


Funding statement: The authors declare no competing financial interest

Data availability statement: The data that support the findings of this study are available from the corresponding author upon reasonable request.

Key messages

- Mast cell activation test (MAT) is a promising diagnostic for IgE-mediated allergies and auto-immune urticaria.
- MAT allows simultaneous analysis of MRGPRX2⁺ and MRGPRX2⁻ subpopulations and silencing of the receptor.
- In combination with BATs, MATs might advance our understanding of effector cell activation and degranulation.

Graphical abstract



Abstract

Since the late nineties, evidence has accumulated that flow-assisted basophil activation test (BAT) might be an accessible and reliable method to explore the mechanisms governing basophil degranulation and diagnostic allowing correct prediction of the clinical outcome following exposure to the offending allergen(s) and cross-reactive structures for different IgE-dependent allergies and particular forms of autoimmune urticaria. Although the BAT offers many advantages over mediator release tests, it is left with some weaknesses that hinder a wider application. It is preferable to perform the BAT analysis within four hours of collection and the technique does not advance diagnosis in patients with non-responsive cells. Besides, the BAT is difficult to standardize mainly because of the difficulty to perform large batch analyses that might span over several days. This article reviews the status on flow cytometric mast cell activation test (MAT) using passively sensitized MCs with patients' sera or plasma (henceforth indicated as passive MAT; pMAT) using both MC lines and cultured MCs in the diagnosis of IgE-dependent allergies. In addition, this paper provides guidance for generating human MCs from peripheral blood CD34⁺ progenitor cells (PBCMCs) and correct interpretation of flow cytometric analyses of activated and/or degranulating cells. With the recent recognition of the mas-related G-protein coupled receptor X2 (MRGPRX2) occupation as a putative mechanism of immediate drug hypersensitivity reactions (IDHRs), we also speculate how direct activation of MCs (dMAT) - that is direct activation by MRGPRX2 agonists without prior passive sensitization - could advance paradigms for this novel endotype of IDHRs.

Keywords: allergy, anaphylaxis, CD63, flow cytometry, human cultured mast cells, IgE, mast cell activation, MRGPRX2

List of abbreviations

aMAT: autoimmune mast cell activation test

ASST: autologous serum skin test

Atra: atracurium

BAT(s): basophil activation test(s)

BM: bone marrow

BMCMC(s): bone marrow cultured mast cell(s)

CB: cord blood

Cipro: ciprofloxacin

Cisatra: cisatracurium

dMAT: direct mast cell activation test (without prior passive sensitization with patients' sera or plasma)

DsRed: *Discosoma* sp. red fluorescent protein

ENPP: ectonucleotide pyrophosphatase/phosphodiesterase

FBS: fetal bovine serum

FcεRI: high affinity receptor for IgE

FCM: flow cytometry

IDHR(s): immediate drug hypersensitivity reaction(s)

IDT(s): intradermal test(s)

IL: interleukin

pMAT: passive mast cell activation tests (after passive sensitization of MCs with patients' sera or plasma)

IMDM: Iscove's Modified Dulbecco's Medium

IPSC: induced pluripotent stem cells

LAMP: lysosome associated molecular protein

Levo: levofloxacin

MAT(s): mast cell activation test(s)

MC(s): mast cell(s)

MC_{TC}: connective tissue mast cells

Moxi: moxifloxacin

MRGPRX2: Mas-related G-protein coupled receptor X2

Mths: months

ND: not determined

NMBA(s): neuromuscular blocking agent(s)

PB: peripheral blood

PBCMC(s): peripheral blood cultured mast cell(s)

RBL: Rat Basophilic Leukaemia

Rocu: rocuronium

SCF: stem cell factor

slgE: specific IgE antibody

SPT(s): skin prick test(s)

ST(s): skin test(s)

THIQ: tetrahydroisoquinoline

Introduction

Although tissue resident mast cells (MCs) have been recognized as primary effector cells in IgE-dependent allergies, evidence about the utility of mast cell activation test (MATs) using passively sensitized MCs (passive MAT; pMAT) is relatively new. Actually, most studies on a functional assay for allergy diagnosis have focused on circulating basophils, as these cells are more accessible for *ex vivo* experiments. Together with the advent of flow cytometers and the discovery of easy to measure basophil identification and activation/degranulation markers that paved the way for the basophil activation test (BAT). The clinical utility of the BAT in the diagnosis and monitoring of allergic diseases has widely been studied and reviewed (1, 2). These studies and reviews show that the BAT, although offering many advantages over mediator release tests, is left with some weaknesses that hinder a wider application. Although the BAT can be executed up to 24 hours of collection (3), it is preferable to perform the analyses within 4 hours (4) and the technique does not advance diagnosis in patients with non-responsive cells. Besides, the BAT is difficult to standardize mainly because of the difficulty to perform large batch analyses that might require repetitive analyses spanning over several days.

As already indicated, MC studies should ideally analyses of tissue-resident cells of interest (5, 6). However, such studies are hindered by technical issues such as isolating sufficient numbers of viable, mature and functionally MCs and poor clonal expansion of such cells *ex vivo*. Furthermore, functional behavior of tissue MCs can be influenced by the applied isolation techniques (7-9). To circumvent the limitations encountered with BAT and tissue MCs, different groups have adopted passive sensitization techniques using specific basophil and MC cell lines (10-12), stripped basophils (13, 14) or primary human MCs cultured from donor progenitor cells (10, 15) (16-18).

This article reviews the status on flow cytometric pMAT using both MC lines and cultured cells in the diagnosis of IgE-dependent allergies. Table 1 summarizes the main differences between traditional flow cytometry (FCM)-assisted BAT and pMAT in their application to explore the effector cell activating potential of IgE antibodies and diagnosis of IgE-dependent allergies. In addition, this paper provides guidance for generating MCs from peripheral blood CD34⁺ progenitor cells (PBCMCs) and correct interpretation of flow cytometric analyses of activated and/or degranulating cells. With the recent recognition of

the mas-related G-protein coupled receptor X2 (MRGPRX2) occupation as a putative mechanism of immediate drug hypersensitivity reactions (IDHRs), we also speculate how direct activation of MCs (dMAT) - that is direct activation by MRGPRX2 agonists without prior passive sensitization - could advance paradigms for this novel endotype of IDHRs. Finally, we provide a proof of concept for application of the MAT to depict the presence of autoantibodies autoimmune urticaria in a technique called autoimmune MAT (aMAT).

Flow cytometric analysis of activated and/or degranulating humanized cells and primary human MCs

For decades, exploration of MC functionality has predominantly relied on quantification of released mediators or proteases (e.g. β -hexosaminidase, histamine, etc.) in the supernatant, collected after centrifugation of stimulated cells (10, 17-53). However, these techniques do not allow to analyze (subtle) responses of individual cells nor do they enable to study (small) subpopulations, as their results represent an average of all stimulated cells. In this respect, multicolor FCM enabling to couple surface alterations (e.g. up-regulation of activation and degranulation markers, exteriorization of granular content) and intracellular changes (e.g. phosphorylation of signal molecules, trapping of chemokines, calcium staining) is a significant asset. As shown in Figure 1, traditional FCM-based MAT relies upon cellular identification and quantification of activation and/or degranulation markers on the surface membrane. These changes are detectable and quantifiable using specific fluorescent-labelled monoclonal antibodies. In most studies, MCs have been identified by positive staining for surface markers such as CD117/CD203c (15, 16, 54-56) or as CD117/Fc ϵ RI (10, 19, 20, 22, 40, 44). Subsequently, after activation, the up-regulation of specific markers, such as CD63 and CD107a or CD203c is measured (10, 11, 15, 16, 21, 22, 44, 54-58). CD63 and CD107a are membrane proteins of the lysosome associated molecular protein (LAMP) tetraspanin family that are barely expressed by resting MCs. Up-regulation of CD63 is closely associated with mediator release during degranulation (6, 10, 59-61). CD203c (or ENPP-3), an ectonucleotide pyrophosphatase/phosphodiesterase family member, is already expressed on the cell surface of quiescent MCs and is up-regulated after stimulation (15, 59, 61, 62). However, the upregulation of CD203c and CD63 is clearly different. Actually, upregulation of CD63, unlike upregulation of CD203c, is generally bimodal with a

subpopulation of degranulating cells that express CD63 with a high intensity versus a resting subpopulation with a significant lower expression. Degranulation of MCs can also be explored by measuring the exteriorization of granule matrix. Briefly, anionic proteoglycans from exteriorized MC granule matrix are stained by cationic fluorescent avidin probes (19-22). MC degranulation can also be measured by flow cytometric techniques to quantify intracellular histamine and its release. In this technique, the intracellular content of histamine is quantified using the affinity of diamine oxidase (DAO) for its substrate histamine (63, 64). By coupling the histaminase DAO to fluorochromes, intracellular histamine content and consequently release of histamine can be measured by FCM at a single cell level (54). Newly synthesized chemokines such as CCL4 or CXCL8 can be trapped and measured intracellularly (22). Next, phosphotyrosine staining, a parameter associated to cell protein kinase function, can be used to study MC activation (22). Finally, MC activation can also be measured by imaging changes of intracellular calcium over time (20, 22, 30, 57, 65).

Basophil and mast cell lines

For a recent comprehensive compendium on humanized basophil and fully human MC lines the reader is referred elsewhere (12). Tables 2 and 3 summarize the different humanized rat leukemia and the fully human MC lines and indicate those applied in pMAT studies. Table 4 shows the cell lines expressing MRGPRX2 that have been used to study MRGPRX2-dependent MC activation and degranulation. As most of these studies have used alternative non-flow cytometric read-outs, they are only touched briefly.

Mast cell lines: utility in IgE-dependent allergy

In an attempt to streamline the functional assessment of sIgE antibodies to trigger effector cells, much effort has been put in the development and optimization of cultured cell lines. Nevertheless, these cell lines still present some important shortcomings that generate a lack of acceptance and skepticism by clinicians toward this experimental tool. The first lineage used in this domain have been humanized or transfected cell lines derived from the Rat Basophilic Leukaemia (RBL) cell line and expressing human FcεRI chains (48-50) such as, RBL SX-38 and RBL-48 (47-52, 65-71) (Table 2). However, as

earlier explained, most of these studies used mediator release as a read out. Only Taudou et al. used FCM to demonstrate that a RBL line transfected with the α -chain of the human Fc ϵ RI produce a calcium response when sensitized with patients' sIgE and subsequently incubated with house dust mite (65).

Although for years some of these humanized RBL cell lines have significantly contributed to our understandings of the mechanisms that govern IgE-dependent effector cell degranulation, their clinical application is limited. First, like the native RBL cells, all the chimeric clones show reduced ability to bind purified IgE after approximately 2 weeks in culture (12). Second, these cell lines often need high serum titers of sIgEs in the donor's serum to reach satisfactory results (e.g. >10 kUA/mL for peanut sIgE in RBL SX-38) and require a high percentage of receptor occupancy (>10% for RBL-2H3 E5.D12.8) (49, 72). Third, human xenoreactive IgGs present in patients' sera can cause cell death and induce background degranulation in up to 25% and laboratory procedures used to limit this shortcoming (i.e. serum dilution) could lead to suboptimal MC sensitization and lead to false negative results (71, 73).

As shown in Table 3, there is only a limited number of fully human MC lineages and only few have been used in IgE-mediated allergy research. Of relevance for this review, is the LAD-2 cell line, generated by Kirshenbaum and co-workers in the early 2000s, and that has been validated to assess peanut allergy and to monitor the effect of peanut oral immunotherapy of in children (11). In a first study, Santos et al (11), have shown that pMAT using passively sensitized LAD-2 cells is comparable to BAT in terms of specificity (98%) but has lower sensitivity (73%) to diagnose peanut allergy (11). However, pMAT provided results for those patients with unresponsive basophils. Indirect MAT also identified patients at risk of severe reactions during food challenges (11) and, along with stripped basophil experiments, enabled to demonstrate the inhibitory capacity of IgG4 antibodies in peanut-tolerant children sensitized to peanut major allergens (58). In a second study, the same group confirmed their previously reported findings on the induction of blocking antibodies with peanut oral immunotherapy and disclose that this treatment does not change the functional characteristics of sIgE, as determined by the ability of LAD-2 cells and basophils to respond to peanut allergen (74).

Igarashi et al. (39) described the use of human induced pluripotent stem cells (iPSC) cell line (201B7) to generate short-lived human iPSC-MC line to use in the pMAT. The authors tested the ability of this cell line to be triggered in a dose-dependent manner by cedar pollen, mite and house dust following appropriate sensitization. iPSC-MC could not be sensitized by ragweed. Due to the human origin of these cells, serum could be used undiluted without cellular damage. However, iPSC-derived cells could be maintained in culture only for maximum eight months, this might limit their routine use as a diagnostic tool.

To the best of our knowledge, humanized or human MC lines have only been used to study the pMAT with proteinaceous allergens and not in the field of IDHRs yet. However, in the clinics seldomly additional diagnostic tests are needed for this proteinaceous allergens, but are more than welcome in IDHRs.

Mast cell lines: utility in MRGPRX2-mediated hypersensitivity

Besides the use of MCs lines to study the IgE-dependent activation, cell lines are also increasingly used to study the IgE-independent activation mechanisms of MCs. Since the seminal description by Mc Neil et al. (40), increasing lines of evidence indicate that MC degranulation via occupation of MRGPRX2 might constitute a new endotype of IDHRs independent from cross-linking of IgE/FcεRI complexes (20, 42-44, 75-78). Examples of drugs that might act via MRGPRX2 engagement are neuromuscular blocking agents (NMBAs), fluoroquinolones, icatibant and opiates such as morphine. Many of these drugs harbor a tetrahydroisoquinoline (THIQ) motif (40). Table 4 summarizes the cell lines, including human MC lines that have been used to study the MRGPRX2-dependent activation in IDHRs per drug class. Note that FCM was only used by Navines-Ferrer et al. (44). Although these *in vitro* studies have nourished our insights in MRGPRX2-dependent IDHRs and anaphylaxis, interpretation of the sometimes-incongruent findings and translation into the clinical setting is difficult. Based on the current models, the MRGPRX2-activating capacity and potency of different drugs is left with some uncertainties. First, it appears that its MRGPRX2-activating capacity is different and difficult, if at all, predictable by its effects on its mouse orthologue. For example, rocuronium is ~ 12 times less potent at the MRGPRX2 receptor in humans than in mice

(40). The difficulties of translating findings with Mrgprb2 – the murine orthologue of MRGPRX2 – into humans and a clinical setting has been addressed elsewhere ⁽⁷⁵⁾ and probably reflect adaptive evolution of the human MRGPRX2 gene ⁽⁷⁹⁾. Second, results in LAD-2 cells are sometimes inconsistent and therefore difficult to interpret or translate (41, 43, 77, 80-82). The reason for these observations could relate to the observation that LAD-2 cells express variable levels of MRGPRX2 as compared to human mature skin MCs (81, 83, 84) and that LAD-2 cells show different responsiveness to MRGPRX2 agonists as compared to primary human MCs (81, 84-86). In addition, the research groups often use different concentrations which make interpretation of the results difficult (43, 81). Besides, the LAD-2 cells demonstrate a relatively slow growth rate and express very low levels of tryptase and chymase (83). Furthermore, the cytokine generation is deficient in the cells, making the LAD-2 cells only intermediately differentiated as compared to human mature skin MC (83, 87). In addition to the disadvantage regarding the MRGPRX2-pathway, the LAD-2 cells are also not always suitable for studying the IgE-dependent activation as LAD-2 cells grown for prolonged periods display excessive clumping or slower growth, may lose responsiveness to biotinylated IgE/FcεRI crosslinking, and have reduced activation and degranulation (87). As recently described, the HMC-1 cell line also expresses the MRGPRX2 although at lower levels compared to LAD-2 and human cultured MCs and with lower reactivity (84). Because of these drawbacks there is a need for a representative human MC model to study simultaneously the sIgE/FcεRI- and MRGPRX2-dependent activation. Finally, another important aspect to mention is the surface expression of particular GTP-binding proteins that might alter ligand affinity of the MRGPRX2 receptor. Not well indicated in the McNeil paper (40) was that their results required a specific co-transfection of HEK cells with a specific GTP-binding protein. This suggest that not only the MRGPRX2 expression levels are important, but also will be the profile of GTP-binding proteins being expressed.

Primary human mast cells

Culture techniques

At the beginning, different research groups developed strategies to culture primary human MCs starting from mononuclear cells isolated out of different source materials such as cord blood (CB), bone marrow (BM), fetal liver cell or rarely out of peripheral

blood (PB) (34-38, 88-94). However, starting from mononuclear cells can be challenging and pose some difficulties. First, the most used sources are CB and fetal liver cells, which are not easily and commonly available. Second, the recovery of MCs out of mononuclear cells is low and cultures are often contaminated with other cells (37, 89). Next, most of the obtained MCs do not express chymase (92-94). Circumventing these disadvantages, several research groups have developed protocols for culturing sizeable numbers of primary MCs from different progenitor cells.

Table 5 summarizes the main culture protocols starting from progenitor cells mainly obtained from BM, CB and PB. Bone marrow and CB contain high amounts of progenitors (95). However, as already indicated, BM and CB are not easily accessible. Furthermore, CB derived MCs (CBDMCs) are often immature with a low expression of FcεRI and CD203c, contain little histamine and are poorly dedicated for IgE-mediated activation (33, 38, 87, 96). On the other hand, PB derived MCs (PBCMCs) are more easily and repeatedly accessible making the (PBCMC) cultures probably the preferred method to study MCs function. As shown in Figure 2, PBCMCs, cultured for 4 weeks in the presence of SCF and IL-3, are characterized by CD117 and CD203c expression (62). Besides, these PBCMCs uniformly express other membrane markers as the FcεRI, CD300a and CD32. MRGPRX2 is partially expressed, creating two distinctive subpopulations, MRGPRX2⁺ and MRGPRX2⁻ cells. PBCMCs also contain intracellular markers like histamine, tryptase and chymase and thereby displaying a MC_{TC}-like phenotype. However, as already indicated in table 5, there exist different culture protocols for PBCMCs and it cannot be excluded that MC obtained from different cultures display phenotypic and functional dissimilarities, mainly because of considerable differences in culture time (3-12 weeks) and in maturation protocols (different growth medium, growth factors and cytokine mixtures). The cultures that are already used to study IgE-mediated allergies or MRGPRX2 mediated IDHRs are indicated with an asterisk in table 5. Note that the outcomes of pMAT and dMAT might differ markedly depending on whether MC_T or MC_{TC}-like cells are used. Our experience is mainly based on MC_{TC}-like cells that includes both a MRGPRX2⁺ and a MRGPRX2⁻ subpopulation enabling comparative analyses. For a comparison between the phenotype of MCs and basophils the reader is referred elsewhere (97).

Passive mast cell activation tests (pMATs): clinical utility in IgE-mediated allergy

The utility of the pMAT that uses passively sensitized human cultured MCs to study allergen-specific IgE-mediated MC activation, although nascent, is promising. In the study by Bahri et al (10), MCs derived from CD117⁺CD34⁺ human progenitor cells and sensitized with sera from patients allergic to peanut, grass pollen, and wasp venom allergy demonstrated allergen-specific degranulation by quantification of CD63 and CD107a, and release of prostaglandin D2 and β -hexosaminidase. Moreover, for peanut, pMAT conferred superior diagnostic accuracy compared to sIgE, SPTs and BAT in distinguishing between patients with and without clinical reactivity. Findings also implied that MC responsiveness is not exclusively dependent on the sIgE titer, but also regulated by additional elements such as avidity or IgE reactivity and specificity patterns. Likely, the sensitivity of the pMAT also depends on the Fc ϵ RI receptor expression levels, the cell's intrinsic sensitivity to aggregation of Fc ϵ RI and the sIgE to total IgE ratio. Two other relevant articles about in vitro MC activation relate to the exploration of interleukin (IL)-33 as a primer of individual cell responsiveness (15, 22). Synthetic analyses reveal that this cytokine – which is produced on epithelial contact by an allergen (review in (98)) – potentiate IgE-mediated MC responses by both increasing the number of responding cells and enhancing the responsiveness of individual cells. A similar, but less pronounced priming effect was seen with IL-6 (15) and SCF (99). Note that IL-33 and IL-6 did not potentiate MRGPRX2-dependent MC activation (15), while SCF can dampening MRGPRX2-dependent activation.

The reference standard for diagnosis of drug allergy is a controlled drug challenge with administration of incremental doses of drug or placebo. However, such challenges are not devoid of risk and sometimes almost impossible to be performed because of the resources required to manage the pharmacologic effects of some drugs (e.g. curarizing neuromuscular blocking agents) (100, 101). Furthermore, their predictive value is not absolute (102). Therefore, drug hypersensitivity reactions are usually diagnosed using detection of drug-reactive sIgE, skin tests (STs), and by BAT (103-105). However, as reviewed elsewhere (106), the sIgE assay is only available for a limited number of drugs and show poor performance for some drug classes (e.g. β -lactams). Besides, skin testing

is still associated with some uncertainties, especially for nonspecific histamine releasers that might act via off-target MRGPRX2 occupation (e.g. opiates and quinolones) (104, 107-109). The BAT has emerged as a performant diagnostic tool for some, but certainly not all, drug(s) (classes) and sensitivity and specificity is highly variable between the different drug(s) (classes) (103, 105, 110-112). As already mentioned, the BAT is left with some limitations, these limitations, together with the search for a better understanding of the mechanisms of IgE-independent IDHRs, has urged us to explore the potential of pMAT in this domain. Although, currently limited to a single proof of concept with the biguanide antiseptic chlorhexidine (16), it appears that the pMAT cannot only be successfully applied to proteinaceous allergens such as pollen (10, 15), food (10, 11) and Hymenoptera venom (10), but also to drugs. Actually, from our proof of concept it appears that the pMAT has a high analytical sensitivity enabling to depict low titers of functionally active drug-reactive sIgE antibodies and that the pMAT discriminates between allergy, and merely sensitization as reflected by an isolated positive sIgE result. Unlike PBCMCs passively sensitized with sera from CHX-allergic patients with positive STs and BATs, PBCMCs passively sensitized with sera from CHX-sensitized individuals with negative STs and BATs, did not respond to subsequent incubation with the antiseptic. Finally, the pMAT can also be used to study the cross-reactivity profiles of CHX-reactive sera. As shown in Figure 3, and already demonstrated by Mueller-Wirth et al (113), this cross-reactivity profile can involve the structural homologues alexidine and octenidine.

Passive mast cell activation tests (pMATs) in IgE-mediated allergy: more than a diagnostic

As indicated in the title, we expect that the pMAT might also become a valuable asset to study many domains of IgE-mediated allergies that remain poorly understood. An example of such an application of the pMAT lies in its capability to detect and identify major allergens and antibody recognition structures, as well as the affinity between epitope and paratope. For example, Hemmings et al (114), showed that Ara h 2, a major allergen from peanut (*Arachis hypogaea*), induce greater maximal MC reactivity and has a lower half maximal effective concentration than Ara h 6 when testing co-sensitized patients. Recently, and as shown in Figure 4, we applied pMAT to identify Can s 2

(profilin) and Can s 5 (a Bet v 1 homologue) as new allergens of *Cannabis sativa* (115). As indicated in the introductory paragraph, basophil and MC activation/degranulation can result from distinct IgE-dependent and IgE-independent mechanisms. Therefore, it is likely that comparative analyses of dMAT and pMAT experiments, along with BATs (110), might advance our understandings in the mechanisms that govern effector cell activation and degranulation. For example, evidence has emerged that IgE-mediated activation is not only achievable by traditional allergens, but can also occur by lectins with a binding specificity that matches the glycosylation of IgE and/or FcεRI, or by other molecules such as super allergens (protein Fv, HIV gp120) or *S. mansoni* IPSE/alpha-1 (116, 117). How dMAT and pMAT can contribute to the exploration of mechanistic endotypes of IDHRs and discriminate between clinically indistinguishable IgE-dependent and MRGPRX2-dependent reactions is discussed in the section below.

The autoimmune MAT (aMATs) in autoimmune urticaria

Acute and chronic histaminergic urticaria and angioedema result from MC and basophil degranulation via diverse innate and adaptive immune responses, including auto-immune processes (118, 119). At present, in chronic spontaneous urticaria, two groups of MC degranulating signals have been identified, that is IgE autoantibodies to auto-allergens and IgG autoantibodies that target FcεRI or IgE/FcεRI complexes present on the MC surface. The presence of such anti-IgE or anti-FcεRI anti-bodies can be assessed functionally using patients' sera in an autologous serum skin test (ASST), via autoimmune BAT and by MAT (18, 120-122). Figure 5 shows our proof of concept of the aMAT in which the presence of anti-FcεRI and/or anti-IgE autoantibodies or IgE autoantibodies towards auto-allergens is depicted via FCM.

Mast cell activation tests: clinical utility in MRGPRX2-mediated allergy

As earlier mentioned, the recognition of off-target occupation of the MRGPRX2 MC receptor heralds a new and attractive domain in our understandings of IgE-independent IDHRs. However, as evidence for this pathway has mainly been gathered in mutated animals and via *in vitro* studies using different cell lines (40, 76), it remains uncertain

whether findings also apply in humans (55). For the time being, there is little direct clinical evidence for a MRGPRX2-dependent mechanism of IDHRs in humans. To the best of our knowledge, description of patients who might possibly have suffered from a MRGPRX2-dependent reaction is restricted to a single series of patients who experienced anaphylaxis from rocuronium (123). In these patients, in contrast to those who had suffered from a documented IgE-dependent rocuronium anaphylaxis, diagnosis could only be established by a positive ST (quantification of sIgE and BAT rocuronium were negative). In other words, MRGPRX2-dependent reactions are undetectable by traditional BAT. The reason for this has to be sought in the fact that traditional BAT uses resting peripheral blood basophils that barely express MRGPRX2 (about 5%) (56, 124). In contrast to isolated basophils that express MRGPRX2 in about 13-23% of the cells (125). Whether the MAT that uses non-passively sensitized cells (LAD-2 or MC_{TC}-like donor MCs) could advance diagnosis of MRGPRX2-dependent reactions seems unlikely, because drugs acting as MRGPRX2 agonists are already active via wild-type MRGPRX2 (Figure 6). In other words, for the time being there is no method that enables to unambiguously and directly document a MRGPRX2-dependent IDHR. However, the combination of positive STs with negative quantification of sIgE, BAT and eventually MAT, albeit indirectly, could point to an MRGPRX2 reaction for drugs known to be potent MRGPRX2 agonists (2, 110, 123). Figure 7 shows an integrated approach using STs, sIgE immunoassays, BAT, MAT/pMAT and MRGPRX2 silencing experiments to deepen our insights in the pathomechanisms of IgE- and MRGPRX2-dependent IDHRs. As cutaneous MCs express significant numbers of MRGPRX2, there is a general belief that STs to drugs that might act via this receptor do not discriminate between both these endotypes of IDHRs. However, we observed that sIgE- and MRGPRX2-dependent reactions might show different ST patterns (123). In patients with a combined positive IgE to morphine and a positive rocuronium BAT suggestive of an IgE-dependent mechanism, skin MCs appeared to be more sensitive. Actually, in 30/36 (83%) such patients STs were generally positive in SPTs or intradermal test (IDT) dilutions beyond the concentrations required to trigger non-specific ST responses. In contrast, in patients with negative sIgE to morphine and negative rocuronium BAT, STs were positive in only 12/25 (48%) of patients in SPT settings, and for IDTs, higher non-irritative concentrations were

required to reach positivity. Whether this observation could reflect topographic differences in MRGPRX2 expression by cutaneous MCs remains elusive.

Conclusions and perspectives

FCM-assisted BATs have emerged as a robust tool to diagnose and monitor allergic diseases. However, the BAT has some limitations; it requires fresh samples and the technique does not advance diagnosis in patients showing a non-responder status of their cells. Besides, resting basophils barely express MRGPRX2, making it difficult to study this mechanism. In contrast, MATs use serum or plasma samples that can be frozen, stored, and shipped to a reference center experienced in MC lines and/or cultures, and allow (deferred) batch testing with necessary quality controls. In addition, the MCs express MRGPRX2, allowing to study the potential of drugs to stimulate MCs through this receptor. Therefore, the MAT can be a promising diagnostic approach, especially in difficult cases where traditional tests are unavailable or yield uncertain results. However, there is far to go from the current experience to the creation of a new robust diagnostic and its entrance in mainstream use. For example, for pMAT, further validation will need exploration of the effects of FcεRI receptor expression levels, the cell's intrinsic sensitivity to aggregation of FcεRI and the sIgE-to-total IgE ratio. All parameters that are likely to depend on MC source and culture conditions. With respect to MRGPRX2 signaling, it will be critical to know more than its expression levels, but also deepen our insights in the effects of GTP-binding protein surface expression on its ligand specificity and affinity. Nevertheless, although currently restricted to a proof of concept and some preliminary results, the pMAT appears to be highly sensitive and has a good discrimination performance between true drug allergy and cases with indeterminate diagnostics (e.g. sensitization as reflected by an isolated positive sIgE result). Admittedly, additional larger studies are required to confirm and critically verify these encouraging results and to assess whether the technique could become a cost-effective new diagnostic. Further automation of data analyses and bioinformatic tools should advance standardization and quality assurance and thus accelerate transition to the clinics. Although the non-diagnostic applications of MAT are still in their infancy, with increasing employment, it is expected that the technique might become an attractive and valuable asset to study various domains of MC activation/degranulation biology, especially when

used in combination with other tools such as IgE binding studies, skin testing and BATs. Actually, the MAT, could become an attractive complementary method to further accelerate our insight into the effector mechanisms of IDHRs, benefit identification of antibody recognition sites, expand our understandings of desensitization and tolerance induction strategies, predict natural disease courses and prognosis.

Table 1: Basophil activation tests vs. passive mast cell activation tests

Characteristic	Basophil activation test (BAT)	Mast cell activation test (pMAT)
Source material(s)	Patients' whole blood Basophils	Donor MCs or MC lines Patients' sera/plasma
Non-responders	Up to 10-15% and unpredictable ^(c)	No (MC responsiveness can be verified in advance) ^(f)
Priming	IL-3 (detrimental for CD203c) ^(d)	IL-33 ^(g)
Used readouts	CD63, CD107a, CD203c, avidin, DAO ^(e)	CD63, CD107a, CD203c, avidin, DAO
Early analyses post index event possible? ^(a)	Yes	Yes
Deferred testing possible? ^(b)	No	Yes
Batch testing	No	Yes ^(h)

^(a) BATs, and likely also pMATs, can be performed early after the index reaction (unpublished data). However, if the test is negative it should be repeated.

^(b) Although BATs can be performed up to 24 hours after sampling, it is recommended to perform analysis within 4 hours after collection (2). In contrast, patients' sera or plasma that are used in the pMAT can easily be stored and shipped and thereby enable deferred analyses.

^(c) Up to 10-15% of the patients have non-responsive basophils that do not react to positive control nor allergen. In these patients the BAT does not advance diagnosis. Note that this unresponsiveness cannot be determined in advance. It is an ad hoc determination. In non-responders in the BAT, the pMAT can be a valuable adjunct (11).

^(d) IL-3 priming induces expression of the ectoenzyme CD203c without exposition to allergen and can therefore blunt the responsiveness of the cells to allergen. This does not apply to the lysosomal associated membrane protein (LAMP) 3 or CD63.

^(e) For a detailed description of the characteristics of the different readout families the reader is referred elsewhere (2) and figure 1.

^(f) In contrast to BAT, that uses patients' basophils, the pMAT uses a MC cell line or cultured donor cells.

Responsiveness of these cells can readily be evaluated before starting the analyses.

^(g) IL-33 has been shown to boost MC responsiveness to sIgE/FcεRI cross-linking but not MRGPRX2-dependent activation (15, 22).

^(h) pMAT allows simultaneous testing of several patients' sera or plasma using the same MC line or donor MC culture.

Table 2: humanized MC lines used for the detection of IgE-mediated activation

Humanized cell line	FcεRI chain expressed	Allergen	Read out	Reference
RBL SX-38	α , β^1 , γ	Peanut, grass pollen, Timothy grass	Radiolabeled serotonin β -hexosaminidase	(49, 67) (47, 51)
RS-ATL8	α , β , γ	Wide variety (e.g. <i>Holcus lantanus</i> , grass pollen)	Chemiluminescence (Luciferase)	(68, 70, 71)
RBL NFAT DsRed	α^2	Wide variety of allergens	Chemiluminescence (DsRed)	(69, 71)
RBL-T8	α^2	House dust mite	Calcium Influx (flow cytometric)	(65)
RBL-48	α^2	House dust mite, latex, ragweed, cedar pollen	Histamine release β -hexosaminidase surface plasmon resonance imaging	(48, 52, 66)
RBL-hEla-2B12	α^2	House dust mite	β -hexosaminidase	(50)
¹ Ali et al showed loss of transfected β -chain in RBL SX-38 (126)				
² Expression of the human α chain is sufficient to confer reactivity toward human IgE (65)				
DsRed: <i>Discosoma</i> sp. red fluorescent protein				

Table 3: Overview of human mast cell lines and their use in the diagnose of an IgE-dependent allergy

	FcεRI	SCF dependence	Mutation in KIT receptor	Used in diagnosis of IgE- dependent allergy	Read out	Allergen	Reference
HMC1	No	No	Yes	No	/	/	
LUVA	Yes*	No	No	No	/	/	
LAD-2	Yes	Yes	No	Yes	Flow cytometry (CD63)	Peanut	(11, 58)
ROSA ^{kit wt}	Yes	Yes	No	No	/	/	
ROSA ^{kit D816V}	Yes	No	Yes	No	/	/	
iPSC-MC	Yes	Yes	No	Yes	Histamine β-hexosaminidase	Cedar pollen, Mite, House dust	(39)

* The expression of FcεRI on LUVA cells is sometimes contradictorily.

Table 4: MRGPRX2-expressing cell lines used to explore the MRGPRX2-activating potential of drugs

Cell line	Read out	Opiates	NMBA	(fluoro)quinolones	Antibiotics	Miscellaneous	Reference
CHO-K1-MRGPRX2	Calcium	+ Morphine	ND	ND	ND	ND	Akuzawa(127)
LAD-2	Histamine release	+ Morphine	+ Rocu + Atra - Succ	+ Moxi	ND	+Octreotide + Mastoparan	McNeil(40)
HEK-293	Cytokine release			+ Cipro			
	Calcium			+ Levo			
	β -hexosaminidase			+ Olfoxacin			
LAD-2	Calcium	ND	+ Atra	+ Cipro	ND	ND	Azimi(41)
	β -hexosaminidase						
HEK-293	Calcium	ND	ND	ND	+ Vancomycin	ND	Azimi(42)
LAD-2	β -hexosaminidase						
LAD-2	Calcium β -hexosaminidase	+ Morphine	- Rocu - Atra	- Moxi	ND	- Octreotide + Mastoparan	Lansu(43)
HEK-293				- Cipro			
HTLA				- Levo			
				- Olfoxacin			
LAD-2	Calcium	ND	ND	+Moxi +Cipro	ND	ND	Han(82)
	Histamine release			+Norfloxacin +Lomefloxacin			
LAD-2	β -hexosaminidase	+ Morphine	- Rocu + Cisatra	ND	- Vancomycin	ND	Navines-Ferrer(44)
	Flow cytometry				- Teicoplanin - Amoxicillin		

(CD63)							
LAD-2	β-hexosaminidase						
	Histamine release		+Cisatra				Che(128, 129)
	TNF-α		+Mivacurim				
	Calcium						
LAD-2 HEK-293	Calcium			+ Moxi			
				+Levo			
				+Sparfloxacin			
				+Gatifloxacin			
	β-hexosaminidase	ND	ND	+Enoxacin	ND	ND	Liu(77)
	Histamine release			+Norfloxacin			
				+Lomefloxacin			
				+Pefloxain			
				+Fleroxacin			
LAD-2 HEK-293	β-hexosaminidase						
	Histamine release	+ Morphine	ND	ND	ND	ND	Zhang(130)
	TNF-α						
	Calcium						
LAD-2	β-hexosaminidase	ND	+Atra	ND	ND	ND	Shtessel(131)
LAD-2	Calcium		+/-Rocu				
	β-hexosaminidase	ND	+Atra	ND	ND	ND	Fernandopulle(80)

CCL2 release

LAD-2	β -hexosaminidase	ND	+ Rocu	ND	ND	ND	Chompunud(81)
+, MCs activation/degranulation; -, no MCs activation/degranulation; ND, not determined; NMBA, neuromuscular blocking agent; rocu, rocuronium; atra, atracurium; cisatra, cisatracurium; moxi, moxifloxacin; Cipro, ciprofloxacin; Levo, levofloxacin; Succ, Succinylcholine							

Table 5: Primary human mast cell protocols

Source	Progenitors	Duration (weeks)	Cytokines	Serum factor	Medium	FcεRI	Tryptase (%)°	Chymase (%)°	Reference
PB	CD34 ⁺	6	SCF, IL-3	Autologous serum	RPMI	+	100	< 3	Rottem (132)
PB	CD34 ⁺	10	SCF, IL-3, IL-6	Serum free	StemPro-34	ND	+	ND	Kirshenbaum (133, 134)
PB	CD34 ⁺	12	SCF, IL-3, IL-6	FBS	Methylcellulose	ND	100	10	Wang (23)
PB	CD34 ⁺	8	SCF, IL-3, IL-6	FBS	Methylcellulose	+	ND	ND	Saito (135)
PB	CD34 ⁺	9	SCF, IL-3, IL-6, IL- 9	Serum-free	StemSpan	+	99	95	Lappalainen (24)
PB	CD133 ⁺	7	SCF, IL-3, IL-6	FBS	StemSpan	+	80	ND	Holm (25)
PB	CD34 ⁺	8	SCF, IL-3, IL-6	Serum free	StemPro-34	ND	ND	ND	Radringer (87)
PB	CD113 ⁺	7	SCF, IL-3, IL-6	Serum free	StemSpan	+	100	< 2	Frandsen (26)
PB	CD133 ⁺	8	SCF, IL-6	Serum free	IMDM + GlutaMAX	+	+	-	Gaudenzio (136)
PB	CD34 ⁺	3	SCF, IL-3, LDL	Serum free	Methylcellulose	+	+	4-20	Schmetzer (27)
PB	CD133 ⁺	8	SCF, IL-3, IL-6, IL- 4	Serum free	StemSpan	+	ND	ND	Hoffman (137)
PB	CD34 ⁺	9	SCF, IL-3, IL-6	Serum free	StemSpan-IMDM	+	95	95	* Joulia (19)
PB	CD34 ⁺	6	SCF, IL-3, IL-6	Serum free	StemPro-34 medium	+	ND	ND	Yin (21)
PB	CD34 ⁺	6	SCF, IL-3, IL-6, IL- 9, IL-4	Serum free	IMDM	+	100	71	** Tam (28)
PB	CD34 ⁺	4	SCF, IL-3, LDL	Serum free	Methylcellulose	+	96	90	*, **,*** Cop

									(54)
PB	CD133 ⁺	7	SCF, IL-3, IL-6	FBS	StemSpan/Collagen Matrix	+	100	97	Derakhshan (29)
PB	CD34 ⁺	5	SCF, IL-3, IL-6	Serum free	StemSpan	ND	ND	ND	Luo (138)
PB	CD117	8	SCF, IL-3, IL-6, LDL	Serum free	StemSpan-IMDM	+	+	+	* Bahri (10, 139)
CB	CD34 ⁺	34	SCF, IL-3	Serum free	IMDM	ND	ND	ND	Durand (140)
CB	CD34 ⁺	10	SCF, IL-3, IL-6, PGE2	FBS	IBL media I	+	99	18	Saito (30)
CB	CD34 ⁺	8	SCF, IL-3, IL-6	FBS	Alfa-medium	ND	+	ND	Kempuraj (141)
CB	CD133 ⁺	12	SCF, IL-3, IL-6	Serum free	StemSpan	+	83	36	Dahl (31)
CB	CD34 ⁺	8	SCF, IL-6	Serum free	QSBF60	+	95	9	Yamaguchi (32)
CB	CD133 ⁺	7	SCF, IL-3, IL-6	FBS	StemSpan	+	75	ND	Anderson (33)
CB	CD133 ⁺	8	SCF, IL-3, IL-6	FBS	StemSpan	ND	ND	ND	Radringer (87)
BM	CD34 ⁺	6	SCF, IL-3	Autologous serum	RPMI	+	100	< 3	Rottem (132)
BM	CD34 ⁺	12	SCF, IL-6	Human serum	Media III	+	100	100	Shimizu (142)
BM	CD34 ⁺	10	SCF, IL-3, IL-6	Serum free	StemPro-34	ND	+	ND	Kirshenbaum (133, 134)

PB, peripheral blood; CB, cord blood; BM, bone marrow; SCF, stem cell factor; IL, interleukine; FBS, fetal bovine serum; IMDM, Iscove's Modified Dulbecco's Medium ;ND, not determined; °, if the number of tryptase or chymase positive cells is not mentioned, it is designated as positive (+) or negative (-); * culture used for pMAT; ** cultured used for aMAT; *** cultured used for dMAT

Figure 1: Mast cell Identification and activation/degranulation markers. Mast cells express various molecules that can be used in isolation or in combination to identify the cells and measure their activation/degranulation status by flow cytometry (FCM) in a technique designated as the MAT. Mast cell activation/degranulation can occur via IgE/FcεRI-dependent and IgE/FcεRI-independent pathways. Individual cell activation/degranulation can be measured by FCM on 5 levels: (1) via appearance or upregulation of surface markers (such as CD63, CD107a, CD63, and avidin binding of membrane-associated exteriorized anionic proteoglycans released from granules); (2) Newly synthesized chemokines such CCL4 or CXCL8 can be trapped and measured intracellularly; (3) phosphotyrosine staining, a parameter associated to cell protein kinase function, can also be used to study MCs activation; (4) via changes in mediator content (such as decrease of histamine); or (5) via increased intracellular calcium staining. Intracellular molecules are denoted in pink.

Figure 2: Representative plot of flow cytometric phenotyping of peripheral blood cultured human mast cells. Cells were selected based on side scatter (SSC) and forward scatter (FSC). MC were defined as CD117 and CD203c positive. These MC uniformly express FcεRI, CD300a and CD32. MRGPRX2 is partially expressed on MC creating two clearly subpopulations MRGPRX2⁺ and MRGPRX2⁻. CD63 is barely expressed on resting MC. The MCs contain also tryptase, chymase and histamine (DAO).

Figure 3: Representative plot of utilization of the indirect mast cell activation test to study cross-reactivity between structural homologues. MCs were passively sensitized with serum of a patient with documented CHX-allergy (sIgE: 0.46 kUa/L, FEIA ImmunoCAP, Thermo Fisher together with positive STs and basophil activation tests CHX). MCs are stimulated with CHX (2.8 μM) or with a structural homologue alexidine (2.8 μM).

Figure 4: Representative plot of a mast cell activation test with recombinant homologue of the major birch pollen allergen Bet v 1 from Cannabis sativa (rCs-Bet v 1 homologue) **(A)** or cannabis sativa profiling **(B)**. Unlike cells not passively sensitized with a rCs-Bet v 1 or Cs-profilin reactive serum (top), passively sensitized show a dose-dependent degranulation to rCs-Bet v 1 or Cs-profilin.

Figure 5: Representative plots of an autoimmune mast cell activation test (aMAT) for suspected autoimmune chronic spontaneous urticaria. After incubation with control serum there is no up-regulation of CD63 (aMAT: negative). After incubation of cells with serum of a patient, 40% of the mast cells up-regulate expression of CD63.

Figure 6: Direct mast cell (MC) activation by MRGPRX2 agonists substance P (natural ligand), the opiate morphine and the fluoroquinolone moxifloxacin. Note that, unlike reported by McNeil (40), rocuronium does not induce degranulation of primary human MCs. Silencing of MRGPRX2-dependent degranulation by DsiRNA (55).

Figure 7: exploring IgE- and MRGPRX2-dependent immediate drug hypersensitivity.

Proposal for an integrated approach using skin tests, IgE-immunoassays, basophil activation tests, direct/passive mast cell activation tests, and selective MRGPRX2 silencing to deepen pathomechanistic insights in IgE- and MRGPRX2-dependent immediate drug hypersensitivity. Note that starting from CD34⁺ progenitors, it is possible to obtain a MRGPRX2⁺ and MRGPRX2⁻ population that can be studied separately using multicolor flow cytometry. Ultimately, this approach could advance development of novel diagnostic.

References

1. Hemmings O, Kwok M, McKendry R, Santos AF. Basophil Activation Test: Old and New Applications in Allergy. *Current allergy and asthma reports*. 2018;18(12):77.
2. Ebo DG, Bridts CH, Mertens CH, Sabato V. Principles, potential and limitations of ex vivo basophil activation by flow cytometry in allergology: a narrative review. *J Allergy Clin Immunol*. 2020.
3. Hoffmann HJ, Santos AF, Mayorga C, Nopp A, Eberlein B, Ferrer M, et al. The clinical utility of basophil activation testing in diagnosis and monitoring of allergic disease. *Allergy*. 2015;70(11):1393-405.
4. Sturm GJ, Kranzelbinder B, Sturm EM, Heinemann A, Groselj-Strele A, Aberer W. The basophil activation test in the diagnosis of allergy: technical issues and critical factors. *Allergy*. 2009;64(9):1319-26.
5. Valent P, Bettelheim P. Cell surface structures on human basophils and mast cells: biochemical and functional characterization. *Adv Immunol*. 1992;52:333-423.
6. Valent P, Schernthaner GH, Sperr WR, Fritsch G, Agis H, Willheim M, et al. Variable expression of activation-linked surface antigens on human mast cells in health and disease. *Immunol Rev*. 2001;179:74-81.
7. Saito H, Ebisawa M, Sakaguchi N, Onda T, Iikura Y, Yanagida M, et al. Characterization of cord-blood-derived human mast cells cultured in the presence of Steel factor and interleukin-6. *Int Arch Allergy Immunol*. 1995;107(1-3):63-5.
8. Sellge G, Bischoff SC. Isolation, culture, and characterization of intestinal mast cells. *Methods Mol Biol*. 2006;315:123-38.
9. Patella V, Marino I, Lamparter B, Arbustini E, Adt M, Marone G. Human heart mast cells. Isolation, purification, ultrastructure, and immunologic characterization. *J Immunol*. 1995;154(6):2855-65.
10. Bahri R, Custovic A, Korosec P, Tsoumani M, Barron M, Wu J, et al. Mast cell activation test in the diagnosis of allergic disease and anaphylaxis. *J Allergy Clin Immunol*. 2018;142(2):485-96.e16.
11. Santos AF, Couto-Francisco N, Bécares N, Kwok M, Bahnson HT, Lack G. A novel human mast cell activation test for peanut allergy. *J Allergy Clin Immunol*. 2018;142(2):689-91.e9.
12. Passante E. Mast Cell and Basophil Cell Lines: A Compendium. *Methods Mol Biol*. 2020;2163:127-44.
13. Kleine Budde I, de Heer PG, van der Zee JS, Aalberse RC. The stripped basophil histamine release bioassay as a tool for the detection of allergen-specific IgE in serum. *Int Arch Allergy Immunol*. 2001;126(4):277-85.

14. Larsen LF, Juel-Berg N, Hansen KS, Clare Mills EN, van Ree R, Poulsen LK, et al. A comparative study on basophil activation test, histamine release assay, and passive sensitization histamine release assay in the diagnosis of peanut allergy. *Allergy*. 2018;73(1):137-44.
15. Cop N, Ebo DG, Bridts CH, Elst J, Hagendorens MM, Mertens C, et al. Influence of IL-6, IL-33, and TNF- α on human mast cell activation: Lessons from single cell analysis by flow cytometry. *Cytometry B Clin Cytom*. 2018;94(3):405-11.
16. Elst J, van der Poorten MM, Faber MA, Van Gasse AL, Garvey LH, Bridts CH, et al. Mast cell activation test in chlorhexidine allergy: a proof of concept. *British journal of anaesthesia*. 2020.
17. Derakhshan T, Bhowmick R, Meinkoth JH, Ritchey JW, Gappa-Fahlenkamp H. Human Mast Cell Development from Hematopoietic Stem Cells in a Connective Tissue-Equivalent Model. *Tissue Eng Part A*. 2019;25(21-22):1564-74.
18. Tam IYS, Lau HYA, Tam SY, Lee TH. Mast cell activation test using patient-derived mast cells exhibits distinct combinatorial phenotypic profiles among allergic patients. *Allergy*. 2020;75(7):1796-9.
19. Joulia R, Gaudenzio N, Rodrigues M, Lopez J, Blanchard N, Valitutti S, et al. Mast cells form antibody-dependent degranulatory synapse for dedicated secretion and defence. *Nat Commun*. 2015;6:6174.
20. Gaudenzio N, Sibilano R, Marichal T, Starkl P, Reber LL, Cenac N, et al. Different activation signals induce distinct mast cell degranulation strategies. *J Clin Invest*. 2016;126(10):3981-98.
21. Yin Y, Bai Y, Olivera A, Desai A, Metcalfe DD. An optimized protocol for the generation and functional analysis of human mast cells from CD34(+) enriched cell populations. *J Immunol Methods*. 2017;448:105-11.
22. Joulia R, L'Faqihi FE, Valitutti S, Espinosa E. IL-33 fine tunes mast cell degranulation and chemokine production at the single-cell level. *J Allergy Clin Immunol*. 2017;140(2):497-509.e10.
23. Wang XS, Sam SW, Yip KH, Lau HY. Functional characterization of human mast cells cultured from adult peripheral blood. *Int Immunopharmacol*. 2006;6(5):839-47.
24. Lappalainen J, Lindstedt KA, Kovanen PT. A protocol for generating high numbers of mature and functional human mast cells from peripheral blood. *Clin Exp Allergy*. 2007;37(9):1404-14.
25. Holm M, Andersen HB, Hetland TE, Dahl C, Hoffmann HJ, Junker S, et al. Seven week culture of functional human mast cells from buffy coat preparations. *J Immunol Methods*. 2008;336(2):213-21.
26. Frandsen PM, Krohn IJ, Hoffmann HJ, Schiotz PO. The Influence of IgE on Cultured Human Mast Cells. *Allergy Asthma Immunol Res*. 2013;5(6):409-14.

27. Schmetzer O, Valentin P, Smorodchenko A, Domenis R, Gri G, Siebenhaar F, et al. A novel method to generate and culture human mast cells: Peripheral CD34+ stem cell-derived mast cells (PSCMCs). *J Immunol Methods*. 2014;413:62-8.
28. Tam IYS, Ng CW, Tam SY, Lau HYA. Novel six-week protocol for generating functional human connective tissue-type (MCTC) mast cells from buffy coats. *Inflamm Res*. 2017;66(1):25-37.
29. Derakhshan T, Bhowmick R, Ritchey JW, Gappa-Fahlenkamp H. Development of Human Mast Cells from Hematopoietic Stem Cells within a 3D Collagen Matrix: Effect of Stem Cell Media on Mast Cell Generation. *Stem Cells Int*. 2018;2018:2136193.
30. Saito H, Ebisawa M, Tachimoto H, Shichijo M, Fukagawa K, Matsumoto K, et al. Selective growth of human mast cells induced by Steel factor, IL-6, and prostaglandin E2 from cord blood mononuclear cells. *J Immunol*. 1996;157(1):343-50.
31. Dahl C, Saito H, Nielsen HV, Schiotz PO. The establishment of a combined serum-free and serum-supplemented culture method of obtaining functional cord blood-derived human mast cells. *J Immunol Methods*. 2002;262(1-2):137-43.
32. Yamaguchi M, Azuma H, Fujihara M, Hamada H, Ikeda H. Generation of a considerable number of functional mast cells with a high basal level of FcεRI expression from cord blood CD34+ cells by co-culturing them with bone marrow stromal cell line under serum-free conditions. *Scand J Immunol*. 2007;65(6):581-8.
33. Andersen HB, Holm M, Hetland TE, Dahl C, Junker S, Schiotz PO, et al. Comparison of short term in vitro cultured human mast cells from different progenitors - Peripheral blood-derived progenitors generate highly mature and functional mast cells. *J Immunol Methods*. 2008;336(2):166-74.
34. Piliponsky AM, Gleich GJ, Nagler A, Bar I, Levi-Schaffer F. Non-IgE-dependent activation of human lung- and cord blood-derived mast cells is induced by eosinophil major basic protein and modulated by the membrane form of stem cell factor. *Blood*. 2003;101(5):1898-904.
35. Furitsu T, Saito H, Dvorak AM, Schwartz LB, Irani AM, Burdick JF, et al. Development of human mast cells in vitro. *Proc Natl Acad Sci U S A*. 1989;86(24):10039-43.
36. Yanagida M, Fukamachi H, Ohgami K, Kuwaki T, Ishii H, Uzumaki H, et al. Effects of T-helper 2-type cytokines, interleukin-3 (IL-3), IL-4, IL-5, and IL-6 on the survival of cultured human mast cells. *Blood*. 1995;86(10):3705-14.
37. Mitsui H, Furitsu T, Dvorak AM, Irani AM, Schwartz LB, Inagaki N, et al. Development of human mast cells from umbilical cord blood cells by recombinant human and murine c-kit ligand. *Proc Natl Acad Sci U S A*. 1993;90(2):735-9.

38. Iida M, Matsumoto K, Tomita H, Nakajima T, Akasawa A, Ohtani NY, et al. Selective down-regulation of high-affinity IgE receptor (FcεRI) alpha-chain messenger RNA among transcriptome in cord blood-derived versus adult peripheral blood-derived cultured human mast cells. *Blood*. 2001;97(4):1016-22.
39. Igarashi A, Ebihara Y, Kumagai T, Hirai H, Nagata K, Tsuji K. Mast cells derived from human induced pluripotent stem cells are useful for allergen tests. *Allergol Int*. 2018;67(2):234-42.
40. McNeil BD, Pundir P, Meeker S, Han L, Undem BJ, Kulka M, et al. Identification of a mast-cell-specific receptor crucial for pseudo-allergic drug reactions. *Nature*. 2015;519(7542):237-41.
41. Azimi E, Reddy VB, Shade KC, Anthony RM, Talbot S, Pereira PJS, et al. Dual action of neurokinin-1 antagonists on Mas-related GPCRs. *JCI Insight*. 2016;1(16):e89362.
42. Azimi E, Reddy VB, Lerner EA. Brief communication: MRGPRX2, atopic dermatitis and red man syndrome. *Itch (Phila)*. 2017;2(1).
43. Lansu K, Karpiak J, Liu J, Huang XP, McCorvy JD, Kroeze WK, et al. In silico design of novel probes for the atypical opioid receptor MRGPRX2. *Nat Chem Biol*. 2017;13(5):529-36.
44. Navines-Ferrer A, Serrano-Candelas E, Lafuente A, Munoz-Cano R, Martin M, Gastaminza G. MRGPRX2-mediated mast cell response to drugs used in perioperative procedures and anaesthesia. *Sci Rep*. 2018;8(1):11628.
45. MacGlashan DW, Jr. Basophil activation testing. *J Allergy Clin Immunol*. 2013;132(4):777-87.
46. Jensen BM, Bartko EA, Baumann K, Skov PS. Measuring Histamine and Cytokine Release from Basophils and Mast Cells. *Methods Mol Biol*. 2020;2163:247-62.
47. Wiegand TW, Williams PB, Dreskin SC, Jouvin MH, Kinet JP, Tasset D. High-affinity oligonucleotide ligands to human IgE inhibit binding to Fc epsilon receptor I. *J Immunol*. 1996;157(1):221-30.
48. Gilfillan AM, Kado-Fong H, Wiggan GA, Hakimi J, Kent U, Kochan JP. Conservation of signal transduction mechanisms via the human Fc epsilon RI alpha after transfection into a rat mast cell line, RBL 2H3. *J Immunol*. 1992;149(7):2445-51.
49. Dibbern DA, Jr., Palmer GW, Williams PB, Bock SA, Dreskin SC. RBL cells expressing human Fc epsilon RI are a sensitive tool for exploring functional IgE-allergen interactions: studies with sera from peanut-sensitive patients. *J Immunol Methods*. 2003;274(1-2):37-45.
50. Takagi K, Nakamura R, Teshima R, Sawada J. Application of human Fc epsilon RI alpha-chain-transfected RBL-2H3 cells for estimation of active serum IgE. *Biol Pharm Bull*. 2003;26(2):252-5.
51. McDermott RA, Porterfield HS, El Mezayen R, Burks AW, Pons L, Schlichting DG, et al. Contribution of Ara h 2 to peanut-specific, immunoglobulin E-mediated, cell activation. *Clin Exp Allergy*. 2007;37(5):752-63.

52. Yuhki Yanase, Takaaki Hiragun, Tetsuji Yanase, Tomoko Kawaguchi, Kaori Ishii, Nobutaka Kumazaki, Takayuki Obara, Michihiro Hide. Clinical diagnosis of type I allergy by means of SPR imaging with less than a microliter of peripheral blood. *Sensing and Bio-Sensin Research*. 2014;2:43-8.
53. Baumann K, Marcelino J, Skov PS, Santos MCP, Wyrosiak I, Scheffel J, et al. Autologous serum skin test reactions in chronic spontaneous urticaria differ from heterologous cell reactions. *J Eur Acad Dermatol Venereol*. 2021.
54. Cop N, Decuyper, II, Faber MA, Sabato V, Bridts CH, Hagendorens MM, et al. Phenotypic and functional characterization of in vitro cultured human mast cells. *Cytometry B Clin Cytom*. 2017;92(5):348-54.
55. Elst J, Sabato V, Faber MA, Bridts CH, Mertens C, Van Houdt M, et al. MRGPRX2 and Immediate Drug Hypersensitivity: Insights from Cultured Human Mast Cells. *Journal of investigational allergology & clinical immunology*. 2020:0.
56. Elst J, Sabato V, Hagendorens MM, van Houdt M, Faber MA, Bridts CH, et al. Measurement and Functional Analysis of the Mas-Related G Protein-Coupled Receptor MRGPRX2 on Human Mast Cells and Basophils. *Methods Mol Biol*. 2020;2163:219-26.
57. Nishida K, Yamasaki S, Ito Y, Kabu K, Hattori K, Tezuka T, et al. Fc ϵ RI-mediated mast cell degranulation requires calcium-independent microtubule-dependent translocation of granules to the plasma membrane. *J Cell Biol*. 2005;170(1):115-26.
58. Santos AF, James LK, Bahnson HT, Shamji MH, Couto-Francisco NC, Islam S, et al. IgG4 inhibits peanut-induced basophil and mast cell activation in peanut-tolerant children sensitized to peanut major allergens. *J Allergy Clin Immunol*. 2015;135(5):1249-56.
59. Koberle M, Kaesler S, Kempf W, Wolbing F, Biedermann T. Tetraspanins in mast cells. *Frontiers in immunology*. 2012;3:106.
60. Grutzkau A, Smorodchenko A, Lippert U, Kirchhof L, Artuc M, Henz BM. LAMP-1 and LAMP-2, but not LAMP-3, are reliable markers for activation-induced secretion of human mast cells. *Cytometry A*. 2004;61(1):62-8.
61. MacGlashan D, Jr. Expression of CD203c and CD63 in human basophils: relationship to differential regulation of piecemeal and anaphylactic degranulation processes. *Clin Exp Allergy*. 2010;40(9):1365-77.
62. Elst J, Sabato V, van der Poorten MM, Faber M, Van Gasse AL, De Puyssseleir LP, et al. Peripheral blood cultured mast cells: Phenotypic and functional outcomes of different culture protocols. *J Immunol Methods*. 2021;492:113003.

63. Dvorak AM, Morgan ES, Schleimer RP, Lichtenstein LM. Diamine oxidase-gold labels histamine in human mast-cell granules: a new enzyme-affinity ultrastructural method. *J Histochem Cytochem.* 1993;41(6):787-800.
64. Dvorak AM. Ultrastructural localization of histamine in human basophils and mast cells; changes associated with anaphylactic degranulation and recovery demonstrated with a diamine oxidase-gold probe. *Allergy.* 1997;52(34 Suppl):14-24.
65. Taudou G, Varin-Blank N, Jouin H, Marchand F, Weyer A, Blank U. Expression of the alpha chain of human Fc epsilon RI in transfected rat basophilic leukemia cells: functional activation after sensitization with human mite-specific IgE. *Int Arch Allergy Immunol.* 1993;100(4):344-50.
66. Williams PB, Akasawa A, Dreskin S, Selner JC. Respirable tire fragments contain specific IgE-binding and bridging latex antigens. *Chest.* 1996;109(3 Suppl):13S.
67. Porterfield HS, Murray KS, Schlichting DG, Chen X, Hansen KC, Duncan MW, et al. Effector activity of peanut allergens: a critical role for Ara h 2, Ara h 6, and their variants. *Clin Exp Allergy.* 2009;39(7):1099-108.
68. Nakamura R, Uchida Y, Higuchi M, Nakamura R, Tsuge I, Urisu A, et al. A convenient and sensitive allergy test: IgE crosslinking-induced luciferase expression in cultured mast cells. *Allergy.* 2010;65(10):1266-73.
69. Wang X, Cato P, Lin HC, Li T, Wan D, Alcocer MJ, et al. Optimisation and use of humanised RBL NF-AT-GFP and NF-AT-DsRed reporter cell lines suitable for high-throughput scale detection of allergic sensitisation in array format and identification of the ECM-integrin interaction as critical factor. *Molecular biotechnology.* 2014;56(2):136-46.
70. Ali EA, Nakamura R, Falcone FH. Use of Humanized RS-ATL8 Reporter System for Detection of Allergen-Specific IgE Sensitization in Human Food Allergy. *Methods Mol Biol.* 2017;1592:147-61.
71. Wan D, Wang X, Nakamura R, Alcocer MJC, Falcone FH. Use of Humanized RBL Reporter Systems for the Detection of Allergen-Specific IgE Sensitization in Human Serum. In: Gibbs BF, Falcone FH, editors. *Basophils and Mast Cells: Methods and Protocols.* New York, NY: Springer US; 2020. p. 145-53.
72. Falcone FH, Alcocer MJ, Okamoto-Uchida Y, Nakamura R. Use of humanized rat basophilic leukemia reporter cell lines as a diagnostic tool for detection of allergen-specific IgE in allergic patients: time for a reappraisal? *Current allergy and asthma reports.* 2015;15(11):67.
73. Palmer GW, Dibbern DA, Jr., Burks AW, Bannon GA, Bock SA, Porterfield HS, et al. Comparative potency of Ara h 1 and Ara h 2 in immunochemical and functional assays of allergenicity. *Clin Immunol.* 2005;115(3):302-12.

74. Santos AF, James LK, Kwok M, McKendry RT, Anagnostou K, Clark AT, et al. Peanut oral immunotherapy induces blocking antibodies but does not change the functional characteristics of peanut-specific IgE. *The Journal of allergy and clinical immunology*. 2020;145(1):440-3 e5.
75. Azimi E, Reddy VB, Shade KC, Anthony RM, Talbot S, Pereira PJ, et al. Dual action of neurokinin-1 antagonists on Mas-related GPCRs. *JCI Insight*. 2016;1(16):e89362.
76. Porebski G, Kwiecien K, Pawica M, Kwitniewski M. Mas-Related G Protein-Coupled Receptor-X2 (MRGPRX2) in Drug Hypersensitivity Reactions. *Frontiers in immunology*. 2018;9:3027.
77. Liu R, Hu S, Zhang Y, Che D, Cao J, Wang J, et al. Mast cell-mediated hypersensitivity to fluoroquinolone is MRGPRX2 dependent. *Int Immunopharmacol*. 2019;70:417-27.
78. Subramanian H, Gupta K, Ali H. Roles of Mas-related G protein-coupled receptor X2 on mast cell-mediated host defense, pseudoallergic drug reactions, and chronic inflammatory diseases. *The Journal of allergy and clinical immunology*. 2016;138(3):700-10.
79. Yang S, Liu Y, Lin AA, Cavalli-Sforza LL, Zhao Z, Su B. Adaptive evolution of MRGX2, a human sensory neuron specific gene involved in nociception. *Gene*. 2005;352:30-5.
80. Fernandopulle NA, Zhang S, Soeding PF, Mackay GA. MRGPRX2 activation in mast cells by neuromuscular blocking agents and other agonists: modulation by sugammadex. *Clin Exp Allergy*. 2020.
81. Chompunud Na Ayudhya C, Amponnawarat A, Roy S, Oskeritzian CA, Ali H. MRGPRX2 Activation by Rocuronium: Insights from Studies with Human Skin Mast Cells and Missense Variants. *Cells*. 2021;10(1).
82. Han S, Lv Y, Kong L, Che D, Liu R, Fu J, et al. Use of the relative release index for histamine in LAD2 cells to evaluate the potential anaphylactoid effects of drugs. *Sci Rep*. 2017;7(1):13714.
83. Guhl S, Babina M, Neou A, Zuberbier T, Artuc M. Mast cell lines HMC-1 and LAD2 in comparison with mature human skin mast cells--drastically reduced levels of tryptase and chymase in mast cell lines. *Exp Dermatol*. 2010;19(9):845-7.
84. Hermans MAW, van Stigt AC, van de Meerendonk S, Schrijver B, van Daele PLA, van Hagen PM, et al. Human Mast Cell Line HMC1 Expresses Functional Mas-Related G-Protein Coupled Receptor 2. *Frontiers in immunology*. 2021;12:625284.
85. Kulka M, Sheen CH, Tancowny BP, Grammer LC, Schleimer RP. Neuropeptides activate human mast cell degranulation and chemokine production. *Immunology*. 2008;123(3):398-410.
86. Jadkauskaite L, Bahri R, Farjo N, Farjo B, Jenkins G, Bhogal R, et al. Nuclear factor (erythroid-derived 2)-like-2 pathway modulates substance P-induced human mast cell activation and degranulation in the hair follicle. *J Allergy Clin Immunol*. 2018;142(4):1331-3 e8.

87. Radinger M, Jensen BM, Kuehn HS, Kirshenbaum A, Gilfillan AM. Generation, isolation, and maintenance of human mast cells and mast cell lines derived from peripheral blood or cord blood. *Curr Protoc Immunol*. 2010;Chapter 7:Unit 7 37.
88. Valent P, Spanblochl E, Sperr WR, Sillaber C, Zsebo KM, Agis H, et al. Induction of differentiation of human mast cells from bone marrow and peripheral blood mononuclear cells by recombinant human stem cell factor/kit-ligand in long-term culture. *Blood*. 1992;80(9):2237-45.
89. Dvorak AM, Mitsui H, Ishizaka T. Ultrastructural morphology of immature mast cells in sequential suspension cultures of human cord blood cells supplemented with c-kit ligand; distinction from mature basophilic leukocytes undergoing secretion in the same cultures. *J Leukoc Biol*. 1993;54(5):465-85.
90. Nilsson G, Butterfield JH, Nilsson K, Siegbahn A. Stem cell factor is a chemotactic factor for human mast cells. *J Immunol*. 1994;153(8):3717-23.
91. Toru H, Eguchi M, Matsumoto R, Yanagida M, Yata J, Nakahata T. Interleukin-4 promotes the development of tryptase and chymase double-positive human mast cells accompanied by cell maturation. *Blood*. 1998;91(1):187-95.
92. Irani AA, Craig SS, Nilsson G, Ishizaka T, Schwartz LB. Characterization of human mast cells developed in vitro from fetal liver cells cocultured with murine 3T3 fibroblasts. *Immunology*. 1992;77(1):136-43.
93. Kambe N, Kambe M, Chang HW, Matsui A, Min HK, Hussein M, et al. An improved procedure for the development of human mast cells from dispersed fetal liver cells in serum-free culture medium. *J Immunol Methods*. 2000;240(1-2):101-10.
94. Xia HZ, Du Z, Craig S, Klisch G, Noben-Trauth N, Kochan JP, et al. Effect of recombinant human IL-4 on tryptase, chymase, and Fc epsilon receptor type I expression in recombinant human stem cell factor-dependent fetal liver-derived human mast cells. *J Immunol*. 1997;159(6):2911-21.
95. Bender JG, Unverzagt K, Walker DE, Lee W, Smith S, Williams S, et al. Phenotypic analysis and characterization of CD34+ cells from normal human bone marrow, cord blood, peripheral blood, and mobilized peripheral blood from patients undergoing autologous stem cell transplantation. *Clin Immunol Immunopathol*. 1994;70(1):10-8.
96. Jensen BM, Frandsen PM, Raaby EM, Schiotz PO, Skov PS, Poulsen LK. Molecular and stimulus-response profiles illustrate heterogeneity between peripheral and cord blood-derived human mast cells. *J Leukoc Biol*. 2014;95(6):893-901.
97. Varricchi G, Raap U, Rivellese F, Marone G, Gibbs BF. Human mast cells and basophils-How are they similar how are they different? *Immunol Rev*. 2018;282(1):8-34.

98. Liew FY, Girard JP, Turnquist HR. Interleukin-33 in health and disease. *Nature reviews Immunology*. 2016;16(11):676-89.
99. Babina M, Guhl S, Artuc M, Zuberbier T. Allergic FcεRI- and pseudo-allergic MRGPRX2-triggered mast cell activation routes are independent and inversely regulated by SCF. *Allergy*. 2018;73(1):256-60.
100. Bousquet PJ, Gaeta F, Bousquet-Rouanet L, Lefrant JY, Demoly P, Romano A. Provocation tests in diagnosing drug hypersensitivity. *Current pharmaceutical design*. 2008;14(27):2792-802.
101. Garvey LH, Ebo DG, Krøigaard M, Savic S, Clarke R, Cooke P, et al. The use of drug provocation testing in the investigation of suspected immediate perioperative allergic reactions: current status. *British journal of anaesthesia*. 2019;123(1):e126-e34.
102. Demoly P, Romano A, Botelho C, Bousquet-Rouanet L, Gaeta F, Silva R, et al. Determining the negative predictive value of provocation tests with beta-lactams. *Allergy*. 2010;65(3):327-32.
103. Mayorga C, Ebo DG, Lang DM, Pichler WJ, Sabato V, Park MA, et al. Controversies in drug allergy: In vitro testing. *J Allergy Clin Immunol*. 2019;143(1):56-65.
104. Brockow K, Garvey LH, Aberer W, Atanaskovic-Markovic M, Barbaud A, Bilo MB, et al. Skin test concentrations for systemically administered drugs -- an ENDA/EAACI Drug Allergy Interest Group position paper. *Allergy*. 2013;68(6):702-12.
105. Decuyper, II, Mangoldt EA, Van Gasse AL, Claesen K, Uyttbroek A, Faber M, et al. In Vitro Diagnosis of Immediate Drug Hypersensitivity Anno 2017: Potentials and Limitations. *Drugs in R&D*. 2017;17(2):265-78.
106. Ebo DG, Faber M, Elst J, Van Gasse AL, Bridts CH, Mertens C, et al. In Vitro Diagnosis of Immediate Drug Hypersensitivity During Anesthesia: A Review of the Literature. *J Allergy Clin Immunol Pract*. 2018;6(4):1176-84.
107. Nasser SM, Ewan PW. Opiate-sensitivity: clinical characteristics and the role of skin prick testing. *Clin Exp Allergy*. 2001;31(7):1014-20.
108. Kelso JM. MRGPRX2 signaling and skin test results. *J Allergy Clin Immunol Pract*. 2020;8(1):426.
109. Uyttbroek AP, Sabato V, Bridts CH, De Clerck LS, Ebo DG. Moxifloxacin hypersensitivity: Uselessness of skin testing. *J Allergy Clin Immunol Pract*. 2015;3(3):443-5.
110. Ebo DG, Elst J, Van Gasse A, De Puyseleir L, Faber MA, Hagendorens MM, et al. Basophil Activation Experiments in Immediate Drug Hypersensitivity: More Than a Diagnostic Aid. *Methods Mol Biol*. 2020;2163:197-211.



111. Takazawa T, Sabato V, Ebo DG. In vitro diagnostic tests for perioperative hypersensitivity, a narrative review: potential, limitations, and perspectives. *British journal of anaesthesia*. 2019;123(1):e117-e25.
112. Mayorga C, Dona I, Perez-Inestrosa E, Fernandez TD, Torres MJ. The Value of In Vitro Tests to Diminish Drug Challenges. *Int J Mol Sci*. 2017;18(6).
113. Mueller-Wirth N, Buenter A, Jorg L, Ebo DG, Glatz M, Fernando SL, et al. IgE-mediated chlorhexidine allergy-Cross-reactivity with other biguanide disinfectants. *Allergy*. 2020;75(12):3237-47.
114. Hemmings O, Du Toit G, Radulovic S, Lack G, Santos AF. Ara h 2 is the dominant peanut allergen despite similarities with Ara h 6. *J Allergy Clin Immunol*. 2020;146(3):621-30.e5.
115. Ebo DG, Decuyper, II, Rihs HP, Mertens CH, Van Gasse AL, Van der Poorten ML, et al. IgE-binding and mast cell activating capacity of the homologue of the major birch pollen allergen and profilin from *Cannabis sativa*. *J Allergy Clin Immunol Pract*. 2021.
116. Pramod SN, Venkatesh YP, Mahesh PA. Potato lectin activates basophils and mast cells of atopic subjects by its interaction with core chitobiose of cell-bound non-specific immunoglobulin E. *Clinical and experimental immunology*. 2007;148(3):391-401.
117. Krithika N, Pramod SN, Mahesh PA, Venkatesh YP. Banana lectin (BanLec) induces non-specific activation of basophils and mast cells in atopic subjects. *European annals of allergy and clinical immunology*. 2018;50(6):243-53.
118. Maurer M, Eyerich K, Eyerich S, Ferrer M, Guterth J, Hartmann K, et al. Urticaria: Collegium Internationale Allergologicum (CIA) Update 2020. *Int Arch Allergy Immunol*. 2020;181(5):321-33.
119. Irinyi B, Gyimesi E, Garaczi E, Bata ZS, Kemeny L, Zeher M, et al. Extended diagnostic value of autologous serum skin test and basophil CD63 expression assay in chronic urticaria. *Br J Dermatol*. 2013;168(3):656-8.
120. Netchiporouk E, Moreau L, Rahme E, Maurer M, Lejtenyi D, Ben-Shoshan M. Positive CD63 Basophil Activation Tests Are Common in Children with Chronic Spontaneous Urticaria and Linked to High Disease Activity. *Int Arch Allergy Immunol*. 2016;171(2):81-8.
121. Chen Q, Zhai Z, Xu J, Chen W, Chen S, Zhong H, et al. Basophil CD63 expression in chronic spontaneous urticaria: correlation with allergic sensitization, serum autoreactivity and basophil reactivity. *J Eur Acad Dermatol Venereol*. 2017;31(3):463-8.
122. Patil SU, Calatroni A, Schneider M, Steinbrecher J, Smith N, Washburn C, et al. Data-driven programmatic approach to analysis of basophil activation tests. *Cytometry B Clin Cytom*. 2018;94(4):667-73.

123. Ebo DG, Van der Poorten ML, Elst J, Van Gasse AL, Mertens C, Bridts C, et al. Immunoglobulin E cross-linking or MRGPRX2 activation: clinical insights from rocuronium hypersensitivity. *British journal of anaesthesia*. 2020.
124. Sabato V, Elst J, Van Houdt M, Bridts C, Mertens C, Ebo DG. Surface expression of MRGPRX2 on resting basophils: An area of controversy. *Allergy*. 2020;75(9):2421-2.
125. Wedi B, Gehring M, Kapp A. The pseudoallergen receptor MRGPRX2 on peripheral blood basophils and eosinophils: Expression and function. *Allergy*. 2020;75(9):2229-42.
126. Ali EA, Kalli M, Wan D, Nakamura R, Onion D, Alanine DGW, et al. Characterization of human FcepsilonRIalpha chain expression and gene copy number in humanized rat basophilic leukaemia (RBL) reporter cell lines. *PLoS One*. 2019;14(8):e0221034.
127. Akuzawa Natsuyo OH, Izumi Takashi, Shigeki Takeda Morphine is an exogenous ligand for MrgX, a G protein-coupled receptor for cortistatin. *J Cell An Biol*. 2007;2 (1), pp.004-009.
128. Che D, Rui L, Cao J, Wang J, Zhang Y, Ding Y, et al. Cisatracurium induces mast cell activation and pseudo-allergic reactions via MRGPRX2. *Int Immunopharmacol*. 2018;62:244-50.
129. Che D, Wang J, Ding Y, Liu R, Cao J, Zhang Y, et al. Mivacurium induce mast cell activation and pseudo-allergic reactions via MAS-related G protein coupled receptor-X2. *Cell Immunol*. 2018;332:121-8.
130. Zhang T, Liu R, Che D, Pundir P, Wang N, Han S, et al. A Mast Cell-Specific Receptor Is Critical for Granuloma Induced by Intrathecal Morphine Infusion. *J Immunol*. 2019;203(7):1701-14.
131. Shtessel M, Limjunyawong N, Oliver ET, Chichester K, Gao L, Dong X, et al. MRGPRX2 Activation Causes Increased Skin Reactivity in Patients with Chronic Spontaneous Urticaria. *J Invest Dermatol*. 2021;141(3):678-81 e2.
132. Rottem M, Okada T, Goff JP, Metcalfe DD. Mast cells cultured from the peripheral blood of normal donors and patients with mastocytosis originate from a CD34+/Fc epsilon RI- cell population. *Blood*. 1994;84(8):2489-96.
133. Kirshenbaum AS, Metcalfe DD. Growth of human mast cells from bone marrow and peripheral blood-derived CD34+ pluripotent progenitor cells. *Methods Mol Biol*. 2006;315:105-12.
134. Bandara G, Metcalfe DD, Kirshenbaum AS. Growth of human mast cells from bone marrow and peripheral blood-derived CD34(+) pluripotent hematopoietic cells. *Methods Mol Biol*. 2015;1220:155-62.
135. Saito H, Kato A, Matsumoto K, Okayama Y. Culture of human mast cells from peripheral blood progenitors. *Nat Protoc*. 2006;1(4):2178-83.
136. Gaudenzio N, Laurent C, Valitutti S, Espinosa E. Human mast cells drive memory CD4+ T cells toward an inflammatory IL-22+ phenotype. *J Allergy Clin Immunol*. 2013;131(5):1400-7 e11.


137. Hoffmann HJ. Generation of a human allergic mast cell phenotype from CD133(+) stem cells. *Methods Mol Biol.* 2014;1192:57-62.
138. Luo Y, Frischbutter S, Scheffel J, Siebenhaar F. Generation and Culture of Peripheral CD34(+) Stem Cell-Derived Mast Cells (PSCMCs). *Methods Mol Biol.* 2020;2163:63-7.
139. Bahri R, Bulfone-Paus S. Mast Cell Activation Test (MAT). *Methods Mol Biol.* 2020;2163:227-38.
140. Durand B, Migliaccio G, Yee NS, Eddleman K, Huima-Byron T, Migliaccio AR, et al. Long-term generation of human mast cells in serum-free cultures of CD34+ cord blood cells stimulated with stem cell factor and interleukin-3. *Blood.* 1994;84(11):3667-74.
141. Kempuraj D, Saito H, Kaneko A, Fukagawa K, Nakayama M, Toru H, et al. Characterization of mast cell-committed progenitors present in human umbilical cord blood. *Blood.* 1999;93(10):3338-46.
142. Shimizu Y, Sakai K, Miura T, Narita T, Tsukagoshi H, Satoh Y, et al. Characterization of 'adult-type' mast cells derived from human bone marrow CD34(+) cells cultured in the presence of stem cell factor and interleukin-6. Interleukin-4 is not required for constitutive expression of CD54, Fc epsilon RI alpha and chymase, and CD13 expression is reduced during differentiation. *Clin Exp Allergy.* 2002;32(6):872-80.

MAST CELL ACTIVATION TESTS


membrane markers






histamine content





intracellular calcium staining








Selection Mast cells





Intracellular markers





Membrane markers









Passively sensitized




Passively sensitized





Negative control DsiRNA



DsiRNA MRGPRX2

