

GATA-1 regulates the generation and function of basophils

Yuichiro Nei^{a,b,1}, Kazushige Obata-Ninomiya^{a,1}, Hidemitsu Tsutsui^a, Kenji Ishiwata^c, Masayuki Miyasaka^d, Kenji Matsumoto^e, Susumu Nakae^{f,g}, Hirotaka Kanuka^c, Naohiko Inase^b, and Hajime Karasuyama^{a,h,2}

Departments of ^aImmune Regulation and ^bIntegrated Pulmonology and ^cJapan Science and Technology Agency, Core Research for Evolutional Science and Technology, Graduate School of Medical and Dental Sciences, Tokyo Medical and Dental University, Tokyo 113-8519, Japan; ^dDepartment of Tropical Medicine, The Jikei University School of Medicine, Tokyo 105-8461, Japan; ^eWorld Premier International Research Center Immunology Frontier Research Center, Osaka University, Suita 565-0871, Japan; ^fDepartment of Allergy and Immunology, National Research Institute for Child Health and Development, Tokyo 157-8535, Japan; and ^gLaboratory of Systems Biology, Center for Experimental Medicine and Systems Biology, The Institute of Medical Science, and ^hJapan Science and Technology Agency, Precursory Research for Embryonic Science and Technology, The University of Tokyo, Tokyo 108-8639, Japan

Edited by Ruslan Medzhitov, Yale University School of Medicine, New Haven, CT, and approved October 8, 2013 (received for review June 20, 2013)

Developmental processes of hematopoietic cells are orchestrated by transcriptional networks. GATA-1, the founding member of the GATA family of transcription factors, has been demonstrated to play crucial roles in the differentiation of erythroid cells, megakaryocytes, eosinophils, and mast cells. However, the role of GATA-1 in basophils remains elusive. Here we show that basophils abundantly express *Gata1* mRNAs, and that siRNA-mediated knockdown of *Gata1* resulted in impaired production of IL-4 by basophils in response to the stimulation with IgE plus antigens. Δ dblGATA mice that carry the mutated *Gata1* promoter and are widely used for functional analysis of eosinophils owing to their selective loss of eosinophils showed a decreased number of basophils with reduced expression of *Gata1* mRNAs. The number of basophil progenitors in bone marrow was reduced in these mice, and the generation of basophils from their bone marrow cells in culture with IL-3 or thymic stromal lymphopoietin was impaired. Δ dblGATA basophils responded poorly *ex vivo* to stimulation with IgE plus antigens compared with wild-type basophils as assessed by degranulation and production of IL-4 and IL-6. Moreover, Δ dblGATA mice showed impaired responses in basophil-mediated protective immunity against intestinal helminth infection. Thus, Δ dblGATA mice showed numerical and functional aberrancy in basophils in addition to the known deficiency of eosinophils. Our findings demonstrate that GATA-1 plays a key role in the generation and function of basophils and underscore the need for careful distinction of the cell lineage responsible for each phenotype observed in Δ dblGATA mice.

The differentiation of hematopoietic cells from pluripotent progenitors is regulated by the coordinated action of transcription factors (1). GATA proteins comprise a family of transcription factors that have highly conserved zinc finger DNA binding domains (2, 3). GATA-1, GATA-2, and GATA-3 among six members play major roles in the hematopoietic and immune systems (4). Each GATA factor shows a tissue- and cell-restricted pattern of expression. GATA-1 is expressed in erythroid cells, megakaryocytes, mast cells, and eosinophils among hematopoietic lineages (5–9), and in Sertoli cells of the testis (10). The critical role for GATA-1 in erythropoiesis has been clearly illustrated by establishing GATA-1–null mice that die during embryogenesis due to severe anemia (11). A series of engineered mice carrying genetic modifications in the promoter region of the *Gata1* gene have been established (12–15), and some of these are not embryonic lethal, despite displaying anemia. Studies using these mutant mice revealed that GATA-1 also plays important roles in the development of platelets, mast cells, and eosinophils (12–18). Δ dblGATA mice were generated by deleting a high-affinity double GATA site in the *Gata1* promoter region (14). The double GATA site is also present in the regulatory regions of eosinophil-specific genes (19). Of note, Δ dblGATA mice show selective loss of the eosinophil lineage, with only mild anemia but no apparent anomaly in platelets and mast cells, and therefore

are widely used as eosinophil-deficient mice for the analysis of eosinophil function *in vivo* (19).

Basophils are the least common granulocytes, and represent less than 1% of peripheral blood leukocytes (20). In addition, they share certain features with tissue-resident mast cells, including the presence of basophilic granules in the cytoplasm, the surface expression of the high-affinity Fc receptor for IgE (Fc ϵ RI), and the release of chemical mediators, such as histamine, after stimulation. Accordingly, basophils have often erroneously been considered as minor and redundant relatives of mast cells or blood-circulating precursors of tissue-resident mast cells, and have long been neglected in immunological studies (21). However, recent development of analytical tools for basophil function *in vivo*, including basophil-deficient mice, has identified pivotal and nonredundant roles for basophils in a variety of immune responses, such as allergic reactions, protective immunity against parasitic infections, and regulation of innate and acquired immunity (22–28). Nevertheless, the origin and developmental pathway of basophils, including transcription factors regulating their differentiation, still remain ill-defined compared with those of other hematopoietic cells. Although a developmental relationship between basophils and eosinophils has been suggested in humans (29), a bipotent progenitor of basophils and mast cells, in addition to a unipotent basophil progenitor, has been identified in mice (30, 31), suggesting a closer lineage

Significance

The GATA-1 transcription factor has been extensively characterized and shown to play crucial roles in the development of erythroid cells, megakaryocytes, eosinophils, and mast cells. However, the role of GATA-1 in basophils remained unidentified. We demonstrate that knockdown of *Gata1* gene expression in basophils results in impaired cytokine production upon allergen-mediated activation. Moreover, Δ dblGATA mice carrying the mutated *Gata1* promoter have reduced numbers of basophils and their progenitors and show impaired responses in basophil-mediated protective immunity against parasitic infections. Thus, GATA-1 plays an important role in both generation and activation of basophils, and Δ dblGATA mice display numerical and functional aberrancy in basophils, in addition to the well-known eosinophil deficiency.

Author contributions: K.O.-N., N.I., and H. Karasuyama designed research; Y.N., K.O.-N., H.T., and K.I. performed research; K.I., M.M., K.M., S.N., and H. Kanuka contributed new reagents/analytic tools; Y.N., K.O.-N., N.I., and H. Karasuyama analyzed data; and Y.N., K.O.-N., and H. Karasuyama wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹Y.N. and K.O.-N. contributed equally to this work.

²To whom correspondence should be addressed. E-mail: karasuyama.mbch@tmd.ac.jp.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1311668110/-DCSupplemental.

relationship of basophils with mast cells in mice. The ordered expression of two transcription factors, GATA-2 and C/EBP α , appears to regulate the commitment of progenitor cells into the basophil lineage (30, 32). Of note, mice deficient in the distal promoter-derived Runt-related transcription factor 1 (P1-Runx1) show severe basophilopenia with no apparent anomaly of mast cell, neutrophils, or eosinophils (33), indicating the requirement of P1-Runx1 for the later stage of basophil development. Recent studies demonstrated that basophils express GATA-1 (30, 32, 34), but its functional significance remains to be determined.

In the present study, we explored the possible involvement of GATA-1 in the ontogeny and function of murine basophils. Knockdown of *Gata1* gene expression in basophils resulted in impaired production of IL-4 in vitro in response to stimulation with IgE plus antigens. Moreover, Δ dblGATA mice had basophilopenia with reduced expression of *Gata1* and poor IL-4 production in basophils, and showed impaired responses in basophil-mediated protective immunity against intestinal helminth infection. Thus, GATA-1 plays an important role in both generation and activation of basophils, and Δ dblGATA mice display aberrancy in basophils, in addition to eosinophil deficiency.

Results

Knockdown of *Gata1* in Basophils Impairs IL-4 Production in Response to Stimulation with IgE Plus Antigens. We first compared the level of *Gata1* mRNA expression in three distinct types of murine granulocytes. Basophils expressed *Gata1* mRNAs at the level that was as high as ~80% of that in eosinophils, whereas neutrophils showed little or no expression (Fig. 1A). To examine the functional significance of *Gata1* expression in basophils, *Gata1*-specific siRNAs were introduced into IL-3-cultured bone marrow-derived basophils to repress *Gata1* expression (Fig. 1B). *Gata1*-knockdown basophils reproducibly showed a slightly lower level of surface Fc ϵ RI α expression compared with control basophils, whereas surface CD200R3 and CD49b expression was comparable between them (Fig. 1C). When stimulated with IgE plus antigens, *Gata1*-knockdown basophils produced significantly lower amounts of IL-4 than did control siRNA-treated basophils at both

mRNA and protein levels (Fig. 1D and E). These results suggested that GATA-1 in basophils contribute to the regulation of Fc ϵ RI expression, and IL-4 production triggered by Fc ϵ RI cross-linking.

Δ dblGATA Mice Show Basophilopenia with Reduced Expression of *Gata1* and Surface Fc ϵ RI in Basophils. The results obtained from the *Gata1* knockdown experiments prompted us to examine Δ dblGATA mice for possible aberrancy of their basophils. In contrast to the nearly complete loss of eosinophils, basophils were detectable in Δ dblGATA BALB/c mice. However, the number of basophils was significantly reduced in these mice, compared with that in wild-type mice, particularly in the bone marrow and peripheral blood, to approximately a half and one-third of normal, respectively (Fig. 2A). Similar basophilopenia was observed in Δ dblGATA C57BL/6 mice compared with control littermates (Fig. S1). Of note, the *Gata1* mRNA expression in Δ dblGATA basophils was not null but reduced to approximately a quarter of that in wild-type basophils, whereas the level of *Gata2* mRNAs was comparable between them (Fig. 2B). Flow cytometric analysis revealed that the expression of cell surface CD200R3 was equivalent between them, whereas the expression of Fc ϵ RI α and CD49b was slightly but significantly reduced in Δ dblGATA basophils (Fig. 2C). Considering the fact that the *Gata1* gene is expressed in basophil progenitors (30), the discrepancy between the reduced and unaltered expression of CD49b on Δ dblGATA basophils (Fig. 2C) and siRNA-treated basophils (Fig. 1C), respectively, could be attributed to the effect of reduced versus normal GATA-1 expression on the process of basophil development.

Δ dblGATA Mice Have a Reduced Number of Basophil Progenitors and Show Poor Generation of Basophils. Basophilopenia in the bone marrow suggested the possible impairment of basophil hematopoiesis in Δ dblGATA mice. Indeed, the number of CD34⁺c-kit⁺Fc ϵ RI⁺ basophil progenitors (30) in the bone marrow of Δ dblGATA mice was only a quarter of that in wild-type mice (Fig. 2D). A recent study demonstrated that thymic stromal lymphopoietin (TSLP) and IL-3 independently promote basophil hematopoiesis (35). We found that the surface expression of




Fig. 1. Knockdown of *Gata1* in basophils impairs their IL-4 production. (A) Eosinophils, basophils, and neutrophils were separately isolated from the bone marrow of BALB/c mice, and subjected to RT-PCR analysis. The relative expression of *Gata1* is shown (mean $\pm$ SEM, $n = 3$ each); the level of expression in eosinophils is set as 1. (B–E) *Gata1*-specific or control siRNAs were introduced into IL-3-cultured basophils generated from BALB/c bone marrow cells. Two days later, siRNA-treated basophils were subjected to RT-PCR analysis for *Gata1* expression (B, mean $\pm$ SEM, $n = 3$ each), and flow cytometric analysis for indicated surface markers (C). In C, representative staining profiles are shown (Upper); gray, black, and shaded histograms indicate those of control and *Gata1* siRNA-treated basophils, and control staining with isotype-matched antibodies, respectively. The mean fluorescence intensity (MFI) of each surface marker is shown (Lower) (mean $\pm$ SEM, $n = 3$ each). In D and E, basophils treated with *Gata1*-specific or control siRNAs were stimulated with TNP-specific IgE plus TNP-OVA or control OVA for 3 h (D) or 6 h (E), and subjected to RT-PCR analysis for *Il4* expression (D, mean $\pm$ SEM, $n = 3$ each) or flow cytometric analysis for IL-4 production (E). In E, representative staining profiles are shown (Upper); black and gray histograms indicate those of TNP-OVA and OVA-treated basophils, respectively. The percentage of IL-4-producing cells among basophils is shown (Lower) (mean $\pm$ SEM, $n = 3$ each). Data in A–E are representative of at least three independent experiments. ** $P < 0.01$; *** $P < 0.001$.




Fig. 3. Δ dblGATA bone marrow cells show impaired generation of basophils when cultured ex vivo with TSLP or IL-3. Bone marrow cells isolated from WT or Δ dbl BALB/c mice were cultured ex vivo with TSLP or IL-3 for 5 d. (A) Basophils were identified as $FSC^{low}SSC^{low}CD200R3^{+}c-kit^{+}$ cells as shown at Left, and the number of basophils in each group is summarized at Right (mean $\pm$ SEM, $n = 4$ each). (B) The cell surface phenotype of bone marrow-derived basophils. Representative staining profiles are shown (TSLP and IL-3 rows); gray, black, and shaded histograms indicate those of WT and Δ dbl basophils and control staining with isotype-matched antibodies, respectively. MFI of each surface marker is shown (MFI rows) (mean $\pm$ SEM, $n = 4$ each). Data shown are representative of at least two independent experiments. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

Discussion

Basophils express GATA-1 and GATA-2 but not GATA-3 among hematopoietic GATA transcription factors (34). GATA-2 has been demonstrated to play a key role in basophil hematopoiesis (30, 32), and positively regulate ST2 (IL-33 receptor) expression in basophils (37). In contrast, the role of GATA-1 in basophils remained to be identified despite the fact that GATA-1 has been extensively characterized in terms of its functions in the development of erythroid cells, megakaryocytes, eosinophils, and mast cells (11–18). We illustrated in the present study that GATA-1 plays an important role in both generation and function of basophils, and that Δ dblGATA mice show numerical and

functional aberrancy in basophils, in addition to the known eosinophil deficiency.

Although several cell types, including Th2 cells and basophils, can produce IL-4, each cell type uses distinct *cis*-regulatory element for the *Il4* gene expression (38). Basophils, unlike Th2 cells, do not express Th2 transcription factors GATA3, RBPJ κ , or c-Maf (34). Instead, basophils express GATA-1, GATA-2, and C/EBP α in contrast to Th2 cells. C/EBP α has been shown to directly regulate *Il4* gene transcription in basophils, in cooperation with NFAT (34). Although GATA-1 and GATA-2 have been reported to bind to the intronic enhancer of the *Il4* gene in mast cells (39), basophils used the HS4 element rather than the intronic enhancer as a lineage-specific *Il4* enhancer (38). Thus, the role of GATA-1 and GATA-2 in the *Il4* gene regulation in basophils remained uncertain. In the present study, we illustrated the key role for GATA-1 in IL-4 production by basophils. *Gata1* knockdown resulted in significantly reduced IL-4 expression at both mRNA and protein levels by basophils when stimulated with IgE plus antigens. In accordance with this, Δ dblGATA basophils expressing reduced amounts of *Gata1* mRNAs also showed impaired production of IL-4. In both *Gata1*-knockdown basophils and Δ dblGATA basophils, the surface Fc ϵ RI expression was slightly but significantly reduced, compared with that in normal counterparts, as observed in mast cells from GATA-1^{low} and GATA-KD mice that are genetically engineered to have mutations in the promoter region of the *Gata1* gene, and hence show reduced expression of GATA-1 (18, 40). However, the extent of MAPK (ERK1/2 and p38) phosphorylation in response to Fc ϵ RI-mediated stimulation was comparable between mutant and wild-type basophils (Fig. S5). Therefore, impaired IL-4 production in mutant basophils may not be attributed to the reduced Fc ϵ RI expression, and is likely due to a reduced amount of GATA-1 that regulates *Il4* gene expression.




Fig. 4. Δ dblGATA basophils show defects in degranulation and cytokine production. (A–C) IL-3-elicited WT or Δ dbl basophils were stimulated with anti-TNP IgE plus TNP-OVA or control OVA for 20 min (A), 3 h (B), or 6 h (C), and subjected to flow cytometric analysis for surface CD63 expression (A), RT-PCR analysis for *Il4* expression (B), or flow cytometric analysis for IL-4 production (C). In A, open and shaded histograms indicate CD63 expression when stimulated with TNP-OVA and control OVA, respectively. (D and E) Primary basophils isolated from the bone marrow and spleen of WT or Δ dbl mice were stimulated with anti-TNP IgE plus TNP-OVA or control OVA for 20 min (D) or 6 h (E), and subjected to flow cytometric analysis for surface CD63 expression (D) or flow cytometric analysis for IL-4 production (E). Data shown are the mean $\pm$ SEM ($n = 3$ or 4 each), and representative of three independent experiments, and displayed as in Fig. 1. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.




Fig. 5. Δ dblGATA mice show impaired acquired protection against intestinal helminths due to basophil anomaly. (A–D) WT and Δ dbl BALB/c mice were infected twice with *N. brasiliensis* larvae, and on day 2 of the second infection, their spleen and skin of larva inoculation site were isolated, and subjected to numerical counts of basophils in the spleen (A), indicated cell types accumulating in the skin (B, Upper and Lower), and skin-trapped worms (D). M2-Mac, M2-type macrophage. In C, basophils were sorted from the skin preparation and subjected to RT-PCR analysis for *Il4* expression. (E) WT and Δ dbl mice were infected once with *N. brasiliensis* larvae, and basophils were sorted from their bone marrow cells on day 18 of infection. These *N. brasiliensis*-sensitized basophils isolated from WT or Δ dbl mice were intraperitoneally transferred into Δ dbl mice (4×10^4 cells per mouse) that had been infected with larvae 18 d before. On the day of cell transfer, the recipient mice were subjected to the second infection, and 2 d later, the skin of the larva inoculation site was isolated, and subjected to a numerical count of larvae. In parallel, twice-infected Δ dbl mice without basophil transfer (–) were analyzed as control. Data shown are the mean $\pm$ SEM ($n = 3$ or 4 each), and representative of three independent experiments. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

GATA-1 has been shown to be involved in the differentiation of mast cells. Morphologically abnormal alcian blue⁺ mast cells and numerous mast cell precursors are detected in connective tissues and peritoneal lavage of GATA-1^{low} and GATA-KD mice (17, 18). Moreover, in vitro generation of c-kit⁺FcεRI⁺ mast cells from bone marrow cells is reduced in these mice (18). Thus, GATA-1 is required for generation and maturation of connective tissue mast cells. Although a close developmental relationship between mast cells and basophils has been suggested in mice (30, 31), the functional significance of GATA-1 expression in basophils in terms of their development remained to be clarified. In the present study, we found that Δ dblGATA mice had a decreased number of basophils with reduced expression of *Gata1* mRNAs. Δ dblGATA bone marrow cells generated fewer basophils ex vivo in culture with IL-3 or TSLP than did wild-type cells, in accordance with a decreased number of basophil progenitors in the bone marrow of Δ dblGATA mice, suggesting that GATA-1 is involved in the development of basophils. This reduced basophilopoiesis does not seem to be linked to eosinophil deficiency, because two other mouse strains with eosinophil deficiency have a normal number of basophils (41). Because

GATA-1 regulates GATA-2 expression (9), and GATA-2 is involved in the regulation of basophil development (30, 32), GATA-1 may contribute to the basophil development through GATA-2 regulation. However, GATA-2 expression in Δ dblGATA basophils remains unaltered despite significant reduction in GATA-1 expression, suggesting that GATA-1 may regulate the expression of a gene(s) involved in basophil differentiation, either independently of GATA-2 or by competing or cooperating with GATA-2. Of note, in Δ dblGATA mice, basophilopoiesis is not completely arrested as in the case of erythropoiesis. This could be attributed to the reduced but not null expression of GATA-1 in Δ dblGATA basophils, even though we cannot formally exclude the possibility that GATA-2 expressed in Δ dblGATA basophils may compensate some of GATA-1 functions. A previous study on Δ dblGATA mice demonstrated that in vitro generation of bone marrow-derived mast cells in culture with IL-3 and stem cell factor seems intact, assessed by toluidine blue staining and flow cytometric analysis of c-kit and FcεRI expression (14). Therefore, the Δ dblGATA mutation appears to impair the generation of eosinophils and basophils but not mast cells.

Δ dblGATA mice have been widely used for functional analysis of eosinophils because they show almost complete loss of the eosinophil lineage, with only mild anemia but no apparent anomaly in platelets and mast cells (14). We illustrated in the present study that Δ dblGATA mice showed impaired acquired protection against helminth infection due to basophil anomaly. Considering recent advances in our understanding of functional significance of basophils in various immune responses (22–28), this finding raises concern about the possibility that certain functions of basophils might have been erroneously interpreted as those of eosinophils in studies using Δ dblGATA mice as eosinophil-deficient mice. Two distinct eosinophil-deficient mouse strains, Δ dblGATA and PHIL, were analyzed to clarify the role of eosinophils in pathogenesis of asthma, resulting in some conflicting results (42, 43). This discrepancy reportedly stems from the different genetic background of these mice (44), but may also in part come from the difference in quantity and quality of basophils between these two strains. Thus, the present study underscores the need for careful distinction of the cell lineage responsible for each phenotype observed in Δ dblGATA mice, by means of parallel analyses with other eosinophil-deficient models and basophil-deficient mice.

Materials and Methods

Mice. Wild-type and Δ dblGATA BALB/c mice (14) were purchased from CLEA Japan and Jackson Laboratory, respectively. Δ dblGATA BALB/c mice were backcrossed to C57BL/6 mice for five generations to obtain Δ dblGATA C57BL/6 mice. All animal studies were approved by the Institutional Animal Care and Use Committee of Tokyo Medical and Dental University.

Antibodies. Biotinylated anti-CD49b (DX5), PE-conjugated anti-CD200R3 (Ba13) and anti-CD63 (NVG-2), PE-Cy7-conjugated anti-c-kit (2B8), APC-conjugated anti-CD200R3 (Ba13), IL-4 (11B11) and CD34 (HM34), and FITC-conjugated CD4 (RM4-5), CD8α (53-6.7), CD11b (M1/70), B220 (RA3-6B2), Gr-1 (RB6-8C5), and anti-FcεRIα (MAR-1) were purchased from BioLegend. Biotinylated anti-CD123 (5B11), PE-conjugated IL-6 (MP5-20F3), APC-conjugated anti-CD49b (HMα2), FITC-conjugated CD11c (HL3), Ax488-conjugated anti-phosphorylated ERK1/2 (20A), Ax488-conjugated anti-phosphorylated p38 MAPK (36/p38), and streptavidin were from BD Pharmingen. PE-conjugated anti-TSLP-Rα was from R&D Systems.

Quantitative RT-PCR. Total mRNAs from cells were isolated by RNeasy RNA cell miniprep system (Promega). The first-strand cDNAs were generated with reverse transcription using oligo-dT, random primers (Life Technologies Corporation), and ReverTra Ace-α (Toyobo). Quantitative PCR of the cDNA was performed on Applied Biosystems StepOnePlus Real-Time PCR system using a Fast SYBR Green Master Mix (Life Technologies Corporation) and the following primer sets: for *Gata1*, forward 5'-CACTCCCCAGTCTTTCAGGTGTA-3' and reverse 5'-GGTGAGCCCCCAGGAATT-3'; for *Gata2*, forward 5'-CACCTGTTGTGCAAATTGTCAGA-3' and reverse 5'-GGATCCCTCTCTTTCATGGT-3'; for *Il4*, forward 5'-ACTTGAGAGAGATCATCGGCA-3' and reverse 5'-AGCTCCATGAGAACTAGAGTT-3'; for *Il6*, forward 5'-CTGCAAGAGACTTCCATCCAG-3'

and reverse 5'-AGTGGTATAGACAGGTCTGTTGG-3'; and for *Hprt*, forward 5'-GGCCAGACTTTGTTGGATTG-3' and reverse 5'-CGCTCATCTTAGGCTTTGATTTG-3'.

Gene expression was analyzed using *Hprt* as an endogenous control in each sample.

Flow Cytometric Analysis and Cell Preparation. For flow cytometric analysis, cells were preincubated with anti-CD16/32 mAb and normal rat serum on ice for 15 min before incubation with the indicated combination of Abs to prevent the nonspecific binding of irrelevant Abs. Stained cells were analyzed using FACSCanto II (BD Biosciences). Basophils were isolated from bone marrow cells using biotinylated anti-CD49b antibody and streptavidin-conjugated magnetic particles (BD Pharmingen), followed by sorting FSC^{low}SSC^{low}CD200R3⁺c-kit⁻ cells with FACSARIA II (BD Bioscience). Eosinophils (FSC^{low}SSC^{high}Siglec-F⁺) and neutrophils (Gr-1^{high}) were directly sorted from bone marrow cells. Basophil progenitors were identified as Lin (CD4, CD8, B220, CD11b, Gr-1, CD11c)~CD34⁺c-kit⁺FcεRI⁺ cells in the bone marrow (30). Bone marrow-derived basophils were generated by culturing bone marrow cells ex vivo with 1 ng/mL of IL-3 or 1 μg/mL of TSLP for 5 d.

Knockdown of *Gata1*. *Gata1*-specific siRNA (Silencer Predesigned siRNA, no. 66474, Life technologies Corporation) and control siRNA (Silencer Negative Control #1 siRNA) were introduced into IL-3-cultured bone marrow-derived basophils by using Neon Transfection System (Life Technologies Corporation).

Stimulation of Basophils. Basophils were enriched from freshly isolated bone marrow cells or IL-3-cultured bone marrow cells by using biotinylated anti-CD49b antibody and streptavidin-conjugated magnetic particles (BD

Pharmingen). For RT-PCR analysis, FSC^{low}SSC^{low}c-kit⁺CD45^{low}CD49b⁺ basophils were further sorted from them. Basophil-enriched CD49b⁺ cells or purified basophils were first sensitized with 1 μg/mL of hapten 2,4,6-trinitrophenol (TNP)-specific IgE, followed by incubation with 100 ng/mL of TNP-conjugated ovalbumin (OVA) or control OVA for 20 min for flow cytometric analysis of the CD63 up-regulation, for 30 min for β-hexosaminidase release assay (Sigma-Aldrich), for 3 h for RT-PCR analysis, and for 6 h in the presence of momesin for cytokine production. For detection of cytokine production, stimulated cells were subjected to fixation and permeabilization using Cytofix/Cytoperm buffer (BD Pharmingen), followed by intracellular staining with antibodies specific to IL-4 or IL-6. In some experiments, basophils were stimulated for 6 h with phorbol 12-myristate 13-acetate (PMA) (0.1 μg/mL, Sigma-Aldrich) plus ionomycin (0.5 μg/mL, Sigma-Aldrich).

Helminth Infection. Mice were first injected s.c. with 500 third-stage larvae (L3) of *N. brasiliensis* in the back, and 18 d later, they were injected intradermally with 500 L3 in the flank. Two days after the second larva inoculation, the skin of the inoculation site was isolated, and subjected to numerical counts of skin-trapped larvae and skin-infiltrating cells as described previously (36).

Statistical Analysis. Statistical analysis was performed using unpaired Student *t* test. A *P* value <0.05 was considered statistically significant.

ACKNOWLEDGMENTS. This work was supported by research fellowships from the Japan Society for the Promotion of Science for Young Scientists (to K.O.-N.) and by research grants from JST, CREST (to H. Karasuyama), JST, PRESTO (to S.N.), and the Japanese Ministry of Education, Culture, Sports, Science, and Technology (to H. Karasuyama).

- Orkin SH, Zon LI (2008) Hematopoiesis: An evolving paradigm for stem cell biology. *Cell* 132(4):631–644.
- Yamamoto M, et al. (1990) Activity and tissue-specific expression of the transcription factor NF-E1 multigene family. *Genes Dev* 4(10):1650–1662.
- Bresnick EH, Lee HY, Fujiwara T, Johnson KD, Keles S (2010) GATA switches as developmental drivers. *J Biol Chem* 285(41):31087–31093.
- Orkin SH (1992) GATA-binding transcription factors in hematopoietic cells. *Blood* 80(3):575–581.
- Romeo PH, et al. (1990) Megakaryocytic and erythrocytic lineages share specific transcription factors. *Nature* 344(6265):447–449.
- Martin DI, Zon LI, Mutter G, Orkin SH (1990) Expression of an erythroid transcription factor in megakaryocytic and mast cell lineages. *Nature* 344(6265):444–447.
- Zon LI, et al. (1993) Expression of mRNA for the GATA-binding proteins in human eosinophils and basophils: Potential role in gene transcription. *Blood* 81(12):3234–3241.
- Migliaccio AR, Rana RA, Vannucchi AM, Manzoli FA (2005) Role of GATA-1 in normal and neoplastic hematopoiesis. *Ann N Y Acad Sci* 1044:142–158.
- Ferreira R, Ohneda K, Yamamoto M, Philipsen S (2005) GATA1 function, a paradigm for transcription factors in hematopoiesis. *Mol Cell Biol* 25(4):1215–1227.
- Ito E, et al. (1993) Erythroid transcription factor GATA-1 is abundantly transcribed in mouse testis. *Nature* 362(6419):466–468.
- Fujiwara Y, Browne CP, Cuniff K, Goff SC, Orkin SH (1996) Arrested development of embryonic red cell precursors in mouse embryos lacking transcription factor GATA-1. *Proc Natl Acad Sci USA* 93(12):5215–5218.
- Shivdasani RA, Fujiwara Y, McDevitt MA, Orkin SH (1997) A lineage-selective knock-out establishes the critical role of transcription factor GATA-1 in megakaryocyte growth and platelet development. *EMBO J* 16(13):3965–3973.
- Takahashi S, et al. (1997) Arrest in primitive erythroid cell development caused by promoter-specific disruption of the GATA-1 gene. *J Biol Chem* 272(19):12611–12615.
- Yu C, et al. (2002) Targeted deletion of a high-affinity GATA-binding site in the GATA-1 promoter leads to selective loss of the eosinophil lineage in vivo. *J Exp Med* 195(11):1387–1395.
- Majewski IJ, et al. (2006) A mutation in the translation initiation codon of Gata-1 disrupts megakaryocyte maturation and causes thrombocytopenia. *Proc Natl Acad Sci USA* 103(38):14146–14151.
- McDevitt MA, Shivdasani RA, Fujiwara Y, Yang H, Orkin SH (1997) A “knockdown” mutation created by *cis*-element gene targeting reveals the dependence of erythroid cell maturation on the level of transcription factor GATA-1. *Proc Natl Acad Sci USA* 94(13):6781–6785.
- Hariigae H, et al. (1998) Differential roles of GATA-1 and GATA-2 in growth and differentiation of mast cells. *Genes Cells* 3(1):39–50.
- Migliaccio AR, et al. (2003) GATA-1 as a regulator of mast cell differentiation revealed by the phenotype of the GATA-1low mouse mutant. *J Exp Med* 197(3):281–296.
- Rothenberg ME, Hogan SP (2006) The eosinophil. *Annu Rev Immunol* 24:147–174.
- Galli SJ (2000) Mast cells and basophils. *Curr Opin Hematol* 7(1):32–39.
- Falcone FH, Haas H, Gibbs BF (2000) The human basophil: A new appreciation of its role in immune responses. *Blood* 96(13):4028–4038.
- Min B (2008) Basophils: What they ‘can do’ versus what they ‘actually do’ *Nat Immunol* 9(12):1333–1339.
- Sullivan BM, Locksley RM (2009) Basophils: A nonredundant contributor to host immunity. *Immunity* 30(1):12–20.
- Schroeder JT (2009) Basophils beyond effector cells of allergic inflammation. *Adv Immunol* 101:123–161.
- Sokol CL, Medzhitov R (2010) Emerging functions of basophils in protective and allergic immune responses. *Mucosal Immunol* 3(2):129–137.
- Karasuyama H, Mukai K, Obata K, Tsujimura Y, Wada T (2011) Nonredundant roles of basophils in immunity. *Annu Rev Immunol* 29:45–69.
- Siracusa MC, Comeau MR, Artis D (2011) New insights into basophil biology: Initiators, regulators, and effectors of type 2 inflammation. *Ann N Y Acad Sci* 1217:166–177.
- Voehringer D (2013) Protective and pathological roles of mast cells and basophils. *Nat Rev Immunol* 13(5):362–375.
- Arock M, Schneider E, Boissan M, Tricottet V, Dy M (2002) Differentiation of human basophils: An overview of recent advances and pending questions. *J Leukoc Biol* 71(4):557–564.
- Arinobu Y, et al. (2005) Developmental checkpoints of the basophil/mast cell lineages in adult murine hematopoiesis. *Proc Natl Acad Sci USA* 102(50):18105–18110.
- Metcalfe D, Ng AP, Baldwin TM, Di Rago L, Mifsud S (2013) Concordant mast cell and basophil production by individual hematopoietic blast colony-forming cells. *Proc Natl Acad Sci USA* 110(22):9031–9035.
- Iwasaki H, et al. (2006) The order of expression of transcription factors directs hierarchical specification of hematopoietic lineages. *Genes Dev* 20(21):3010–3021.
- Mukai K, et al. (2012) Critical role of P1-Runx1 in mouse basophil development. *Blood* 120(1):76–85.
- Qi X, Nishida J, Chaves L, Ohmori K, Huang H (2011) CCAAT/enhancer-binding protein alpha (C/EBPalpha) is critical for interleukin-4 expression in response to FcεpsilonRI receptor cross-linking. *J Biol Chem* 286(18):16063–16073.
- Siracusa MC, et al. (2011) TSLP promotes interleukin-3-independent basophil hematopoiesis and type 2 inflammation. *Nature* 477(7363):229–233.
- Obata-Ninomiya K, et al. (2013) The skin is an important bulwark of acquired immunity against intestinal helminths. *J Exp Med* 210:1084–1094.
- Baba Y, et al. (2012) GATA2 is a critical transactivator for the human IL1RL1/ST2 promoter in mast cells/basophils: Opposing roles for GATA2 and GATA1 in human IL1RL1/ST2 gene expression. *J Biol Chem* 287(39):32689–32696.
- Yagi R, Tanaka S, Motomura Y, Kubo M (2007) Regulation of the IL4 gene is independently controlled by proximal and distal 3′ enhancers in mast cells and basophils. *Mol Cell Biol* 27(23):8087–8097.
- Henkel G, Brown MA (1994) PU.1 and GATA: Components of a mast cell-specific interleukin 4 intronic enhancer. *Proc Natl Acad Sci USA* 91(16):7737–7741.
- Nishiyama C, et al. (2005) GATA-1 is required for expression of FcεpsilonRI on mast cells: Analysis of mast cells derived from GATA-1 knockdown mouse bone marrow. *Int Immunol* 17(7):847–856.
- Doyle AD, et al. (2013) Expression of the secondary granule proteins major basic protein 1 (MBP-1) and eosinophil peroxidase (EPX) is required for eosinophilopoiesis in mice. *Blood* 122(5):781–790.
- Lee JJ, et al. (2004) Defining a link with asthma in mice congenitally deficient in eosinophils. *Science* 305(5691):1773–1776.
- Humbles AA, et al. (2004) A critical role for eosinophils in allergic airways remodeling. *Science* 305(5691):1776–1779.
- Walsh ER, et al. (2008) Strain-specific requirement for eosinophils in the recruitment of T cells to the lung during the development of allergic asthma. *J Exp Med* 205(6):1285–1292.

Supporting Information

Nei et al. 10.1073/pnas.1311668110




Fig. S1. Δ dblGATA C57BL/6 mice show basophilopenia. Numbers of basophils, defined as $FSC^{low}SSC^{low}CD200R3^{+}c-kit^{-}$ cells, in the bone marrow, peripheral blood, and spleen of Δ dbl C57BL/6 mice and their control littermates (LMs) are shown (mean $\pm$ SEM, $n = 5$ each), representative of two independent experiments. * $P < 0.05$; ** $P < 0.01$.




Fig. S2. β -Hexosaminidase release and IL-4 production in wild-type (WT) versus Δ dblGATA basophils. (A) IL-3-elicited WT or Δ dbl BALB/c basophils were stimulated with anti-2,4,6-trinitrophenol (TNP) IgE plus TNP-ovalbumin (OVA) or control OVA for 30 min, and subjected to β -hexosaminidase release assay. (B) IL-3-elicited WT or Δ dbl basophils were stimulated with PMA plus ionomycin (PMA-Iono) or control medium for 6 h, and subjected to flow cytometric analysis for IL-4 production. Data shown are the mean $\pm$ SEM ($n = 3$ each), and representative of three independent experiments. Med, control medium. * $P < 0.05$; n.s., not significant.




Fig. S3. Δ dblGATA basophils show impaired IL-6 production. (A and B) IL-3-elicited WT or Δ dbl BALB/c basophils were stimulated with anti-TNP IgE plus TNP-OVA or control OVA for 3 h (A) or 6 h (B), and subjected to RT-PCR analysis for *I/6* expression (A) or flow cytometric analysis for IL-6 production (B). (C and D) Primary basophils isolated from the bone marrow and spleen of WT or Δ dbl mice were stimulated with anti-TNP IgE plus TNP-OVA or control OVA for 3 h (C) or 6 h (D), and subjected to RT-PCR analysis for *I/6* expression (C) or flow cytometric analysis for IL-6 production (D). Data shown are the mean $\pm$ SEM ($n = 3$ or 4 each), and representative of three independent experiments. *** $P < 0.001$.




Fig. S4. Primary Δ dblGATA basophils show impaired *I/4* up-regulation. Primary basophils ($FSC^{low}SSC^{low}c-kit^{-}CD45^{low}CD49b^{+}$) sorted from bone marrow cells of WT or Δ dbl BALB/c mice were stimulated with anti-TNP IgE plus indicated concentration (conc.) of TNP-OVA or control OVA for 3 h and subjected to RT-PCR analysis for *I/4* expression (mean $\pm$ SEM, $n = 3$ each). Data shown are representative of two independent experiments. * $P < 0.05$; *** $P < 0.001$.




Fig. 55. Δ dblGATA basophils show intact phosphorylation of ERK1/2 and p38 when stimulated with IgE plus antigens. Basophils isolated from the bone marrow and spleen of WT and/or Δ dbl BALB/c mice were stimulated with anti-TNP IgE plus TNP-OVA or control OVA for 5 min, and subjected to flow cytometric analysis for phosphorylation of ERK1/2 (A) and p38 MAPK (B). Representative staining profiles are shown (Upper), in that black and gray histograms indicate those of TNP-OVA and OVA-treated basophils, respectively. Mean fluorescence intensity of each staining is shown (Lower) (mean $\pm$ SEM, $n = 3$ each). Data shown are representative of three independent experiments; n.s., not significant.