



Contents lists available at ScienceDirect

Allergology International

journal homepage: <http://www.elsevier.com/locate/alit>

Original Article

Analysis of factors associated with cedar pollen sensitization and development of pollinosis in a young Japanese adult population

Satoshi Nakamura ^{a,1}, Sachiko Tsunoda ^{b,1}, Hiroshi Sakaida ^a, Sawako Masuda ^c, Ahamad Shah Said ^a, Kazuhiko Takeuchi ^{a,*}^a Department of Otorhinolaryngology, Head & Neck Surgery, Mie University Graduate School of Medicine, Mie, Japan^b School of Medicine, Mie University, Mie, Japan^c Department of Otorhinolaryngology, Mie National Hospital, Mie, Japan

ARTICLE INFO

Article history:

Received 29 January 2018

Received in revised form

2 May 2018

Accepted 13 May 2018

Available online xxx

Keywords:

Association

Factors

Japanese cedar pollinosis

Sensitization

Specific IgE antibody

Abbreviations:

FEIA fluorescent enzyme immunoassay

CpG ODN cytosine phosphate-guanosine

oligodeoxynucleotides

ABSTRACT

Background: Genetic and environmental factors are proposed to be involved in cedar pollen allergy sensitization and onset. The impact of these factors will provide key information for the prevention of cedar pollen sensitization and allergy onset, which we investigated in this cross-sectional study.

Methods: Subjects were 382 young adult volunteers who completed a self-administered questionnaire on self-reported subjective symptoms of pollinosis, physician-diagnosed pollinosis, and background factors. We also measured their serum IgE antibody titers specific for cedar, cypress, and mites. Factors associated with subjective symptoms, physician diagnosis, and the three specific antigens were determined using both univariate and multivariate analyses.

Results: Sensitization to cedar, cypress, and mites, defined as specific IgE levels of class 1 or above, was found in 78.8%, 64.4%, and 56.0% of subjects, respectively. The prevalence of cedar pollinosis was 41.2% based on subjective symptoms and 22.2% based on physician diagnosis. Factors associated with increased cedar pollen sensitization were mite sensitization, comorbid allergic rhinitis, and family history of cedar pollinosis. Risk-reducing factors for cedar pollen sensitization were keeping a cat, number of common colds, and hours of sleep. Risk-increasing factors for both subjective pollinosis symptoms and physician-diagnosed pollinosis were comorbid allergic rhinitis and family history of cedar pollinosis.

Conclusions: Sensitization to cedar pollen in this population was extremely high. Both common and distinct factors were associated with sensitization to pollen and with the development of pollinosis. The distinct factors were associated with sensitization to cedar and cypress antigens.

Copyright © 2018, Japanese Society of Allergy. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Seasonal allergic rhinitis caused by Japanese cedar pollen is the most common allergic disease in Japan and is considered a national affliction.¹ The prevalence of Japanese cedar pollinosis increased 2.6-fold between 1980 and 2000, and the prevalence differs considerably according to age² and degree of urbanization.³ In a 2006–2007 survey of adults aged 20–49 years in Fukui Prefecture, the positive rate of serum Japanese cedar pollen specific IgE level above 0.7 (CAP RAST score of 2) was 55.5% and the prevalence of cedar pollinosis reported on a questionnaire was 36.7%.⁴ Among

adolescents, the prevalence of pollinosis, defined as the percentage of the population who were sensitized and had one or more seasonal rhinoconjunctival symptoms, was reported to be 28.7% in metropolitan areas in the year 2004.³ In younger children, a 2003 epidemiological survey of allergic diseases among first-year junior high school students (age 11–12 years) in Wakayama Prefecture found that 48.6% of children had specific IgE antibodies against Japanese cedar pollen by MAST-26 (Hitachi Chemical, Tokyo, Japan)⁵ and in 2015 sensitization to Japanese cedar pollen, defined as specific IgE class 1 or above by ImmunoCAP (Phadia, Uppsala, Sweden), was detected in 39% of lower-grade schoolchildren (aged 6–9 years).⁶

Japanese cedar pollinosis is a multifactorial disease, and genetic⁷ and environmental factors^{8–10} have been known to influence disease development. Familial clustering and intra-individual clustering suggest that overlapping genetic factors influence the development of allergic diseases. Many environmental factors have

* Corresponding author. Department of Otorhinolaryngology, Mie University Graduate School of Medicine, 2-174 Edobashi, Tsu, Mie 514-8507, Japan.

E-mail address: kazuhiko@clin.medic.mie-u.ac.jp (K. Takeuchi).

Peer review under responsibility of Japanese Society of Allergy.

¹ These two authors equally contributed to this project.

<https://doi.org/10.1016/j.alit.2018.05.006>

1323–8930/Copyright © 2018, Japanese Society of Allergy. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

been investigated as influential factors for the development of allergic rhinitis. It has been reported that increased amounts of antigens,⁸ living environment,¹⁰ history of pet ownership,¹¹ smoking,^{12–16} dietary habits,^{17,18} and house dust mite sensitization¹⁹ are associated with allergic rhinitis. Allergic rhinitis, and cedar pollen allergy in particular, imposes a high socioeconomic burden. Given the recent increases in cedar pollen allergy in Japan, effective preventive measures as well as effective treatments are important to determine. The elucidation of factors associated with the development of cedar pollen allergy will help effective strategies for prevention to be drawn up.

The development of allergic rhinitis occurs in two steps, sensitization to allergens and development of symptoms, and each step involves different mechanisms. Therefore, the influential factors for sensitization and symptomatic onset must be identified separately. This study sought to clarify the factors that influence the sensitization and symptomatic development of cedar pollinosis in a single population. Two parameters, self-reported subjective symptoms of cedar pollinosis and physician-diagnosed cedar pollinosis, were used to evaluate the development of pollinosis. Moreover, sensitization to Japanese cypress and house dust mites was evaluated, and the factors associated with sensitization to each of the 3 allergens were determined.

Methods

Subjects and ethical considerations

Initially, the subjects in this cross-sectional study were 590 volunteers (260 males and 330 females; age range 5–80 years) who were students, workers, or residents living in Mie Prefecture, Japan, regardless of whether they were symptomatic for cedar pollen allergy. We mounted posters to recruit volunteers. The study was conducted from August 2013 to March 2017 and adhered to the tenets of the Declaration of Helsinki. The study protocol was approved by Mie University Ethics Committee (No.2590), and written informed consent was obtained from all subjects or their legal guardians.

Questionnaire

The questionnaire included questions to collect data in two broad categories: (1) general data on subject profiles and (2) data associated with allergies, and more specifically with cedar pollen allergies. The questionnaire (Supplementary Methods 1) included items on age, sex, occupation, and height and weight of the subject. The questions on allergies surveyed (1) self-reported subjective symptoms such as sneezing, nasal discharge and/or congestion, and itchy and/or teary eyes lasting for at least 2 weeks during the cedar pollen dispersal period (February–March) in the absence of a cold and (2) physician-diagnosed cedar pollen allergy. Data were collected on family history of cedar pollen allergy (among the subject's father, mother, siblings, grandfathers, grandmothers, and children) and on any concomitant and family history of allergic diseases (e.g., allergic rhinitis other than cedar pollen allergy, food allergies, hives, atopic dermatitis, oral allergy syndrome, and asthma). Also recorded was any past or concomitant history of sinusitis, angina, hyperlipidemia, diabetes, obstructive sleep apnea syndrome, cerebral infarction, or gastroesophageal reflux disease. Subjects also answered questions on the number of colds per year, history of smoking and passive smoking, number of cohabitants, current living environment, main site of daytime activity, frequency of playing outside during childhood, stress, regular exercise habits, and history of owning pets such as dogs and cats.

Questions about diet included the frequency of consumption of yogurt, meat, fish, fast food, beer, Japanese sake, wine, carbonated soft drinks, coffee, butter, eggs, fruit, margarine, milk, nuts, noodles and pasta, bread, rice, potatoes, vegetables, and soy products. There were three possible answers for frequency of consumption: (1) very rarely, (2) 1–2 times per week, and (3) ≥ 3 times per week.

Measurement of IgEs

Levels of specific IgE antibodies against cedar (*Cryptomeria japonica*), cypress (*Chamaecyparis obtusa*), and mites (*Dermatophagoides pteronyssinus*) and total IgE antibody levels were measured by CAP-FEIA (fluorescent enzyme immunoassay) (SRL, Tokyo, Japan). Sensitization to cedar pollen was defined as specific IgE levels of class 1 or above based on blood test results alone. Development of cedar pollen allergy was defined as (1) having pollinosis symptoms such as sneezing, nasal discharge and/or congestion, and itchy and/or teary eyes lasting for at least 2 weeks during the cedar pollen dispersal period every year (February–March) in the absence of a cold and (2) previous diagnosis of cedar pollen allergy by a physician.

Statistical analysis

Univariate analysis with chi-square and Mann–Whitney *U* tests were used to determine sensitization to Japanese cedar pollen and development of pollinosis evaluated by subjective symptoms or physician diagnosis. Factors that exhibited strong correlations ($p < 0.2$) were further examined using multiple logistic regression analysis. Analyses were performed with SPSS statistical software version 21 (IBM, Chicago, IL). The same models were applied to identify factors associated with mites or cypress sensitization. A *p* value of less than 0.05 was considered statistically significant.

Results

Subjects

Questionnaires were received for a total of 590 subjects, and blood samples were taken from all subjects. Figure 1 shows the age distribution of the 590 subjects. Given the high number of subjects in their twenties, a subanalysis was performed among subjects in




Fig. 1. Age distribution of the 590 subjects in this study. Given the large number of subjects in their twenties, a subanalysis was performed among subjects in the age range 20–29 years.

the age range 20–29 years ($n = 382$; 188 men, 194 women; mean age 23.6 years; age range 20–29 years). Mean height, body weight, and BMI of these 382 subjects were 166 cm, 59.4 kg, and 21.2, respectively. The findings reported below are confined to this young adult population only.

Prevalence

Among these 382 young adult subjects, 41.2% reported symptoms in the cedar pollen scattering period, and 22.2% had physician-diagnosed cedar pollinosis (Table 1). The prevalence rates of comorbidities and family history of comorbidities such as allergic rhinitis other than cedar pollinosis, food allergy, urticarial rash, atopic dermatitis, oral allergy syndrome, and asthma are shown in Table 1. Of the 382 subjects, 9.9% reported they have the common cold ≥ 5 times per year. Smokers constituted 8.6%. Dogs and cats were kept as a pet by 37.2% and 13.9%, respectively. Sensitization to cedar, cypress, and mites, defined as specific IgE levels of class 1 or above, was present in 78.8%, 64.4%, and 56.0%, respectively (Table 1, Fig. 2).

Factors associated with cedar pollinosis

The results of univariate analysis are shown in Table 2, 3. Only those items that were statistically significant ($p < 0.05$) are shown.

The results of multiple logistic regression analysis are shown in Table 4, 5. As shown in Figure 3A, the factors associated with increased cedar pollen sensitization were mite sensitization, comorbid allergic rhinitis other than cedar pollinosis, and family history of cedar pollinosis (Fig. 3A). Three factors were associated with reduced risk of cedar pollen sensitization: keeping a pet cat, hours of sleep, and number of common colds per year. The factors associated with increased cypress pollen sensitization were serum total IgE level and family history of cedar pollinosis. Those associated with reduced risk of cypress pollen sensitization were number of common colds per year, BMI, stress, smoking, consumption of margarine, and consumption of fast foods (Fig. 3A).

Table 1

Prevalence of antigen sensitization, cedar pollinosis, comorbidities, family history, and lifestyle among young Japanese adults aged 20–29 years ($n = 382$).

		%	Present	Absent
Sensitization	<i>Dermatophagoides pteronyssinus</i>	56.0	214	168
	<i>Cryptomeria japonica</i>	78.8	301	81
	<i>Chamaecyparis obtusa</i>	64.4	246	136
	Self-reported symptoms of cedar pollinosis	41.2	160	222
Physician-diagnosed cedar pollinosis		22.2	85	297
	Allergic rhinitis	26.2	100	282
	Food allergy	7.9	30	352
	Urticaria	17.0	65	317
Comorbidity	Atopic dermatitis	21.2	81	301
	Oral allergy syndrome	4.5	21	361
	Asthma	12.8	49	333
	Cedar pollinosis	68.8	263	119
Family history	Allergic rhinitis	32.3	125	257
	Asthma	15.2	58	324
	Atopic dermatitis	18.6	71	311
	Conjunctivitis	4.7	18	364
Catching the common cold ≥ 5 times per year		16.2	62	320
	Food allergy	2.9	11	371
	Oral allergy syndrome	9.9	38	344
	Smoking	8.6	33	349
Keeping a pet dog		37.2	142	240
Keeping a pet cat		13.9	53	329

The factors associated with increased house dust mite sensitization were serum total IgE level and comorbid asthma, while those associated with a reduced risk of mite sensitization were consumption of eggs and milk (Fig. 3A).

The factors associated with increased subjective symptoms of pollinosis were concomitant allergic rhinitis other than cedar pollen allergy and family history of cedar pollen allergy (Fig. 3B). Consumption of butter was a risk-reducing factor for subjective pollinosis symptoms (Fig. 3B).

The factors associated with physician diagnosis were concomitant allergic rhinitis other than cedar pollen allergy, family history of cedar pollinosis, and family history of conjunctivitis. History of consuming carbonated soft drinks was a risk-reducing factor for physician diagnosis (Fig. 3B).

Discussion

The present study has been performed in Mie Prefecture. Geographically, Mie is situated close to the center of Japan, along the Pacific Ocean. The Prefecture has a long, thin shape, being longer in the north-south direction than in the east-west direction. It takes approximately 3 h to reach Tsu City (the capital of Mie) from Tokyo and 50 min from Nagoya. The annual average temperature for the Ise Plain is 15 °C which is relatively warm for Japan. As a result, the annual cedar pollen count is larger than the average and the prevalence of cedar pollinosis is higher in Mie (33.3%) than the national average (26.5%).²⁰

The rates of sensitization to cedar pollen and the prevalence of pollinosis differ considerably according to age, period, and degree of urbanization. In this study that focused on a young adult population in their twenties, the sensitization rate to Japanese cedar pollen was 78.8% and subjective symptoms were reported by 41.2% of subjects during the pollen season. The sensitization rate in the present study is higher than that in any other previous report.^{3–6} The rate of sensitization and prevalence of cedar pollinosis depends on the age groups. This is partly because the cedar pollen count increased recently and sensitization started to occur in the younger ages than before.²⁰

Sensitization to cedar pollen was defined as specific IgE levels of class 1 or above. Before we conducted statistics, we calculated percentage of subjects having pollinosis symptoms according to seven classes of cedar pollen specific IgE. As a result, subjects with class 1 had higher percentage of pollinosis than those with class 0. Thus we defined class 1 or more as sensitization.

The factors associated with increased cedar pollen sensitization were mite sensitization, comorbid allergic rhinitis other than cedar pollinosis, and family history of cedar pollinosis. According to a survey of first-grade schoolchildren conducted by Kanazawa et al.,²¹ children who were sensitized to mites but not to cedar had significantly elevated cedar pollen specific IgE levels compared with the group not sensitized to mites. This finding indicates that mite sensitization is very strongly associated with cedar pollen sensitization. Sensitization to house dust mites is the only factor associated with sensitization to cedar pollen in asymptomatic subjects.¹⁹ Mite sensitization can potentially cause minimal persistent inflammation^{22,23}; thus, repeated exposure to mite antigens at amounts that may not trigger allergies may still cause inflammation and a subsequent deficit in the defensive function of the nasal mucosa.

Two risk factors were common to both cedar pollen sensitization and the development of cedar pollinosis, regardless of the two different diagnostic criteria. The first risk factor was concomitant allergic rhinitis other than cedar pollen allergy. Inflammation of the nasal mucosa lowers the defensive function of the nasal epithelium and facilitates the entry of cedar pollen allergens into the body;




Fig. 2. Distribution of specific IgE classes of the three antigens cedar, cypress, and mite among 382 subjects. Sensitization to cedar (*Cryptomeria japonica*), cypress (*Chamaecyparis obtusa*), and mites (*Dermatophagoides pteronyssinus*) was found in 78.8%, 64.4%, and 56.0% of subjects, respectively.

Table 2
Result of univariate analysis on sensitization to three antigens.

	Mites (P value)	Cedar (P value)	Cypress (P value)
Total IgE	0.000	0.000	0.000
Mite IgE class	NS	0.000	0.004
Cedar IgE class	0.000	NS	NS
Age	NS	NS	0.020
Body weight	NS	NS	0.045
BMI	NS	NS	0.014
Asthma	0.000	0.006	NS
Food allergy	0.006	0.013	0.024
Atopic dermatitis	0.000	0.005	0.001
Allergic rhinitis other than cedar pollinosis	0.000	0.001	NS
GERD	0.023	NS	NS
Otitis media with effusion	NS	0.006	NS
Family history	NS	0.004	0.000
Cedar pollinosis	NS	NS	NS
Asthma	0.015	NS	NS
Atopic dermatitis	0.029	NS	NS
Smoking	0.043	NS	0.017
Keeping a pet cat	NS	0.037	NS
Number of colds	NS	0.004	0.002
Fast food	NS	NS	0.042
Margarine	NS	NS	0.025
Carbonated drinks	NS	NS	0.006
Vegetables	NS	0.013	NS
Eggs	0.047	NS	NS
Milk	0.006	NS	NS

BMI, body mass index; GERD, gastroesophageal reflux disease; NS, not significant.

Table 3
Result of univariate analysis on pollinosis symptoms and diagnosis.

	Subjective symptoms (P value)	Physician diagnosis (P value)
Food allergy	0.036	NS
Allergic rhinitis other than cedar pollinosis	0.000	0.000
OAS	NS	0.004
Family history	NS	0.001
Cedar pollinosis	0.000	0.001
Atopic dermatitis	NS	0.023
Conjunctivitis	0.008	0.000
OAS	0.035	0.009
Keeping dogs as pets	0.024	NS
Meat	0.043	NS
Butter	0.014	NS
Carbonated drinks	NS	0.003

OAS, oral allergy syndrome; NS, not significant.

smaller in the allergic mucosa than that required to trigger allergy in normal mucosa.²² The second risk factor common to sensitization and development was family history of cedar pollinosis, which has previously been reported to be an important risk factor for both.^{24,25} One reason may be the greater likelihood of genetic polymorphisms shared by blood relatives. Various genetic polymorphisms are associated with allergic rhinitis. These specifically include interleukin-33⁷, which causes allergic inflammation by inducing eosinophil chemotaxis and cytokine production, and the expression of filaggrin,²⁶ which is important for skin barrier function.

The present study revealed some protective factors against sensitization to cedar and cypress. These included the number of common colds, hours of sleep, keeping a cat, and smoking. One of the most striking findings in this study was that the number of colds inversely correlated with sensitization to cedar (OR: 0.186, 95% CI: 0.078–0.448) and cypress (OR: 0.324, 95% CI: 0.141–0.743). The sensitization rate of those who have the common cold ≥ 5 times

increased allergen levels within the body are hypothesized to increase the likelihood of sensitization. In addition, cells such as eosinophils and neutrophils infiltrate the nasal mucosa, thereby increasing sensitivity in patients with allergic rhinitis. This is suggested to increase the likelihood of a cedar pollen allergy developing because the amount of cedar pollen allergens would be

Table 4

Result of multiple regression analysis on sensitization to three antigens.

	House dust mites		Japanese cedar		Japanese cypress	
	Odd ratio (95%CI)	P value	Odd ratio (95%CI)	P value	Odd ratio (95%CI)	P value
Total IgE	1.005 (1.003–1.007)	0.000		NS	1.005 (1.003–1.007)	0.000
Mite IgE class		NA	1.571 (1.311–1.882)	0.000		NS
BMI		NS		NS	0.900 (0.821–0.987)	0.025
Asthma	8.067 (2.504–25.988)	0.000		NS		NS
Allergic rhinitis other than cedar pollinosis		NS	3.182 (1.351–7.494)	0.008		NS
Family history of cedar pollinosis		NS	2.8769 (1.608–5.142)	0.000	2.474 (1.456–4.203)	0.001
Keeping cats as pets		NS	0.418 (0.196–0.892)	0.024		NS
Hours of sleep		NS	0.666 (0.480–0.925)	0.015		NS
Number of colds		NS	0.186 (0.078–0.448)	0.000	0.324 (0.141–0.743)	0.008
Smoking		NS		NS	0.338 (0.139–0.821)	0.017
Stress		NS		NS	0.514 (0.309–0.854)	0.010
Fast food ≥ 3 per week		NS		NS	0.025 (0.001–0.683)	0.029
Margarine 1–2 per week		NS		NS	0.321 (0.107–0.961)	0.042
Eggs 1–2 per week	0.286 (0.094–0.871)	0.028		NS		NS
Milk ≥ 3 per week	0.446 (0.236–0.843)	0.013		NS		NS

BMI, body mass index; CI, confidence interval; NS, not significant; NA, not applicable.

Table 5

Result of multiple regression analysis on pollinosis symptoms and diagnosis.

	Subjective symptoms		Physician diagnosis	
	Odds ratio (95%CI)	P value	Odds ratio (95%CI)	P value
Allergic rhinitis other than cedar pollinosis	2.799 (1.696–4.619)	0.000	4.204 (2.536–7.256)	0.000
Family history				
Cedar pollinosis	3.539 (2.132–5.876)	0.000	2.482 (1.282–4.802)	0.007
Allergic conjunctivitis		NS	4.289 (1.481–12.416)	0.007
Butter ≥ 3 per week	0.294 (0.126–0.685)	0.005		NS
Carbonated drinks 1–2 per week		NS	0.483 (0.261–0.896)	0.021

CI, confidence interval; NS, not significant.

per year was 60.5%, while it was 80.8% in those who have colds <5 times. We defined susceptibility to cold as having cold five times or more per year, because it is reported that the common cold is estimated to occur an average of 2–5 times for adults in the general healthy population.²⁷ To our knowledge, the inverse correlation between the sensitization to pollen antigens and the number colds has not been reported before in the literature. It is believed that wheezing associated with viral infections is a pivotal risk factor for the development of asthma.²⁸ However, some viral infections are

reported to protect against persistent IgE sensitization,²⁹ and certain viral infections caught during the sensitization and challenge phases reduce the development of delayed eosinophilic allergic rhinitis in BALB/c mice.³⁰ Allergic rhinitis was reportedly associated with decreased expression of TLR9 in sinonasal epithelial cells.³¹

The above results have implications for the treatment of cedar pollinosis. Microbial factors likely modulate nasal innate immunity to maintain homeostasis. An approach targeting TLR9 by synthetic




Fig. 3. Factors linked to sensitization to house dust mites, cedar pollen, and cypress (A). Factors linked to self-reported subjective symptoms and physician diagnosis of Japanese cedar pollinosis (B). Upward arrows indicate positively correlated factors and downward arrows indicate negatively correlated factors.

cytosine phosphate-guanosine oligodeoxynucleotides (CpG ODN) is a promising new treatment paradigm for modulating the immune response.³² A clinical trial of immunotherapy with a ragweed-TLR9 agonist vaccine in adults allergic to ragweed has been proven successful.³³ Animal models have elucidated the mechanism of action of this therapy. CpG ODN is likely recognized by TLR9 on conventional dendritic cells and alveolar macrophages, which produce mRNA encoding IL-12. IL-12 is necessary for the subsequent production of innate and adaptive interferon- γ .³⁴ Mice that received CpG showed reduced IgE antibody production at both neonatal and adult stages.³⁵

In this study, keeping a cat was found to have a protective effect against sensitization to cedar pollen (OR: 0.418, 95% CI 0.196–0.892). Current cat ownership was related to significantly lower rates of allergic rhinitis (adjusted OR [aOR]: 0.71, 95% CI 0.57–0.89) and Japanese cedar pollinosis (aOR: 0.57, 95% CI 0.44–0.75). However, there was no association between the prevalence of allergies and dog ownership.¹¹ It remains unclear why the association observed with cats did not extend to dogs. However, owning a cat has been shown to result in the accumulation of large amounts of endotoxins in the living environment (OR: 1.91; 95% CI: 1.43–2.55),¹⁰ and there is a potentially larger bacterial load, including that of colon bacilli, indoors. The reason why keeping a cat has a protective effect only to cedar pollen but not mite is currently unknown and should be investigated in the future study.

In this study, smoking was a protective factor for sensitization to cypress (OR: 0.338, 95% CI: 0.139–0.821). Current smokers were at a significantly lower risk of cedar pollinosis among men (hazard ratio [HR]: 0.64, 95% CI: 0.50–0.83) and women (HR: 0.64, 95% CI: 0.47–0.88).¹⁵ Strikingly, it has been reported that passive smoking was also related to a significantly lower rate of allergic rhinitis (aOR: 0.83, 95% CI: 0.77–0.89) and Japanese cedar pollinosis (aOR: 0.81, 95% CI: 0.74–0.88).¹¹ The underlying mechanism is likely to be cigarette smoke-mediated inhibition of inflammatory cytokine production and T cell suppression, which would also trigger sensitization.^{12,13} In the present study, we were unable to detect an association between allergy onset and history of smoking; however, Eriksson *et al.* previously reported that the prevalence of allergic rhinitis was significantly lower in males with a history of smoking.¹⁴ Although smoking has a protective effect against allergic rhinitis, active, passive, and electronic cigarette smoking is associated with asthma in adolescents.¹⁶

Duration of sleeping hours was also a protective factor. The present result means the negative correlation between the duration of sleeping hours and the sensitization rate, which indicates that short sleeping hours may increase risk of cedar pollen sensitization. Although the precise mechanism is unclear, it is known that sleep deprivation has detrimental effects on metabolic and immune regulation.³⁶ Zhang *et al.* reported that short sleep duration is associated with the risk of sensitization to food and aero allergens in rural Chinese adolescents.³⁷ Thus, it is probable that risk of allergic sensitization can be reduced by appropriate guidance on sleep duration.

This study showed an association between consuming several foods and decreased sensitization and development of pollinosis. Eggs and milk were inversely associated with sensitization to mites, and fast foods and margarine were inversely associated with sensitization to cypress. Butter and carbonated drinks were inversely associated with subjective symptoms and physician-diagnosed pollinosis, respectively. Reportedly, an increased risk of rhinoconjunctivitis was associated with the consumption of fast foods, margarine, and butter and a decreased risk with eggs and milk.¹⁷ The reason why fast foods, margarine, and butter were associated with reduced rates of sensitization and pollinosis is

unclear. It is possible that individuals sensitized to pollen and those with pollinosis avoided consuming these foods.

Although our results showed that the consumption of carbonated soft drinks was a protective factor against pollinosis based on physician diagnosis, no previous reports support this finding. Carbonated water may promote sympathetic nerve activity and ease the symptoms of allergic rhinitis such as nasal discharge and/or congestion, which are associated with parasympathetic nerve activation, thus delaying the onset of pollinosis. However, it will be a challenge to determine the mechanism underlying the apparent protective factor of consuming soft drinks 1–2 times per week.

Total IgE level, which was not identified as a risk factor for cedar pollen sensitization, was found to be associated with mite sensitization. Total IgE level can be used to estimate the contribution of an individual's characteristics to IgE antibody production and the strength of that individual's predisposition to atopy. Thus, our results demonstrate that although predisposition to atopy was not associated with the development of cedar pollen sensitization, mite sensitization was more likely to occur in people who exhibited a stronger predisposition to atopy. Our results showed that there is a causal relationship between mite IgE and cedar pollen sensitization. This was previously confirmed by studies showing that mite sensitization, which appears to be associated with a predisposition to atopy, occurs at a younger age than cedar pollen sensitization.^{38,39}

One of the limitations of this study is that we did not consider the specialties of physicians who made diagnosis of pollinosis. Diagnosis of cedar pollinosis may be different between specialized ENT physicians and general family physicians.

Skin prick test is commonly used for the detection of causative allergen because of the high sensitivity, rapidity and inexpensiveness. However, allergen-specific IgE blood assay is widely used for clinics in our country and we also used serum IgE in this study. The major reason for this is no risk of severe allergic reaction in serum IgE measurement. However, serum IgE measurement may result in underestimation of sensitization status to allergens.

The sensitization rate to cedar pollen in a young Japanese adult population was extremely high. Both common and distinct factors were associated with sensitization to pollen and the development of pollinosis. Distinct factors were associated with sensitization to cedar and cypress antigens.

Acknowledgements

We thank Ms. Mayu Nakatani for assistance with this study. This study was supported in part by a grant from the Ministry of Health, Labor and Welfare in Japan (Prevention and treatment of immunology and allergy disease; Chief: Yoshitaka Okamoto, H23-006), Takeda Science Foundation, and by a discretionary budget allocation from the director of Mie University Hospital.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.alit.2018.05.006>.

Conflict of interest

The authors have no conflict of interest to declare.

Authors' contributions

SM, HS, and KT designed the study. KT wrote the manuscript. SAS contributed to data collection. ST, SN, and KT performed the statistical analysis and interpretation of the results. All authors read and approved the final manuscript.

References

1. Yamada T, Saito H, Fujieda S. Present state of Japanese cedar pollinosis: the national affliction. *J Allergy Clin Immunol* 2014;**133**:632–9.
2. Ozasa K, Hama T, Dejima K, Watanabe Y, Hyo S, Terada T, et al. A 13-year study of Japanese cedar pollinosis in Japanese schoolchildren. *Allergol Int* 2008;**57**:175–80.
3. Kaneko Y, Motohashi Y, Nakamura H, Endo T, Eboshida A. Increasing prevalence of Japanese cedar pollinosis: a meta-regression analysis. *Int Arch Allergy Immunol* 2005;**136**:365–71.
4. Sakashita M, Hirota T, Harada M, Nakamichi R, Tsunoda T, Osawa Y, et al. Prevalence of allergic rhinitis and sensitization to common aeroallergens in a Japanese population. *Int Arch Allergy Immunol* 2010;**151**:255–61.
5. Yoda S, Enomoto T, Dake Y, Ikeda H, Shibano A, Sakoda T, et al. [Epidemiological survey of allergic diseases in first-year junior high school students in Wakayama Prefecture in 2003]. *Nihon Jibiinkoka Gakkai Kaiho* 2006;**109**:742–8 (in Japanese).
6. Yamazaki S, Shima M, Nakadate T, Ohara T, Omori T, Ono M, et al. Patterns of sensitization to inhalant allergens in Japanese lower-grade schoolchildren and related Factors. *Int Arch Allergy Immunol* 2015;**167**:253–63.
7. Sakashita M, Yoshimoto T, Hirota T, Harada M, Okubo K, Osawa Y, et al. Association of serum interleukin-33 level and the interleukin-33 genetic variant with Japanese cedar pollinosis. *Clin Exp Allergy* 2008;**38**:1875–81.
8. Honda K, Saito H, Fukui N, Ito E, Ishikawa K. The relationship between pollen count levels and prevalence of Japanese cedar pollinosis in Northeast Japan. *Allergol Int* 2013;**62**:375–80.
9. Nakamura K, Nagata C, Wada K, Fujii K, Kawachi T, Takatsuka N, et al. Association of farming with the development of cedar pollinosis in Japanese adults. *Ann Epidemiol* 2010;**20**:804–10.
10. Bischof W, Koch A, Gehring U, Fahlbusch B, Wichmann HE, Heinrich J. Indoor Exposure and Genetics in Asthma Study Group. Predictors of high endotoxin concentrations in the settled dust of German homes. *Indoor Air* 2002;**12**:2–9.
11. Kurosaka F, Nakatani Y, Terada T, Tanaka A, Ikeuchi H, Hayakawa A, et al. Current cat ownership may be associated with the lower prevalence of atopic dermatitis, allergic rhinitis, and Japanese cedar pollinosis in schoolchildren in Himeji, Japan. *Pediatr Allergy Immunol* 2006;**17**:22–8.
12. Sopori M. Effects of cigarette smoke on the immune system. *Nat Rev Immunol* 2002;**2**:372–7.
13. Ouyang Y, Virasch N, Hao P, Aubrey MT, Mukerjee N, Brier BE, et al. Suppression of human IL-1 β , IL-2, IFN- γ , and TNF- α production by cigarette smoke extracts. *J Allergy Clin Immunol* 2000;**106**:280–7.
14. Eriksson J, Ekerljung L, Sundblad BM, Lötval J, Torén K, Rönmark E, et al. Cigarette smoking is associated with high prevalence of chronic rhinitis and low prevalence of allergic rhinitis in men. *Allergy* 2013;**68**:347–54.
15. Nagata C, Nakamura K, Fujii K, Kawachi T, Takatsuka N, Oba S, et al. Smoking and risk of cedar pollinosis in Japanese men and women. *Int Arch Allergy Immunol* 2008;**147**:117–24.
16. Kim SY, Sim S, Choi HG. Active, passive, and electronic cigarette smoking is associated with asthma in adolescents. *Sci Rep* 2017;**7**:17789.
17. Ellwood P, Asher MI, García-Marcos L, Williams H, Keil U, Robertson C, et al. ISAAC Phase III Study Group. Do fast foods cause asthma, rhinoconjunctivitis and eczema? Global findings from the International Study of Asthma and Allergies in Childhood (ISAAC) phase three. *Thorax* 2013;**68**:351–60.
18. Miyake Y, Sasaki S, Tanaka K, Ohya Y, Miyamoto S, Matsunaga I, et al. Osaka maternal and child Health study group. Fish and fat intake and prevalence of allergic rhinitis in Japanese females: the Osaka maternal and child Health study. *J Am Coll Nutr* 2007;**26**:279–87.
19. Sakaida H, Masuda S, Takeuchi K. Analysis of factors influencing sensitization of Japanese cedar pollen in asymptomatic subjects. *Auris Nasus Larynx* 2013;**40**:543–7.
20. Okubo K, Kurono Y, Ichimura K, Enomoto M, Okamoto Y, Kawauchi Y, et al., editors. [Practical Guideline for the Management of Allergic Rhinitis in Japan]. 8th ed. Tokyo: Life Science; 2016 (in Japanese).
21. Kanazawa A, Terada T, Ozasa K, Hyo S, Araki N, Kawata R, et al. Continuous 6-year follow-up study of sensitization to Japanese cedar pollen and onset in schoolchildren. *Allergol Int* 2014;**63**:95–101.
22. Canonica GW, Compalati E. Minimal persistent inflammation in allergic rhinitis: implications for current treatment strategies. *Clin Exp Immunol* 2009;**158**:260–71.
23. Ricca V, Landi M, Ferrero P, Bairo A, Tazzer C, Canonica GW, et al. Minimal persistent inflammation is also present in patients with seasonal allergic rhinitis. *J Allergy Clin Immunol* 2000;**105**:54–7.
24. Warm K, Backman H, Lindberg A, Lundbäck B, Rönmark E. Low incidence and high remission of allergic sensitization among adults. *J Allergy Clin Immunol* 2012;**129**:136–42.
25. Li CW, Chen DH, Zhong JT, Lin ZB, Peng H, Lu HG, et al. Epidemiological characterization and risk factors of allergic rhinitis in the general population in Guangzhou City in China. *PLoS One* 2014;**9**:e114950.
26. Imoto Y, Enomoto H, Fujieda S, Okamoto M, Sakashita M, Susuki D, et al. S2554X mutation in the flaggrin gene is associated with allergen sensitization in the Japanese population. *J Allergy Clin Immunol* 2010;**125**:498–500.e2.
27. Monto AS. Epidemiology of viral respiratory infections. *Am J Med* 2002;**112**:4S–12.
28. Jackson DJ, Gern JE, Lemanske Jr RF. Lessons learned from birth cohort studies conducted in diverse environments. *J Allergy Clin Immunol* 2017;**139**:379–86.
29. Saghaian-Hedengren S, Sverremark-Ekström E, Linde A, Lilja G, Nilsson C. Early-life EBV infection protects against persistent IgE sensitization. *J Allergy Clin Immunol* 2010;**125**:433–8.
30. Sasaki Y, Hayashi T, Hasegawa K. Lactate dehydrogenase-elevating virus infection at the sensitization and challenge phases reduces the development of delayed eosinophilic allergic rhinitis in BALB/c mice. *Scand J Immunol* 2007;**66**:628–35.
31. Melvin TA, Nguyen MT, Lane AP, Lin SY. Allergic rhinitis is associated with decreased expression of Toll-like receptor 9 by sinonasal epithelial cells. *Int Forum Allergy Rhinol* 2011;**1**:153–6.
32. Ihler F, Canis M. Ragweed-induced allergic rhinoconjunctivitis: current and emerging treatment options. *J Asthma Allergy* 2015;**8**:15–24.
33. Creticos PS, Schroeder JT, Hamilton RG, Balcer-Whaley SL, Khattignavong AP, Lindblad R, et al. Immune tolerance network group. *N Engl J Med* 2006;**355**:1445–55.
34. Lee LM, Ji M, Sinha M, Dong MB, Ren X, Wang Y, et al. Determinants of divergent adaptive immune responses after airway sensitization with ligands of toll-like receptor 5 or toll-like receptor 9. *PLoS One* 2016;**11**:e0167693.
35. de Brito CA, Fusaro AE, Victor JR, Rigato PO, Goldoni AL, Muniz BP, et al. CpG-induced Th1-type response in the downmodulation of early development of allergy and inhibition of B7 expression on T cells of newborn mice. *Int Forum Allergy Rhinol* 2011;**1**:153–6.
36. Bryant PA, Trinder J, Curtis N. Sick and tired: does sleep have a vital role in the immune system? *Nat Rev Immunol* 2004;**4**:457–67.
37. Zhang S, Liu X, Kim JS, Ouyang F, Wang B, Li Z, et al. Association between short sleep duration and the risk of sensitization to food and aero allergens in rural Chinese adolescents. *Clin Exp Allergy* 2011;**41**:547–55.
38. Mori A. [Sensitization and onset of Japanese cedar pollinosis in children]. *Arerugi* 1995;**44**:7–15 (in Japanese).
39. Takano S, Aramaki H, Kitajima S, Ueda N, Shigi K. [Relation between the development of nasal allergy]. *Practica Oto-Rhino-Laryngologica* 1997;**90**:1013–7 (in Japanese).

Medical interview sheet

This medical interview sheet is for the clinical examination: "Research on Sensitization and development of cedar pollinosis"

Please fill out the form with ☒ and also write something if necessary

Personal information will be protected..

Date		year		month		Day
age		Age	work		Have work	
gender		Male			No work (housework etc)	
		Female			student	
height		cm	weight		kg	Residing city

I. Asking about your health condition including allergy

1. Have you ever had symptoms of sneezing, watery rhinorrhea, nose blocking, itchy eyes, tearing for more than two weeks continuously in February to March (time of cedar pollen scattering) without having cold?

	No
	Every year
	sometimes
	This year for the first time

2. Have you ever been told that you have cedar allergy by a medical doctor?

	No
	I didn't see a doctor but I think I have cedar allergy.
	I was told so by a doctor.※

※When did you see a doctor and doctor told you that you have cedar allergy? Please write number below.

	When I was () years old		() years ago
--	--------------------------	--	---------------

3. Have you ever told that you have allergic rhinitis other than cedar pollinosis?

	No
	Yes※

※If yes; please write the cause if you know it.

--

Supplementary Methods 1

4. Have you ever had below mentioned allergies? Please mark ☒.

☐	allergic conjunctivitis	☐	Food allergy
☐	hives	☐	atopic dermatitis
☐	Oral Allergy Syndrome※	☐	asthma

※Oral Allergy Syndrome: When you eat fruits and vegetables, you will feel itchy or uncomfortable in your mouth.

5. Have you ever had any diseases below or do you have any? Please mark ☒.

☐	sinusitis	☐	angina
☐	hypertension	☐	dyslipidemia
☐	diabetes mellitus	☐	sleep apnea syndrome
☐	brain stroke	☐	gastroesophageal regurgitation

6. How often do you have cold?

☐	5 times or more per year
☐	Less than 5 times a year

II. Asking other things.

1. Does your family have cedar pollinosis?

☐	no
☐	yes※

※If yes, who has one?

☐	father	☐	mother	☐	brother
☐	grandfather	☐	grandmother	☐	child

2. Does your family have any other allergic rhinitis other than cedar pollinosis?

☐	No
☐	Yes ※

※If yes, who it is?

☐	father	☐	mother	☐	brother
☐	grandfather	☐	grandmother	☐	child

3. Does your family have asthma?

☐	no
☐	yes※

※If yes, who it is?

Supplementary Methods 1

	father		mother		brother
	grandfather		grandmother		child

4. Does your family have atopic dermatitis?

	no
	yes✕

✕If yes, who it is?

	father		mother		brother
	grandfather		grandmother		child

5. Does your family have allergic conjunctivitis?

	no
	yes✕

✕If yes, who it is?

	father		mother		brother
	grandfather		grandmother		child

6. Does your family have food allergy?

	no
	yes✕

✕If yes, who it is?

	father		mother		brother
	grandfather		grandmother		child

7. Does your family have aral allergy syndrome?

	no
	yes✕

✕If yes, who it is?

	father		mother		brother
	grandfather		grandmother		child

8. Do you smoke?

	no
	yes
	Not now, before

Supplementary Methods 1

9. Does your family smoke in your house?

	no
	yes※

※If yes, who it is?

	father		mother		brother
	grandfather		grandmother		child
	partner		others		

1 0. How many brothers and sisters do you have including yourself? Please write number. And please write in which order are you in your brothers and sisters?

	Numbers of brothers and sisters		I am the ()th
--	---------------------------------	--	----------------

1 1. How many people do you live together including yourself? Please write number.

	persons
--	---------

1 2. Tell me about surroundings of your house

	suburb (residence area)
	village (surrounded by field, forest and nature)
	others

1 3. While you are working or in a daytime, do you spend your time inside or outside?

	mainly inside
	mainly outside

1 4. When you were a child, where were you playing around?

	often played outside
	seldom spend time outside

1 5. Are you vulnerable to stress?

	Yes
	Not really.

1 6. How many hours do you work a day? (if you work).

	hours
--	-------

1 7. How long is your average sleeping hours?

Supplementary Methods 1

	hours
--	-------

- 1 8. Do you exercise more than 30 minutes a day for twice a week or more, and continue for more than a year?

	Yes
	No

- 1 9. Do you have pets? Or have you had pets? Please mark ☒.

	no	Yes I have now.	Not now but before I had
dog			
cat			
other (write what)			

- 2 0. How often do you eat below mentioned items a week? Please mark ☒.

	seldom	1-2 a week	More than 3 times a week
Yogurt			
Meat			
Fish			
Fast food like burgers			
Beer			
Japanese sake			
Wine			
Soda			
Green tea, black tea, coffee			
Butter			
Egg			
Fruit			
Margarine			
Milk			
Nuts			
Noodles			
Bread			
Rice			
Potatoes			
Vegetables			
Bean product (Miso soup, tofu)			

This is the end of the interview. Thank you very much.