

Advance Publication

The Journal of Veterinary Medical Science

Accepted Date: 25 November 2019

J-STAGE Advance Published Date: 25 December 2019

©2019 The Japanese Society of Veterinary Science

Author manuscripts have been peer reviewed and accepted for publication but have not yet been edited.

Internal Medicine

Full paper

Title:

Hydrolyzed diets may stimulate food-reactive lymphocytes in dogs

RUNNING HEAD: PROTEINS IN CANINE HYDROLYZED DIETS

Kenichi Masuda¹, Atsushi Sato¹, Atsushi Tanaka¹, Akiko Kumagai¹

¹Animal Allergy Clinical Laboratories Inc., SIC-2 #301, 5-4-30, Nishihashimoto, Midori-ku,
Sagamihara-city, Kanagawa 252-0131, Japan

Correspondence: Kenichi Masuda, Animal Allergy Clinical Laboratories Inc., SIC-2 #301, 5-4-30,
Nishihashimoto, Midori-ku, Sagamihara, Kanagawa 252-0131, Japan Tel.: +81-042-770-9437; fax:
+81-042-770-9438

E-mail address: masuda@aacl.co.jp

ABSTRACT

Hydrolyzed proteins are often prescribed for dogs with food hypersensitivity in food elimination programs. However, the potential of these diets to stimulate lymphocyte-mediated hypersensitivity is currently unknown. In this study, two commercially available hydrolyzed diets for dogs, D-1 (Aminopeptide Formula Dry, Royal Canin Japan, Tokyo, Japan), and D-2 (Canine z/d Ultra Dry, Hill's-Colgate (Japan) Ltd., Tokyo, Japan), were analyzed to identify residual proteins or peptides, as well as activated helper T-lymphocyte reactions in dogs with suspected food hypersensitivity. Proteins and peptides with molecular weights > 1 kDa (majority 1.5 - 3.5 kDa) were detected in both diet extracts with sodium dodecyl sulfate polyacrylamide gel electrophoresis, and size exclusion chromatography. When peripheral blood mononuclear cells (PBMC's) from 316 dogs with suspected food allergies were cultured with hydrolyzed diet extracts, flow cytometry analysis revealed detectable levels of CD25^{low} helper T-lymphocytes stimulated by D-1 and D-2 in 91 of 316, (28.8%), and 75 of 316 (23.7%) samples, respectively. These data indicated that the extracts contained proteins or peptides large enough to activate the lymphocytes. The percentages of CD25^{low} helper T-lymphocytes stimulated by D-1 and D-2 extracts increased to 38.7% and 29.6%, respectively, in 186 of the original 316 samples (186/316, 58.9%), also reactive to poultry-related antigens. Thus, both poultry-related antigens, and D-1 and D-2 diet extracts may activate helper T-lymphocytes. These results demonstrate that hydrolyzed diets may contain proteins that stimulate helper T-lymphocytes, and may not be effective for treating all dogs with food hypersensitivity.

KEYWORDS: diets, dogs, food allergy, hydrolyzation, lymphocytes

INTRODUCTION

Elimination diets often contain low molecular weight (MW) proteins and peptides resulting from enzymatic digestion (hydrolysis) of proteins from source materials [2, 10, 20]. Hydrolyzed protein diets are considered therapeutic for companion animals since they may prevent allergic reactions due to food hypersensitivity [2]. In humans, IgE may recognize protein allergens with MW ranging from 5 - 50 kDa, therefore, hydrolyzed proteins in elimination diets must have MW lower than 5 kDa to prevent allergic reactions [29]. A recent study in dogs demonstrated that a canine diet made from hydrolyzed poultry-feather meal containing 95% hydrolyzed proteins with $MW \leq 1$ kDa did not induce clinical reactions in any of the dogs allergic to chicken. However, another diet containing 78% hydrolyzed chicken liver proteins with $MW \leq 1$ kDa did induce allergic reactions in 40% of the dogs [2].

A clinical study of dogs diagnosed with food hypersensitivity after food restriction and provocation found type IV hypersensitivity lymphocyte-mediated reactions in 82% of the dogs, however, type I hypersensitivity was rarely detected [8]. Another report evaluating clinical cases of dogs with chronic dermatitis and suspected food hypersensitivity confirmed this trend [5]. In addition, a study conducted on dogs with gastrointestinal symptoms clinically diagnosed as inflammatory bowel disease, found peripheral lymphocytes reactive to at least one food antigen in all the animals [12]. Flow cytometry analysis revealed that the type IV lymphocytes associated with hypersensitivity were $CD4^+$ helper T-lymphocytes [5, 21]. Theoretically, these types of lymphocytes are reactive to proteins or peptides that do not induce IgE-mediated reactions [10, 14]. Therefore, it is clear that clinical use of hydrolyzed diets, as part of food elimination programs, may not be appropriate for all dogs with food hypersensitivity.

Small peptides (T cell epitopes) recognized by helper T-lymphocytes are displayed by MHC (Major Histocompatibility Complex) class II molecules found on antigen-presenting cells, such as macrophages, dendritic cells, and B-lymphocytes, which engulf antigens via phagocytosis. In previous studies, Wu et al., 2002, and Masuda et al. 2004, experimentally determined the amino acid (AA) sequences of common allergens by mapping with overlapping peptides containing *ca.* 15 AA's covering

all the protein allergen sequences [17, 30]. The estimated MW of the 15 AA length peptides ranged from 1–3 kDa [6, 17, 23], depending on AA composition. In a 2007 study, Kabuki and Joh showed that a peptide MW range of 1-3 kDa was critical for induction of a T-lymphocyte reaction in a human infant with casein-reactive T-lymphocytes. The study results demonstrated that helper T-lymphocytes were stimulated by mildly hydrolyzed casein with MW of ≥ 2 kDa, but not highly hydrolyzed casein with MW ≤ 1 kDa [10]. Moreover, in another study, a specific T-lymphocyte helper clone could recognize a peptide consisting of only 5 AA, with a MW < 1 kDa [7]. These results suggest that helper T-lymphocytes may recognize hydrolyzed proteins in a clinical setting, therefore, lower hydrolyzed-protein MW are desirable to prevent reactions.

Since T-cell epitopes are linear peptides, allergen AA sequences may be conserved in similar species, and therefore cause allergic reactions due to T-lymphocyte cross-reactivity. In a recent study, the major allergens from Japanese cedar and cypress Cry j 1 and Cha o 1, contained AA sequences with approximately 80% homology [26], and also contained identical T-cell epitope sequences recognized by human T-lymphocytes reactive to both Cry j 1 and Cha o 1 [23]. The Cry j 1 and Cha o 1 T-cell epitopes were also identical to various tree pollen and vegetable species epitopes [9, 31]. Alpha-parvalbumin, a major allergen from chicken meat, contains AA sequences that are highly conserved (greater than 80% homology) in poultry and livestock animals, and in a previous study resulted in induction of cross-reactive human IgE [15]. In 2018, Bexley et al. reported that dog serum IgE cross-reactivity between chicken and fish was related to nine proteins common to both species [1]. Alpha-actin, a highly conserved gene in vertebrates, was identified as one of the nine proteins [3]. Consequently, since T-lymphocytes may be widely cross-reactive to low MW allergens from taxonomically similar species, it is possible that T-lymphocytes could react to proteins or peptides even after hydrolysis.

In this study, the MW and antigenicity of protein extracts from two types of hydrolyzed diets, Amino peptide Formula Dry (Royal Canin Japon) and Canine z/d Ultra Dry (Hill's-Colgate (Japan) Ltd.), were examined to determine whether hydrolyzed proteins could stimulate helper T-lymphocytes in dogs

with suspected food hypersensitivity.

MATERIALS AND METHODS

Protein extraction from hydrolyzed diets

Two types of commercially available hydrolyzed diets (D-1, Aminopeptide Formula Dry, Royal Canin Japan, known as Anallergenic, Royal Canin, Aimargues, France and D-2, Hill's Prescription Diet, Canine z/d Ultra Dry, Hill's-Colgate (Japan) Ltd.) were purchased. When the study was initiated, three product lots of D-1, and two lots of D-2 were available. The protein extraction protocol was similar to the one previously reported [11]. Briefly, one gram of crushed kibbles from each diet was minced and suspended in 10 ml of physiological saline, and subsequently ground with glass beads under shaking conditions at a speed of 6000 rpm for 1 min, in an ULTRA-TURRAX® Tube Drive (IKA® Werke GmbH & Co. KG., Staufen, Germany). This procedure was repeated three times with one-minute intervals. The suspension was maintained at 4°C for 30 min to remove sediments after homogenization, and the supernatant was collected and centrifuged at $10,000 \times g$ for 10 min at 4°C. The supernatant was filtered and sterilized after centrifugation, with a 0.45 μ m pore size PES membrane, and protein concentrations were measured with ultraviolet (UV) light absorbance at a wavelength of 280 nm. Finally, the extracts were analyzed with sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and size-exclusion chromatography, and tested for lymphocyte stimulation.

SDS-PAGE

The hydrolyzed diet and chicken meat extracts (Greer Laboratories, Inc., Lenoir, NC, U.S.A.) were separated by SDS-PAGE in a 5–20% gradient polyacrylamide gel (ATTO Corporation, Tokyo, Japan) using a previously described method [11]. The total protein amounts in D-1, D-2, and chicken meat extract were adjusted to 10 μ g per lane. After electrophoresis, the gel was stained with Coomassie brilliant blue, (CBB) (FUJIFILM Wako Pure Chemical Corporation, Osaka, Japan), or silver stain reagent (Cosmo Bio Co., Ltd., Tokyo, Japan) for protein band visualization.

120

121 *Size-exclusion chromatography*

122 Size-exclusion chromatography of the extracts was performed using slight modifications of a
123 previously reported procedure [11]. A Superdex 200 10/300 GL column (GE Healthcare Bio-Sciences
124 AB, Uppsala, Sweden) was equilibrated with phosphate buffered saline (PBS). Each sample was eluted
125 isocratically at a flow rate of 0.5 ml/min in PBS, and protein elution was monitored with UV absorbance
126 at 280 nm. The column was calibrated with a Gel Filtration Low MW Kit (GE Healthcare Bio-Sciences
127 AB, Uppsala, Sweden), containing different MW protein standards: ferritin (MW 440 kDa), aldolase
128 (MW 158 kDa), conalbumin (MW 75 kDa), ovalbumin (MW 43 kDa), carbonic anhydrase (MW 29
129 kDa), ribonuclease A (MW 13.7 kDa), and aprotinin (MW 6.5 kDa).

130

131 *Lymphocyte stimulation*

132 As previously reported [5, 12, 25], lymphocyte proliferation of PBMCs isolated (by veterinarians
133 in private animal hospitals) from dogs with suspected food hypersensitivity, was investigated in a
134 routine assay with 18 types of food antigens in a commercial laboratory (Animal Allergy Clinical
135 Laboratories, Inc, Sagamihara, Japan). The food allergens for the standard lymphocyte assay were
136 extracted from beef, pork, chicken, egg white, egg yolk, milk, wheat, soybean, corn, lamb, turkey, duck,
137 salmon, cod fish, catfish, potato, and rice, and all were prepared and purchased from Greer Laboratories
138 Inc. (Lenoir, NC, U.S.A.) as previously described [5, 12, 25]. However, capelin was prepared by
139 homogenization of the dried fish body in PBS. Detailed clinical symptoms and dog history were
140 unknown, however, dermatological and/or gastrointestinal symptoms were suspected. Additionally, no
141 information on the status of allergen exposure and treatment history was available for the dogs providing
142 the blood samples. The blood samples were kept at 4 °C and shipped to the laboratory by refrigerated
143 courier services within 1-3 days after collection from local veterinarians. Briefly, PBMC's were purified
144 from whole blood with Ficoll-Hypaque (Lymphoprep, Axis-shield, Oslo, Norway) under centrifugation.
145 The cell viability was determined by exclusion of trypan blue (Sigma-Aldrich, Inc., St Louis, MO,

U.S.A.). When more than 95% of cells were viable, the PBMC's were cultured for seven days in RPMI-1640 medium (FUJIFILM Wako Pure Chemical Corporation), containing 10% fetal bovine serum, 100 μ g/ml streptomycin, and 100 U/ml penicillin (Sigma-Aldrich), at 1 μ g/ml antigen concentration for each allergen. Following the standard assay with 18 types of food allergens, the remaining PBMC's were also tested in a similar manner with extracts from D-1 and D-2. On Day 4 of the culture, recombinant human IL-2 (Pepro Tech Inc., Rocky Hill, NJ, U.S.A.) was added at a concentration of 100 U/ml. Subsequently, the cultured PBMC's were stained with antibodies, and analyzed via flow cytometry.

Flow cytometry

As previously described [5, 14], the antibodies used in the flow cytometry analysis included anti-dog CD4 rat antibody labeled with Alexa Fluor 647 (Clone YKIX302.9) (Bio-Rad Laboratories, Inc., Hercules, CA, U.S.A.), and anti-human CD25 antibody labeled with phycoerythrin (Clone ACT-1) (DAKO Cytomation, Glostrup, Denmark), as well as isotype controls such as Alexa Fluor 647-labeled purified rat IgG2a (Bio-Rad Laboratories), and PE-labeled mouse IgG1 (Bio-Rad Laboratories). Stained dead cells were excluded by propidium iodide incorporation. Finally, CD4⁺CD25⁺ cells were measured in a lymphocyte gate with FACS Canto II and FACS Diva software (Becton Dickinson, San Jose, CA, U.S.A.). We determined the number of activated helper T-lymphocytes against each antigen by calculating the percentage of CD4⁺ CD25^{low} lymphocytes in 2500 CD4⁺ lymphocytes [18, 19]. We defined activation as $\geq 0.4\%$ CD4-positive T-lymphocytes, since at least ten positive cells (0.4% of 2500 cells) confirmed the presence of a distinct cell population.

When considering clinical use of hydrolyzed diets in a food elimination program, induction of a lymphocyte response in dogs allergic to poultry-related antigens should be a major concern, since the diets examined in this study contained poultry-related proteins, such as feather meal or chicken meat. In this study, the positive lymphocyte ($\geq 0.4\%$) response to poultry-related antigens was compared to the response induced with hydrolyzed proteins. When a positive lymphocyte response to two or more

poultry-related antigens was detected in a sample, the maximum antigen value was designated as representative of a lymphocyte response. Positive lymphocyte responses to poultry-related antigens and hydrolyzed diets were statistically analyzed with Spearman's rank correlation coefficient. In addition, since positive lymphocyte responses might not always be detected, qualitative comparisons were also conducted. The Chi-squared test was utilized to analyze categorical data from samples with positive ($\geq 0.4\%$) and negative ($< 0.4\%$) lymphocyte responses, and from poultry-related antigens and hydrolyzed diets. Statistical analyses were conducted with XLSTAT software (Addinsoft, Paris, France).

A value of 1.2% has been considered a threshold of clinical manifestation in dogs with food hypersensitivity [5]. Therefore, we counted the number of samples with hydrolyzed diet response values of $\geq 1.2\%$, to calculate a lymphocyte response potentially associated with clinical manifestation.

RESULTS

Hydrolyzed diet protein molecular weights

SDS PAGE gels of the D-2 extract processed with CBB and silver stain revealed multiple broad bands with MW from 3.5 - 24 kDa, and faint bands at approximately 24 kDa and 52 kDa. No bands were detected in the D-1 extract processed with CBB, and silver staining revealed only one faint band with a MW slightly larger than 24 kDa (Fig. 1). Since the protein amounts in all lanes (D-1, D-2 and chicken meat) were adjusted to 10 μ g, the differences in visualized bands were due to protein component differences in the samples.

Size-exclusion chromatography revealed 6-8 protein waveform peaks with different molecular weights in both D-1 and D-2 (Fig. 2 and Fig. 3). Based on the standard protein reagents, the two major peaks had an estimated MW > 1 kDa, approximately 1.5 kDa and 3.5 kDa. A peak with a MW > 440 kDa (the largest standard protein) was detected in the D-2 extracts. All three D-1 product lots, and two D-2 product lots displayed similar wave patterns (Fig. 3). Size-exclusion chromatography data obtained in a different laboratory (Toray Research Center, Inc., Tokyo, Japan), revealed waveform peaks similar to our results (data not shown).

Lymphocyte responses to hydrolyzed proteins and poultry-related antigens

Of 316 PBMC total samples examined in the study, positive lymphocyte responses ($\geq 0.4\%$) against D-1 and D-2 extracts were detected in 91 (91/316, 28.8%), and 75 (75/316, 23.7%) samples, respectively (Table 1).

We applied Spearman's rank correlation coefficient (due to lack of normality) to determine if there was a correlation between positive lymphocyte responses in the D-1 or D-2 extracts, and the maximum response with poultry-related antigens. However, no correlations were observed, (D-1, $r_s=0.256$; D-2, $r_s=0.239$). Positive lymphocyte responses ($\geq 0.4\%$) were observed with at least one poultry-related antigen, such as chicken, duck, turkey, or chicken egg, in 186 of 316 total samples (186/316, 58.9%). The D-1 extracts elicited positive lymphocyte responses in 72 samples (72/186, 38.7%), and the D-2 extracts elicited responses in 55 (55/186, 29.6%) samples (Table 1). D-1 and D-2 extracts elicited positive responses in 130 samples that did not exhibit positive lymphocyte responses to any poultry-related antigens; 20 samples (20/130, 15.4%) responded to D-1, and 19 samples (19/130, 14.6%) to D-2. Application of the Chi-squared test demonstrated that there was no association between the maximum percentage of positive lymphocyte responses to poultry-related antigens, and the positive responses to the D-1 or D-2 extract.

Only seven (7/316, 2.2 %) D-1 samples and six D-2 samples (6/316, 1.9 %) elicited lymphocyte reactions with values $\geq 1.2\%$.

DISCUSSION

Hydrolyzed diets are considered effective for food elimination programs in companion animals with food hypersensitivity, since they contain proteins or peptides with molecular weights that are typically too low to stimulate allergic reactions. However, clinical cases of dogs with food hypersensitivity that did not respond positively to food elimination programs with hydrolyzed diets have been reported [2]. In addition, a previous report revealed that several carbohydrate proteins with molecular weights ranging from 21 - 67 kDa, and those recognized by serum IgE from individual cases

of dogs with suspected food hypersensitivity, were detected in the same commercial diets examined in this study [22]. In this study, SDS-PAGE gels of D-1 and D-2 extracts revealed bands with slightly higher MW than 24 kDa. In addition, a peak with an extremely high MW (>440 kDa) was detected only in D-2 extracts with size exclusion chromatography. These results suggest that type I hypersensitivity may occur in clinical cases where serum IgE is reactive to the proteins.

Previous reports showed that a type IV hypersensitivity allergic reaction mediated by T-lymphocytes recognizing small peptides not recognized by IgE, was observed in dogs with food hypersensitivity [5, 8]. In this study, the lower MW proteins and peptides in the hydrolyzed diet extracts were tested for induction of lymphocyte-mediated allergic reactions. Size-exclusion chromatography of D-1 and D-2 extracts revealed proteins and peptides approximately 1.5 kDa and 3.5 kDa MW (Fig 2). Previous studies suggested that proteins or peptides of this size were large enough to induce helper T-lymphocyte reactions, since synthetic peptides between *ca.* 1-1.5 kDa MW were used in experiments to detect helper T-lymphocyte reactions to T-cell epitopes [13, 18]. In addition, flow cytometry results in this study demonstrated a lymphocyte response to low MW proteins and peptides in the hydrolyzed diets. These results confirm that even low MW proteins and peptides may not be small enough to avoid a T-lymphocyte response in dogs with food hypersensitivity. Consequently, although hydrolyzed diets have been clinically effective in food elimination programs for many dogs with food hypersensitivity as previously reported [2], it is possible that T-lymphocyte allergic reactions might be prompted by residual proteins in hydrolyzed diets, and therefore these diets should not be considered appropriate for all dogs with food hypersensitivity.

Although potential induction of T-lymphocyte reactions against hydrolyzed protein diets has been suspected, induction of clinical symptoms related to food hypersensitivity in dogs was considered unlikely for three reasons. First, in this study, T-lymphocyte responses with values ≥ 1.2 % were detected in only seven of 316 (2.2%), and six of 316 (1.9%) cases stimulated by D-1 and D-2, respectively. Since 1.2% has been considered the threshold for clinical manifestation of canine food hypersensitivity [5], hydrolyzed diets should be safe in most cases. Second, in dogs fed hydrolyzed

diets, gastrointestinal digestion should decrease the molecular size of proteins or peptides, possibly preventing antigenicity against T-lymphocytes. For example, frequent gastrointestinal digestion of collagen *in vitro* results in small peptides with MW < 1 kDa [24], a molecular size consistent with common T-cell epitopes [17, 23]. Therefore, any T-cell epitopes remaining in hydrolyzed protein diets should be further reduced in size during gastrointestinal digestion. Third, since the protein sources in the hydrolyzed diets examined in this study were poultry-feather meal (D-1), and chicken meat (D-2) [2], they may have included T-cell epitopes taxonomically similar to many poultry proteins. In general, T-cell epitopes are conserved among major allergens with taxonomic similarity [28], and cross-reactivity against multiple allergens can occur [9, 23, 31]. T-cell epitopes should be conserved in poultry-related allergens; this may explain the “bird-egg syndrome” phenomenon of human allergic reactions to some avian meats [13] and feathers [27]. However, in this study, we found no significant correlation between positive lymphocyte responses against D-1 or D-2 extracts, and poultry-related allergens, suggesting that cross-reactivity of T-lymphocyte reactions may not be very common. Although further study on T-cell epitopes of poultry-related antigens in dogs is necessary, at present, hydrolyzed diets are considered to have a low risk for T-lymphocyte cross-reactivity. Consequently, a food elimination clinical program using hydrolyzed diets should be successful in most dogs with food hypersensitivity.

We detected one faint but clear protein band slightly larger than 24 kDa MW in silver stained SDS-PAGE gels of D-1 and D-2 extracts. This was inconsistent with results from a previous report where wide ranges of residual proteins from 21-67 kDa MW were found in the same hydrolyzed diets examined in this study [22]. Possibly, this may be due to differences in protein extraction procedures. The previous report described a homogenization procedure with a tissue homogenizer, *versus* the easy-to-use tabletop homogenizer utilized in the present study. In addition, protein extraction efficiency may be increased when the starting material is pulverized rather than homogenized [16], therefore an optimized method for protein extraction from hydrolyzed diets should be developed.

In this study, protein concentrations were measured with UV light absorption at 280 nm because

dye-staining methods, such as the Bradford method, were unsuccessful for the D-1 extracts (data not shown). This result was consistent with the SDS-PAGE results since CBB staining did not visualize any D-1 extract proteins in the lower MW ranges (3.5-24 kDa), whereas broad protein bands were visualized with both CBB and silver stain in the D-2 extracts. It is likely that proteins were not stained in the D-1 extracts due to structural damage of AA. CBB reacts with many types of AA, including aromatic and basic, as well as with the AA N-terminus [4], likely destroyed in the D-1 extract. In comparison, during silver staining, silver ions bind to AA sulfhydryl and hydroxyl groups, subsequently visualized with paraformaldehyde in a basic pH solution that reduces silver ions to metallic silver [17]. Notably, the D-1 extract proteins were also not visualized with silver stain, suggesting that the AA sulfhydryl and/or hydroxyl groups might have been damaged in the extraction procedure. In addition, regardless of the negative staining results, the D-1 extract protein concentrations were easily measured with UV light, which is absorbed by AA benzene rings. These results suggest that AA conformation in the D-1 extracts might have been partially damaged, but not destroyed. However, since proteins in the D-2 extracts were detected after both CBB and silver staining; the protein extraction procedure could not have affected protein conformation in the D-1 extracts. Therefore, proteins in the D-1 source material, *i.e.* hydrolyzed feather meal, should be analyzed to determine whether hydrolyzation could potentially cause changes in protein conformation.

Since the lymphocyte response is influenced by several factors in cell culture conditions such as reactive antigen or peptide concentrations, duration of antigen stimulation, cell concentrations, and ratios of reactive lymphocytes and antigen-presenting cells, T-lymphocyte reactions would not necessarily always be detected in this study. False-negative results might occur depending on cell culture conditions; although a lymphocyte response $\geq 1.2\%$ was designated as the food hypersensitivity detection threshold in dogs [5, 9]. Therefore, the significance of this study was the *in vitro* detection of lymphocyte responses in dogs stimulated by hydrolyzed diet extracts, previously reported in humans [10], but not in dogs. Lymphocyte responses would be controllable in constant culture conditions, and comparable to lymphocyte responses stimulated by different antigens in other studies. These results

could be obtained in dogs, (as in studies of human helper T-lymphocyte proliferation), by determination of the specific co-culture ratio between antigen-presenting cells, and CD4 positive lymphocytes, autologously isolated from blood samples [1, 3].

The PBMC's utilized in this study were sent by veterinarians for a previously described standard lymphocyte proliferation assay [5, 12, 25], and after the assay was completed, the remaining cells were used to examine T-lymphocyte responses to the hydrolyzed diets. Therefore, we did not have access to the history and clinical features of the dogs and blood samples. Ideally, before blood samples are submitted for lymphocyte proliferation assays, dogs with suspected allergies should be clinically diagnosed with food hypersensitivity based on their responses to food elimination programs and provocation [8]. Typically, only a limited number of dogs may be accepted for this type of analysis since the diagnostic procedure is complex, and may not always be successfully completed. In this study, we examined 316 samples, and although the sample inclusion criteria were not strictly defined, the large sample size permitted a comprehensive analysis of T-lymphocyte responses to hydrolyzed diets in dogs with potential food hypersensitivity. However, further studies should be conducted to confirm T-lymphocyte reactions to hydrolyzed diets in dogs with food hypersensitivity.

In summary, the current study demonstrated that hydrolyzed diets contain proteins and peptides with sufficiently high molecular weights to stimulate T-lymphocytes. Therefore, it is important to avoid hydrolyzed diets in food elimination programs when treating dogs previously diagnosed with lymphocytes reactive to poultry-related antigens.

CONFLICT OF INTEREST

K. Masuda is the Chief Executive Officer and a stockholder of Animal Allergy Clinical Laboratories, Inc.

ACKNOWLEDGMENTS

We want to thank veterinarians at various animal hospitals for sending blood samples for

lymphocyte proliferation assay analyses. We would also like to thank Dr. Makio Hayashiya (Hayashiya Animal Hospital, Kyoto, Japan) for their critical comments and suggestions on clinical aspects of the study.

REFERENCES

1. Bexley, J., Kingswell, N. and Olivry, T. 2018. Serum IgE cross-reactivity between fish and chicken meats in dogs. *Vet Dermatol* **30**: 25-e28.
2. Bizikova, P. and Olivry, T. 2016. A randomized, double-blinded crossover trial testing the benefit of two hydrolysed poultry-based commercial diets for dogs with spontaneous pruritic chicken allergy. *Vet Dermatol* **27**: 289-e270.
3. Chang, K.S., Rothblum, K.N. and Schwartz, R.J. 1985. The complete sequence of the chicken alpha-cardiac actin gene: a highly conserved vertebrate gene. *Nucleic Acids Res* **13**: 1223-1237.
4. Compton, S.J. and Jones, C.G. 1985. Mechanism of dye response and interference in the Bradford protein assay. *Anal Biochem* **151**: 369-374.
5. Fujimura, M., Masuda, K., Hayashiya, M. and Okayama, T. 2011. Flow cytometric analysis of lymphocyte proliferative responses to food allergens in dogs with food allergy. *J Vet Med Sci* **73**: 1309-1317.
6. Greer, F.R., Sicherer, S.H. and Burks, A.W. 2008. Effects of early nutritional interventions on the development of atopic disease in infants and children: the role of maternal dietary restriction, breastfeeding, timing of introduction of complementary foods, and hydrolyzed formulas. *Pediatrics* **121**: 183-191.
7. Hemmer, B., Kondo, T., Gran, B., Pinilla, C., Cortese, I., Pascal, J., Tzou, A., McFarland, H.F., Houghten, R. and Martin, R. 2000. Minimal peptide length requirements for CD4(+) T cell clones--implications for molecular mimicry and T cell survival. *Int Immunol* **12**: 375-383.
8. Ishida, R., Masuda, K., Kurata, K., Ohno, K. and Tsujimoto, H. 2004. Lymphocyte blastogenic responses to inciting food allergens in dogs with food hypersensitivity. *J Vet Intern Med* **18**:

- 354 25-30.
- 355 9. Jahn-Schmid, B., Radakovics, A., Luttkopf, D., Scheurer, S., Vieths, S., Ebner, C. and Bohle,
 356 B. 2005. Bet v 1142-156 is the dominant T-cell epitope of the major birch pollen allergen and
 357 important for cross-reactivity with Bet v 1-related food allergens. *J Allergy Clin Immunol* **116**:
 358 213-219.
- 359 10. Kabuki, T. and Joh, K. 2007. Extensively hydrolyzed formula (MA-mi) induced exacerbation
 360 of food protein-induced enterocolitis syndrome (FPIES) in a male infant. *Allergol Int* **56**: 473-
 361 476.
- 362 11. Kainuma, K., Ookura, T. and Kawamura, Y. 1995. Purification and characterization of protein
 363 disulfide isomerase from soybean. *J Biochem* **117**: 208-215.
- 364 12. Kawano, K., Oumi, K., Ashida, Y., Horiuchi, Y. and Mizuno, T. 2013. The prevalence of dogs
 365 with lymphocyte proliferative responses to food allergens in canine allergic dermatitis. *Pol J*
 366 *Vet Sci* **16**: 735-739.
- 367 13. Kelso, J.M., Cockrell, G.E., Helm, R.M. and Burks, A.W. 1999. Common allergens in avian
 368 meats. *J Allergy Clin Immunol* **104**: 202-204.
- 369 14. Knipping, K., Simons, P.J., Buelens-Sleumer, L.S., Cox, L., den Hartog, M., de Jong, N.,
 370 Teshima, R., Garssen, J., Boon, L. and Knippels, L.M. 2014. Development of beta-
 371 lactoglobulin-specific chimeric human IgE kappa monoclonal antibodies for *in vitro* safety
 372 assessment of whey hydrolysates. *PLoS One* **9**: e106025.
- 373 15. Kuehn, A., Lehnert, C., Hilger, C. and Hentges, F. 2009. Food allergy to chicken meat with IgE
 374 reactivity to muscle alpha-parvalbumin. *Allergy* **64**: 1557-1558.
- 375 16. Mahbod, M., Shahhoseini, S., Khabazkhoob, M., Beheshtnejad, A.H., Bakhshandeh, H., Atyabi,
 376 F. and Hashemi, H. 2015. Amniotic Membrane Extract Preparation: What is the Best Method?
 377 *J Ophthalmic Vis Res* **9**: 314-319.
- 378 17. Masuda, K., Sakaguchi, M., Saito, S., Yasueda, H., Iwabuchi, S., Tsukui, T., Hayashi, N., Nakao,
 379 Y., Kurata, K., Maeda, S., Ohno, K. and Tsujimoto, H. 2004. Identification of peptides

containing T-cell epitopes of Japanese cedar (*Cryptomeria japonica*) pollen allergen (Cry j 1) in dogs. *Vet Immunol Immunopathol* **102**: 45-52.

18. Masuda, K. and Yasuda, N. 2008. The antibody against human CD25, ACT-1, recognizes canine T-lymphocytes in the G2/M and G0/G1 phases of the cell cycle during proliferation. *J Vet Med Sci* **70**: 1285-1287.

19. Mizuno, T., Suzuki, R., Umeki, S. and Okuda, M. 2009. Crossreactivity of antibodies to canine CD25 and Foxp3 and identification of canine CD4+CD25 +Foxp3+ cells in canine peripheral blood. *J Vet Med Sci* **71**: 1561-1568.

20. Olivry, T., Bexley, J. and Mougeot, I. 2017. Extensive protein hydrolyzation is indispensable to prevent IgE-mediated poultry allergen recognition in dogs and cats. *BMC Vet Res* **13**: 251.

21. Olivry, T., Kurata, K., Paps, J.S. and Masuda, K. 2007. A blinded randomized controlled trial evaluating the usefulness of a novel diet (aminoprotect care) in dogs with spontaneous food allergy. *J Vet Med Sci* **69**: 1025-1031.

22. Roitel, O., Bonnard, L., Stella, A., Schiltz, O., Maurice, D., Douchin, G., Jacquenet, S., Favrot, C., Bihain, B.E. and Couturier, N. 2017. Detection of IgE-reactive proteins in hydrolysed dog foods. *Vet Dermatol* **28**: 589.

23. Sone, T., Dairiki, K., Morikubo, K., Shimizu, K., Tsunoo, H., Mori, T. and Kino, K. 2005. Identification of human T cell epitopes in Japanese cypress pollen allergen, Cha o 1, elucidates the intrinsic mechanism of cross-allergenicity between Cha o 1 and Cry j 1, the major allergen of Japanese cedar pollen, at the T cell level. *Clin Exp Allergy* **35**: 664-671.

24. Sun, L., Chang, W., Ma, Q. and Zhuang, Y. 2016. Purification of Antioxidant Peptides by High Resolution Mass Spectrometry from Simulated Gastrointestinal Digestion Hydrolysates of Alaska Pollock (*Theragra chalcogramma*) Skin Collagen. *Mar Drugs* **14**.

25. Suto, A., Suto, Y., Onohara, N., Tomizawa, Y., Yamamoto-Sugawara, Y., Okayama, T. and Masuda, K. 2015. Food allergens inducing a lymphocyte-mediated immunological reaction in canine atopic-like dermatitis. *J Vet Med Sci* **77**: 251-254.

- 406 26. Suzuki, M., Komiyama, N., Itoh, M., Itoh, H., Sone, T., Kino, K., Takagi, I. and Ohta, N. 1996.
 407 Purification, characterization and molecular cloning of Cha o 1, a major allergen of
 408 Chamaecyparis obtusa (*Japanese cypress*) pollen. *Mol Immunol* **33**: 451-460.
- 409 27. Szepefalusi, Z., Ebner, C., Pandjaitan, R., Orlicek, F., Scheiner, O., Boltz-Nitulescu, G., Kraft,
 410 D. and Ebner, H. 1994. Egg yolk alpha-livetin (chicken serum albumin) is a cross-reactive
 411 allergen in the bird-egg syndrome. *J Allergy Clin Immunol* **93**: 932-942.
- 412 28. Westernberg, L., Schulten, V., Greenbaum, J.A., Natali, S., Tripple, V., McKinney, D.M.,
 413 Frazier, A., Hofer, H., Wallner, M., Sallusto, F., Sette, A. and Peters, B. 2016. T-cell epitope
 414 conservation across allergen species is a major determinant of immunogenicity. *J Allergy Clin*
 415 *Immunol* **138**: 571-578 e577.
- 416 29. Woodfolk, J.A., Commins, S.P., Schuyler, A.J., Erwin, E.A. and Platts-Mills, T.A. 2015.
 417 Allergens, sources, particles, and molecules: Why do we make IgE responses? *Allergol Int* **64**:
 418 295-303.
- 419 30. Wu, B., Elst, L.V., Carlier, V., Jacquemin, M.G. and Saint-Remy, J.M. 2002. The
 420 Dermatophagoides pteronyssinus group 2 allergen contains a universally immunogenic T cell
 421 epitope. *J Immunol* **169**: 2430-2435.
- 422 31. Zulehner, N., Nagl, B., Briza, P., Roulias, A., Ballmer-Weber, B., Zlabinger, G.J., Ferreira, F.
 423 and Bohle, B. 2017. Characterization of the T-cell response to Dau c 1, the Bet v 1-homolog in
 424 carrot. *Allergy* **72**: 244-251.

425





Figure 1. SDS-PAGE of D-1 (Lane 1), D-2 (Lane 2), and chicken meat (Lane 3) extracts with Coomassie brilliant blue (CBB) and silver staining. The total protein amount applied to each lane was 10 μ g. Note that there were no visualized bands in Lane 1. SDS-PAGE, sodium dodecyl sulphate-polyacrylamide gel electrophoresis



432

433 Figure 2. Size-exclusion chromatography of D-1 extracts. The assay was repeated with three different

434 product lots. Similar wave patterns were detected in all product lots.

435




Figure 3. Size-exclusion chromatography of D-2 extracts. The assay was repeated with two different product lots. Similar wave patterns were detected in all product lots.

450

451 Table 1. In each sample, the presence or absence of helper T-lymphocytes reactive to hydrolyzed foods
 452 (D-1 and D-2) and poultry proteins was compared. Samples were stratified at 0.4% of reactive helper
 453 T-lymphocytes, since it was considered in this study that 0.4% was the minimum percentage to show
 454 reactive helper T-lymphocytes.

	Poultry $\geq 0.4\%$ ^{a)}	Poultry $< 0.4\%$ ^{b)}	Total
D-1 $\geq 0.4\%$	72	19	91
D-1 $< 0.4\%$	114	111	225
Total	186	130	316

	Poultry $\geq 0.4\%$ ^{a)}	Poultry $< 0.4\%$ ^{b)}	Total
D-2 $\geq 0.4\%$	55	20	75
D-2 $< 0.4\%$	131	110	241
Total	186	130	316

a) When helper T-lymphocyte response to any of chicken, turkey, duck, egg white, and egg yolk was 0.4% or more, the sample was counted as $\geq 0.4\%$.

b) When helper T-lymphocyte response to any of chicken, turkey, duck, egg white, and egg yolk was less than 0.4%, the sample was counted as $0.4\% <$.

455