

# Allergy Practice Worldwide: A Report by the World Allergy Organization Specialty and Training Council

John O. Warner<sup>a</sup> Michael A. Kaliner<sup>b</sup> Carlos D. Crisci<sup>c</sup> Sergio Del Giacco<sup>d</sup> Anthony J. Frew<sup>a</sup>  
G.H. Liu<sup>e</sup> Jorge Maspero<sup>f</sup> Hee-Bom Moon<sup>g</sup> Takemasa Nakagawa<sup>h</sup> Paul C. Potter<sup>i</sup>  
Lanny J. Rosenwasser<sup>j</sup> Anand B. Singh<sup>k</sup> Erkka Valovirta<sup>l</sup> Paul Van Cauwenberge<sup>m</sup>

<sup>a</sup> Southampton General Hospital Centre, Southampton, UK; <sup>b</sup> Institute for Asthma and Allergy, Bethesda, Md., USA; <sup>c</sup> Department of Allergy and Immunology, National University of Rosario, Rosario, Argentina;

<sup>d</sup> Dipartimento Di Scienze Mediche Internistiche, Policlinico Universitario, Cagliari, Italy; <sup>e</sup> China;

<sup>f</sup> Buenos Aires San Jose Cine a Foundation, Buenos Aires, Argentina; <sup>g</sup> Department of Allergy/Rheumatology, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea; <sup>h</sup> St. Marianna University, Toyoko Hospital, Department of Internal Medicine, Kawasaki, Japan; <sup>i</sup> Cape Town University, UTC Lung Institute, Cape Town, South Africa; <sup>j</sup> National Jewish Medical & Research Center, Denver, Colo., USA;

<sup>k</sup> Institute of Genomics and Integrative Biology, Delhi University Campus, New Delhi, India;

<sup>l</sup> Turku Allergy Center, Turku, Finland; <sup>m</sup> University Hospital Gent, ENT Department, Gent, Belgium

## Key Words

Allergy, specialty and training · Allergy disease prevalence · Allergy practice

diseases would be seen by a pediatrician with appropriate training. The survey results highlight a pressing need for the development of allergy services worldwide.

Copyright © 2006 S. Karger AG, Basel

## Abstract

In 2004 the World Allergy Organization's Specialty and Training Council conducted a survey of World Allergy Organization (WAO) member societies to obtain information about the status of the specialty of allergy worldwide. Responses were received from 33 countries, representing a population of 1.39 billion people, of whom it was estimated that 22% may suffer from some form of allergic disease. Allergy was reported by 23 respondents to be a certified or accredited specialty in their country, and the number of certified allergists per head of population ranged from 1:25 million to 1:16,000. Allergists were ranked as the fifth most likely clinicians to see cases of allergic asthma, third most likely to see allergic rhinitis, and fourth most likely to see eczema or sinusitis. Nine countries only reported that children with allergic

## Introduction

Allergic diseases constitute the most common causes of chronic illness in developed countries. Although rates may appear to be lower in developing countries, wherever prevalence studies have been conducted there is evidence of a steady increase over time in the proportion of populations suffering from allergies [1–7]. It has been proposed that there is a worldwide epidemic of allergic diseases which is likely to be a consequence of the changing environment and improved general health, superimposed on a range of genetic susceptibilities. Therefore, the treatment of allergy should have high priority in most countries, and the specialty of allergy should constitute an important branch of medicine.

In order to treat properly the large number of patients with allergic diseases, there needs to be a highly trained workforce specifically competent to handle their problems. However, around the world, allergy practice differs enormously. For some countries it does not even exist, either as a specialty or subspecialty. So-called 'allergy care pathways' also differ enormously, resulting in allergy patients being referred for care to organ-based specialists, clinical immunologists or generalists.

The World Allergy Organization (WAO) is a global federation representing 74 regional and national allergy and clinical immunology societies. WAO exists to build a global alliance to advance excellence in clinical care, research, education and training. WAO supports the rights of individuals with allergies worldwide to be seen by trained and certified allergists and for children to be seen by pediatricians with appropriate allergy training. To advance these objectives, the Allergy Specialty and Training Council was established to define the global outreach of, and need for, the specialty of allergy and clinical immunology. The Council's aim is to strengthen the specialty by providing guidance on training standards and certification requirements and to assist in the development of allergy training and accreditation programs.

The Council's first task was to define the current state of allergy training and services in the countries represented within the organization. A survey was sent to all WAO national society member organizations to obtain their subjective view of the state of the specialty of allergy within their countries. The questions were completed by allergists knowledgeable about their own countries, but in some countries the data were based on impressions rather than on published data. Some of the responses should be interpreted with caution, in that they represent the perspective of practicing allergists who may or may not have had access to accurate data. Nevertheless, the returns have provided a qualitative evaluation that will aid WAO in directing its energies to supporting the development of the specialty in collaboration with national organizations, to ensure that in the future patients suffering from allergic disorders receive a high standard of care.

## Methods

Each WAO member society was sent a questionnaire soliciting general information about the country's population and the pattern of allergy services within the country. It requested information about the form of training and certification for adult and pediatric practice. Some information was obtained on care pathways in relation to specific allergic disorders.

Questionnaires were sent to 61 national member organizations. Thirty-four responses were received, representing 33 countries (table 1). A number of respondents were unable to provide information on all questions. While the data solicited has not been cross-checked by reference to external published information on exact population figures and numbers of allergists in each country, it was decided to compile a preliminary report on the data received in order to give a broad indication of the current state of allergy services. Some of the highly populated regions of the world are not represented within this report, such as China, India and Indonesia. Additional information on numbers of trained allergists was obtained from a regional survey completed in 2005 by the Latin American Society of Allergy, Asthma and Immunology, and from the WAO Emerging Societies Meeting held in Russia in 2004.

## Results

The total population of the world represented by the returns is 1.39 billion, with a total world population in 2000 of 6.1 billion. The prevalence rates for allergy in the countries where physicians responded to the survey ranged from 7.5 to 40%, suggesting an average mean of 22% across the populations surveyed (fig. 1, 2). However, these two extreme figures were significant outliers. The overwhelming majority of countries reported between 15 and 35% incidence rates (22 of the 30 countries that gave relevant figures). The countries reporting the lowest rates were Colombia, Italy, Philippines, South Africa (although rates are rapidly increasing in South Africa, particularly with the migration of rural populations to urban areas [8]) and Turkey. The highest rates were from Ukraine, Japan, Bulgaria, Sweden and the United Kingdom. While some of the figures quoted accurately reflect output from large prevalence studies, such as ISAAC, others show discrepancies with published findings. This may represent a misunderstanding of whether the question about allergy patients related to the percentage of the population with allergic sensitization, as distinct from allergic disease. The figures quoted for individual atopic illnesses much more clearly reflected published figures, but were only elaborated by a minority of respondents.

The number of certified allergists in each country varied enormously from zero in Bangladesh to 5,000 in Germany. Based on those countries that had at least one certified allergist, as was the case in Malaysia, the number of allergists per head of population ranged from 1 in 25 million in Malaysia to 1 in 16,000 in Germany (table 2). Twenty-three out of the 30 countries giving information had some type of formal certification procedure, with a mixture of certification either in allergy alone, or combined as allergy and clinical immunology. Where allergy


**Table 1.** Responses received

| Country           | Respondent                        | WAO Member Society                                              |
|-------------------|-----------------------------------|-----------------------------------------------------------------|
| Argentina         | Edgardo Jares                     | Argentine Association of Allergy and Clinical Immunology        |
| Bangladesh        | Asif Mujtaba Mahmud               | Bangladesh Society of Allergy and Immunology                    |
| Belgium           | Wim Stevens                       | Belgian Society of Allergy and Clinical Immunology              |
| Bulgaria          | Vasil Dimitrov                    | Bulgarian National Society of Allergology                       |
| Colombia          | Luis Caraballo                    | Colombian Association of Allergy, Asthma and Immunology         |
| Czech Republic    | Vit Petru                         | Czech Society of Allergology and Clinical Immunology            |
| Denmark           | Ronald Dahl                       | Danish Society of Allergology                                   |
| Ecuador           | Manuel Viteri-Acaiturri           | Ecuadorian Society of Allergy and Immunology                    |
| Finland           | Iikka Annila                      | Finnish Society of Allergology and Immunology                   |
| France            | Daniel Vervloet                   | French Society of Allergology and Clinical Immunology           |
| Germany           | Gerhard Schulze-Werninghaus       | German Society of Allergology and Clinical Immunology           |
| Greece            | Dimitrios Papaioannou             | Hellenic Society of Allergy and Clinical Immunology             |
| Hungary           | Kristof Nekam                     | Hungarian Society of Allergology and Clinical Immunology        |
| Israel            | Meir Shalit                       | Israel Society of Allergy and Clinical Immunology               |
| Italy             | Sergio del Giacco                 | Italian Society of Allergology and Clinical Immunology          |
| Japan             | Yoichi Kohno                      | The Japanese Society of Allergology                             |
| Lebanon           | Fares Zaitoun                     | Lebanese Society of Allergy and Immunology                      |
| Malaysia          | Ranbir Kaulsay                    | Malaysian Society of Allergy and Immunology                     |
| Mongolia          | Munkhbayarlah Sonomjamts          | Mongolian Society of Allergology                                |
| Netherlands       | Anthony Dubois                    | Netherlands Society of Allergology                              |
| Philippines       | Madeleine Sumpaico                | Philippine Society of Allergy, Asthma and Immunology            |
| Portugal          | Celso Perreira/Angela Gaspar      | Sociedade Portuguesa de Alergologia e Imunologia Clinica        |
| Romania           | Radu Jeana-Rodica                 | Romanian Society of Allergology and Clinical Immunology         |
| Serbia/Montenegro | Vojislav Djuric                   | Association for Allergy and Immunology of Serbia and Montenegro |
| South Africa      | Cas Motala and Paul Potter        | Allergy Society of South Africa                                 |
| Sweden            | Mona Palmqvist                    | Swedish Association of Allergology                              |
| Switzerland       | Prof. U. Müller, Prof. A. Bircher | Swiss Society for Allergology and Immunology                    |
| Thailand          | Pakit Vichyanond                  | The Allergy and Immunology Society of Thailand                  |
| Turkey            | Sevim Baybek                      | Turkish National Society of Allergy and Clinical Immunology     |
| United Kingdom    | Chris Corrigan                    | British Society for Allergy and Clinical Immunology             |
| Ukraine           | Lyudmyla Yashyna                  | Ukrainian Association of Allergologists and Immunology          |
| USA (AAAAI)       | Lanny Rosenwasser                 | American Academy of Allergy, Asthma and Immunology              |
| USA (ACAAI)       | Michael Blaiss                    | American College of Allergy, Asthma and Immunology              |
| Venezuela         | Mario Sanchez-Borges              | Venezuelan Society of Allergy, Asthma and Immunology            |

AAAAI = American Academy of Allergy, Asthma and Immunology; ACAAI = American College of Allergy, Asthma and Immunology.

and clinical immunology were combined, the responsibilities of the certified clinicians included autoimmune disease and might also include infectious diseases, such as HIV, and primary immune deficiency. Additional information from the WAO Emerging Societies Meeting held with representatives from member countries of the Commonwealth of Independent States (CIS) indicates that allergy is a specialty, or a subspecialty of internal medicine and/or pediatrics in Belarus, Tajikistan, Moldova, Azerbaijan, Russia, Turkmenistan and Armenia, with formal certification reported by all countries except Tajikistan.


It is clear that allergy constitutes a component of post-graduate training in other specialties. In 16 countries, allergy is taught as part of respiratory medicine, in 15 as part of pediatrics, in 12 as part of dermatology, in 14 as part of ENT (otorhinolaryngology), and in 14 as part of internal medicine. In only 1 country was allergy part of the training for general practice, and in 1 country it was part of gastroenterology. These latter two figures are perhaps the most remarkable. Given the prevalence of allergy, the need to include allergy training in primary care training programs would appear to be overwhelming. Furthermore, food allergy constitutes a significant per-



1

**Fig. 1.** Estimated percentage of allergy patients per country surveyed. ACAAI = American College of Allergy, Asthma and Immunology.

**Fig. 2.** Allergy patients in the population surveyed (total population = 1,389,309,100).



2

centage of a gastroenterologist's clinical work. This information alone indicates the desperate need to improve training and expertise in order to ensure a good standard of care for patients with allergy. South Africa does not have a formal system for certification of allergists, but local allergists have taken a pragmatic approach to the need to create a level of competence amongst the clinicians who are the primary interface with allergy patients by creating an allergy diploma course. Points are accrued for

attending specialist clinics, conferences, and participation in a wide range of educational activities related to allergy, and an examination for the award of the diploma can be taken when 1,000 points have been attained.

With regard to specific clinical allergic problems, it is likely that the allergist may not be the primary referral in the overwhelming majority of countries. For allergic rhinitis in many countries, there is a rank order of the type of doctor most likely to be managing the patient, and the

**Table 2.** Allergists per head of population

| Country                  | Allergists per head of population |
|--------------------------|-----------------------------------|
| Argentina                | 1:46,353                          |
| Belgium                  | 1:900,000                         |
| Brazil <sup>a</sup>      | 1:100,000                         |
| Bulgaria                 | 1:106,250                         |
| Chile <sup>a</sup>       | 1:500,000                         |
| Colombia                 | 1:571,428                         |
| Czech Republic           | 1:17,543                          |
| Denmark                  | 1:135,000                         |
| Ecuador                  | 1:2,400,000                       |
| El Salvador <sup>a</sup> | 1:470,000                         |
| Finland                  | 1:94,545                          |
| France                   | 1:1,240,000                       |
| Germany                  | 1:16,000                          |
| Greece                   | 1:183,333                         |
| Honduras <sup>a</sup>    | 1:1,380,000                       |
| Hungary                  | 1:50,000                          |
| Israel                   | 1:52,000                          |
| Italy                    | 1:43,200                          |
| Japan                    | 1:61,200                          |
| Lebanon                  | 1:121,000                         |
| Malaysia                 | 1:25,000,000                      |
| Mexico <sup>a</sup>      | 1:175,000                         |
| Mongolia                 | 1:2,250,001                       |
| Paraguay <sup>a</sup>    | 1:200,000                         |
| Peru <sup>a</sup>        | 1:1,360,000                       |
| Philippines              | 1:66,115                          |
| Portugal                 | 1:63,334                          |
| Romania                  | 1:197,577                         |
| Serbia/Montenegro        | 1:500,000                         |
| South Africa             | 1:1,666,666                       |
| Spain <sup>a</sup>       | 1:44,000                          |
| Sweden                   | 1:42,857                          |
| Switzerland              | 1:36,649                          |
| Thailand                 | 1:1,000,000                       |
| Turkey                   | 1:1,076,923                       |
| United Kingdom           | 1:1,083,333                       |
| Ukraine                  | 1:94,441                          |
| Uruguay                  | 1:110,000                         |
| USA (ACAAI)              | 1:65,546                          |
| Venezuela                | 1:109,090                         |

ACAAI = American College of Allergy, Asthma and Immunology.

<sup>a</sup> Figures from LSAI Survey.

most likely is the primary care doctor, and if referral occurs, it may be to an otorhinolaryngology specialist before the allergist (fig. 3). For asthma, the most likely clinician is the primary care physician or internist followed by the pulmonologist, and then pediatricians, and only fifth on

**Table 3.** Immunotherapy practice

| Country           | Is immuno-therapy performed? | By specialists only | Specialists and GPs | Untrained practitioners |
|-------------------|------------------------------|---------------------|---------------------|-------------------------|
| Argentina         | ✓                            | ✓                   |                     |                         |
| Bangladesh        | Very rarely                  |                     |                     | ✓                       |
| Belgium           | ✓                            | ✓                   |                     |                         |
| Bulgaria          | ✓                            | ✓                   |                     |                         |
| Colombia          | ✓                            | ✓                   |                     |                         |
| Czech Republic    | Not recorded                 |                     |                     |                         |
| Denmark           | ✓                            |                     | ✓                   |                         |
| Ecuador           | ✓                            |                     | ✓                   |                         |
| Finland           | ✓                            |                     | ✓                   |                         |
| France            | ✓                            |                     | ✓ <sup>a</sup>      |                         |
| Germany           | ✓                            | ✓                   | ✓                   |                         |
| Greece            | ✓                            |                     |                     |                         |
| Hungary           | ✓                            | ✓                   |                     |                         |
| Israel            | ✓                            |                     | ✓                   |                         |
| Italy             | ✓                            |                     | ✓                   |                         |
| Japan             | ✓                            |                     | ✓                   |                         |
| Lebanon           | ✓                            |                     | ✓                   |                         |
| Malaysia          | ✓                            | ✓                   |                     |                         |
| Mongolia          | X                            |                     |                     |                         |
| Philippines       | Not recorded                 |                     |                     |                         |
| Portugal          | ✓                            | ✓                   |                     |                         |
| Romania           | ✓                            | ✓                   |                     |                         |
| Serbia/Montenegro | ✓                            | ✓                   |                     |                         |
| South Africa      | ✓                            |                     |                     |                         |
| Sweden            | ✓                            | ✓                   | ✓                   |                         |
| Switzerland       | ✓                            |                     |                     |                         |
| Thailand          | ✓                            | ✓                   | ✓                   |                         |
| Turkey            | ✓                            |                     |                     |                         |
| United Kingdom    | ✓                            | ✓                   |                     |                         |
| Ukraine           | ✓                            | ✓                   |                     |                         |
| USA (ACAAI)       | ✓                            | ✓                   |                     |                         |
| Venezuela         | ✓                            | ✓                   |                     |                         |

ACAAI = American College of Allergy, Asthma and Immunology.

<sup>a</sup> France: maintenance therapy by GPs.

the list, the allergist. For eczema, the dermatologist is the most likely to be managing the disease, followed by general practitioner, pediatrician and then allergist. For urticaria and angioedema, again, the primary care physician or internist is primarily involved, followed by the dermatologist, and fourthly the allergist. For sinusitis it is primary care/internal medicine, followed by ENT, and then pediatrics, and finally the allergist. It is remarkable that these figures seem to be irrespective of the number of certified allergists in the country. Even in Germany with a very large number of allergists representing a large percentage of the total physicians in the country, it is the




**Fig. 3.** Who sees rhinitis patients?

internist, system specialist and pediatricians who manage asthma rather than allergists.

Immunotherapy is performed in all but 1 country that responded (table 3) and is predominantly handled either by a specialist alone or by the specialist in collaboration with a primary care physician.

While the majority of countries have pediatricians who specialize in allergy, only 8 reported certified training programs. Remarkably, from a pediatrician's perspective, in the majority of countries, children with allergic problems could be seen by either a pediatrician or an adult specialist, whether or not the specialist had allergy or organ-based specialty training. Only 9 countries reported that children with allergic problems will be seen by someone with appropriate pediatric training (table 4).

## Discussion

There is overwhelming evidence that allergic diseases constitute the most common cause of chronic problems in developed countries. These observations are consistent with other global surveys [1–7]. Despite these remarkable figures, health services worldwide have failed to respond to the need to help allergic patients with a change in provision of specialists suitable to handle this ever-increasing problem. Even in Germany, which has a high rate of allergy training, studies of health care utilization suggest that the system is under substantial strain, with half of the children with asthma symptoms making unplanned visits to health care facilities [9]. Similar figures come from the USA [10]. Indeed, studies for the USA have shown that emergency department visits and hospital admissions account for around three quarters of the direct costs of asthma [11]. One might have expected that such

**Table 4.** Pediatric allergy practice

| Country           | Number of pediatricians specializing in allergy <sup>a</sup> | Is pediatric allergy a certified specialty | Pediatric referrals to allergists <sup>b</sup> | Pediatric referrals to organ-based physicians <sup>b</sup> |
|-------------------|--------------------------------------------------------------|--------------------------------------------|------------------------------------------------|------------------------------------------------------------|
| Argentina         | <i>160</i>                                                   | No                                         | Both                                           | Pediatric                                                  |
| Bangladesh        |                                                              | No                                         | Both                                           | Both                                                       |
| Belgium           | 5                                                            | No                                         | Both                                           | Both                                                       |
| Bulgaria          | 25                                                           | No                                         | Both                                           | Pediatric                                                  |
| Colombia          | 7                                                            | No                                         | Both                                           | Both                                                       |
| Czech Republic    | 180                                                          | Not recorded                               | Not recorded                                   | Not recorded                                               |
| Denmark           | 30                                                           |                                            | Pediatric                                      | Pediatric                                                  |
| Ecuador           | 5 (accredited overseas)                                      | No                                         | Both                                           | Pediatric                                                  |
| Finland           | 35                                                           | Yes, until 2005                            | Pediatric                                      | Pediatric                                                  |
| France            | 20                                                           | No                                         | Both                                           | Both                                                       |
| Germany           | <i>11,000<sup>c</sup></i>                                    | No                                         | Both                                           | Both                                                       |
| Greece            |                                                              | No                                         | Adult                                          | Both                                                       |
| Hungary           | 60                                                           | No                                         | Pediatric                                      | Both                                                       |
| Israel            | Not known                                                    | Yes                                        | Pediatric                                      | Pediatric                                                  |
| Italy             | Not known                                                    | Not recorded                               | Pediatric                                      | Both                                                       |
| Japan             | 253                                                          | No                                         | Both                                           | Neither                                                    |
| Lebanon           | 3–5                                                          | Yes                                        | Both                                           | Both                                                       |
| Malaysia          | 0                                                            | No                                         | Neither                                        | Adult                                                      |
| Mongolia          | 3                                                            | No                                         | Both                                           | Pediatric                                                  |
| Philippines       | 49                                                           | Not recorded                               | Not recorded                                   | Not recorded                                               |
| Portugal          |                                                              | No                                         | Both                                           | Both                                                       |
| Romania           | 110 <sup>d</sup>                                             |                                            | Not recorded                                   | Not recorded                                               |
| Serbia/Montenegro |                                                              | No                                         | Pediatric                                      | Pediatric                                                  |
| South Africa      | 20                                                           | No                                         | Pediatric                                      | Pediatric                                                  |
| Sweden            | 10–20                                                        | Yes                                        | Pediatric                                      | Pediatric                                                  |
| Switzerland       | 20                                                           | No                                         | Both                                           | Both                                                       |
| Thailand          | 20                                                           | Yes                                        | Neither                                        | Pediatric                                                  |
| Turkey            | 3–4                                                          | Yes                                        | Pediatric                                      | Pediatric                                                  |
| United Kingdom    | 5                                                            | No                                         | Neither                                        | Pediatric                                                  |
| Ukraine           | 364                                                          | Yes                                        | Both                                           | Both                                                       |
| USA (ACAAI)       | 2,590                                                        | No                                         | Both                                           | Pediatric                                                  |
| Venezuela         | 50–60                                                        | No                                         | Both                                           | Neither                                                    |

ACAAI = American College of Allergy, Asthma and Immunology.

<sup>a</sup> Italicized numbers represent estimated numbers, numbers in regular print represent actual ones.

<sup>b</sup> Both refers to both pediatrician and adult specialist, neither refers to neither pediatrician nor adult specialist.

<sup>c</sup> In Germany, all pediatricians will see some allergy, hence the high figure of 11,000.

<sup>d</sup> In Romania, the specialty of allergy and clinical immunology training program covers adult and pediatric allergy.

figures would have prompted health care planners to evolve a system to improve this state of affairs; that it has not occurred has prompted a number of responses from the profession. In the UK, the Royal College of Physicians published a report in 2003, entitled Allergy: the Unmet Need [12]. This report subsequently led to a UK parliamentary response with the publication of a document entitled The Provision of Allergy Services [13]. Sadly, hitherto this has not led to any changes in health care policies.

It is a remarkable paradox that in the UK, a country which has an outstanding record in allergy research, there is a remarkably poor clinical service for allergy sufferers [14]. These depressing facts appear to be reflected in all the countries surveyed and highlight the very pressing need for WAO to support national societies in furthering the interests of allergy sufferers.

In the absence of appropriately trained specialists, patients are predominantly seen by relatively poorly trained

or allergy-naïve system specialists, generalists or primary care physicians. It is not surprising, therefore, that morbidity statistics are unacceptably high and health costs considerable. It is likely that a significant number of individuals with allergic problems are not seen by any physician, but either tolerate their symptoms or purchase over-the-counter remedies. Where the patient is seen by a nonallergist, they are presumably being primarily given pharmacotherapy with little probability of having appropriate triggers of their disease identified. In some countries, patients and their families resort to a range of complementary diagnostic and therapeutic procedures of dubious provenance.

Many allergy sufferers have more than one manifestation of their disease. Thus, the overwhelming majority of patients with asthma also have allergic rhinitis and up to 40% have had or continue to have eczema. Similarly, sinusitis occurs at an alarmingly high rate in asthmatics, especially in the more severely affected asthmatic. This constellation of allergic manifestations often means that patients are required to see 2 or 3 different system specialists in order to handle the range of their allergic problems. This inefficiency increases health care costs considerably [15]. The one-stop service provided by an allergist should certainly reduce these costs, although there are no data available to confirm this suspicion. Furthermore, there are significant dangers implicit in a lack of coordination of treatment between the different system specialties. One possible consequence of multiple physicians caring for individual aspects of allergic diseases is overprescription of topical steroids to the skin, nose and lungs with consequent potential for major unwanted effects. Overreliance on pharmacotherapy without attempting to identify potentially avoidable triggers of the disease is likely to increase health care costs, morbidity and potential for side effects.

The situation for pediatrics appears even more depressing. The number of countries with specific specialization in pediatric allergy and appropriate certification of training is small, and children with allergic diseases may be seen by adult physicians, primary care physicians or general pediatricians. All countries would assert the importance of children with medical problems being seen by an appropriately trained pediatrician. The special needs of children can only be addressed by someone with complete pediatric training. However, in addition to this, there must be appropriate specialty training within pediatrics so that the specific problem is handled effectively. This is well iterated in a recently published document from the UK on a National Service Framework for Chil-

dren, Young People and Maternity Services, 'All children and young people who are ill or thought to be ill or injured, should have timely access to appropriate advice and to effective services which address their health, social, educational and emotional needs throughout the period of their illness' [16]. Implicit in this is that children need to be cared for by people with appropriate training and understanding of their specific needs.

It is important to appreciate the shortcomings of this survey. It did not include full information from some of the most populous areas of the world including Brazil, India, China, Russia and Indonesia, and responses were not received from some WAO Member Societies in westernized countries where allergy rates are known to be high. It has not been possible to substantiate all the figures quoted by individuals returning the questionnaire and in some cases, it is clear that the estimates do not match up with published data. The survey provides the subjective impressions of allergy specialists about the situation in their countries, and the data must be viewed as qualitative rather than accurately quantitative. There is no doubting, however, the fact that the prevalence of allergic disease is not matched by an appropriate development of clinical services to support sufferers. Even in countries with a well-developed system of training, there are still significant problems of morbidity and health economic costs accruing as a consequence of inadequacies in the care of patients [17]. WAO is in a position to provide strong support to all national organizations attempting to improve this state of affairs. There is the opportunity to define the levels of competence required to handle allergic problems and then to define the training requirements at the various levels of care. Given the prevalence of allergic problems, much attention must focus on training undergraduate medical students, primary care physicians and generalists, as well as ensuring that system specialists, who are likely to be dealing with allergic patients, have a higher level of training. Finally there must be a sufficient number of specialist (tertiary) care centers in each country that are able to set the standards, prosecute research, organize training at a local level and support secondary and primary care by establishing appropriate networks.

## References

- ▶ 1 Worldwide variation in prevalence of symptoms of asthma, allergic rhinoconjunctivitis and atopic eczema: ISAAC. The International Study of Asthma and Allergies in Childhood (ISAAC) Steering Committee. *Lancet* 1998; 351:1225–1232.
- ▶ 2 European Community Respiratory Health Survey: Variations in the prevalence of respiratory symptoms, self reported asthma attacks and use of asthma medication in the European community respiratory health survey ECRHS. *Eur Respir J* 1996;9:687–695.
- ▶ 3 Grundy J, Matthews S, Bateman B, et al: Rising prevalence of allergy to peanut in children: data from two sequential cohorts. *J Allergy Clin Immunol* 2002;110:784–789.
- ▶ 4 Sheikh A, Alves B: Hospital admissions for anaphylaxis: time trend study. *Br Med J* 2000; 320:1441.
- ▶ 5 Garabrant DH, Schweitzer S: Epidemiology of latex sensitisation and allergies in health care workers. *J Allergy Clin Immunol* 2002;110: 582–595.
- ▶ 6 Demoly P, Bousquet J: Epidemiology of drug allergy. *Curr Opin Allergy Clin Immunol* 2001; 1:305–310.
- ▶ 7 Bousquet J: Allergy as a global problem: think globally act globally. *Allergy* 2000;57:661–662.
- ▶ 8 Steinman H, Donson H, Kawalski M, Toerien A, Potter PC: Bronchial hyper-responsiveness in urban, periurban and rural South African children. *Pediatr Allergy Immunol* 2003;14: 383–393.
- ▶ 9 Maziak W, von Mutius E, Keil U, et al: Predictors of health care utilisation of children with asthma in the community. *Pediatr Allergy Immunol* 2004;14:166–171.
- 10 Glaxo Wellcome: Asthma in America National Survey. Washington, Glaxo Wellcome, 1998.
- ▶ 11 Weiss KB, Gergen PJ, Hogson TA: An economic evaluation of asthma in the United States. *N Engl J Med* 1992;326:862–866.
- 12 Allergy: The Unmet Need. A Blueprint for Better Patient Care. A report of the Royal College of Physicians Working Party on the Provision of Allergy Services in the UK. London, Royal College of Physicians, 2003.
- 13 House of Commons Health Committee: The Provision of Allergy Services 6th Report of Session 2003/2004. House of Commons. London, The Stationery Office, HC696-1.
- ▶ 14 Warner JO: The allergy paradox. *Pediatr Allergy Immunol* 2004;15:287–288.
- ▶ 15 Grupp-Phelan J, Lozars P, Fishman P: Health care utilisation and cost in children with asthma and selected comorbidities. *J Asthma* 2001; 38:363–373.
- 16 NSF for Children, Young People and Maternity Services. London, Department of Health Publications, 2004.
- ▶ 17 van den Akker-van Marle ME, Bruil J, Detmar SB: Evaluation of cost disease: assessing the burden to society of asthma in children in the European Union. *Allergy* 2005;60:140–149.