

REVIEW

Herbal Medicines Prevent the Development of Atopic Dermatitis by Multiple Mechanisms*

MAN Mao-qiang (蔺茂强)^{1,2}, HU Li-zhi (胡立志)³, and Peter M. Elias^{1,2}

ABSTRACT Atopic dermatitis (AD) is among the most common skin disorders in humans. Although a variety of regimens are available for the treatment of AD, preventive approaches are limited. Recent studies have demonstrated that certain naturally-occurring herbal medicines are effective in preventing the development of AD via divergent mechanisms, such as inhibiting cytokine and chemokine expression, IgE production, inflammatory cell infiltration, histamine release, and/or enhancement of epidermal permeability barrier function. Yet, they exhibit few adverse effects. Since herbal medicines are widely available, inexpensive and generally safe, they could represent an ideal approach for preventing the development of AD, in both highly developed and developing countries.

KEYWORDS herbal medicines, T cells, cytokines, permeability barrier, dermatitis

Various types of eczematous dermatitis are among the most common skin disorders in human. These disorders include atopic dermatitis (AD), nummular eczema, seborrheic dermatitis and contact dermatitis.^(1,2) A variety of anti-inflammatory approaches are available to treat these disorders. For example, topical glucocorticoids are commonly used to treat allergic contact dermatitis, irritant contact dermatitis and AD. Moreover, oral cyclosporin A reportedly increases regulatory T cells in AD, and is effective in treating both AD^(3,4) and allergic contact dermatitis.⁽⁵⁾ Furthermore, immunomodulators, such as tacrolimus and pimecrolimus, also are effective in treating AD in both adults and children.⁽⁶⁻⁹⁾ Finally, tumor necrosis factor α (TNF- α) inhibitors, such as infliximab, are effective in treating AD.^(10,11) The effects of these anti-inflammatory approaches in treating eczematous dermatoses are not always sufficient. Moreover, each of these anti-inflammatory approaches carries a substantial risk of side effects. These adverse effects provide a rationale for alternative approaches to both treat and/or prevent these disorders. Though both systemic and topical administrations of herbal medicines alleviate AD, allergic and irritant contact dermatitis,⁽¹²⁻¹⁶⁾ herbal regimens that prevent the development of dermatitis, including AD, are not greatly appreciated by the medical community. Certain topical medicinal herbal preparations reportedly prevent the development of contact dermatitis in human,⁽¹⁷⁻²⁰⁾ and the preventive effects of herbal medicines on the development of contact dermatitis and AD-like dermatitis has also

been convincingly demonstrated in murine disease models.^(21,22) Most studies on the preventive effects of herbal medicines on the development of AD-like lesions have been carried out in murine models, including repeated topical applications of a hapten or allergen to pre-sensitized murine skin,^(1,23) or in NC/Nga and filaggrin deficient mice, which develop AD-like skin lesions with AD-like, Th2-dominant immunophenotype under appropriate condition.⁽²⁴⁾ Numerous studies have demonstrated that either systemic or topical administration of herbal medicines prevents the development of AD as evaluated by clinical symptoms, stratum corneum biophysical properties, cytokine expression and inflammatory cell infiltration in these murine AD models.

In a review of the patients' quality of lives, as well as cumulative medical costs, prevention is more important than treatment of these eczematous diseases. This paper updates current knowledge

©The Chinese Journal of Integrated Traditional and Western Medicine Press and Springer-Verlag Berlin Heidelberg 2015

*Supported by the National Institute of Health, USA (No. AR061106), Veterans Affairs Medical Center, USA (No. VA Merit-1 101BX000608), National Eczema Association, and National Science Foundation of China (No. 81301360)

1. Dermatology Service, Veterans Affairs Medical Center, CA (94121), USA; 2. Department of Dermatology, University of California San Francisco, CA (94121), USA; 3. School of Basic Medical Sciences, Tianjin Medical University, Tianjin (300070), China

Correspondence to: Dr. MAN Mao-qiang, E-mail: mqman@hotmail.com, or Dr. HU Li-zhi, E-mail: lizhihuleopard@yahoo.com
DOI: 10.1007/s11655-015-2438-1

about the utility of herbal medicines in preventing the development of AD, and the mechanisms that appear to account for their benefits. These studies demonstrate the potential utility of herbal medicines for the prevention of AD.

Clinical Symptoms

The benefits of herbal medicines in preventing the development of clinical symptoms have been assessed primarily in AD mouse models. For example, mice topically treated repeatedly with house dust mites, or haptens, such as 2,4-dinitrochlorobenzene (DNCB) or oxazolone develop AD symptoms. In addition, NC/Nga mice develop AD like lesions in a conventional (non-specific pathogen free) environment. It has been reported that mice pre-fed with azuki bean extract ($250 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}$) for 2 weeks display a greater than 70% reduction in dermatitis score in a repeated DNBCB-induced model of AD.⁽²⁵⁾ Likewise, mice pre-fed with apigenin for 4 weeks also demonstrated a dramatic reduction in dermatitis scores during the development of AD induced by repeated topical 2,4,6-trinitrochlorobenzene (TNCB) challenges.⁽²⁶⁾ Moreover, either oral *Rumex japonicus* houtt, or *Actinidia arguta* or persimmon leaf extract dramatically reduced the amount of scratching behavior in different AD models.⁽²⁷⁻²⁹⁾ Orally given *Alpinia katsumadai* Hayata extract, along with topical allergen challenges for 4 weeks, also significantly lowered dermatitis scores in NC/Nga mice with spontaneous AD.⁽²²⁾ However, when 6-week old NC/Nga mice were fed with *Lithospermum erythrorhizon* extract for 10 weeks, the appearance of their skin, as well as the extent of their scratching behavior were nearly normalized.⁽³⁰⁾

In most comparative studies, the effect of oral herbal extracts in the prevention of AD development was compared favorably to potent topical or systemic glucocorticoids.⁽³¹⁻³⁶⁾ For example, oral herbal extracts display similar or superior effect to oral prednisolone in preventing the increase in ear inflammation⁽³³⁾ and dermatitis score during the development of AD.^(28,34) In another comparative study, oral Yokukansan (Yi-Gan-San in Chinese) was more effective than the H1 blocker, fexofenadine hydrochloride, in reducing the amount of scratching, as well as dermatitis scores in NC/Nga mice.⁽³⁷⁾ Similarly, oral ginseng extract displayed similar effect to cyclosporin A for dermatitis symptoms in an AD mouse model.⁽³⁸⁾

Not only do oral, but also topical herbal medicines prevent the increase in dermatitis scores of AD. For instance, topical *Angelicae Dahuricae Radix* extract effectively decreased dermatitis scores and ear thickness, with an comparable effect as topical tacrolimus in a dust mite-induced AD model.⁽³⁹⁾ Additionally, intraperitoneally administrations of herbal extracts prevented the expected increase in AD severity scores and ear thickness induced by topical DNFB.^(40,41) All these studies indicate that either systemic or topical herbal medicines prevent the development of clinical symptoms of AD. Herbal medicines shown to be effective in preventing the development of AD-like symptoms are shown in Table 1.

Stratum Corneum Function

Stratum corneum (SC) biophysical properties reflect different epidermal functions. Studies indicate that elevated transepidermal water loss (TEWL) levels, an indicator of epidermal permeability barrier function, correlate positively with the severity of AD phenotype,^(64,65) while SC hydration correlates negatively with the severity of AD.⁽⁶⁶⁾ Therefore, changes in both barrier function and SC hydration have been used to evaluate the therapeutic and preventive effect of herbal medicines on AD. A number of studies have demonstrated that orally administrations of herbal extracts attenuate the expected deterioration in barrier function in murine AD models. Funakushi, et al⁽³⁷⁾ reported that oral administrations of herbal extract, Yokukansan, for 3 weeks significantly inhibited the increase in TEWL in NC/Nga mice. Likewise, oral administration of an extract from *actinidia arguta* at a dosage of either 50 or 100 mg/kg for 16 days prevented the expected deterioration in barrier function in a rat AD model induced by magnesium deficiency.⁽²⁷⁾ Similarly, an oral extract from persimmon leaves also normalized barrier function in NC/Nga mice.⁽²⁹⁾ Moreover, orally administrated *Lithospermum erythrorhizon* extract, started one day prior to oxazolone challenge not only reduced basal TEWL rates, but also accelerated barrier recovery rates in an AD model.⁽⁶⁷⁾ Furthermore, topically co-applications of bamboo extract with DNBCB lowered TEWL rates in a repeated DNBCB-treated mouse AD model.⁽⁶⁸⁾ As stated above, decrease in SC hydration is another feature of AD.^(1,69,70) Oral Yokukansan prevented the usual reduction of SC hydration in NC/Nga mice, in parallel with an inhibition in the development of AD.⁽⁴³⁾ Notably, the effect

Table 1. List of Herbal Medicines that Display Preventive Effects on the Development of AD in Murine AD Models

No.	Herbal extract	Route of administration	AD model	Dermatitis score	Ear thickness	Number of scratching	Reference
1	<i>Alpinia katsumadai</i> Hayata (草豆蔻)	Orally	House dust mite-treated NC/Nga mice	↓	↓	N/D	22
2	<i>Vigna angularis</i> (红豆)	Orally	DNCB-treated NC/Nga mice	↓	↓	N/D	25
3	Apigenin from <i>Flos Chrysanthemi</i> (菊花)	Orally	PCI-treated NC/Nga mice	↓	N/D	N/D	26
4	<i>Lithospermum erythrorhizon</i> (紫草)	Orally	NC/Nga Mice	N/D	N/D	↓	30
5	<i>Radix Glycyrrhizae</i> (甘草), <i>Zingiber officinale</i> (生姜), <i>Citrus sunki</i> (酸橘), <i>Magnolia officinale</i> (厚朴), <i>Atractylodes japonica</i> (苍术), <i>Zizyphus jujuba</i> (大枣)	Orally	DNFB-treated NC/Nga mice	↓	N/D	↓	31
6	<i>Atractylodes Japonica</i> (苍术)	Orally	TNCB-treated NC/Nga mice	↓	N/D	↓	32
7	<i>Astragalus membranaceus</i> (黄芪)	Orally	DNFB-treated NC/Nga mice	↓	↓	N/D	33
8	Fruits of <i>Cudrania tricuspidata</i> (柘树)	Orally	House dust mite-treated NC/Nga mice	↓	N/D	N/D	34
9	<i>Actinidia arguta</i> (猕猴桃)	Orally	TNCB-treated NC/Nga mice; NC/Nga mice; magnesium deficient rat	↓	N/D	↓	35, 36, 27, 42
10	<i>Rumex japonicus</i> houtt (土大黄)	Orally	PCI-treated NC/Nga mice	↓	↓	↓	28
11	<i>Atractylodes lancea</i> rhizome (苍术), <i>Poria</i> (茯苓), <i>Cnidium officinale</i> (川芎), <i>Radix Angelicae sinensis</i> (当归), <i>Radix Bupleuri</i> (柴胡), <i>Radix Glycyrrhizae</i> (甘草), and <i>Ramulus uncaria</i> cum <i>uncis</i> (钩藤)	Orally	NC/Nga mice	↓	N/D	↓	37,43
12	Persimmon leaf (柿叶)	Orally	NC/Nga mice	↓	↓	↓	29
13	<i>Panax red ginseng</i> (红参)	Orally	TNCB-treated NC/Nga mice	↓	↓	N/D	38
14	<i>Ramulus uncaria</i> cum <i>uncis</i> (钩藤)	Orally	DNFB-treated NC/Nga mice	↓	↓	N/D	44
15	<i>Saururus chinensis</i> (LOUR.) BAILL (三白草)	Orally	PCI-treated NC/Nga mice	↓	↓	↓	45
16	<i>Betula platyphylla</i> var. <i>japonica</i> (白桦)	Orally	PCI-treated NC/Nga mice	↓	↓	↓	46
17	Ginsenoside Rh1 from <i>Radix ginseng</i> (人参)	Orally	OX-treated hairless mice	↓	↓	N/D	47
18	<i>Radix Angelicae sinensis</i> (当归), <i>Moutan Cortex</i> (牡丹皮), <i>Rehmannia glutinosa</i> <i>Libosch</i> (地黄), <i>Glycyrrhizae Radix</i> (甘草), <i>Flos Lonicerae</i> (金银花), <i>Radix Ledebouriellae Divaricatae</i> (防风), <i>Spirodelae Herba</i> (浮萍), <i>Forsythiae Fructus</i> (连翘), <i>Radix Paeoniae rubra</i> (赤芍), <i>Rhizoma Ligusticum wallichii</i> (川芎), <i>Schizonepetae Herba</i> (荆芥), <i>Bombyx Batryticatus</i> (僵蚕), <i>Menthae Herba</i> (薄荷), <i>Bubali Cornu</i> (水牛角) and <i>Radix Scutellariae</i> (黄芩)	Orally	DNCB-treated NC/Nga mice	↓	↓	N/D	48
19	<i>Radix Platycodi</i> (桔梗)	Orally	DNCB-treated NC/Nga mice	↓	↓	N/D	49
20	<i>Chelidonium majus</i> (白屈菜)	Orally	DNCB-treated NC/Nga mice	↓	↓	↓	50
21	<i>Euglena gracilis</i> Z (小眼虫)	Orally	TNCB-treated NC/Nga mice	↓	↓	N/D	51
22	Naringenin	Intraperitoneally	DNFB-treated NC/Nga mice	↓	↓	N/D	40
23	Rosmarinic acid	Intraperitoneally	DNFB-treated NC/Nga mice	↓	↓	N/D	41
24	<i>Radix Angelicae Dahuricae</i> (白芷)	Topically	House dust mite-treated NC/Nga mice	↓	↓	N/D	39
25	Silymarin	Topically	House dust mite-treated NC/Nga mice	↓	↓	N/D	52
26	<i>Lindera obtusiloba</i> (三桠乌药)	Topically	DNCB-treated BALB/c; House dust mite-treated BALB/c	↓	↓	N/D	53
27	<i>Broussonetia kazinoki</i> Sieb (小构树, 楮)	Topically	House dust mite-treated NC/Nga mice	↓	↓	N/D	54

(To Be Continued)

(Continued)

No.	Herbal extract	Route of administration	AD model	Dermatitis score	Ear thickness	Number of scratching	Reference
28	<i>Illicium verum</i> (八角)	Topically	House dust mite-treated NC/Nga mice	↓	↓	N/D	55
29	<i>Drynaria fortunei</i> (骨碎补)	Topically	House dust mite-treated NC/Nga mice	↓	↓	N/D	56
30	<i>Vernonia amygdalina</i> leaf (南非叶)	Topically	DNCB-treated NC/Nga mice	↓	↓	↓	57
31	<i>Sophora alopecuroides</i> L. (苦参)	Topically	DNFB-treated NC/Nga mice	↓	↓	N/D	58
32	<i>Chelidonium majus</i> (白屈菜)	Topically	DNCB-treated NC/Nga mice	↓	↓	↓	50
33	<i>Pleurotus eryngii</i> (灵芝)	Topically	DNCB-treated NC/Nga mice	↓	↓	N/D	59
34	<i>Radix ginseng</i> (人参), <i>Astragalus membranaceus</i> (黄芪), <i>Radix Paeoniae rubra</i> (赤芍), <i>Angelica gigantis Radix</i> (当归), <i>Atractylodes Rhizoma Alba</i> (白术), <i>Citri Unshius Pericarpium</i> (陈皮), <i>Forsythiae Fructus</i> (连翘), <i>Lonicerae Flos</i> (金银花), <i>Poria cocos</i> (茯苓), <i>Radix Angelicae Dahuricae</i> (白芷), <i>Glycyrrhiza uralensis</i> (甘草)	Topically	DNCB-treated BALB/c	↓	↓	N/D	60
35	<i>Cortex Cinnamomi</i> (肉桂)	Topically	House dust mite-treated NC/Nga mice	↓	N/D	N/D	61
36	<i>Rehmannia glutinosa</i> Libosch (地黄)	Topically	House dust mite-treated NC/Nga mice	↓	↓	N/D	62
37	<i>Radix ginseng</i> (人参)	Topically	DNCB-treated NC/Nga mice	↓	↓	N/D	63

Notes: N/D: non-determined; DNFB: 2,4-dinitrofluorobenzene; DNCB: 1-chloro-2,4-dinitrobenzene; TNCB: trinitrochlorobenzene; OX: oxazolone; PCI: 1-chloro-2,4,6-trinitrobenzene (picryl chloride)

of oral red ginseng extract in preventing a TNCB-induced increase in TEWL was comparable to that of oral cyclosporin A.⁽³⁸⁾ The mechanisms accounting for improving barrier function are not clear. One study showed that *Lithospermum erythrorhizon* increased epidermal ceramide content resulting from reduction of ceramidase expression in NC/Nga mice.⁽³⁰⁾ Together, these studies demonstrate that either systemic or topical herbal medicines prevent the development of AD, as indicated by prevention in expected AD-associated changes in epidermal permeability barrier function and SC hydration. These studies further suggest that assessment of SC biophysical properties could comprise a useful non-invasive approach to evaluate the potential efficacy of herbal medicines in preventing the development of AD. Herbal medicines that prevent

the increase in TEWL during the development of AD-like lesions are shown in Table 2.

Inflammation

Inflammatory Cell Infiltration

Although AD is generally considered as a Th2 cytokines mediated immune disorder,⁽⁷¹⁾ other inflammatory cells such as Th1, Th17, mast cells and B cells also are involved in the pathogenesis of AD.⁽⁷¹⁻⁷⁵⁾ Although AD is a Th2 dominant allergic disorder, characterized by elevated interleukin (IL)-4, 5 and 13 levels, mast cells also produce Th2 cytokines⁽⁷⁶⁾ and release histamine upon stimulation,^(77,78) all of which contribute to the development of AD.^(79,80) Accordingly, the preventive effects of herbal medicines on the development of AD also reflect an inhibition in inflammatory cell activation,

Table 2. Herbal Medicines that Prevent the Deterioration in Barrier Function During AD Development

Herbal extract	Route of administration	AD model	TEWL	Reference
<i>Actinidia arguta</i>	Orally	Magnesium deficient rats	↓	27
<i>Atractylodes lancea</i> rhizome, <i>Poria</i> , <i>Cnidium officinale</i> , <i>Radix Angelicae sinensis</i> , <i>Radix Bupleuri</i> , <i>Radix Glycyrrhizae</i> , and <i>Ramulus uncaria cum uncis</i>	Orally	NC/Nga mice	↓	37
Persimmon leaf	Orally	NC/Nga mice	↓	29
Panax red ginseng	Orally	TNCB-treated NC/Nga mice	↓	38
<i>Lithospermum erythrorhizon</i> (柴草)	Orally	OX-treated hairless mice	↓	67
<i>Bambusae caulis in Liquamen</i> (竹沥)	Topically	DNCB-treated hairless mice	↓	68

differentiation and infiltration.

Studies have shown that orally administrated extracts of herbal medicines inhibit inflammatory cell infiltration by eosinophils and mast cells in various murine AD models, including magnesium deficiency, NC/Nga mice or repeated DNFB applications.^(27,28,37,43,44-47) Pokharel, et al⁽⁴⁸⁾ showed that oral Sopungyangjae-Tang extract at a dosage of 40 mg/day for 12 weeks completely diminished inflammatory infiltration of the epidermis in a NC/Nga model. Even intraperitoneally administrated naringenin significantly reduced the number of CD4⁺ T cells, CD8⁺ T cells, total mast cells and degranulated mast cells in a NC/Nga model.⁽⁴⁰⁾ Oral herbal medicines not only reduce inflammatory cell infiltration in skin, but also decrease circulating inflammatory cells. It has been demonstrated that oral extract of *Actinidia arguta* prevented the increase of activated B (CD45RA⁺ positive) and CD11b⁺ cells in peripheral blood mononuclear cells (PBMC) and the skin of AD rats.⁽²⁷⁾ Furthermore, *Actinidia arguta* extract induced a 30% reduction in CD4⁺IL-4⁺ cells and over 40% reduction in CD19⁺IgE⁺ B cells in the splenocytes of ovalbumin-sensitized mice.⁽⁸¹⁾ Oral *Rheum rhabarbarum* extract caused an almost 200% reduction in neutrophil, and an over 85% reduction in eosinophil counts in the blood during the development of AD.⁽⁴²⁾ Moreover, oral extracts from the fruit of *Cudrania tricuspidata* exhibited similar effect to oral prednisolone in inhibiting mast cell infiltration in the NC/Nga model.⁽³⁴⁾ Another study showed that efficacy of oral ginsenoside Rh1 was comparable to prednisolone on inflammatory cell infiltration in an oxazolone-induced AD model.⁽⁴⁷⁾ Moreover, there was no difference in the extent of inhibition of cutaneous CD11a⁺ cell infiltration between oral *Panax ginseng* extract and oral cyclosporin A.⁽³⁸⁾

Not only do systemic administration, but also topical applications of herbal medicines reduce inflammatory cell infiltration during the development of AD. Qi, et al⁽⁶⁸⁾ showed that topical bamboo extract inhibited the expected increase in neutrophils and eosinophils in peripheral blood mononuclear cell (PBMC). Reduction in mast cell infiltration, induced by topical extracts from *Silybum marianum* or *Lindera obtusiloba*, has been reported in NC/Nga mice and in an DNCB-induced AD model.^(52,53) Interestingly, the inhibitory effect of topical extracts from *Broussonetia kazinoki* Sieb. on mast cell infiltration is superior to conventional immunosuppressor, tacrolimus, in NC/Nga mice.⁽⁵⁴⁾ Again, the efficacies of topical *Illicium verum* extract and *Drynaria fortunei* extracts on inhibition

of mast cell infiltration were also comparable to tacrolimus during the development of house dust mite-induced AD.^(55,56) In addition to inflammatory cell infiltration, the epidermal thickness normalized in *Angelicae Dahuricae Radix* extract-treated mice to an extent that was comparable to tacrolimus-treated skin in an AD model.⁽³⁹⁾ Taken together, these studies reveal that either systemic or topical administrations of herbal medicines prevent the development of AD by inhibiting inflammatory cell infiltration.

Cytokine/Chemokine Expression

T cells, eosinophils, mast cells, cytokines and chemokines play crucial role in the development of AD.⁽⁸²⁻⁸⁴⁾ Thus, blockade of cytokine production or action has become a common approach in preventing and treating AD.^(85,86) Alternatively, approaches that stimulate the production of anti-inflammatory cytokines, such as IL-10, also could be beneficial in AD. Several studies have shown that herbal medicines, that prevent the development of AD, also down-regulated pro-inflammatory cytokine expression. The pro-inflammatory cytokine, TNF- α binds to its receptors on T cells, B cells and antigen-presenting cells thereby regulating inflammatory responses. It is reported that oral ginseng extract is as effective as cyclosporin A in the reduction of cutaneous levels of both protein and mRNA for TNF- α , as well as thymic stromal lymphopoietin (TSLP), an epidermal cytokine that stimulates T cell maturation, in a hapten-induced AD model.⁽³⁸⁾ Pretreatment with topical *Vernonia amygdalina* leaf extracts remarkably lowered serum TNF- α levels and even normalized serum IL-4 levels in NC/Nga AD model.⁽⁵⁷⁾ In contrast, IL-10 is anti-inflammatory cytokine. Accordingly, inhibition of IL-10 action increases TNF- α and IL-1 β production.⁽⁸⁷⁾ Oral or topical applications of some herbal extracts increase IL-10 levels in serum⁽⁴⁹⁾ and skin⁽⁵⁸⁾ while decreasing cutaneous mRNA levels of IL-4 and TNF- α .⁽⁴⁹⁾

Although most herbal medicines, that simultaneously exhibit preventive benefit on the development of AD, lower pro-inflammatory cytokine levels, the effects of herbal medicines on cytokine and its mRNA expression varies among the herbal ingredients (Table 3). Majority of herbal medicines down-regulate Th2 cytokines, such as IL-4, IL-5 and IL-13. The influence of herbal medicines on Th1 cytokines such as interferon- γ (IFN)- γ varies dramatically. Some herbs increase IFN- γ ,^(31,88) while others decrease IFN- γ ,^(33,40,44) or cause a little changes

in IFN- γ .^(26,28,38) Yet, it is worth noting that both oral and topical apigenin inhibit the development of AD in murine models,^(26,89) but oral administrations of apigenin did not change serum levels of a wide range of cytokines,⁽²⁶⁾ suggesting that the inhibitory effect of some herbal medicine, such as apigenin, on the development of AD could be independent of cytokines. The influence of herbal medicines on cytokine mRNA expression also varies among tissues. It has been shown that topical Taglisodog-eum extract lowers mRNA levels of IL-31, IFN- γ and TNF- α in skin lesion, but not CD4⁺ T cells in draining lymph nodes.⁽⁶⁰⁾ Some herbal medicines induce down-regulation of some cytokine mRNA expression is attributable to inhibition of NF- κ B activation.^(60,63,64)

In addition to cytokines, chemokines also play crucial role in the development of AD.⁽⁹⁰⁻⁹²⁾ For example, macrophage inflammatory protein 1 (MIP-1/CCL3) can activate granulocytes (neutrophils, eosinophils and basophils) and induce the synthesis and release of other pro-inflammatory cytokines such as IL-1, IL-6 and TNF- α .^(93,94) Regulated upon activation normal T cell expressed and presumably secreted (RANTES, CCL5) can activate and recruit inflammatory cells into inflamed skin sites.⁽⁹⁵⁾ Both thymus activation regulated chemokine (TARC/CCL17) and the macrophage-derived chemokine, MDC/CCL22, are AD associated chemokines that recruit Th2 cells.⁽⁹⁶⁾ Increased levels of these chemokines have been observed in AD patients.^(94,97-100) Accordingly, some herbal medicines that prevent the development of AD also down-regulate chemokine expression. A number of studies have demonstrated that both oral and topical applications of some herbal extracts decrease both protein and mRNA levels for these chemokines (Table 3). Moreover, the effect of oral administration of *Cudrania tricuspidata* extract on serum TARC levels was comparable to oral prednisolone in NC/Nga mouse model of AD.⁽³⁴⁾ Topical applications of either *Illicium verum* extract or tacrolimus induce a comparable extent of inhibition in mRNA expression for TARC and RANTES.⁽⁵⁵⁾ Taken together, these evidence suggest that either oral or topical applications of herbal medicines can inhibit cytokine and/or chemokine expression, resulting attenuation of AD development.

IgE Levels

IgE is synthesized by B cells and further modulated by Th2 cytokines, such as IL-4 and IL-13,^(101,102) but a large portion of allergen-specific IgE

is also produced by CD138⁺ plasma cells.⁽¹⁰³⁾ IgE plays a crucial role in the development of AD. IgE molecules bind to the high affinity IgE receptor (Fc ϵ RI) on the surface of mast cells and dendritic cells, stimulating histamine release.⁽¹⁰⁴⁻¹⁰⁶⁾ Histamine in turn enhances dendritic cell function and cytokine production by T cells.⁽¹⁰⁷⁻¹⁰⁹⁾ Both AD patients and mice exhibit higher serum IgE levels.^(2,110,111) Studies have shown that most herbal medicines that inhibit AD development lower serum IgE levels. For example, Topical herbal medicines such as silymarin and aloperine also reduce serum IgE levels in NC/Nga AD model.⁽⁵²⁾ Some herbal medicines lower both total serum IgE and allergen specific IgE.^(53,81) Mice pre-fed with paramylon, an extract from *Euglena gracilis* Z, for 1 week before induction of AD significantly reduced serum IgE levels in NC/Nga mice.⁽⁵¹⁾ Finally, oral ginseng extract lowered serum IgE levels more than did cyclosporin A.⁽²⁹⁾

Yet, other herbal medicines inhibit the development of AD without altering serum IgE levels,^(30,44,112,113) and one study even showed that topical *Rehmannia glutinosa* extract increased serum IgE levels.⁽⁶²⁾ Together, these results suggest that the effects of herbal medicines on serum IgE levels vary among specific ingredients. The effects of herbal medicines on IgE levels are summarized in Table 3.

Other Pathogenic Mechanisms Impacted by Herbal Medicines

Colonization of *S. aureus* is another common and provocative feature of AD.⁽¹¹⁵⁾ Lee, et al⁽²⁸⁾ reported that oral *Rumex japonicus* houtt extract, beginning 3 days before TNCB challenges prevented the development of AD, and induced an over 60% reduction in the density of *S. aureus* on the skin. Toll-like receptors, such as TLR-2, 4, and 9 not only are involved in the pathogenesis of AD,⁽¹¹⁶⁻¹¹⁹⁾ but also protect against *S. aureus* infection.⁽¹²⁰⁻¹²²⁾ Oral administrations of *Actinidia arguta* extract at dose of 100 mg·kg⁻¹·day⁻¹ for 7 weeks during TNCB challenges remarkably increased the expression levels of mRNA for TLR-2,4,9 in skin.⁽³⁵⁾ During the development of cutaneous inflammation, migration of inflammatory cells into the skin is mediated by adhesion molecules, such as intercellular adhesion molecule 1 (ICAM-1) and vascular cell adhesion molecule 1 (VCAM-1).⁽¹²³⁻¹²⁵⁾ The cutaneous mRNA levels of ICAM-1 and VCAM-1 are significantly decreased by topical herbal extract treatment for 23 days.^(55,62) However, topical *Drynaria fortune*

Table 3. Influence of Herbal Medicines on Cytokine and Chemokine Expression during the Development of AD in Murine Models

Herbal extracts [△]	Cytokines and Chemokines (protein)	Cytokines and Chemokines (mRNA)	IgE	Histamine
1	*TARC ↓	N/D	↓	↓
2	N/D	IL-4 ↓, TNF-α ↓, IFN-γ ↓	↓	N/D
3	IL-4(-), IL-13(-), IFN-γ (-), Botaxin(-), MIP-1(-)	IL-4(-), IFN-γ ↓	↓	N/D
5	IL-4 ↓, IFN-γ ↑	IL-4 ↓, TNF-α ↓, IFN-γ ↓	↓	↓
7	IFN-γ ↓, IL-4(-)	N/D	(-)	N/D
8	IL-4 ↓, TARC ↓	N/D	↓	↓
9	IL-4 ↓, IL-5 ↓, IL-10 ↓, IL-13, IFN-γ ↑, IL-12 ↑	IL-4 ↓, IFN-γ ↑	↓	N/D
10	IL-4 ↓, IFN-γ (-)	N/D	↓	N/D
13	TNF-α ↓, TSLP ↓, IL-4(-), IL-17(-), and IFN-γ (-)	TNF-α (-), TSLP ↓	↓	N/D
14	IFN-γ ↓, IL-4(-)	N/D	(-)	N/D
15	N/D	IL-4(-), IFN-γ ↑	↓	N/D
16	N/D	IL-4 ↓, IFN-γ (-)	↓	N/D
17	IL-6 ↓	IL-4(-), IFN-γ ↑	N/D	N/D
18	*TARC ↓	*TARC ↓	↓	N/D
19	TARC ↓, IL-10 ↑	IL-4 ↓, TNF-α ↓	↓	N/D
20	TNF-α ↓, IL-4	N/D	↓	↓
21	IL-4 ↓, IL-18 ↓, IL-12 ↓, IFN-γ (-)	N/D	↓	N/D
Oral fermented barley extract ⁽¹¹²⁾	IL-4(-), IL-17 ↓, IFN-γ ↓	N/D	(-)	N/D
Oral hesperidin ⁽¹¹⁴⁾	IL-4(-), IL-10(-), IL-17 ↓, IFN-γ ↓	N/D	↓	N/D
22	*IL-4 ↓, *IFN-γ ↓	N/D	↓	N/D
23	*IL-4 ↓, *IFN-γ ↓	N/D	↓	
24	*TARC ↓	*TARC ↓, *MDC ↓	↓	↓
Topical <i>Bambusae caulis</i> in Liquamen ⁽⁶⁸⁾	N/D	IL-4 ↓, IL-13 ↓, TNF-α ↓, IFN-γ ↑	↓	N/D
25	IL-4 ↓	IL-4 ↓, TNF-α ↓, IFN-γ ↑	↓	N/D
26	IL-4 ↓, IL-13 ↓, IL-31, TNF-α ↓	N/D	↓	↓
27	IL-4 ↓, *TARC ↓	*TARC ↓, *MDC ↓, *RANTES ↓	↓	N/D
28	*TNF-α ↓, *IL-4 ↓, IL-6 ↓	IL-4 ↓, IL-6 ↓, TNF-α ↓, TARC ↓, RANTES ↓, IFN-γ (-)	↓	↓
29	IL-6 ↓	IL-4 ↓, IL-6 ↓, TNF-α ↓, IFN-γ (-)	↓	N/D
30	IL-4 ↓, IL-5 ↓, MIP-1 ↓, TNF-α ↓, IFN-γ ↓, Eotaxin ↓	N/D	↓	N/D
31	IL-1 β ↓, IL-4 ↓, IL-6 ↓, IL-10 ↑, IL-13 ↓, TNF-α ↓, IFN-γ ↓	N/D	↓	N/D
32	TNF-α ↓, IL-4 ↓	N/D	↓	↓
33	TARC ↓	IL-4 ↓, IL-5 ↓, IL-13 ↓, TNF-α ↓, IFN-γ ↓	↓	N/D
34	IL-4 ↓, IL-13 ↓	IL-4 ↓, IL-5 ↓, IL-13 ↓, IL-17 ↓, IL-31 ↓, TNF-α ↓, IFN-γ ↓	↓	N/D
35	TNF-α ↓, TARC ↓	IL-4, TNF-α ↓, TARC ↓, *MDC ↓, *RANTES ↓	↓	↓
36	*TARC ↓, *MDC ↓, *RANTES ↓	IL-4 ↓, TNF-α ↓, TARC ↓, MDC ↓, RANTES ↓	↑	↓
37	TARC ↓	TARC ↓, IL-4 ↓, IL-5 ↓, IL-13 ↓, TNF-α ↓, IFN-γ ↓	↓	N/D

Notes: [△]Numbers and route of administration are from Table 1. ACD: allergic contact dermatitis; ICD: irritant contact dermatitis; TSLP: thymic stromal lymphopoietin; TARC: thymus and activation-regulated chemokine; MDC: macrophage-derived chemokine; MIP-1: Macrophage inflammatory protein 1; RANTES: regulated upon activation normal T cell expressed and presumably secreted; (-): no changes, **In vitro* study.

extract treatment for the same period of time does not alter the cutaneous mRNA levels of ICAM-1 and VCAM-1.⁽⁷¹⁾ These studies indicate that herbal medicines that prevent the development of AD also regulate cutaneous innate immune response and cell migration.

Concluding Remarks

Before or during the hapten or allergen challenge status, either systemic or topical herbal medicines can inhibit the development of AD. These AD preventive herbs improve stratum corneum function, AD severity score and scratching via divergent mechanisms including improving epidermal permeability barrier function, lowering IgE levels, inhibiting inflammatory cell infiltration, cytokine expression and immune defense depending on the type of herbal ingredients utilized. The effect of herbs in preventing AD development is comparable or superior to conventional immune modulators, antihistamine agents and glucocorticoids, but with fewer adverse effects. Herbal medicines could be optional alternatives for preventing the development of AD.

Conflict of Interest

Authors have no conflict of interest.

REFERENCES

- Man MQ, Hatano Y, Lee SH, Man M, Chang S, Feingold KR, et al. Characterization of a hapten-induced, murine model with multiple features of atopic dermatitis: structural, immunologic, and biochemical changes following single versus multiple oxazolone challenges. *J Invest Dermatol* 2008;128:79-86.
- Scharschmidt TC, Man MQ, Hatano Y, Crumrine D, Gunathilake R, Sundberg JP, et al. Filaggrin deficiency confers a paracellular barrier abnormality that reduces inflammatory thresholds to irritants and haptens. *J Allerg Clin Immunol* 2009;124:496-506.
- Brandt C, Pavlovic V, Radbruch A, Worm M, Baumgrass R. Low-dose cyclosporine A therapy increases the regulatory T cell population in patients with atopic dermatitis. *Allergy* 2009;64:1588-1596.
- Hijnen DJ, ten Berge O, Timmer-de Mik L, Bruijnzeel-Koomen CA, de Bruin-Weller MS. Efficacy and safety of long-term treatment with cyclosporin A for atopic dermatitis. *J Eur Acad Dermatol Venereol* 2007;21:85-89.
- Jacob SE. Cyclosporin ophthalmic emulsion—a novel therapy for benzyl alcohol-associated eyelid dermatitis. *Contact Dermatitis* 2008;58:169-170.
- Ruzicka T, Bieber T, Schöpf E, Rubins A, Dobozy A, Bos JD, et al. A short-term trial of tacrolimus ointment for atopic dermatitis. *European Tacrolimus Multicenter Atopic Dermatitis Study Group. N Engl J Med* 1997;337:816-821.
- Hanifin JM, Ling MR, Langley R, Breneman D, Rafal E. Tacrolimus ointment for the treatment of atopic dermatitis in adult patients: part I, efficacy. *J Am Acad Dermatol* 2001;44(1 Suppl):S28-S38.
- Boguniewicz M, Fiedler VC, Raimier S, Lawrence ID, Leung DY, Hanifin JM. A randomized, vehicle-controlled trial of tacrolimus ointment for treatment of atopic dermatitis in children. *Pediatric Tacrolimus Study Group. J Allergy Clin Immunol* 1998;102(4 Pt 1):637-644.
- Kang S, Lucky AW, Pariser D, Lawrence I, Hanifin JM. Long-term safety and efficacy of tacrolimus ointment for the treatment of atopic dermatitis in children. *J Am Acad Dermatol* 2001;44(1 Suppl):S58-S64.
- Jacobi A, Antoni C, Manger B, Schuler G, Hertl M. Infliximab in the treatment of moderate to severe atopic dermatitis. *J Am Acad Dermatol* 2005;2(3 Pt 1):522-526.
- Cassano N, Loconsole F, Coviello C, Vena GA. Infliximab in recalcitrant severe atopic eczema associated with contact allergy. *Int J Immunopathol Pharmacol* 2006;19:237-240.
- Yang XG, Chen DY, Lin J, Du Y. Modulatory effect of the oral liquid expelling wind and stopping itching on Th1/Th2 cytokines in patients with dermatitis and eczema. *Chin J Dermatovenereol (Chin)* 2005;19:307-308.
- Shang JW. Surveying the curative effect on the Ointment of Qi Shen Lian to treat the eczema of the dermatitis. *Chin J Dermatovenereol (Chin)* 2007;21:448-448.
- Zhang XM, Lv DH, Zhang XM. Clinical observation of the therapeutic effect of Jinzhi Jieyin Gargle on patients with dermatitis and eczema. *Chin J Dermatovenereol (Chin)* 2006;20:220-221.
- Man MQ, Shi Y, Man M, Lee SH, Demerjian M, Chang S, et al. Chinese herbal medicine (Tuhui extract) exhibits topical anti-proliferative and anti-inflammatory activity in murine disease models. *Exp Dermatol* 2008;17:681-687.
- Man W, Man M, Hupe M, Martin-Ezquerria G, Feingold KR, Elias PM, et al. Topical herbal extract (Huangdang mixture) exhibits both preventive and therapeutic effects in murine acute irritant contact dermatitis. *Int J Dermatol* 2011;50:1421-1427.
- Li YE, Huang HJ. Clinical observation on the efficacy of topical aloe on the prevention of contact dermatitis in patients with arteriovenous internal fistula. *Attend Pract Res (Chin)* 2009;6(22):97.
- Xie ZM, Xie ZJ. In-house preparation of Fu'an Oil prevents the development of contact dermatitis: a study of 250 cases. *Clin J Trad Chin Med (Chin)* 2004;16:567-568.
- Liao RY, He Y, Meng W, Peng HT. Clinical observation on the efficacy of *Radix Arnebiae* ointment in the prevention of contact dermatitis-induced by drug: a 38 case studies. *Hebei J Tradit Chin Med (Chin)* 2013;35:517-518

20. Ye JQ, Zheng CX. Clinical observation of MEBO in the prevention of infant diaper dermatitis in 165 cases. *Zhejiang J Trad Chin Med (Chin)* 2010;45:272.
21. Park DK, Lee YG, Park HJ. Extract of *Rhus verniciflua* Bark suppresses 2,4-dinitrofluorobenzene-induced allergic contact dermatitis. *Evid Based Complement Alternat Med* 2013;2013:879696.
22. Lim HS, Seo CS, Ha H, Lee H, Lee JK, Lee MY, et al. Effect of *Alpinia katsumadai* Hayata on house dust mite-induced atopic dermatitis in NC/Nga mice. *Evid Based Complement Alternat Med* 2012;2012:705167.
23. Jin H, He R, Oyoshi M, Geha RS. Animal models of atopic dermatitis. *J Invest Dermatol* 2009;129:31-40.
24. Suto H, Matsuda H, Mitsuishi K, Hira K, Uchida T, Unno T, et al. NC/Nga mice: a mouse model for atopic dermatitis. *Int Arch Allergy Immunol* 1999;120(S1):70-75.
25. Collantes TM, Rho MC, Kwon HJ, Jung BG, Alfajaro MM, Kim DS, et al. Azuki bean (*Vigna angularis*) extract inhibits the development of experimentally induced atopic dermatitis-like skin lesions in NC/Nga mice. *Food Chem* 2012;132:1269-1275.
26. Yano S, Umeda D, Yamashita S, Yamada K, Tachibana H. Dietary apigenin attenuates the development of atopic dermatitis-like skin lesions in NC/Nga mice. *J Nutr Biochem* 2009;20:876-881.
27. Choi JJ, Park B, Kim DH, Pyo MY, Choi S, Son M, et al. Blockade of atopic dermatitis-like skin lesions by DA-9102, a natural medicine isolated from *Actinidia arguta*, in the Mg-deficiency induced dermatitis model of hairless rats. *Exp Biol Med (Maywood)* 2008;233:1026-1234.
28. Lee HS, Kim SK, Han JB, Choi HM, Park JH, Kim EC, et al. Inhibitory effects of *Rumex japonicus* houtt on the development of atopic dermatitis-like skin lesions in NC/Nga mice. *Br J Dermatol* 2006;155:33-38.
29. Matsumoto M, Kotani M, Fujita A, Higa S, Kishimoto T, Suemura M, et al. Oral administration of persimmon leaf extract ameliorates skin symptoms and transepidermal water loss in atopic dermatitis model mice, NC/Nga. *Br J Dermatol* 2002;146:221-227.
30. Kim J, Kim Y, Seo D, Kim S, Lee S, Cho Y. Oral supplementation of *Lithospermum erythrorhizon* prevents the development of atopic dermatitis with reducing ceramide degradation in the epidermis of NC/Nga mice. *Phytother Res* 2009;23:1250-1256.
31. Han NR, Moon PD, Kim HM, Jeong HJ. Effect of Pyeongwee-San (KMP6) on 2,4-dinitrofluorobenzene-induced atopic dermatitis-like skin lesions in NC/Nga mice. *Life Sci* 2012;90:147-153.
32. Lim H, Lee JH, Kim J, Kim YS, Kim HP. Effects of the rhizomes of *Atractylodes japonica* and atractylenolide I on allergic response and experimental atopic dermatitis. *Arch Pharm Res* 2012;35:2007-2012.
33. Lee SJ, Oh SG, Seo SW, Ahn HJ, Geum D, Cho JJ, et al. Oral administration of *Astragalus membranaceus* inhibits the development of DNFB-induced dermatitis in NC/Nga mice. *Biol Pharm Bull* 2007;30:1468-1471.
34. Lee H, Ha H, Lee JK, Seo CS, Lee NH, Jung DY, et al. The fruits of *Cudrania tricuspidata* suppress development of atopic dermatitis in NC/Nga mice. *Phytother Res* 2012;26:594-599.
35. Kim JY, Lee IK, Son MW, Kim KH. Effects of orally administered *Actinidia arguta* (Hardy Kiwi) fruit extract on 2-chloro-1,3,5-trinitrobenzene-induced atopic dermatitis-like skin lesions in NC/Nga mice. *J Med Food* 2009;12:1004-1015.
36. Park EJ, Park KC, Eo H, Seo J, Son M, Kim KH, et al. Suppression of spontaneous dermatitis in NC/Nga murine model by PG102 isolated from *Actinidia arguta*. *J Invest Dermatol* 2007;127:1154-1160.
37. Funakushi N, Yamaguchi T, Jiang J, Imamura S, Kuhara T, Suto H, et al. Ameliorating effect of Yokukansan on the development of atopic dermatitis-like lesions and scratching behavior in socially isolated NC/Nga mice. *Arch Dermatol Res* 2011;303:659-667.
38. Cho E, Cho SH. Effects of Korean red ginseng extract on the prevention of atopic dermatitis and its mechanism on early lesions in a murine model. *J Ethnopharmacol* 2013;145:294-302.
39. Lee H, Lee JK, Ha H, Lee MY, Seo CS, Shin HK. *Angelicae Dahuricae Radix* inhibits dust mite extract-Induced atopic dermatitis-like skin lesions in NC/Nga Mice. *Evid Based Complement Alternat Med* 2012;2012:743075.
40. Kim TH, Kim GD, Ahn HJ, Cho JJ, Park YS, Park CS. The inhibitory effect of naringenin on atopic dermatitis induced by DNFB in NC/Nga mice. *Life Sci* 2013;93:516-524.
41. Jang AH, Kim TH, Kim GD, Kim JE, Kim HJ, Kim SS, et al. Rosmarinic acid attenuates 2,4-dinitrofluorobenzene-induced atopic dermatitis in NC/Nga mice. *Int Immunopharmacol* 2011;11:1271-1277.
42. Park EJ, Kim B, Eo H, Park K, Kim Y, Lee HJ, et al. Control of IgE and selective T(H)1 and T(H)2 cytokines by PG102 isolated from *Actinidia arguta*. *J Allergy Clin Immunol* 2005;116:1151-1157.
43. Jiang J, Yamaguchi T, Funakushi N, Kuhara T, Fan PS, Ueki R, et al. Oral administration of Yokukansan inhibits the development of atopic dermatitis-like lesions in isolated NC/Nga mice. *J Dermatol Sci* 2009;56:37-42.
44. Kim DY, Jung JA, Kim TH, Seo SW, Jung SK, Park CS. Oral administration of *Uncariae rhynchophylla* inhibits the development of DNFB-induced atopic dermatitis-like skin lesions via IFN-gamma down-regulation in NC/Nga mice. *J Ethnopharmacol* 2009;122:567-572.
45. Choi MS, Kim EC, Lee HS, Kim SK, Choi HM, Park JH, et

- al. Inhibitory effects of *Saururus chinensis* (LOUR.) BAILL on the development of atopic dermatitis-like skin lesions in NC/Nga mice. *Biol Pharm Bull* 2008;31:51-56.
46. Kim EC, Lee HS, Kim SK, Choi MS, Lee S, Han JB, et al. The bark of *Betula platyphylla* var. *japonica* inhibits the development of atopic dermatitis-like skin lesions in NC/Nga mice. *J Ethnopharmacol* 2008;116:270-278.
47. Zheng H, Jeong Y, Song J, Ji GE. Oral administration of ginsenoside Rh1 inhibits the development of atopic dermatitis-like skin lesions induced by oxazolone in hairless mice. *Int Immunopharmacol* 2011;11:511-518.
48. Pokharel YR, Lim SC, Kim SC, Heo TH, Choi HK, Kang KW. Sopungyangjae-tang inhibits development of dermatitis in nc/nga mice. *Evid Based Complement Alternat Med* 2008;5:173-180.
49. Choi JH, Han EH, Park BH, Kim HG, Hwang YP, Chung YC, et al. *Platycodi Radix* suppresses development of atopic dermatitis-like skin lesions. *Environ Toxicol Pharmacol* 2012;33:446-452.
50. Yang G, Lee K, Lee MH, Kim SH, Ham IH, Choi HY. Inhibitory effects of *Chelidonium majus* extract on atopic dermatitis-like skin lesions in NC/Nga mice. *J Ethnopharmacol* 2011;138:398-403.
51. Sugiyama A, Hata S, Suzuki K, Yoshida E, Nakano R, Mitra S, et al. Oral administration of paramylon, a beta-1,3-D-glucan isolated from *Euglena gracilis* Z inhibits development of atopic dermatitis-like skin lesions in NC/Nga mice. *J Vet Med Sci* 2010;72:755-763.
52. Kang JS, Yoon WK, Han MH, Lee H, Lee CW, Lee KH, et al. Inhibition of atopic dermatitis by topical application of silymarin in NC/Nga mice. *Int Immunopharmacol* 2008;8:1475-1480.
53. Choi EJ, Lee S, Kim HH, Singh TS, Choi JK, Choi HG, et al. Suppression of dust mite extract and 2,4-dinitrochlorobenzene-induced atopic dermatitis by the water extract of *Lindera obtusiloba*. *J Ethnopharmacol* 2011;137:802-807.
54. Lee JK, Ha H, Lee HY, Park SJ, Jeong SL, Choi YJ, et al. Inhibitory effects of heartwood extracts of *Broussonetia kazinoki* Sieb on the development of atopic dermatitis in NC/Nga mice. *Biosci Biotechnol Biochem* 2010;74:1802-1806.
55. Sung YY, Yang WK, Lee AY, Kim DS, Nho KJ, Kim YS, et al. Topical application of an ethanol extract prepared from *Illicium verum* suppresses atopic dermatitis in NC/Nga mice. *J Ethnopharmacol* 2012;144:151-159.
56. Sung YY, Kim DS, Yang WK, Nho KJ, Seo HS, Kim YS, et al. Inhibitory effects of *Drynaria fortunei* extract on house dust mite antigen-induced atopic dermatitis in NC/Nga mice. *J Ethnopharmacol* 2012;144:94-100.
57. Ngatu NR, Okajima MK, Yokogawa M, Hirota R, Takaishi M, Eitoku M, et al. Anti-allergic effects of *Vernonia amygdalina* leaf extracts in hapten-induced atopic dermatitis-like disease in mice. *Allergol Int* 2012;61:597-607.
58. Yuan XY, Ma HM, Li RZ, Wang RY, Liu W, Guo JY. Topical application of aloperine improves 2,4-dinitrofluorobenzene-induced atopic dermatitis-like skin lesions in NC/Nga mice. *Eur J Pharmacol* 2011;658:263-269.
59. Choi JH, Kim HG, Jin SW, Han EH, Khanal T, Do MT, et al. Topical application of *Pleurotus eryngii* extracts inhibits 2,4-dinitrochlorobenzene-induced atopic dermatitis in NC/Nga mice by the regulation of Th1/Th2 balance. *Food Chem Toxicol* 2013;53:38-45.
60. Hwang JS, Kim JE, Kim HT, Yu YB, Im SH. Topical application of Taglisodog-eum inhibits the development of experimental atopic dermatitis. *J Ethnopharmacol* 2013;145:536-546.
61. Sung YY, Yoon T, Jang JY, Park SJ, Jeong GH, Kim HK. Inhibitory effects of *Cinnamomum cassia* extract on atopic dermatitis-like skin lesions induced by mite antigen in NC/Nga mice. *J Ethnopharmacol* 2011;133(2):621-628.
62. Sung YY, Yoon T, Jang JY, Park SJ, Kim HK. Topical application of *Rehmannia glutinosa* extract inhibits mite allergen-induced atopic dermatitis in NC/Nga mice. *J Ethnopharmacol* 2011;134:37-44.
63. Choi JH, Jin SW, Park BH, Kim HG, Khanal T, Han HJ, et al. Cultivated ginseng inhibits 2,4-dinitrochlorobenzene-induced atopic dermatitis-like skin lesions in NC/Nga mice and TNF- α /IFN- γ -induced TARC activation in HaCaT cells. *Food Chem Toxicol* 2013;56:195-203.
64. Kim DW, Park JY, Na GY, Lee SJ, Lee WJ. Correlation of clinical features and skin barrier function in adolescent and adult patients with atopic dermatitis. *Int J Dermatol* 2006;45:698-701.
65. Addor FA, Takaoka R, Rivitti EA, Aoki V. Atopic dermatitis: correlation between non-damaged skin barrier function and disease activity. *Int J Dermatol* 2012;51:672-676.
66. Knor T, Meholfić-Fetahović A, Mehmedagić A. Stratum corneum hydration and skin surface pH in patients with atopic dermatitis. *Acta Dermatovenerol Croat* 2011;19:242-247.
67. Lee JH, Jung KM, Bae IH, Cho S, Seo DB, Lee SJ, et al. Anti-inflammatory and barrier protecting effect of *Lithospermum erythrorhizon* extracts in chronic oxazolone-induced murine atopic dermatitis. *J Dermatol Sci* 2009;56:64-66.
68. Qi XF, Kim DH, Yoon YS, Li JH, Jin D, Deung YK, et al. Effects of *Bambusae caulis* in Liguamen on the development of atopic dermatitis-like skin lesions in hairless mice. *J Ethnopharmacol* 2009;123:195-200.
69. Simpson E, Böhlting A, Bielfeldt S, Bosc C, Kerrouche N. Improvement of skin barrier function in atopic dermatitis patients with a new moisturizer containing a ceramide precursor. *J Dermatolog Treat* 2013;24:122-125.
70. Sugawara T, Kikuchi K, Tagami H, Aiba S, Sakai S. Decreased lactate and potassium levels in natural moisturizing factor from the stratum corneum of mild atopic

- dermatitis patients are involved with the reduced hydration state. *J Dermatol Sci* 2012;66:154-159.
71. Romagnani S. Biology of human TH1 and TH2 cells. *J Clin Immunol* 1995;15:121-129.
 72. Grewe M, Bruijnzeel-Koomen CA, Schöpf E, Thepen T, Langeveld-Wildschut AG, Ruzicka T, et al. A role for Th1 and Th2 cells in the immunopathogenesis of atopic dermatitis. *Immunol Today* 1998;19:359-361.
 73. Di Cesare A, Di Meglio P, Nestle FO. A role for Th17 cells in the immunopathogenesis of atopic dermatitis? *J Invest Dermatol* 2008;128:2569-2571.
 74. Jee HM, Kim KW, Hong JY, Sohn MH, Kim KE. Increased serum B cell-activating factor level in children with atopic dermatitis. *Clin Exp Dermatol* 2010;35:593-598.
 75. Yanaba K, Kamata M, Asano Y, Tada Y, Sugaya M, Kadono T, et al. CD19 expression in B cells regulates atopic dermatitis in a mouse model. *Am J Pathol* 2013;182:2214-2222.
 76. Masuda A, Yoshikai Y, Aiba K, Matsuguchi T. Th2 cytokine production from mast cells is directly induced by lipopolysaccharide and distinctly regulated by c-Jun N-terminal kinase and p38 pathways. *J Immunol* 2002;169:3801-3810.
 77. Wei JF, Wei XL, Mo YZ, He SH. Induction of mast cell accumulation, histamine release and skin edema by N49 phospholipase A2. *BMC Immunol* 2009;10:21.
 78. Kim HH, Choi PH, Yoo JS, Jeon H, Chae BS, Park JS, et al. Ripe fruit of *Rubus coreanus* inhibits mast cell-mediated allergic inflammation. *Int J Mol Med* 2012;29:303-310.
 79. Suárez-Fariñas M, Dhingra N, Gittler J, Shemer A, Cardinale I, de Guzman Strong C, et al. Intrinsic atopic dermatitis shows similar TH2 and higher TH17 immune activation compared with extrinsic atopic dermatitis. *J Allergy Clin Immunol* 2013;132:361-370.
 80. Liu FT, Goodarzi H, Chen HY. IgE, mast cells, and eosinophils in atopic dermatitis. *Clin Rev Allergy Immunol* 2011;41:298-310.
 81. Jin JH, Ngoc TM, Bae K, Kim YS, Kim HP. Inhibition of experimental atopic dermatitis by rhubarb (rhizomes of *Rheum tanguticum*) and 5-lipoxygenase inhibition of its major constituent, emodin. *Phytother Res* 2011;25:755-759.
 82. Lee J, Noh G, Lee S, Youn Y, Rhim J. Atopic dermatitis and cytokines: recent patents in immunoregulatory and therapeutic implications of cytokines in atopic dermatitis—part I : cytokines in atopic dermatitis. *Recent Pat Inflamm Allergy Drug Discov* 2012;6:222-247.
 83. Yamanaka K, Mizutani H. The role of cytokines/chemokines in the pathogenesis of atopic dermatitis. *Curr Probl Dermatol* 2011;41:80-92.
 84. Chen L, Martinez O, Overbergh L, Mathieu C, Prabhakar BS, Chan LS. Early up-regulation of Th2 cytokines and late surge of Th1 cytokines in an atopic dermatitis model. *Clin Exp Immunol* 2004;138:375-387.
 85. Numerof RP, Asadullah K. Cytokine and anti-cytokine therapies for psoriasis and atopic dermatitis. *Bio Drugs* 2006;20:93-103.
 86. Stevceva L. Cytokines and their antagonists as therapeutic agents. *Curr Med Chem* 2002;9:2201-2207.
 87. Isomaki P, Luukkainen R, Saario R, Toivanen P, Punnonen J. Interleukin-10 functions as an antiinflammatory cytokine in rheumatoid synovium. *Arthritis Rheum* 1996;39:386-395.
 88. Makino T, Hamanaka M, Yamashita H, Mizukami H. Effect of bakumijogan, an herbal formula in traditional Chinese medicine, on atopic dermatitis-like skin lesions induced by mite antigen in NC/Jic mice. *Biol Pharm Bull* 2008;31:2108-2113.
 89. Man MQ, Hupe M, Sun R, Man G, Mauro TM, Elias PM. Topical apigenin alleviates cutaneous inflammation in murine models. *Evid Based Complement Alternat Med* 2012;2012:912028.
 90. Gros E, Bussmann C, Bieber T, Förster I, Novak N. Expression of chemokines and chemokine receptors in lesional and nonlesional upper skin of patients with atopic dermatitis. *J Allergy Clin Immunol* 2009;124:753-760.
 91. Pivarsci A, Homey B. Chemokine networks in atopic dermatitis: traffic signals of disease. *Curr Allergy Asthma Rep* 2005;5:284-290.
 92. Ying S, Meng Q, Barata LT, Kay AB. Macrophage inflammatory protein-1alpha and C-C chemokine receptor-1 in allergen-induced skin late-phase reactions: relationship to macrophages, neutrophils, basophils, eosinophils and T lymphocytes. *Clin Exp Allergy* 2001;31:1724-1731.
 93. Kaburagi Y, Shimada Y, Nagaoka T, Hasegawa M, Takehara K, Sato S. Enhanced production of CC-chemokines (RANTES, MCP-1, MIP-1alpha, MIP-1beta, and eotaxin) in patients with atopic dermatitis. *Arch Dermatol Res* 2001;293:350-355.
 94. Bisset LR, Schmid-Grendelmeier P. Chemokines and their receptors in the pathogenesis of allergic asthma: progress and perspective. *Curr Opin Pulm Med* 2005;11:35-42.
 95. Belperio JA, Dy M, Murray L, Burdick MD, Xue YY, Strieter RM, et al. The role of the Th2 CC chemokine ligand CCL17 in pulmonary fibrosis. *J Immunol* 2004;173:4692-4698.
 96. Yamada H, Chihara J, Matsukura M, Yasuba H, Yudate T, Tezuka T. Elevated plasma RANTES levels in patients with atopic dermatitis. *J Clin Lab Immunol* 1996;48:87-91.
 97. Shimada Y, Takehara K, Sato S. Both Th2 and Th1 chemokines (TARC/CCL17, MDC/CCL22, and Mig/CXCL9) are elevated in sera from patients with atopic dermatitis. *J Dermatol Sci* 2004;34:201-208.
 98. Fujisawa T, Nagao M, Hiraguchi Y, Katsumata H, Nishimori H, Iguchi K, et al. Serum measurement of thymus and activation-regulated chemokine/CCL17 in children with atopic dermatitis: elevated normal levels in infancy and age-specific analysis in atopic dermatitis. *Pediatr Allergy Immunol* 2009;20:633-641.
 99. Hayashida S, Uchi H, Takeuchi S, Esaki H, Moroi Y, Furue M. Significant correlation of serum IL-22 levels with CCL17

- levels in atopic dermatitis. *J Dermatol Sci* 2011;61:78-79.
100. Hokazono H, Omori T, Ono K. Effects of single and combined administration of fermented barley extract and gamma-aminobutyric acid on the development of atopic dermatitis in NC/Nga mice. *Biosci Biotechnol Biochem* 2010;74:135-139.
 101. Angeli V, Staumont D, Charbonnier A, Hammad H, Gosset P, Pichavant M, et al. Activation of the D prostanoid receptor 1 regulates immune and skin allergic responses. *J Immunol* 2004;172:3822.
 102. Eckl-Dorna J, Niederberger V. What is the source of serum allergen-specific IgE? *Curr Allergy Asthma Rep* 2013;13:281-287.
 103. Eckl-Dorna J, Pree I, Reisinger J, Marth K, Chen KW, Vrtala S, et al. The majority of allergen-specific IgE in the blood of allergic patients does not originate from blood-derived B cells or plasma cells. *Clin Exp Allergy* 2012;42:1347-1355.
 104. Frandsen PM, Krohn IJ, Hoffmann HJ, Schiøtz PO. The influence of IgE on cultured human mast cells. *Allergy Asthma Immunol Res* 2013;5:409-414.
 105. Ring J, Bieber T, Vieluf D, Kunz B, Przybilla B. Atopic eczema, Langerhans cells and allergy. *Int Arch Allergy Appl Immunol* 1991;94:194-201.
 106. Bieber T. The pro- and anti-inflammatory properties of human antigen-presenting cells expressing the high affinity receptor for IgE (Fc epsilon RI). *Immunobiology* 2007;212:499-503.
 107. Amaral MM, Davio C, Ceballos A, Salamone G, Cañones C, Geffner J, et al. Histamine improves antigen uptake and cross-presentation by dendritic cells. *J Immunol* 2007;179:3425-3433.
 108. Jutel M, Watanabe T, Klunker S, Akdis M, Thomet OA, Malolepszy J, et al. Histamine regulates T-cell and antibody responses by differential expression of H1 and H2 receptors. *Nature* 2001;413:420-425.
 109. Gutzmer R, Mommert S, Gschwandtner M, Zwingmann K, Stark H, Werfel T. The histamine H4 receptor is functionally expressed on T(H)2 cells. *J Allergy Clin Immunol* 2009;123:619-625.
 110. Schäfer T, Heinrich J, Wjst M, Adam H, Ring J, Wichmann HE. Association between severity of atopic eczema and degree of sensitization to aeroallergens in schoolchildren. *J Allergy Clin Immunol* 1999;104:1280-1284.
 111. Hatano Y, Man MQ, Uchida Y, Crumrine D, Scharschmidt TC, Kim EG, et al. Maintenance of an acidic stratum corneum prevents emergence of murine atopic dermatitis. *J Invest Dermatol* 2009;129:1824-1835.
 112. Iguchi T, Kawata A, Watanabe T, Mazumder TK, Tanabe S. Fermented barley extract suppresses the development of atopic dermatitis-like skin lesions in NC/Nga mice, probably by inhibiting inflammatory cytokines. *Biosci Biotechnol Biochem* 2009;73:489-493.
 113. Abeck D, Mempel M. *Staphylococcus aureus* colonization in atopic dermatitis and its therapeutic implications. *Br J Dermatol* 1998;139(S53):13-16.
 114. Nagashio Y, Matsuura Y, Miyamoto J, Kometani T, Suzuki T, Tanabe S. Hesperidin inhibits development of atopic dermatitis-like skin lesions in NC/Nga mice by suppressing Th17 activity. *J Funct Foods* 2013;5:1633-1641.
 115. Gong JQ, Lin L, Lin T, Hao F, Zeng FQ, Bi ZG, et al. Skin colonization by *Staphylococcus aureus* in patients with eczema and atopic dermatitis and relevant combined topical therapy: a double-blind multicentre randomized controlled trial. *Br J Dermatol* 2006;155:680-687.
 116. Lesiak A, Smolewski P, Sobolewska-Sztychny D, Sysa-Jedrzejowska A, Narbutt J. The role of T-regulatory cells and Toll-like receptors 2 and 4 in atopic dermatitis. *Scand J Immunol* 2012;76:405-410.
 117. Sakai T, Kogiso M, Mitsuya K, Komatsu T, Yamamoto S. Defect of toll-like receptor 9-mediated activation in NC/Nga mouse macrophages. *Immunol Lett* 2006;106:91-95.
 118. Kuo IH, Carpenter-Mendini A, Yoshida T, McGirt LY, Ivanov AI, Barnes KC, et al. Activation of epidermal toll-like receptor 2 enhances tight junction function: implications for atopic dermatitis and skin barrier repair. *J Invest Dermatol* 2013;133:988-998.
 119. Yimin, Kohanawa M, Zhao S, Ozaki M, Haga S, Nan G, et al. Contribution of Toll-like receptor 2 to the innate response against *staphylococcus aureus* infection in mice. *PLoS One* 2013;8:e74287.
 120. Fournier B. The function of TLR2 during staphylococcal diseases. *Front Cell Infect Microbiol* 2012;2:167.
 121. Stenzel W, Soltek S, Sanchez-Ruiz M, Akira S, Miletic H, Schlüter D, et al. Both TLR2 and TLR4 are required for the effective immune response in *Staphylococcus aureus*-induced experimental murine brain abscess. *Am J Pathol* 2008;172:132-145.
 122. Parker D, Prince A. *Staphylococcus aureus* induces type I IFN signaling in dendritic cells via TLR9. *J Immunol* 2012;189:4040-4046.
 123. Singer KH, Le PT, Denning SM, Whichard LP, Haynes BF. The role of adhesion molecules in epithelial-T-cell interactions in thymus and skin. *J Invest Dermatol* 1990;94(6 S):85S-90S.
 124. Goebeler M, Gutwald J, Roth J, Meinardus-Hager G, Sorg C. Expression of intercellular adhesion molecule-1 in murine allergic contact dermatitis. *Int Arch Allergy Appl Immunol* 1990;93:294-299.
 125. Johnson LA, Clasper S, Holt AP, Lalor PF, Baban D, Jackson DG. An inflammation-induced mechanism for leukocyte transmigration across lymphatic vessel endothelium. *J Exp Med* 2006;203:2763-2777.

(Received July 6, 2014)
 Edited by YUAN Lin