

REVIEW ARTICLE

Skin disorders are prominent features in primary immunodeficiency diseases: A systematic overview of current data

Jill de Wit¹  | Romke J. K. Brada¹ | Joyce van Veldhuizen¹ | Virgil A. S. H. Dalm² | Suzanne G. M. A. Pasmans¹

¹Department of Dermatology, Erasmus MC University Medical Center, Rotterdam, the Netherlands

²Department of Internal Medicine, Division of Clinical Immunology and Department of Immunology, Erasmus MC University Medical Center, Rotterdam, the Netherlands

Correspondence

Jill de Wit, Department of Dermatology, Erasmus MC University Medical Center, Rotterdam, the Netherlands.

Email: j.dewit.1@erasmusmc.nl

and

Suzanne G. M. A. Pasmans, Department of Pediatric Dermatology, Sophia Children's Hospital, Erasmus MC University Medical Center, Rotterdam, the Netherlands.

Email: s.pasmans@erasmusmc.nl

Funding information

The Department of Dermatology of the Erasmus MC University Medical Center Rotterdam received an unrestricted grant of Micros Human Health, the Netherlands.

Abstract

Primary immunodeficiency diseases (PIDs) are characterized by an increased risk of infections, autoimmunity, autoinflammation, malignancy, and allergic disorders. Skin disorders are also common clinical features in PIDs and may be among the presenting manifestations. Recognition of specific PID-associated skin conditions in combination with other clinical features as described in the currently used warning signs could raise suspicion of an underlying PID. We aimed to provide a systematically obtained overview of skin disorders and their prevalence in PIDs. Secondary, the prevalence of *Staphylococcus* (*S.*) *aureus*-associated skin disorders and atopy was reviewed, as these are the most prominent skin features in PIDs. A systematic search was performed in EMBASE, MEDLINE, Web of Science, Cochrane, and Google Scholar (up to May 9, 2018). All original observational and experimental human studies that address the presence of skin disorders in PIDs were selected. We rated study quality using the Institute of Health Economics Quality Appraisal Checklist for Case Series Studies. Sixty-seven articles (5030 patients) were included. Study quality ranged from 18.2% to 88.5%. A broad spectrum of skin disorders was reported in 30 PIDs, mostly in single studies with a low number of included patients. An overview of associated PIDs per skin disorder was generated. Data on *S. aureus*-associated skin disorders and atopy in PIDs were limited. In conclusion, skin disorders are prominent features in PIDs. Through clustering of PIDs per skin disorder, we provide a support tool to use in clinical practice that should raise awareness of PIDs based on presenting skin manifestations.

KEYWORDS

atopy, primary immunodeficiency disease, skin manifestation, *Staphylococcus aureus*

1 | INTRODUCTION

Primary immunodeficiency diseases (PIDs) represent a heterogeneous group of inherited disorders caused by mutations in genes encoding functional proteins of the immune cells. Based on registries and epidemiologic surveys, it has been suggested that six

million people are living with a PID worldwide, whereas only 27,000–60,000 patients have been identified to date.¹ PIDs are usually characterized by recurrent and/or severe infections as well as an increased risk of autoimmunity, autoinflammation, malignancy, and allergic disorders.^{2–4} Moreover, both infectious and noninfectious skin disorders are common in PIDs and may be among the presenting clinical manifestations.^{5–8} *Staphylococcus* (*S.*) *aureus*-induced skin infections are the most common infectious

Brada and van Veldhuizen equally contributed to this work.

skin disorders reported in PIDs, including leukocyte adhesion defects (LAD), chronic granulomatous disease (CGD), severe congenital neutropenia, and hyper-IgE syndrome (HIES).⁹⁻¹¹ On the other hand, dermatitis is one of the most prominent noninfectious skin manifestations in PIDs and may be part of the atopic syndrome.¹² Patients with an atopic constitution show next to atopic dermatitis (AD) tendency toward development of food allergies, asthma, and rhinoconjunctivitis.¹³

Based on previous narrative reviews without a systematic approach, *S. aureus* skin infections, dermatitis, and other skin disorders as well as atopy seem to be all fairly common in patients with a PID, but are also frequently described in the general population.¹² Therefore, it is of importance to realize that presence of specific skin symptoms alone does not necessarily point toward a PID. However, recognition of specific skin conditions in combination with other clinical features suggestive of an immunodeficiency should raise awareness to an underlying PID and may facilitate earlier diagnosis of PIDs.¹⁴

The aim of this review was to provide a systematically obtained overview of skin disorders and their prevalence in patients with PIDs. Focusing on two prevalent skin disorders in PIDs, the relation between PIDs and *S. aureus*-related skin disorders and atopy will be reviewed in more detail.

2 | METHODS

2.1 | Studies

This review with a systematic approach was conducted and reported according to the Meta-analysis Of Observational Studies in Epidemiology (MOOSE) guidelines, where applicable.¹⁵ All original observational and experimental human studies were included. We selected both articles reporting skin disorders in patients with PIDs and articles presenting a differential diagnosis of a specific skin disorder that includes a PID. No restrictions were made with respect to publication date and language. We excluded case reports (<5 patients), conference abstracts, letters, and editorials as the quality of these types of articles can be highly variable. Also articles describing acquired immunodeficiencies, articles reporting skin disorders in PIDs that developed after or during treatment/intervention, and articles in which the description of skin disorders in PIDs was not part of the results section were excluded. Data on skin disorders were only extracted if at least five patients per PID were reported.

2.2 | Study participants

Patients of all ages with a PID according to Picard et al¹⁶ from both hospital setting and general population were included.

2.3 | Study outcomes

The primary outcome is the presence of skin disorders in PIDs. Secondary outcomes include the prevalence of skin disorders in PIDs, *S. aureus*-associated skin disorders in PIDs, and PIDs associated with

an atopic constitution (ie, atopic dermatitis, food allergy, asthma, rhinoconjunctivitis).

2.4 | Search strategy

The electronic search was conducted in EMBASE, MEDLINE, Web of Science, Cochrane, and Google Scholar up to May 9, 2018 (Table S1). The search was composed of terms of the categories primary immunodeficiency, skin disorder, *S. aureus*, and atopy supplemented by specific PIDs and skin disorders based on recent literature.^{12,17-19}

2.5 | Study selection and data extraction

All studies identified in the systematic search were screened for relevance by title and abstract. Duplicates and studies that did not meet our inclusion criteria were excluded (Appendix S1). Remaining articles were assessed for eligibility by full-text review. Furthermore, a cross-reference check was performed to identify other eligible studies based on the reference lists of all included articles and relevant review articles. Translation of non-English studies was conducted officially. Study selection and data extraction were performed independently by two researchers (JdW and JvV, JdW and RB or JvV and RB). Disagreements were resolved and consensus was reached. If one population was described in different articles, we included the study with the most detailed description of the results. The methodological quality of the individual articles was rated using the Institute of Health Economics (IHE) Quality Appraisal Checklist for Case Series Studies (Appendix S1).²⁰

2.6 | Analysis of data

The prevalence of skin disorders in PIDs was extracted from the included studies. If required, the prevalence was calculated with the available raw data. Because the reported number of patients with a PID was mainly low, the proportion of patients with a PID and skin disorders was descriptively presented. Proportions of skin disorders in PIDs were compared to the prevalence of skin disorders in the general population.²¹⁻²⁶ Data from the general population were based on a birth cohort in Finland ($n = 1932$, age 45-47 years) and a Dermatology outpatient clinic in Turkey ($n = 11\,040$, age 1-99 years).^{21,22} In addition, a nationwide study of Furue et al²³ reported the prevalence of cutaneous disorders in 67 448 Japanese patients of all ages. In the study of Verhoeven et al,²⁴ the skin disease prevalence per 1000 patient-years in family practices in the Netherlands was converted to a point prevalence in the general population ($n = 501$, age 18-97 years). Finally, two studies from the United States of America and the United Kingdom performed in 1978 and 1976 showed the prevalence of skin disorders in community studies in, respectively, 20 749 (age 1-74 years) and 614 (age 15-74 years) patients.^{25,26}

3 | RESULTS

3.1 | Study characteristics

The literature search identified 15 871 studies. Removal of duplicates resulted in 12 834 studies. Screening on title and abstract yielded 86 full-text articles of which 36 articles remained after full-text screening. Finally, after cross-reference check, a total of 67 articles (5030 patients) were included for further analysis (Figure 1). Skin disorders in patients with PIDs were described in 67 articles, and three articles reported PIDs as part of the differential diagnosis of a specific skin disorder. Fifty-seven studies showed a mean percentage of males of 62.2%. Both children and adults were included with a mean age of 15.8 years, reported in 26 articles. The IHE Quality Appraisal Checklist for Case Series Studies ranged from 18.2% to 88.5% (Table S1).

3.2 | Skin disorders and their prevalence in primary immunodeficiency diseases

Thirty individual PIDs and their related skin manifestations were found. We categorized the skin disorders in 15 main groups and in 20 more specific subgroups (Table 1). The skin disorders per PID were mainly reported in single studies. Therefore, meta-analysis was not possible. The presence of skin telangiectasia, café au lait macules, and hypopigmented macules in ataxia-telangiectasia (AT); skin abscesses in HIES; atopic dermatitis in hypogammaglobulinemia; atopic dermatitis, alopecia (areata), vitiligo, and psoriasis in selective IgA deficiency (SIgAD); alopecia, vitiligo, and nail dystrophy in autoimmune polyendocrinopathy candidiasis ectodermal dystrophy (APECED); and abscesses and granuloma in CGD were confirmed in at least three articles. All reported skin disorders per PID were used to provide an overview of PIDs per skin disorder group (Figure 2).




FIGURE 1 Flow chart of search strategy and study selection

TABLE 1 Skin disorders and their prevalence in primary immunodeficiency diseases

Primary immunodeficiency disease					General population	
Main groups of skin disorders	Subgroups of skin disorders	Skin disorders as reported in included articles	Number of reported cases with skin disorder (proportion)	Prevalence of skin disorder (%)	Prevalence of skin disorder (%) [†]	
Immunodeficiencies affecting cellular and humoral immunity						
Severe combined immunodeficiency						
Dermatitis-like lesions		Seborrheic dermatitis	2/9 ⁷	22.2	2.2-11.7 ²¹⁻²⁵	
Skin infections	Fungal skin infections	Candidiasis	4/9 ⁷	44.4	0.6-1.0 ^{22,23}	
Omenn syndrome						
Hair abnormalities	Hair loss disorders	Severe alopecia	5/7 ⁵²	71.4	0.4-2.5 ²¹⁻²³	
		Alopecia of eyelashes and eyebrows	3/7 ⁵²	42.9	-	
Erythematous skin lesions		Exfoliative erythroderma ^a	7/7 ⁵²	100	0.1 ²³	
Other skin disorders		Skin induration	6/7 ⁵²	85.7	-	
Combined immunodeficiencies with associated or syndromic features						
Ataxia-telangiectasia						
Dermatitis-like lesions		Dermatitis	1/62 ⁶	1.6	32.4 ²⁴	
		Eczema	2/22 ⁵³	9.1	9.0-27.4 ^{21,26}	
		Nummular eczema	1/22 ⁵³	4.5	1.9-2.2 ^{21,22}	
		Seborrheic rash	2/32 ⁵⁴	6.3	2.2-11.7 ²¹⁻²⁵	
Hair abnormalities	Excessive hair growth disorders	Hypertrichosis	7/32 ⁵⁴	21.9	-	
		Hirsutism	2/12 ³³	16.7	0.4 ²²	
	Hair pigmentation disorders	Poliosis	5/12 ³³	41.7	-	
Skin infections	Fungal skin infections	Oral candidiasis	1/12 ³³	8.3	-	
		Coccidioidomycosis	1/22 ⁵³	4.5	-	
	Viral skin infections	Viral warts	2/32, ⁵⁴ 8/22 ⁵³	6.3-36.4	3.4-4.5 ^{22,23,26}	
		Herpes simplex	2/12 ³³	16.7	0.8-1.0 ^{22,23}	
	Bacterial skin infections	Chronic impetigo	1/22 ⁵³	4.5	0.8-1.6 ^{22,23}	
		Impetigo	1/12 ³³	8.3	0.8-1.6 ^{22,23}	
Erythematous skin lesions		Pinpoint erythematous macules	2/12 ³³	16.7	-	
Vascular disorders	Telangiectasia	Skin telangiectasia	6/62, ⁶ 16/26, ⁷ 4/12 ³³	9.7-61.5	-	
		Telangiectasia on cheeks or nose	4/32, ⁵⁴ 18/22 ⁵³	12.5-81.8	-	
		Telangiectasia on ears	15/32 ⁵⁴	46.9	-	
		Telangiectasia on back/shoulders/neck	5/32 ⁵⁴	15.6	-	
	Vasculitis	Allergic vasculitis	1/22 ⁵³	4.5	-	
	Pigmentation disorders	Hyperpigmentation disorders	Café au lait macules	27/32, ⁵⁴ 3/22, ⁵³ 4/12 ³³	13.6-84.4	12.4 ²¹
			Pigmented nevi (>5 mm)	12/32 ⁵⁴	37.5	-
Hyperpigmentation			1/62 ⁶	1.6	-	
Acanthosis nigricans			3/12 ³³	25.0	-	
Hypopigmentation disorders		Hypopigmented macules	3/62, ⁶ 14/32, ⁵⁴ 2/12 ³³	4.8-43.8	-	
	Albinism	1/32, ⁵⁴ 8/22 ⁵³	3.1-36.4	-		
	Vitiligo	1/12 ³³	8.3	1.2-1.7 ²¹⁻²³		

(Continues)

TABLE 1 (Continued)

Primary immunodeficiency disease					General population	
Main groups of skin disorders	Subgroups of skin disorders	Skin disorders as reported in included articles	Number of reported cases with skin disorder (proportion)	Prevalence of skin disorder (%)	Prevalence of skin disorder (%) [†]	
	Other pigmentation disorders	Blue nevus	1/22 ⁵³	4.5	1.3 ²¹	
		Freckles	1/22 ⁵³	4.5	-	
	Neoplastic disorders	Others neoplastic disorders	Basal cell carcinoma	1/22 ⁵³	4.5	0.4-0.5 ^{21,23}
			Juvenile melanoma	1/22 ⁵³	4.5	-
Rash		Facial papulosquamous rash	13/32 ⁵⁴	40.6	-	
Nail disorders	Noninfectious nail disorders	Congenital nail dystrophy	2/12 ³³	16.7	-	
Granulomatous disorders		Skin granulomas	8/8 ⁵⁵	100	0.3 ²³	
Other skin disorders		Lichen simplex chronicus	1/32 ⁵⁴	3.1	3.0 ²²	
		Sclerodermoid changes	1/22 ⁵³	4.5	-	
		Senile keratosis (actinic keratosis)	1/22 ⁵³	4.5	0.4-0.6 ^{21,23}	
		Aged skin	2/22 ⁵³	9.1	-	
		Shagreen patch	1/12 ³³	8.3	-	
		Lipoatrophy	1/12 ³³	8.3	-	
		Hydroa vacciniforme	1/12 ³³	8.3	-	
		Dermatofibroma	1/12 ³³	8.3	0.2-22.2 ^{21,23}	
		Purpura	5/26 ⁷	19.2	-	
Wiskott-aldrich syndrome						
Dermatitis-like lesions		Eczema	5/5 ⁷	100	9.0-27.4 ^{21,26}	
Skin infections	Fungal skin infections	Yeast/fungi	18/154 ⁵⁶	11.7	-	
		Viral skin infections	Varicella	25/154 ⁵⁶	16.2	-
	Bacterial skin infections	HSV I/HSV II	24/154 ⁵⁶	15.6	0.8-1.0 ^{22,23}	
		Molluscum contagiosum	13/154 ⁵⁶	8.4	0.8-0.9 ^{22,23}	
		Warts	10/154 ⁵⁶	6.5	3.4-4.5 ^{22,23,26}	
		Impetigo	19/154 ⁵⁶	12.3	0.8-1.6 ^{22,23}	
		Abscesses	19/154 ⁵⁶	12.3	1.7 ²²	
		Cellulitis	19/154 ⁵⁶	12.3	0.9-1.5 ^{22,23}	
		Vascular disorders	Vasculitis	Henoch-Schönlein purpura	8/154 ⁵⁶	5.2
Skin vasculitis	12/55 ⁵⁷			7.8	-	
Hyper-IgE syndrome unspecified						
Dermatitis-like lesions		Atopic dermatitis	5/6, ⁶ 5/5, ⁵⁸ 28/43, ³⁴ 5/8 ²⁷	62.5-100	2.2-12.4 ²¹⁻²⁴	
		Eczema	30/30 ⁵⁹	100	9.0-27.4 ^{21,26}	
Skin infections	Fungal skin infections	Candida	4/6, ⁶ 25/30 ⁵⁹	66.7-83.3	0.6-1.0 ^{22,23}	
	Bacterial skin infections	Skin abscesses	6/6, ⁶ 36/43, ³⁴ 3/8, ²⁷ 26/30 ⁵⁹	37.5-100	1.7 ²²	
	Other skin infections	Skin infection	3/5 ⁵⁸	60.0	4.6-43.5 ^{21,25,26}	
Ulcers	Oral ulcers	Oral aphthous ulcers	1/5 ⁵⁸	20.0	0.7 ²²	
		Oral ulceration	8/11 ⁶⁰	72.7	0.7 ²²	
Rash		Newborn rash	35/43 ³⁴	81.4	-	
		Maculopapular rash	1/6 ⁶	16.7	-	
		Papulopustular eruption	8/8 ²⁷	100	-	

(Continues)

TABLE 1 (Continued)

Primary immunodeficiency disease					General population
Main groups of skin disorders	Subgroups of skin disorders	Skin disorders as reported in included articles	Number of reported cases with skin disorder (proportion)	Prevalence of skin disorder (%)	Prevalence of skin disorder (%) [†]
Acne-like lesions		Neonatal acne	6/43 ³⁴	14.0	-
Other skin disorders		Coarse face	3/6, ⁶ 2/8 ²⁷	25.0-50.0	-
Autosomal dominant hyper-IgE syndrome					
Dermatitis-like lesions		Ecematous dermatitis	20/21 ²⁸	95.2	32.4 ²⁴
		Ecema	47/82, ³⁵ 17/17 ²⁹	57.3-100	9.0-27.4 ^{21,26}
Skin infections	Fungal skin infections	Oral candidiasis	4/21 ²⁸	19.0	-
		Genitalia fungal infection	1/17 ²⁹	5.9	-
		Varicella zoster virus infection	9/21 ²⁸	42.9	-
	Viral skin infections	Herpes simplex virus infection	4/21 ²⁸	19.0	0.8-1.0 ^{22,23}
		Herpes infection	5/67 ³⁵	7.5	0.8-1.0 ^{22,23}
		Molluscum contagiosum	1/21, ²⁸ 1/82 ³⁵	1.2-4.8	0.8-0.9 ^{22,23}
		Cold abscesses	20/21, ²⁸ 9/17 ²⁹	52.9-95.2	1.7 ²²
		Skin abscesses	61/82 ³⁵	74.4	1.7 ²²
		Cellulitis	15/82 ³⁵	18.3	0.9-1.5 ^{22,23}
		Pustulosis	14/17 ²⁹	82.4	-
		Folliculitis	7/17 ²⁹	41.2	1.1-6.0 ^{21,23}
	Other skin infections	Recurrent skin infections	17/17 ²⁹	100	4.6-43.5 ^{21,25,26}
Ulcers	Oral ulcers	Oral ulcer	1/17 ²⁹	2.4	0.7 ²²
Neoplastic disorders	Cutaneous lymphomas	Pilotropic cutaneous T-cell lymphoma	1/21 ²⁸	4.8	-
	Other neoplastic disorders	Squamous cell carcinoma	1/82 ³⁵	1.2	0.3-0.7 ^{23,24}
Rash		Papulopustular rash (<2 months)	14/21 ²⁸	66.7	-
Nail disorders	Infectious nail disorders	Chronic paronychia	8/21 ²⁸	38.1	-
		Onychomycosis	23/82, ³⁵ 4/17 ²⁹	23.5-28.0	9.4 ²¹
Urticaria		Urticaria	13/82 ³⁵	15.9	0.5-8.3 ²¹⁻²³
Other skin disorders		Lichenification	1/21 ²⁸	4.8	-
		Coarse facies	10/21 ²⁸	47.6	-
		Dry skin	18/21 ²⁸	85.7	-
		Thrush	17/82, ³⁵ 6/17 ²⁹	20.7-35.3	-
		Angioedema	9/82 ³⁵	11.0	-
Autosomal recessive hyper-IgE syndrome					
Dermatitis-like lesions		Ecema	19/21, ³⁶ 7/10 ³⁷	70.0-90.5	9.0-27.4 ^{21,26}
		Atopic dermatitis	7/10 ³⁷	70.0	2.2-12.4 ²¹⁻²⁴
Skin infections	Fungal skin infections	Mucocutaneous candidiasis	9/21 ³⁶	42.9	0.6-1.0 ^{22,23}
		Chronic candidiasis of mucosal sites	10/13 ³⁰	76.9	-
	Viral skin infections	Viral warts	13/21 ³⁶	61.9	3.4-4.5 ^{22,23,26}
		Verruca plana	1/10 ³⁷	10.0	3.4-4.5 ^{22,23,26}

(Continues)

TABLE 1 (Continued)

Primary immunodeficiency disease					General population	
Main groups of skin disorders	Subgroups of skin disorders	Skin disorders as reported in included articles	Number of reported cases with skin disorder (proportion)	Prevalence of skin disorder (%)	Prevalence of skin disorder (%) [†]	
	Bacterial skin infections	Herpes simplex virus	12/21, ³⁶ 8/13 ³⁰	57.1-61.5	0.8-1.0 ^{22,23}	
		Recurrent herpes	1/10 ³⁷	10.0	0.8-1.0 ^{22,23}	
		Molluscum contagiosum	10/21, ³⁶ 4/13 ³⁰	30.8-47.6	0.8-0.9 ^{22,23}	
		Severe primary varicella zoster	7/21, ³⁶ 2/13 ³⁰	15.4-33.3	-	
		Herpes zoster	5/21 ³⁶	23.8	1.4-2.4 ^{22,23}	
		Bacterial skin infection	17/21 ³⁶	81.0	-	
		Skin abscesses	11/13 ³⁰	84.6	1.7 ²²	
		MRSA wound infected eczema	1/10 ³⁷	10.0	-	
		Other skin infections	Recurrent stomatitis	1/10 ³⁷	10.0	-
		Neoplastic disorders	Cutaneous lymphomas	Cutaneous T-cell lymphoma	1/21 ³⁶	4.8
Other neoplastic disorders	Squamous cell carcinoma		4/21 ³⁶	19.0	0.3-0.7 ^{23,24}	
Rash		Severe eczematoid rash	13/13 ³⁰	100	9.0-27.4 ^{21,26}	
		Newborn rash	5/21 ³⁶	23.8	-	
Other skin disorders		Immune thrombocytopenic purpura	1/10 ³⁷	10.0	-	
Nijmegen breakage syndrome						
Skin infections	Fungal skin infections	Candidiasis	6/21 ⁶¹	28.6	0.6-1.0 ^{22,23}	
	Viral skin infections	Herpes virus lip infection	2/21 ⁶¹	9.5	0.8-1.0 ^{22,23}	
	Other skin infections	Angular cheilitis	2/21 ⁶¹	9.5	0.1-0.3 ^{22,23}	
Vascular disorders	Telangiectasia	Cutaneous telangiectasia	3/32 ⁶²	9.4	-	
Pigmentation disorders	Hyperpigmentation disorders	Café au lait spots	18/21 ⁶²	85.7	12.4 ²¹	
	Hypopigmentation disorders	Vitiligo	14/21 ⁶²	66.7	1.2-1.7 ²¹⁻²³	
Granulomatous disorders		Skin granuloma	5/35 ⁶³	14.3	0.3 ²³	
Other skin disorders		Hyperkeratosis	1/21 ⁶¹	4.8	-	
		Gingivitis	19/21 ⁶¹	90.5	-	
DiGeorge syndrome						
Rash		Rash	5/5 ⁶⁴	100	-	
Comèl-Netherton syndrome						
Dermatitis-like lesions		Eczema ^b	8/9 ³¹	88.9	9.0-27.4 ^{21,26}	
Hair abnormalities	Hair loss disorders	Severe alopecia	9/9 ⁵²	100	0.4-2.5 ²¹⁻²³	
		Alopecia of eyelashes and eyebrows	5/9 ⁵²	55.6	-	
	Other hair abnormalities	Bamboo hair	9/9 ³¹	100	-	
Skin infections	Bacterial skin infections	Recurrent/persistent <i>S. aureus</i> skin infections	9/9 ³¹	100	-	

(Continues)

TABLE 1 (Continued)

Primary immunodeficiency disease					General population	
Main groups of skin disorders		Subgroups of skin disorders	Skin disorders as reported in included articles	Number of reported cases with skin disorder (proportion)	Prevalence of skin disorder (%)	Prevalence of skin disorder (%) [†]
	Erythematous skin lesions		Exfoliative erythroderma ^c	9/9 ⁵²	100	0.1 ²³
	Other skin disorders		Congenital ichthyosis	9/9 ³¹	100	0.1 ^{22,23}
Predominantly antibody deficiencies						
X-linked agammaglobulinemia						
	Dermatitis-like lesions		Dermatitis	4/23 ⁶	17.4	32.4 ²⁴
	Hair abnormalities	Hair loss disorders	Alopecia	1/110 ⁶⁵	0.9	0.4-2.5 ²¹⁻²³
Skin infections	Bacterial skin infections	Abscesses	3/23 ⁶	13.0	1.7 ²²	
		Furunculosis	2/23 ⁶	8.7	1.7 ²²	
		Impetigo	2/23 ⁶	8.7	0.8-1.6 ^{22,23}	
Pigmentation disorders	Hypopigmentation disorders	Vitiligo	0/110 ⁶⁵	0.0	1.2-1.7 ²¹⁻²³	
Rash		Maculopapular rash	3/23 ⁶	13.0	-	
Psoriasis-like lesions		Psoriasis	0/110 ⁶⁵	0.0	1.4-8.0 ²¹⁻²⁶	
Urticaria		Urticaria	2/23 ⁶	8.7	0.5-8.3 ²¹⁻²³	
Other skin disorders		Pyoderma	1/10 ⁷	10.0	1.0 ²²	
		Lichen planus	0/110 ⁶⁵	0.0	0.3-14.1 ^{21,23,25}	
Hypogammaglobulinemia						
	Dermatitis-like lesions		Atopic dermatitis	28/28, ⁶⁶ 0/12, ³⁸ 46/78 ³⁹	0.0-100	2.2-12.4 ²¹⁻²⁴
Common variable immunodeficiency						
Dermatitis-like lesions		Dermatitis	6/28 ⁶	21.4	32.4 ²⁴	
		Atopic dermatitis	9/47 ⁶⁷	19.1	2.2-12.4 ²¹⁻²⁴	
		Eczema	4/15 ⁶⁸	26.7	9.0-27.4 ^{21,26}	
Hair abnormalities	Hair loss disorders	Alopecia areata	1/28, ⁶ 1/47 ⁶⁷	2.1-3.6	0.4-2.5 ²¹⁻²³	
		Alopecia	4/244 ⁶⁵	1.6	0.4-2.5 ²¹⁻²³	
Skin infections	Fungal skin infections	Candida	4/28 ⁶	14.3	0.6-1.0 ^{22,23}	
		Pseudomembranous candidiasis	4/15 ⁶⁸	26.7	-	
	Viral skin infections	Recurrent herpes labialis	1/15 ⁶⁸	6.7	0.8-1.0 ^{22,23}	
		Bacterial skin infections	Recurrent skin abscesses	2/31 ⁶⁹	6.5	1.7 ²²
		Other skin infections	Skin infections	7/47 ⁶⁷	14.9	4.6-43.5 ^{21,25,26}
Ulcers	Oral ulcers	Recurrent aphthosis	5/47 ⁶⁷	10.6	0.7 ²²	
		Oral ulcers	9/15 ⁶⁸	60.0	0.7 ²²	
		Oral aphthae	10/31 ⁶⁹	32.3	0.7 ²²	
Pigmentation disorders	Hypopigmentation disorders	Vitiligo	8/244, ⁶⁵ 2/47 ⁶⁷	3.3-4.3	1.2-1.7 ²¹⁻²³	
Rash		Maculopapular rash	1/28 ⁶	3.6	-	
Psoriasis-like lesions		Psoriasis	2/244, ⁶⁵ 9/47 ⁶⁷	0.8-19.1	1.4-8.0 ²¹⁻²⁶	
Acne-like lesions		Acne	6/47 ⁶⁷	12.8	0.8-13.1 ²²⁻²⁴	

(Continues)

TABLE 1 (Continued)

Primary immunodeficiency disease					General population	
Main groups of skin disorders	Subgroups of skin disorders	Skin disorders as reported in included articles	Number of reported cases with skin disorder (proportion)	Prevalence of skin disorder (%)	Prevalence of skin disorder (%) [†]	
Urticaria		Urticaria ^d	1/28 ⁶	3.6	0.5-8.3 ²¹⁻²³	
Other skin disorders		Pyoderma	2/5 ⁷	40.0	1.0 ²²	
		Lichen planus	1/244 ⁶⁵	0.4	0.3-14.1 ^{21,23,25}	
Selective IgA deficiency						
Dermatitis-like		Dermatitis	5/17 ⁶	29.4	32.4 ²⁴	
		Eczema	2/39 ⁷⁰	5.1	9.0-27.4 ^{21,26}	
		Atopic dermatitis	13/13, ⁶⁶ 4/12, ³⁸ 59/102, ⁶⁷ 12/159, ⁴⁰ 12/23, ⁴² 9/123, ⁴¹ 1/8, ⁷¹ 9/81, ⁴³ 16/347 ⁷²	4.6-100	2.2-12.4 ²¹⁻²⁴	
		Allergic contact dermatitis	11/347 ⁷²	3.2	1.0-8.5 ²¹⁻²⁵	
		Seborrheic dermatitis	4/347 ⁷²	1.2	2.2-11.7 ²¹⁻²⁵	
Hair abnormalities	Hair loss disorders	Alopecia	2/60, ⁶⁵ 1/102, ⁶⁷ 1/8 ⁷¹	1.0-12.5	0.4-2.5 ²¹⁻²³	
		Alopecia areata	1/8, ⁷³ 1/123, ⁴¹ 2/347 ⁷²	0.6-12.5	0.4-2.5 ²¹⁻²³	
		Infection-related alopecia areata	1/8 ⁷³	12.5	0.4-2.5 ²¹⁻²³	
Skin infections	Fungal skin infections	Candida	2/17 ⁶	11.8	0.6-1.0 ^{22,23}	
		Pseudomembranous candidiasis	10/39 ⁷⁰	25.6	-	
	Viral skin infections	Recurrent herpes labialis	10/39, ⁷⁰ 2/123 ⁴¹	1.6-25.6	0.8-1.0 ^{22,23}	
		Herpes simplex	4/347 ⁷²	1.2	0.8-1.0 ^{22,23}	
		Herpes zoster	1/347 ⁷²	0.3	1.4-2.4 ^{22,23}	
	Molluscum contagiosum	2/347 ⁷²	0.9	0.8-0.9 ^{22,23}		
		Bacterial skin infections	Folliculitis	1/17 ⁶	5.9	1.1-6.0 ^{21,23}
			Erysipelas recidivans	1/8 ⁷³	12.5	0.1 ²³
			Chronic recurrent furunculosis	1/8 ⁷³	12.5	1.7 ²²
	Cellulitis	5/347 ⁷²	1.4	0.9-1.5 ^{22,23}		
		Other skin infections	Skin infections	2/102 ⁶⁷	2.0	4.6-43.5 ^{21,25,26}
			Angular stomatitis	1/39 ⁷⁰	2.6	-
			Scabies	3/347 ⁷²	0.9	0.2-1.5 ^{22,23}
		Ulcers	Oral ulcers	Recurrent aphthosis	5/102 ⁶⁷	4.9
Oral ulcers	24/39 ⁷⁰			61.5	0.7 ²²	
Aphthosis recidivans	1/8 ⁷³			12.5	0.7 ²²	
Erythematous skin lesions		Erythroderma	1/8 ⁷³	12.5	0.1 ²³	
		Erythema nodosum	1/123 ⁴¹	0.8	0.2-0.3 ^{22,23}	
Vascular disorders	Vasculitis	Vasculitis	3/7 ⁷	42.9	-	
		Kawasaki disease	1/8 ⁷¹	12.5	-	
	Other vascular disorders	Raynaud syndrome	1/8 ⁷¹	12.5	-	
Pigmentation disorder	Hypopigmentation disorders	Vitiligo	1/7, ⁷ 3/60, ⁶⁵ 3/102, ⁶⁷ 1/8, ⁷³ 7/159, ⁴⁰ 3/123, ⁴¹ 1/81, ⁴³ 2/347 ⁷²	0.6-14.3	1.2-1.7 ²¹⁻²³	
Psoriasis-like disorders		Psoriasis	0/60, ⁶⁵ 2/102, ⁶⁷ 7/159, ⁴⁰ 1/123, ⁴¹ 7/347 ⁷²	0.0-4.4	1.4-8.0 ²¹⁻²⁶	

(Continues)

TABLE 1 (Continued)

Primary immunodeficiency disease					General population
Main groups of skin disorders	Subgroups of skin disorders	Skin disorders as reported in included articles	Number of reported cases with skin disorder (proportion)	Prevalence of skin disorder (%)	Prevalence of skin disorder (%) [†]
Acne-like lesions		Acne	6/102, ⁶⁷ 69/347 ⁷²	5.9-19.9	0.8-13.1 ²²⁻²⁴
Urticaria		Urticaria	5/23 ⁴²	21.7	0.5-8.3 ²¹⁻²³
		Atopic urticaria	4/123 ⁴¹	3.3	0.5-8.3 ²¹⁻²³
		Chronic spontaneous urticaria	17/347 ⁷²	4.9	0.5-8.3 ²¹⁻²³
Other skin disorders		Lichen planus	0/60, ⁶⁵ 2/159 ⁴⁰	0.0-1.3	0.3-14.1 ^{21,23,25}
		Immune thrombocytopenic purpura	1/17 ⁶	5.9	-
		Idiopathic thrombocytopenic purpura	2/123 ⁴¹	1.6	-
		Chronic idiopathic thrombocytopenic purpura	1/81 ⁴³	1.2	-
		Epidermolysis bullosa dystrophica	1/8 ⁷³	12.5	-
		Local skin scleroderma	1/123 ⁴¹	0.8	-
		Dermatitis herpetiformis	1/123 ⁴¹	0.8	0.2-0.3 ^{21,22}
		Ichthyosis and keratoderma of handpalms and footsoles in epileptic patients (Rud syndrome)	1/123 ⁴¹	0.8	-
IgM deficiency					
Dermatitis-like lesions		Atopic dermatitis	14/14, ⁶⁶ 11/53 ³⁸	20.8-100	2.2-12.4 ²¹⁻²⁴
IgG deficiency					
Dermatitis-like lesions		Atopic dermatitis	11/11 ⁶⁶	100	2.2-12.4 ²¹⁻²⁴
Diseases of immune dysregulation					
Autoimmune polyendocrinopathy candidiasis ectodermal dystrophy					
Dermatitis-like lesions		Recurrent and troublesome napkin dermatitis	5/18 ⁷⁴	27.8	-
Hair abnormalities	Hair loss disorders	Alopecia	2/15, ⁷¹ 6/18, ⁷⁴ 20/68, ⁷⁵ 6/35, ⁷⁶ 6/22 ⁷⁷	13.3-33.3	0.4-2.5 ²¹⁻²³
	Hair pigmentation disorders	Poliosis	1/18 ⁷⁴	5.6	-
Skin infections	Fungal skin infections	Oral candidiasis	41/68 ⁷⁵	60.3	-
		Dermal candidiasis	6/68, ⁷⁵ 6/35 ⁷⁶	8.8-17.1	0.6-1.0 ^{22,23}
		Chronic mucocutaneous candidiasis	18/18, ⁷⁴ 30/35 ⁷⁶	85.7-100	0.6-1.0 ^{22,23}
		Mucocutaneous candidiasis	21/22 ⁷⁷	95.5	0.6-1.0 ^{22,23}
		Lifelong genital moniliasis	1/18 ⁷⁴	5.6	-

(Continues)

TABLE 1 (Continued)

Primary immunodeficiency disease					General population
Main groups of skin disorders	Subgroups of skin disorders	Skin disorders as reported in included articles	Number of reported cases with skin disorder (proportion)	Prevalence of skin disorder (%)	Prevalence of skin disorder (%) [†]
	Other skin infections	Angular cheilitis	13/18 ⁷⁴	72.2	0.1-0.3 ^{22,23}
Vascular disorders	Vasculitis	Cutaneous vasculitis	2/68 ⁷⁵	2.9	-
Pigmentation disorders	Hypopigmentation disorders	Vitiligo	1/15, ⁷¹ 2/18, ⁷⁴ 9/68, ⁷⁵ 13/35, ⁷⁶ 6/22 ⁷⁷	6.7-37.1	1.2-1.7 ²¹⁻²³
		Halo naevi	1/18 ⁷⁴	5.6	0.8 ²¹
Nail disorders	Infectious nail disorders	Ungual candidiasis	33/50 ⁷⁵	66.0	-
		Candidal paronychia and/or onychomycosis	13/18 ⁷⁴	72.2	9.4 ²¹
		Nail candidiasis	12/35 ⁷⁶	34.3	-
	Noninfectious nail disorders	Nail dystrophy	26/50, ⁷⁵ 6/35, ⁷⁶ 4/22 ⁷⁷	17.1-52.0	-
Urticaria		Urticarial eruption	23/35 ⁷⁶	8.6	0.5-8.3 ²¹⁻²³
		Urticarial rash	2/22 ⁷⁷	9.1	0.5-8.3 ²¹⁻²³
Other skin disorders		Oral thrush	35/35 ⁷⁶	100	-
Immunodysregulation polyendocrinopathy enteropathy X-linked syndrome					
Dermatitis-like		Atopic dermatitis	7/10 ⁷⁸	70.0	2.2-12.4 ²¹⁻²⁴
		Severe eczema	5/14 ⁴⁴	35.7	9.0-27.4 ^{21,26}
		Mild eczema	4/14 ⁴⁴	28.6	9.0-27.4 ^{21,26}
		Eczema	5/5, ⁷⁹ 2/5 ⁸⁰	40.0-100	9.0-27.4 ^{21,26}
Hair abnormalities	Hair loss disorders	Alopecia ^e	2/14 ⁴⁴	14.3	0.4-2.5 ²¹⁻²³
Skin infections	Fungal skin infections	Chronic mucocutaneous candidiasis	4/5 ⁷⁹	80.0	0.6-1.0 ^{22,23}
	Viral skin infections	HSV skin infection	1/5 ⁷⁹	20.0	0.8-1.0 ^{22,23}
		Herpes zoster	1/5 ⁷⁹	20.0	1.4-2.4 ^{22,23}
	Bacterial skin infections	Staphylococcal superinfection of eczema	1/5 ⁷⁹	20.0	-
	Other skin infections	Cheilitis	4/10 ⁷⁸	40.0	0.1-0.3 ^{22,23}
Erythematous skin lesions		Exfoliative erythroderma	2/10 ⁷⁸	20.0	0.1 ²³
Nail disorders	Infectious nail disorders	Onychodystrophy ^f	2/10 ⁷⁸	20.0	-
		(Peri-)onyxis	1/10 ⁷⁸	10.0	-
Psoriasis-like lesions		Psoriasiform rash	2/10 ⁷⁸	20.0	1.4-8.0 ²¹⁻²⁶
Other skin disorders		Palmar keratoderma	1/10 ⁷⁸	10.0	-
		Inflammatory edema (lips and perioral) or Quincke edema	3/10 ⁷⁸	30.0	-
		Mild xerosis	1/14 ⁴⁴	7.1	-
		Acrodermatitis enteropathica	1/5 ⁷⁹	20.0	-
Adenosine deaminase 2 deficiency					
Dermatitis-like lesions		Eczema	5/9 ⁸¹	55.6	9.0-27.4 ^{21,26}
Ulcers	Oral ulcers	Oral aphthae	3/9 ⁸¹	33.3	0.7 ²²

(Continues)

TABLE 1 (Continued)

Primary immunodeficiency disease					General population
Main groups of skin disorders	Subgroups of skin disorders	Skin disorders as reported in included articles	Number of reported cases with skin disorder (proportion)	Prevalence of skin disorder (%)	Prevalence of skin disorder (%) [†]
	Other ulcers	Livedoid vasculopathy/ulcers	1/9 ⁸¹	11.1	-
Erythematous skin lesions		Erythema nodosum	2/9, ⁸¹ 1/8 ⁸²	12.5-22.2	0.2-0.3 ^{22,23}
Vascular disorders	Vasculitis	Cutaneous vasculitis	3/9 ⁸¹	33.3	-
		Polyarteritis nodosa	1/9 ⁸¹	11.1	-
	Other vascular disorders	Raynaud's phenomenon	1/8 ⁸²	12.5	-
Neoplastic disorder	Other neoplastic disorders	AML chloroma	1/9 ⁸¹	11.1	-
Rash		Rash	3/9 ⁸¹	33.3	-
Urticaria		Urticaria-like rash	1/8 ⁸²	12.5	0.5-8.3 ²¹⁻²³
Other skin disorders		Livedo reticularis	3/9, ⁸¹ 6/8 ⁸²	33.3-75.0	0.1 ²³
		Livedo racemose	1/9 ⁸¹	11.1	0.1 ²³
		Aspecific skin induration	4/9 ⁸¹	44.4	-
<i>Congenital defects of phagocyte number or function</i>					
Leukocyte adhesion defect unspecified					
Skin infections	Bacterial skin infections	Cellulitis	2/6 ⁶	33.3	0.9-1.5 ^{22,23}
		Folliculitis	2/6 ⁶	33.3	1.1-6.0 ^{21,23}
Other skin disorders		Periodontitis	2/6 ⁶	33.3	-
Leukocyte adhesion defect type 1					
Skin infections	Fungal skin infections	Candida infection	8/15 ⁸³	53.3	0.6-1.0 ^{22,23}
	Bacterial skin infections	Skin abscesses	12/15 ⁸³	80.0	1.7 ²²
		Cellulitis	4/15 ⁸³	26.7	0.9-1.5 ^{22,23}
Ulcers	Oral ulcers	Oral ulcers	13/15 ⁸³	86.7	0.7 ²²
Other skin disorders		Gingivitis	9/15 ⁸³	60.0	-
Chronic granulomatous disease					
Dermatitis-like lesions		Dermatitis	nm/429, ⁸⁴ nm/39 ⁸⁵	-	32.4 ²⁴
		Eczema	8/48 ⁴⁵	16.7	9.0-27.4 ^{21,26}
Skin infections	Bacterial skin infections	Abscesses	23/34, ⁶ nm/429, ⁸⁴ nm/39, ⁸⁵ 1/48, ⁴⁵ 156/368, ⁸⁶ 11/49 ³²	2.1-67.6	1.7 ²²
		Folliculitis	5/34 ⁶	14.7	1.1-6.0 ^{21,23}
		Impetigo	3/34, ⁶ 16/95 ³²	8.8-16.8	0.8-1.6 ^{22,23}
		Cellulitis	18/368 ⁸⁶	4.9	0.9-1.5 ^{22,23}
		Furunculosis	nm/429 ⁸⁴	-	-
		Pustular eruption	10/48 ⁴⁵	20.8	-
		Chronic cutaneous infections	1/6 ⁸⁷	16.7	4.6-43.5 ^{21,25,26}
	Other skin infections	Cutaneous/subcutaneous infections	22/48, ⁴⁵ 43/84 ⁸⁸	45.8-51.2	4.6-43.5 ^{21,25,26}
		Skin infection	59/130 ³²	45.4	4.6-43.5 ^{21,25,26}
Ulcers	Oral ulcers	Oral ulceration	7/9, ⁶⁰ 2/6 ⁸⁷	33.3-77.8	0.7 ²²
	Other ulcers	Nose ulcers	5/34 ⁶	14.7	-

(Continues)

TABLE 1 (Continued)

Primary immunodeficiency disease					General population
Main groups of skin disorders	Subgroups of skin disorders	Skin disorders as reported in included articles	Number of reported cases with skin disorder (proportion)	Prevalence of skin disorder (%)	Prevalence of skin disorder (%) [†]
Vascular disorders	Vasculitis	Kawasaki disease	2/48 ⁴⁵	4.2	-
Nail disorders	Infectious nail disorders	Paronychia	2/34 ⁶	5.9	-
Granulomatous disorders		Granuloma	2/11, ⁷¹ nm/429, ⁸⁴ 4/48 ⁴⁵	8.3-18.2	0.3 ²³
Acne-like lesions		Acne	nm/429 ⁸⁴	-	0.8-13.1 ²²⁻²⁴
Urticaria		Urticaria	1/48 ⁴⁵	2.1	0.5-8.3 ²¹⁻²³
Other skin disorders		Discoid lupus erythematosus ⁸	30/340 ⁸⁶	8.8	0.1-0.3 ^{21,22}
		Thrush	11/48 ⁴⁵	22.9	-
Severe congenital neutropenia					
Skin infections	Fungal skin infections	Mucocutaneous candidiasis	5/18 ⁸⁹	27.8	0.6-1.0 ^{22,23}
	Bacterial skin infections	Cutaneous abscesses	10/18 ⁸⁹	55.6	1.7 ²²
	Other skin infections	Cutaneous infections	7/18 ⁸⁹	38.9	4.6-43.5 ^{21,25,26}
Ulcers	Oral ulcers	Oral ulcers	13/18 ⁸⁹	72.2	0.7 ²²
Papillon-Lefèvre syndrome					
Nail disorders	Noninfectious nail disorders	Nail changes (mainly slight thickening of nails)	13/47 ⁹⁰	27.7	3.3-3.4 ^{24,26}
Psoriasis-like lesions		Extensive psoriasiform plaques	3/47 ⁹⁰	6.4	1.4-8.0 ²¹⁻²⁶
Other skin disorders		Punctate hyperkeratosis on palms and soles	8/47 ⁹⁰	17.0	-
		Well-demarcated hyperkeratosis of knees and elbows	23/47 ⁹⁰	48.9	-
		Ichthyosis	2/47 ⁹⁰	4.3	0.1 ^{22,23}
GATA2 deficiency					
Neoplastic disorders	Other neoplastic disorders	Cutaneous melanoma	1/71 ⁹¹	1.4	0.1-1.2 ^{21,23}
Defects in intrinsic and innate immunity					
Chronic mucocutaneous candidiasis					
Skin infections	Other skin infections	Perleche (angular cheilitis)	3/7 ⁶	42.9	0.1-0.3 ^{22,23}
Other skin disorders		Thrush ^h	7/7 ⁶	100	-
Autoinflammatory disorders					
PLCG2 associated antibody deficiency and immune dysregulation					
Ulcers	Other ulcers	Neonatal-onset ulcerative lesions (cold-sensitive regions)	8/36 ⁴⁶	22.2	2.0 ²³
Erythematous skin lesions		Recurrent red papules and patches	1/36 ⁴⁶	2.8	-
Pigmentation disorders	Hypopigmentation disorders	Vitiligo	1/36 ⁴⁶	2.8	1.2-1.7 ²¹⁻²³
Granulomatous disorders		Granulomatous inflammation	4/36 ⁴⁶	11.1	0.3 ²³

(Continues)

TABLE 1 (Continued)

Primary immunodeficiency disease					General population
Main groups of skin disorders	Subgroups of skin disorders	Skin disorders as reported in included articles	Number of reported cases with skin disorder (proportion)	Prevalence of skin disorder (%)	Prevalence of skin disorder (%) [†]
Urticaria		Cold urticaria	36/36 ⁴⁶	100	0.5-8.3 ²¹⁻²³
Muckle-Wells syndrome					
Ulcers	Oral ulcers	Oral ulcers	7/29 ⁹²	24.1	0.7 ²²
Pigmentation disorders	Hyperpigmentation disorders	Hyperpigmented, sclerotic and hypertrichotic plaques	6/6 ⁹³	100	-
Rash		Skin rash	15/15 ⁹²	100	-
Urticaria		Attacks of recurrent urticaria	2/6 ⁹³	33.3	0.5-8.3 ²¹⁻²³
		Urticaria	8/8 ⁹⁴	100	0.5-8.3 ²¹⁻²³
		Cold-induced urticaria	14/29 ⁹²	48.3	0.5-8.3 ²¹⁻²³
Other skin disorders		Weals caused by cold	16/16 ⁹⁵	100	-
Neonatal-onset multisystem inflammatory disease					
Urticaria		Urticaria	8/8 ⁹⁴	100	0.5-8.3 ²¹⁻²³
Complement deficiencies					
C2 deficiency					
Urticaria		Chronic urticaria	2/47 ⁹⁶	4.3	0.5-8.3 ²¹⁻²³
Other skin disorders		Subacute cutaneous lupus erythematosus	2/47 ⁹⁶	4.3	0.3 ²²
		Dermatitis herpetiformis	1/47 ⁹⁶	2.1	0.2-0.3 ^{21,22}

Abbreviation: nm, not mentioned.

[†]Data on the prevalence of skin disorders in the general population were based on a birth cohort in Finland (n = 1932, age 45-47 years) and a Dermatology outpatient clinic in Turkey (n = 11 040, age 1-99 years).^{21,22} In addition, a nationwide study of Furue et al²³ reported the prevalence of cutaneous disorders in 67 448 Japanese patients of all ages. In the study of Verhoeven et al,²⁴ the skin disease prevalence per 1000 patient-years in family practices in the Netherlands was converted to a point prevalence in the general population (n = 501, age 18-97 years). Last, two studies from the United States of America and the United Kingdom performed in 1978 and 1976 showed the prevalence of skin disorders in community studies in, respectively, 20 749 (age 1-74 years) and 614 (age 15-74 years) patients.^{25,26}

^{a-h}See Figure S1.

3.3 | *Staphylococcus aureus*-associated skin disorders in primary immunodeficiency diseases

Skin disorders associated with *S. aureus* in PIDs were reported in six articles (Table S1). In HIES unspecified, 4/7 patients with a papulopustular eruption had a positive *S. aureus* culture.²⁷ *S. aureus* was also found positive in patients with AD-HIES and a papulopustular rash (2/5), eczematous dermatitis (20/20), cold abscesses (20/20), or wounds (3/4).^{28,29} Renner et al³⁰ described that skin abscesses were frequently due to *S. aureus* infections in AR-HIES. In Comèl-Netherton syndrome, 8/9 described patients showed recurrent or persistent *S. aureus* skin infections once skin lesions had developed.³¹ Lastly, *S. aureus* was isolated in 1/4 patients with CGD and suppurative dermatitis.³²

3.4 | Primary immunodeficiency diseases associated with atopy

The prevalence of at least two atopic symptoms (ie, eczema, food allergy, asthma, and/or rhinoconjunctivitis) in PIDs was described in 17

articles (Table S1). Cohen et al³³ found no atopy in patients with AT. In HIES unspecified, AD-HIES and AR-HIES, all of the atopic symptoms, if reported, were present in about half of the patients.^{28,29,34-37} Renner et al³¹ described the presence of atopy in most Comèl-Netherton patients. The number of patients with eczema, food allergy, asthma, and rhinoconjunctivitis was presented in two studies with hypogammaglobulinemia patients.^{38,39} Eight studies reported the prevalence of atopy in a total of 398 patients with SIgAD, in which 11.6% (46/398) patients had eczema, 3.4% (9/263) had food allergy, 37.1% (43/116) had asthma, and 20.0% (55/275) had rhinoconjunctivitis.^{38,40-43} In IgM-deficient patients, eczema, asthma, and rhinoconjunctivitis were prevalent symptoms.³⁸ Immunodysregulation polyendocrinopathy enteropathy X-linked syndrome (IPEX) patients were mainly positive for eczema.⁴⁴ Eight of 48 CGD patients had eczema and one out of 18 had rhinoconjunctivitis.⁴⁵ Finally, Aderibigbe et al⁴⁶ have shown that one to three out of eight patients with phospholipase C gamma 2 (PLCG2) gene associated antibody deficiency and immune dysregulation (PLAID) had eczema, food allergy, asthma, and/or rhinoconjunctivitis.




FIGURE 2 Differential diagnosis of primary immunodeficiency diseases per skin disorder. nm, not mentioned[†]. [†]The exact prevalence of acne-like lesions in chronic granulomatous disease was not reported in the included article. The prevalence of skin disorders per primary immunodeficiency disease was based on the reported proportion of patients with the skin disorders in the total group of patients with the primary immunodeficiency disease. In case of multiple studies reporting a skin disorder in a primary immunodeficiency disease, the general prevalence was calculated by dividing the total number of affected patients (skin disorder) by the total number of patients with the primary immunodeficiency disease. nm, not mentioned.

4 | DISCUSSION

This review demonstrates that skin disorders are common symptoms in both children and adult patients with PIDs based on data from 67 systematically selected studies. Only a few PIDs related to *S. aureus*-associated skin disorders or atopy were reported in mainly single studies.

This is the first review using a systematic approach without limitations on skin disorders or PIDs. Therefore, we managed to obtain a complete spectrum of skin disorders in PIDs. A recent study of Ettinger et al⁴⁷ that focused on PIDs and the respective gene defects included an overview of PIDs per skin disorder in a nonsystematic approach. Although some PIDs are characterized by skin disorders, such as telangiectasia in AT and granuloma in CGD, the novelty of this review is showing an overview of all skin disorders in PIDs including skin disorders of which an association with a PID was not yet known.

Furthermore, we succeeded in composing an overview of PIDs per skin disorder that could serve as a valuable support tool for PID awareness in clinical practice and for registries. In the Netherlands, the diagnostic delay in PIDs (ie, time period between the date of onset of first symptoms and the date of diagnosis) ranges from 0 to 14.5 years and is dominated by the defects in innate immunity (14.5 years), HIES (10.5 years), and hypogammaglobulinemia (10.0 years).⁴⁸ Moin et al⁶ and Berron-Ruiz et al⁷ reported that in, respectively, 31.8% and 78.9% of the PIDs, the cutaneous alterations preceded and were the basis for the clinical immunological diagnosis. Increased attention for these cutaneous manifestations as signal function of PIDs in combination with presence of the current warning signs for the suspicion of PIDs might improve earlier diagnosis of PIDs. These warning signs include (a) four or more new ear infections within 1 year; (b) two or more serious sinus infections within 1 year; (c) two or more months on antibiotics with little effect; (d) two or more pneumonias within 1 year; (e) failure of an infant to gain weight or grow normally; (f) recurrent deep skin or organ abscesses; (g) persistent thrush in mouth or fungal infection on skin; (h) need for intravenous antibiotics to clear infections; (i) two or more deep-seated infections including septicemia; and (j) a family history of PID.⁴⁹ In addition, narrowing the number of eligible PIDs through clustering of skin disorders could further reduce the diagnostic delay in PIDs. Using the multistage diagnostic protocol of de Vries⁴ or the phenotypic approach for PID classification and diagnosis by Bousfiha et al,⁵⁰ the diagnosis of suspected PIDs or PID classes based on clinical symptoms could be confirmed with laboratory tests. For example, a first diagnostic step in case of a supposed antibody deficiency or neutropenia could be blood count and differentiation, IgG, IgA, IgM, and IgE. In case of a possible combined immunodeficiency disease, these tests should be supplemented by lymphocyte subpopulations.

Our review has some limitations. First of all, exclusion of case reports describing fewer than five cases in our analysis might have resulted in loss of information about skin disorders in rare PIDs. However, the quality of case reports is highly variable, potential

publication bias plays a role, and thus, exclusion of these case reports might have improved the reliability of this review. Furthermore, through the addition of selected PIDs and skin disorders to our electronic search we could have caused a selection bias. Although we used a cross-reference check, we cannot exclude that we might have missed some articles. Thirdly, demonstrating that specific skin disorders are characteristic for PIDs was not possible. We could only compare the presence of a number of skin disorders in patients with a PID with the prevalence of skin disorders in the general population based on six studies varying in publication year and age of the studied population.²¹⁻²⁶ However, most of these skin disorders were more prevalent in patients with a PID compared with the general population. Lastly, the reliability of the description of skin disorders might be questioned since in only 27 of the 67 included articles the Department of Dermatology was involved. Probably, the described skin disorders were not all diagnosed by a dermatologist, but by an immunologist or pediatrician. Moreover, the majority of studies did not use skin biopsy to confirm the diagnosis of the cutaneous manifestations histopathologically. In severe combined immunodeficiency (SCID) patients, it was shown by Denianke et al⁵¹ that clinically comparable skin lesions could demonstrate different histopathological images, possibly due to an altered immune system. Subsequently, the reported clinical diagnosis of skin disorders reported in articles included in this review might not correlate with the corresponding histopathological diagnosis as well.

Because most PIDs are rare, reliable prevalence of skin disorders in PIDs can only be obtained by reporting skin disorders on an international basis. The international PID database of The European Society for Immunodeficiencies (ESID) registers, among others, data on warning signs of PIDs. These warning signs give only attention to infectious skin disorders. Noninfectious cutaneous symptoms are not included. Based on data of this review, we suggest to start to collect more detailed data on all skin disorders in the ESID registry.

Future research is needed to validate these data and support an association between specific cutaneous symptoms and PIDs. Given the low number of articles reporting *S. aureus*-associated skin disorders and atopy in PIDs, more data have to be collected to further improve earlier recognition of PIDs. In addition, data on *S. aureus* associated skin disorders might provide new treatment options for skin disorders, such as targeted therapy directed against *S. aureus*.

5 | CONCLUSION

This review with a systematic approach shows that skin disorders are a prominent feature in PIDs. Earlier diagnosis of PIDs can be facilitated by recognition of specific skin conditions as signal function of PIDs in combination with the current warning signs for PIDs or by recognizing PID specific clusters of skin conditions. We provide a support tool to use in clinical practice that should raise awareness of PIDs based on the presenting skin manifestations. Limited data are available on *S. aureus*-associated skin disorders and atopy in PIDs.

ACKNOWLEDGMENTS

We thank Wichor M. Bramer (BSc), biomedical information specialist, who assisted us in conducting the systematic digital search.

CONFLICTS OF INTEREST

JdW, RB, JvV, VD, and SP have no conflict of interest to declare.

ORCID

Jill de Wit  <https://orcid.org/0000-0001-5080-1453>

REFERENCES

- Bousfiha AA, Jeddane L, Ailal F, et al. Primary immunodeficiency diseases worldwide: more common than generally thought. *J Clin Immunol*. 2013;33:1-7.
- Samarghitean C, Ortutay C, Vihinen M. Systematic classification of primary immunodeficiencies based on clinical, pathological, and laboratory parameters. *J Immunol*. 2009;183:7569-7575.
- Notarangelo L, Casanova JL, Fischer A, et al. Primary immunodeficiency diseases: an update. *J Allergy Clin Immunol*. 2004;114:677-687.
- de Vries E, European Society for Immunodeficiencies m. Patient-centred screening for primary immunodeficiency, a multi-stage diagnostic protocol designed for non-immunologists: 2011 update. *Clin Exp Immunol*. 2012;167:108-119.
- Al-Herz W, Nanda A. Skin manifestations in primary immunodeficient children. *Pediatr Dermatol*. 2011;28:494-501.
- Moin A, Farhoudi A, Moin M, Pourpak Z, Bazargan N. Cutaneous manifestations of primary immunodeficiency diseases in children. *Iran J Allergy Asthma Immunol*. 2006;5:121-126.
- Berron-Ruiz A, Berron-Perez R, Ruiz-Maldonado R. Cutaneous markers of primary immunodeficiency diseases in children. *Pediatr Dermatol*. 2000;17:91-96.
- Sillevis Smitt JH, Kuijpers TW. Cutaneous manifestations of primary immunodeficiency. *Curr Opin Pediatr*. 2013;25:492-497.
- Johnston SL. Clinical immunology review series: an approach to the patient with recurrent superficial abscesses. *Clin Exp Immunol*. 2008;152:397-405.
- Rosenzweig SD, Holland SM. Phagocyte immunodeficiencies and their infections. *J Allergy Clin Immunol*. 2004;113:620-626.
- Slatter MA, Gennery AR. Clinical immunology review series: an approach to the patient with recurrent infections in childhood. *Clin Exp Immunol*. 2008;152:389-396.
- Pichard DC, Freeman AF, Cowen EW. Primary immunodeficiency update: part I. Syndromes associated with eczematous dermatitis. *J Am Acad Dermatol*. 2015;73:355-364.
- Spergel JM, Paller AS. Atopic dermatitis and the atopic march. *J Allergy Clin Immunol*. 2003;112(6 suppl):S118-S127.
- European Society for Immunodeficiencies Clinical Working Party. 10 Warnings Signs of PID - General. <https://esid.org/Working-Parties/Clinical-Working-Party/Resources/10-Warning-Signs-of-PID-General>. Accessed December 3, 2018.
- Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Int J Surg*. 2010;8:336-341.
- Picard C, Bobby Gaspar H, Al-Herz W, et al. International Union of Immunological Societies: 2017 Primary immunodeficiency diseases committee report on inborn errors of immunity. *J Clin Immunol*. 2018;38:96-128.
- Pichard DC, Freeman AF, Cowen EW. Primary immunodeficiency update: part II. Syndromes associated with mucocutaneous candidiasis and noninfectious cutaneous manifestations. *J Am Acad Dermatol*. 2015;73:367-381.
- Gniadecki R. In: Griffiths CEM, Barker J, Bleiker TO, Chalmers RJG, Creamer D, eds. *Rook's Textbook of Dermatology, Chapter 148: The Skin and Disorders of the Haematopoietic and Immune Systems*. 9th ed. Hoboken, New Jersey, USA: John Wiley & Sons Inc; 2016.
- Schaffer JV, Paller AS, In: Callen JP, Cerroni L, Heymann WR, Hruza GJ, Mancini AJ, Patterson JW, Röcken M, Schwarz T, eds. *Bologna Dermatology, Chapter 60: Primary Immunodeficiencies*. 3rd ed. Amsterdam, The Netherlands: Elsevier Limited; 2012.
- Huizing M, Anikster Y, Gahl WA. Hermansky-Pudlak syndrome and related disorders of organelle formation. *Traffic*. 2000;1:823-835.
- Sinikumpu SP, Huilaja L, Jokelainen J, et al. High prevalence of skin diseases and need for treatment in a middle-aged population. A Northern Finland Birth Cohort 1966 study. *PLoS One*. 2014;9:e99533.
- Bilgili ME, Yildiz H, Sarici G. Prevalence of skin diseases in a dermatology outpatient clinic in Turkey. A cross-sectional, retrospective study. *J Dermatol Case Rep*. 2013;7:108-112.
- Furue M, Yamazaki S, Jimbow K, et al. Prevalence of dermatological disorders in Japan: a nationwide, cross-sectional, seasonal, multicenter, hospital-based study. *J Dermatol*. 2011;38:310-320.
- Verhoeven EW, Kraaijaat FW, van Weel C, et al. Skin diseases in family medicine: prevalence and health care use. *Ann Fam Med*. 2008;6:349-354.
- Johnson M-LT, Roberts J. Skin conditions and related need for medical care among persons 1-74 years, United States, 1971-1974. 1978.
- Rea JN, Newhouse ML, Halil T. Skin disease in Lambeth. A community study of prevalence and use of medical care. *Br J Prev Soc Med*. 1976;30:107-114.
- Chamlin SL, McCalmont TH, Cunningham BB, et al. Cutaneous manifestations of hyper-IgE syndrome in infants and children. *J Pediatr*. 2002;141:572-575.
- Olaiwan A, Chandesris MO, Fraitag S, et al. Cutaneous findings in sporadic and familial autosomal dominant hyper-IgE syndrome: a retrospective, single-center study of 21 patients diagnosed using molecular analysis. *J Am Acad Dermatol*. 2011;65:1167-1172.
- Wu J, Chen J, Tian ZQ, et al. Clinical manifestations and genetic analysis of 17 patients with autosomal dominant hyper-IgE syndrome in Mainland China: new reports and a literature review. *J Clin Immunol*. 2017;37:166-179.
- Renner ED, Puck JM, Holland SM, et al. Autosomal recessive hyper-immunoglobulin E syndrome: a distinct disease entity. *J Pediatr*. 2004;144:93-99.
- Renner ED, Hartl D, Rylaarsdam S, et al. Comel-Netherton syndrome defined as primary immunodeficiency. *J Allergy Clin Immunol*. 2009;124:536-543.
- Zhou Q, Hui X, Ying W, et al. A cohort of 169 chronic granulomatous disease patients exposed to BCG vaccination: a Retrospective Study from a single center in Shanghai, China (2004-2017). *J Clin Immunol*. 2018;38:260-272.
- Cohen LE, Tanner DJ, Schaefer HG, Levis WR. Common and uncommon cutaneous findings in patients with ataxia-telangiectasia. *J Am Acad Dermatol*. 1984;10:431-438.
- Eberting CL, Davis J, Puck JM, Holland SM, Turner ML. Dermatitis and the newborn rash of hyper-IgE syndrome. *Arch Dermatol*. 2004;140:1119-1125.
- Gernez Y, Freeman AF, Holland SM, et al. Autosomal dominant hyper-IgE syndrome in the USIDNET registry. *J Allergy Clin Immunol Pract*. 2018;6:996-1001.
- Chu EY, Freeman AF, Jing H, et al. Cutaneous manifestations of DOCK8 deficiency syndrome. *Arch Dermatol*. 2012;148:79-84.

37. Broides A, Mandola AB, Levy J, et al. The clinical and laboratory spectrum of dedicator of cytokinesis 8 immunodeficiency syndrome in patients with a unique mutation. *Immunol Res*. 2017;65:651-657.
38. Altun D, Akpınar M, Haskoğlu ZS, et al. Immunoglobulin isotype deficiency together with allergic diseases. *Asim Allerji Immunoloji*. 2016;14:164-169.
39. Szczawińska-Popłonyk A, Komasińska P, Bręborowicz A. IgA deficiency: a risk factor for food allergy-related atopic dermatitis in infants and young children. *Postepy Dermatol Alergol*. 2016;33:369-374.
40. Koskinen S. Long-term follow-up of health in blood donors with primary selective IgA deficiency. *J Clin Immunol*. 1996;16:165-170.
41. Patrizi A, Ricci G, Cassoli C, Specchia F, Neri I, Masi M. Dermatologic diseases associated with IgA deficiency. *G Ital Dermatol Venerol*. 1992;127(7-8):325-329.
42. Aghamohammadi A, Cheraghi T, Gharagozlou M, et al. IgA deficiency: correlation between clinical and immunological phenotypes. *J Clin Immunol*. 2009;29:130-136.
43. Erkoçoğlu M, Metin A, Kaya A, et al. Allergic and autoimmune disorders in families with selective IgA deficiency. *Turk J Med Sci*. 2017;47:592-598.
44. Gambineri E, Perroni L, Passerini L, et al. Clinical and molecular profile of a new series of patients with immune dysregulation, polyendocrinopathy, enteropathy, X-linked syndrome: inconsistent correlation between forkhead box protein 3 expression and disease severity. *J Allergy Clin Immunol*. 2008;122:1105-1112.
45. Wu J, Wang WF, Zhang YD, Chen TX. Clinical features and genetic analysis of 48 patients with chronic granulomatous disease in a single center study from Shanghai, China (2005-2015): new studies and a literature review. *J Immunol Res*. 2017;2017:8745254.
46. Aderibigbe OM, Priel DL, Lee CC, et al. Distinct cutaneous manifestations and cold-induced leukocyte activation associated with PLCG2 mutations. *JAMA Dermatol*. 2015;151:627-634.
47. Ettinger M, Schreml J, Wirsching K, Berneburg M, Schreml S. Skin signs of primary immunodeficiencies - how to find the genes to check. *Br J Dermatol*. 2018;178:335-349.
48. Jonkman-Berk BM, van den Berg JM, Ten Berge IJ, et al. Primary immunodeficiencies in the Netherlands: national patient data demonstrate the increased risk of malignancy. *Clin Immunol*. 2015;156:154-162.
49. Jeffrey Modell Foundation. 10 Warning Signs of Primary Immunodeficiency. <http://downloads.info4pi.org/pdfs/10-Warning-Signs-Genetic-Text-2-.pdf>. Accessed December 3, 2018.
50. Bousfiha AA, Jeddane L, Ailal F, et al. A phenotypic approach for IUIS PID classification and diagnosis: guidelines for clinicians at the bedside. *J Clin Immunol*. 2013;33:1078-1087.
51. Denianke KS, Frieden IJ, Cowan MJ, Williams ML, McCalmont TH. Cutaneous manifestations of maternal engraftment in patients with severe combined immunodeficiency: a clinicopathologic study. *Bone Marrow Transplant*. 2001;28:227-233.
52. Pruszkowski A, Bodemer C, Fraïtag S, Teillac-Hamel D, Amoric JC, de Prost Y. Neonatal and infantile erythrodermas: a retrospective study of 51 patients. *Arch Dermatol*. 2000;136:875-880.
53. Reed WB, Epstein WL, Boder E, Sedgwick R. Cutaneous manifestations of ataxia-telangiectasia. *JAMA*. 1966;195:746-753.
54. Greenberger S, Berkun Y, Ben-Zeev B, Levi YB, Barzilai A, Nissenkorn A. Dermatologic manifestations of ataxia-telangiectasia syndrome. *J Am Acad Dermatol*. 2013;68:932-936.
55. Paller AS, Massey RB, Curtis MA, et al. Cutaneous granulomatous lesions in patients with ataxia-telangiectasia. *J Pediatr*. 1991;119:917-922.
56. Sullivan KE, Mullen CA, Blaese RM, Winkelstein JA. A multiinstitutional survey of the Wiskott-Aldrich syndrome. *J Pediatr*. 1994;125(6 Pt 1):876-885.
57. Dupuis-Girod S, Medioni J, Haddad E, et al. Autoimmunity in Wiskott-Aldrich syndrome: risk factors, clinical features, and outcome in a single-center cohort of 55 patients. *Pediatrics*. 2003;111:622-7.
58. Aghamohammadi A, Moghaddam ZG, Abolhassani H, et al. Investigation of underlying primary immunodeficiencies in patients with severe atopic dermatitis. *Allergol Immunopathol (Madr)*. 2014;42:336-341.
59. Grimbacher B, Holland SM, Gallin JI, et al. Hyper-IgE syndrome with recurrent infections—an autosomal dominant multisystem disorder. *N Engl J Med*. 1999;340:692-702.
60. Charon JA, Mergenhagen SE, Gallin JI. Gingivitis and oral ulceration in patients with neutrophil dysfunction. *J Oral Pathol*. 1985;14:150-155.
61. Gregorek H, Olczak-Kowalczyk D, Dembowska-Baginska B, et al. Oral findings in patients with Nijmegen breakage syndrome: a preliminary study. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2009;108:39-45.
62. Hiel JA, Weemaes CM, van Engelen BG, et al. Nijmegen breakage syndrome in a Dutch patient not resulting from a defect in NBS1. *J Med Genet*. 2001;38:19.
63. Deripapa E, Balashov D, Rodina Y, et al. Prospective study of a cohort of Russian Nijmegen breakage syndrome patients demonstrating predictive value of low kappa-deleting recombination excision circle (KREC) numbers and beneficial effect of hematopoietic stem cell transplantation (HSCT). *Front Immunol*. 2017;8:807.
64. Markert ML, Alexieff MJ, Li J, et al. Complete DiGeorge syndrome: development of rash, lymphadenopathy, and oligoclonal T cells in 5 cases. *J Allergy Clin Immunol*. 2004;113:734-741.
65. Azizi G, Tavakol M, Rafiemanesh H, et al. Autoimmunity in a cohort of 471 patients with primary antibody deficiencies. *Expert Rev Clin Immunol*. 2017;13:1099-1106.
66. Celiksoy MH, Topal E, Sancak R, Catal F, Sogut A. Relationship between hypogammaglobulinemia and severity of atopic dermatitis. *Ann Allergy Asthma Immunol*. 2014;113:467-469.
67. Gualdi G, Lougaris V, Baronio M, et al. Burden of skin disease in selective IgA deficiency and common variable immunodeficiency. *J Invest Allergol Clin Immunol*. 2015;25:369-371.
68. Porter SR, Scully C. Orofacial manifestations in primary immunodeficiencies: common variable immunodeficiencies. *J Oral Pathol Med*. 1993;22:157-158.
69. Muşabak UH, Demirel F, Yeşilliik S, et al. Adults with common variable immunodeficiency: a single-center experience. *Turk J Med Sci*. 2017;47:1-12.
70. Porter SR, Scully C. Orofacial manifestations in primary immunodeficiencies involving IgA deficiency. *J Oral Pathol Med*. 1993;22:117-119.
71. Blazina Š, Markelj G, Jeverica AK, et al. Autoimmune and inflammatory manifestations in 247 patients with primary immunodeficiency—a report from the Slovenian National registry. *J Clin Immunol*. 2016;36:764-773.
72. Magen E, Masalha A, Waitman DA, et al. Prevalence of dermatologic diseases among patients with selective immunoglobulin A deficiency. *Allergy Asthma Proc*. 2017;38:70-77.
73. Goring HD. [IgA deficiency in the dermatologic clinic. Frequency, clinical relevance, diagnosis and therapy]. *Hautarzt*. 1981;32:505-511.
74. Collins SM, Dominguez M, Ilmarinen T, Costigan C, Irvine AD. Dermatological manifestations of autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy syndrome. *Br J Dermatol*. 2006;154:1088-1093.
75. Ahonen P, Myllarniemi S, Sipila I, Perheentupa J. Clinical variation of autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED) in a series of 68 patients. *N Engl J Med*. 1990;322:1829-1836.
76. Ferre EM, Rose SR, Rosenzweig SD, et al. Redefined clinical features and diagnostic criteria in autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy. 2016.

77. Zaidi G, Bhatia V, Sahoo SK, et al. Autoimmune polyendocrine syndrome type 1 in an Indian cohort: a longitudinal study. *Endocr Connect*. 2017;6:289-296.
78. Halabi-Tawil M, Ruemmele FM, Fraitag S, et al. Cutaneous manifestations of immune dysregulation, polyendocrinopathy, enteropathy, X-linked (IPEX) syndrome. *Br J Dermatol*. 2009;160:645-651.
79. Uzel G, Sampaio EP, Lawrence MG, et al. Dominant gain-of-function STAT1 mutations in FOXP3 wild-type immune dysregulation-polyendocrinopathy-enteropathy-X-linked-like syndrome. *J Allergy Clin Immunol*. 2013;131:1611-1623.
80. Barış HE, Kıyıkım A, Nain E, Özen AO, Karakoç-Aydiner E, Barış S. The plethora, clinical manifestations and treatment options of autoimmunity in patients with primary immunodeficiency. *Turk Pediatr Ars*. 2016;51:186-192.
81. Van Montfrans JM, Hartman EAR, Braun KPJ, et al. Phenotypic variability in patients with ADA2 deficiency due to identical homozygous R169Q mutations. *Rheumatology*. 2016;55:902-910.
82. Sahin S, Adrovic A, Barut K, et al. Clinical, imaging and genotypical features of three deceased and five surviving cases with ADA2 deficiency. *Rheumatol Int*. 2018;38:129-136.
83. Movahedi M, Entezari N, Pourpak Z, et al. Clinical and laboratory findings in Iranian patients with leukocyte adhesion deficiency (study of 15 cases). *J Clin Immunol*. 2007;27:302-307.
84. van den Berg JM, van Koppen E, Ahlin A, et al. Chronic granulomatous disease: the European experience. *PLoS One*. 2009;4:e5234.
85. Liese J, Kloos S, Jendrossek V, et al. Long-term follow-up and outcome of 39 patients with chronic granulomatous disease. *J Pediatr*. 2000;137:687-693.
86. Winkelstein JA, Marino MC, Johnston RB Jr, et al. Chronic granulomatous disease. Report on a national registry of 368 patients. *Medicine (Baltimore)*. 2000;79:155-169.
87. Cohen MS, Leong PA, Simpson DM. Phagocytic cells in periodontal defense. Periodontal status of patients with chronic granulomatous disease of childhood. *J Periodontol*. 1985;56:611-617.
88. Wolach B, Gavrieli R, de Boer M, et al. Chronic granulomatous disease: clinical, functional, molecular, and genetic studies. The Israeli experience with 84 patients. *Am J Hematol*. 2017;92:28-36.
89. Rezaei N, Moin M, Pourpak Z, et al. The clinical, immunohematological, and molecular study of Iranian patients with severe congenital neutropenia. *J Clin Immunol*. 2007;27:525-533.
90. Ullbro C, Crossner CG, Nederfors T, Alfadley A, Thestrup-Pedersen K. Dermatologic and oral findings in a cohort of 47 patients with Papillon-Lefevre syndrome. *J Am Acad Dermatol*. 2003;48:345-351.
91. Nguyen J, Alexander T, Jiang H, et al. Melanoma in patients with GATA2 deficiency. *Pigment Cell Melanoma Res*. 2018;31:337-340.
92. Sobolewska B, Angermair E, Deuter C, Doycheva D, Kuemmerle-Deschner J, Zierhut M. NLRP3 A439V Mutation in a Large Family with Cryopyrin-associated Periodic Syndrome: description of Ophthalmologic Symptoms in Correlation with Other Organ Symptoms. Erratum appears in *J Rheumatol*. 2016;43:1451; PMID: 27371654. *J Rheumatol*. 2016;43:1101-1106.
93. El-Darouti MA, Marzouk SA, Abdel-Halim MR. Muckle-Wells syndrome: report of six cases with hyperpigmented sclerodermoid skin lesions. *Int J Dermatol*. 2006;45:239-244.
94. Mehr S, Allen R, Boros C, et al. Cryopyrin-associated periodic syndrome in Australian children and adults: epidemiological, clinical and treatment characteristics. *J Paediatr Child Health*. 2016;52:889-895.
95. Haas N, Kuster W, Zuberbier T, Henz BM. Muckle-Wells syndrome: clinical and histological skin findings compatible with cold air urticaria in a large kindred. *Br J Dermatol*. 2004;151:99-104.
96. Lipsker DM, Schreckenberg-Gilliot C, Uring-Lambert B, et al. Lupus erythematosus associated with genetically determined deficiency of the second component of the complement. *Arch Dermatol*. 2000;136:1508-1514.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: de Wit J, Brada RJK, van Veldhuizen J, Dalm VASH, Pasmans SGMA. Skin disorders are prominent features in primary immunodeficiency diseases: A systematic overview of current data. *Allergy*. 2019;74:464-482. <https://doi.org/10.1111/all.13681>

















SUPPORTING INFORMATION

Table E1

Electronic search

Database	Search
Embase	((('immune deficiency'/exp OR (((immun* OR antibod*) NEAR/3 (deficien* OR deficit* OR incompeten*)) OR immunodeficien* OR immunocomprom*):ab,ti) AND ('congenital disorder'/de OR 'genetic and familial disorders'/de OR 'familial disease'/de OR 'genetic disorder'/de OR 'immune deficiency'/exp/dm_cn OR (((primar* OR congenital* OR hereditar* OR genetic* OR familial*) NEAR/6 (immunodeficien*)) OR ((primar* OR congenital* OR hereditar* OR genetic* OR familial*) NEAR/6 (immun* OR antibod*) NEAR/3 (deficien* OR deficit* OR incompeten*))):ab,ti)) OR ('severe combined immunodeficiency'/exp OR 'ataxia telangiectasia'/de OR 'Chediak Higashi syndrome'/de OR 'common variable immunodeficiency'/de OR 'complement deficiency'/de OR 'DiGeorge syndrome'/de OR 'hyper IgE syndrome'/de OR 'hyper IgM syndrome'/de OR 'immunoglobulin A deficiency'/de OR 'immunoglobulin M deficiency'/de OR 'leukocyte adhesion deficiency'/de OR 'X linked agammaglobulinemia'/de OR 'chronic granulomatous disease'/de OR ('mucocutaneous candidiasis'/de AND 'chronic disease'/de) OR 'Griscelli syndrome'/de OR 'ocular albinism'/de OR 'Wiskott Aldrich syndrome'/de OR 'Netherton disease'/de OR ((severe* NEAR/3 combin* NEAR/3 immun*) OR (ataxi* NEAR/3 (telangiect* OR teleangiect*)) OR (Chediak NEAR/3 (Higashi OR anomal* OR syndrom*)) OR (common NEAR/3 variable NEAR/3 immun*) OR (complement* NEAR/3 deficienc*) OR (Congenital* NEAR/3 CD4 NEAR/3 deficien*) OR (Cyclic* NEAR/3 haematopoie*) OR DiGeorge OR (hyper NEXT/1 (IgE OR igm)) OR 'job syndrome' OR (hyperimmunoglobul* NEXT/1 (E or M)) OR (Idiopathic NEAR/3 CD4 NEAR/3 lymphocytopenia) OR ((immunoglobulin-A OR iga OR immunoglobulin-M OR igm OR irak OR IL-1-receptor-associated-kinase OR leukocyte-adhesion OR tap OR transporter-associated-with-antigen*) NEAR/3 deficien*) OR whim OR (Wart* NEAR/6 Hypogammaglobulin* NEAR/6 Immun NEAR/6 Myelokath*) OR ((linked OR bruton) NEXT/1 agammaglobulin*) OR (chronic NEAR/3 granulomat*) OR (chronic* NEAR/3 mucocutan* NEAR/3

	candidias*) OR Griscelli OR (ocular NEAR/3 albin*) OR (Hermansky NEAR/3 Pudlak) OR (Wiskott NEAR/3 Aldrich) OR Netherton OR IPEX-syndrome OR APECED OR PLAID OR ((DOCK8 OR Phosphoglucomutase-3 OR PGM3 OR Mammalian-sterile OR MST1 OR GATA2 OR CARD9 OR IL-17RA OR IL-17F OR ACT1) NEAR/3 deficien*) OR (mutation NEAR/3 (STAT1 OR Dectin-1))):ab,ti)) AND ('skin disease'/exp OR 'small vessel vasculitis'/de OR 'skin abscess'/de OR 'atopy'/de OR 'Staphylococcus aureus'/de OR 'Staphylococcus infection'/de OR (((skin OR cutaneous OR dermal OR mucocutan* OR pigment*) NEAR/3 (disease* OR disorder* OR manifestat* OR infect* OR candid* OR abscess* OR granulom* OR lymphoma OR carcinoma* OR tumor* OR cancer* OR neoplas*)) OR eczem* OR verruca* OR Wart* OR dermatit* OR (lupus NEAR/3 erythemat*) OR (small-vessel* NEAR/3 vasculit*) OR Ulcer* OR decubit* OR pyoderm* OR atop* OR ((Staphylococcus OR s) NEXT/1 aureus) OR ((Basal OR squamous) NEXT/1 cell* NEAR/6 carcinom*) OR hyperpigment* OR hypopigment* OR Café-au-lait OR albinis* OR vitiligo OR depigment* OR erythroderma* OR Hyperkeratosis OR Ecthyma OR Melanoma OR Panniculitis):ab,ti) AND ('observational study'/exp OR 'cohort analysis'/exp OR 'longitudinal study'/exp OR 'retrospective study'/exp OR 'prospective study'/exp OR 'health survey'/de OR 'health care survey'/de OR 'epidemiological data'/de OR 'case control study'/de OR 'cross-sectional study'/de OR 'correlational study'/de OR 'population research'/de OR 'family study'/de OR 'major clinical study'/de OR 'multicenter study'/de OR 'comparative study'/de OR 'follow up'/de OR 'open study'/de OR 'review'/exp OR 'systematic review'/exp OR 'meta analysis'/de OR (((observation* OR epidemiolog* OR famil* OR comparativ* OR communit*) NEAR/6 (stud* OR data OR research)) OR cohort* OR longitudinal* OR retrospectiv* OR prospectiv* OR population* OR (national* NEAR/3 (stud* OR survey)) OR (health* NEAR/3 survey*) OR ((case OR cases OR match*) NEAR/3 (control*)) OR (cross NEXT/1 section*) OR correlation* OR multicenter* OR multi-center* OR follow-up* OR followup* OR review* OR meta-analy*):ab,ti) NOT ([Conference Abstract]/lim OR [Letter]/lim OR [Note]/lim OR [Editorial]/lim) NOT ([animals]/lim NOT [humans]/lim)
Medline	((exp "Immunologic Deficiency Syndromes"/ OR (((immun* OR antibod*) ADJ3 (deficien* OR deficit* OR incompeten*)) OR immunodeficien* OR immunocomprom*).ab,ti.) AND ("Congenital, Hereditary, and Neonatal Diseases and Abnormalities"/ OR exp "Immunologic Deficiency Syndromes"/co OR "Genetic

	Diseases, Inborn"/ OR (((primar* OR congenital* OR hereditar* OR genetic* OR familial*) ADJ6 (immunodeficien*)) OR ((primar* OR congenital* OR hereditar* OR genetic* OR familial*) ADJ6 (immun* OR antibod*) ADJ3 (deficien* OR deficit* OR incompeten*))).ab,ti.)) OR ("Severe Combined Immunodeficiency"/ OR "Ataxia Telangiectasia"/ OR "Chediak-Higashi Syndrome"/ OR "Common Variable Immunodeficiency"/ OR Complement C3/df OR "DiGeorge Syndrome"/ OR "Job Syndrome"/ OR exp "Hyper-IgM Immunodeficiency Syndrome"/ OR "IgA Deficiency"/ OR Immunoglobulin M/df OR "Leukocyte-Adhesion Deficiency Syndrome"/ OR (Agammaglobulinemia/ AND "Genetic Diseases, X-Linked"/) OR "Granulomatous Disease, Chronic"/ OR "Candidiasis, Chronic Mucocutaneous"/ OR "Cryopyrin-Associated Periodic Syndromes"/ OR "Albinism, Ocular"/ OR "Wiskott-Aldrich Syndrome"/ OR "Netherton Syndrome"/ OR ((severe* ADJ3 combin* ADJ3 immun*) OR (ataxi* ADJ3 (telangiect* OR teleangiect*)) OR (Chediak ADJ3 (Higashi OR anomal* OR syndrom*)) OR (common ADJ3 variable ADJ3 immun*) OR (complement* ADJ3 deficienc*) OR (Congenital* ADJ3 CD4 ADJ3 deficien*) OR (Cyclic* ADJ3 haematopoie*) OR DiGeorge OR (hyper ADJ (IgE OR igm)) OR "job syndrome" OR (hyperimmunoglobul* ADJ (E or M)) OR (Idiopathic ADJ3 CD4 ADJ3 lymphocytopenia) OR ((immunoglobulin-A OR iga OR immunoglobulin-M OR igm OR irak OR IL-1-receptor-associated-kinase OR leukocyte-adhesion OR tap OR transporter-associated-with-antigen*) ADJ3 deficien*) OR whim OR (Wart* ADJ6 Hypogammaglobulin* ADJ6 Immun ADJ6 Myelokath*) OR ((linked OR bruton) ADJ agammaglobulin*) OR (chronic ADJ3 granulomat*) OR (chronic* ADJ3 mucocutan* ADJ3 candidas*) OR Griscelli OR (ocular ADJ3 albin*) OR (Hermansky ADJ3 Pudlak) OR (Wiskott ADJ3 Aldrich) OR Netherton OR IPEX-syndrome OR APECED OR PLAID OR ((DOCK8 OR Phosphoglucomutase-3 OR PGM3 OR Mammalian-sterile OR MST1 OR GATA2 OR CARD9 OR IL-17RA OR IL-17F OR ACT1) ADJ3 deficien*) OR (mutation ADJ3 (STAT1 OR Dectin-1))).ab,ti.)) AND (exp "Skin Diseases"/ OR "Staphylococcus aureus"/ OR "Staphylococcal Infections"/ OR (((skin OR cutaneous OR dermal OR mucocutan* OR pigment*) ADJ3 (disease* OR disorder* OR manifestat* OR infect* OR candid* OR abscess* OR granulom* OR lymphoma OR carcinoma* OR tumor* OR cancer* OR neoplas*)) OR eczem* OR verruca* OR Wart* OR dermatit* OR (lupus ADJ3 erythemat*) OR (small-vessel* ADJ3 vasculit*) OR Ulcer* OR decubit* OR pyoderm* OR atop* OR ((Staphylococcus OR s) ADJ aureus) OR ((Basal OR
--	---

	squamous) ADJ cell* ADJ6 carcinom*) OR hyperpigment* OR hypopigment* OR Cafe-au-lait OR albinis* OR vitiligo OR depigment* OR erythroderma* OR Hyperkeratosis OR Ecthyma OR Melanoma OR Panniculitis).ab,ti.) AND ("observational study"/ OR exp "Cohort Studies"/ OR "Health Surveys"/ OR "Epidemiologic Studies"/ OR "Case-Control Studies"/ OR "Cross-Sectional Studies"/ OR "multicenter study"/ OR "comparative study"/ OR "review"/ OR "Meta-Analysis"/ OR (((observation* OR epidemiolog* OR famil* OR comparativ* OR communit*) ADJ6 (stud* OR data OR research)) OR cohort* OR longitudinal* OR retrospectiv* OR prospectiv* OR population* OR (national* ADJ3 (stud* OR survey)) OR (health* ADJ3 survey*) OR ((case OR cases OR match*) ADJ3 (control*)) OR (cross ADJ section*) OR correlation* OR multicenter* OR multicenter* OR follow-up* OR followup* OR review* OR meta-analy*).ab,ti.) NOT (letter OR news OR comment OR editorial OR congresses OR abstracts).pt. NOT (exp animals/ NOT humans/)
Web of Science	TS=(((((((immun* OR antibod*) NEAR/2 (deficien* OR deficit* OR incompeten*)) OR immunodeficien* OR immunocomprom*)) AND (((primar* OR congenital* OR hereditar* OR genetic* OR familial*) NEAR/5 (immunodeficien*)) OR ((primar* OR congenital* OR hereditar* OR genetic* OR familial*) NEAR/5 (immun* OR antibod*) NEAR/2 (deficien* OR deficit* OR incompeten*))))) OR (((severe* NEAR/2 combin* NEAR/2 immun*) OR (ataxi* NEAR/2 (telangiect* OR teleangiect*)) OR (Chediak NEAR/2 (Higashi OR anomal* OR syndrom*)) OR (common NEAR/2 variable NEAR/2 immun*) OR (complement* NEAR/2 deficienc*) OR (Congenital* NEAR/2 CD4 NEAR/2 deficien*) OR (Cyclic* NEAR/2 haematopoie*) OR DiGeorge OR (hyper NEAR/1 (IgE OR igm)) OR "job syndrome" OR (hyperimmunoglobul* NEAR/1 (E or M)) OR (Idiopathic NEAR/2 CD4 NEAR/2 lymphocytopenia) OR ((immunoglobulin-A OR iga OR immunoglobulin-M OR igm OR irak OR IL-1-receptor-associated-kinase OR leukocyte-adhesion OR tap OR transporter-associated-with-antigen*) NEAR/2 deficien*) OR whim OR (Wart* NEAR/5 Hypogammaglobulin* NEAR/5 Immun NEAR/5 Myelokath*) OR ((linked OR bruton) NEAR/1 agammaglobulin*) OR (chronic NEAR/2 granulomat*) OR (chronic* NEAR/2 mucocutan* NEAR/2 candidias*) OR Griscelli OR (ocular NEAR/2 albin*) OR (Hermansky NEAR/2 Pudlak) OR (Wiskott NEAR/2 Aldrich) OR Netherton OR IPEX-syndrom OR APECED OR PLAID OR ((DOCK8 OR Phosphoglucomutase-3 OR PGM3 OR

	Mammalian-sterile OR MST1 OR GATA2 OR CARD9 OR IL-17RA OR IL-17F OR ACT1) NEAR/2 deficien*) OR (mutation NEAR/2 (STAT1 OR Dectin-1))))) AND (((((skin OR cutaneous OR dermal OR mucocutan* OR pigment*) NEAR/2 (disease* OR disorder* OR manifestat* OR infect* OR candid* OR abscess* OR granulom* OR lymphoma OR carcinoma* OR tumor* OR cancer* OR neoplas*)) OR eczem* OR verruca* OR Wart* OR dermatit* OR (lupus NEAR/2 erythemat*) OR (small-vessel* NEAR/2 vasculit*) OR Ulcer* OR decubit* OR pyoderm* OR atop* OR ((Staphylococcus OR s) NEAR/1 aureus) OR ((Basal OR squamous) NEAR/1 cell* NEAR/5 carcinom*) OR hyperpigment* OR hypopigment* OR Café-au-lait OR albinis* OR vitiligo OR depigment* OR erythroderma* OR Hyperkeratosis OR Ecthyma OR Melanoma OR Panniculitis)) AND (((((observation* OR epidemiolog* OR famil* OR comparativ* OR communit*) NEAR/5 (stud* OR data OR research)) OR cohort* OR longitudinal* OR retrospectiv* OR prospectiv* OR population* OR (national* NEAR/2 (stud* OR survey)) OR (health* NEAR/2 survey*) OR ((case OR cases OR match*) NEAR/2 (control*)) OR (cross NEAR/1 section*) OR correlation* OR multicenter* OR multi-center* OR follow-up* OR followup* OR review* OR meta-analy*))) AND DT=(article)
Cochrane	(((((immun* OR antibod*) NEAR/3 (deficien* OR deficit* OR incompeten*)) OR immunodeficien* OR immunocomprom*):ab,ti) AND (((((primar* OR congenital* OR hereditar* OR genetic* OR familial*) NEAR/6 (immunodeficien*)) OR ((primar* OR congenital* OR hereditar* OR genetic* OR familial*) NEAR/6 (immun* OR antibod*) NEAR/3 (deficien* OR deficit* OR incompeten*))):ab,ti)) OR (((severe* NEAR/3 combin* NEAR/3 immun*) OR (ataxi* NEAR/3 (telangiect* OR teleangiect*)) OR (Chediak NEAR/3 (Higashi OR anomal* OR syndrom*)) OR (common NEAR/3 variable NEAR/3 immun*) OR (complement* NEAR/3 deficienc*) OR (Congenital* NEAR/3 CD4 NEAR/3 deficien*) OR (Cyclic* NEAR/3 haematopoie*) OR DiGeorge OR (hyper NEXT/1 (IgE OR igm)) OR 'job syndrome' OR (hyperimmunoglobul* NEXT/1 (E or M)) OR (Idiopathic NEAR/3 CD4 NEAR/3 lymphocytopenia) OR ((immunoglobulin-A OR iga OR immunoglobulin-M OR igm OR irak OR IL-1-receptor-associated-kinase OR leukocyte-adhesion OR tap OR transporter-associated-with-antigen*) NEAR/3 deficien*) OR whim OR (Wart* NEAR/6 Hypogammaglobulin* NEAR/6 Immun NEAR/6 Myelokath*) OR ((linked OR bruton) NEXT/1 agammaglobulin*) OR (chronic NEAR/3 granulomat*) OR (chronic* NEAR/3 mucocutan* NEAR/3

	candidias*) OR Griscelli OR (ocular NEAR/3 albin*) OR (Hermansky NEAR/3 Pudlak) OR (Wiskott NEAR/3 Aldrich) OR Netherton OR IPEX-syndrome OR APECED OR PLAID OR ((DOCK8 OR Phosphoglucomutase-3 OR PGM3 OR Mammalian-sterile OR MST1 OR GATA2 OR CARD9 OR IL-17RA OR IL-17F OR ACT1) NEAR/3 deficien*) OR (mutation NEAR/3 (STAT1 OR Dectin-1))):ab,ti)) AND (((skin OR cutaneous OR dermal OR mucocutan* OR pigment*) NEAR/3 (disease* OR disorder* OR manifestat* OR infect* OR candid* OR abscess* OR granulom* OR lymphoma OR carcinoma* OR tumor* OR cancer* OR neoplas*)) OR eczem* OR verruca* OR Wart* OR dermatit* OR (lupus NEAR/3 erythemat*) OR (small-vessel* NEAR/3 vasculit*) OR Ulcer* OR decubit* OR pyoderm* OR atop* OR ((Staphylococcus OR s) NEXT/1 aureus) OR ((Basal OR squamous) NEXT/1 cell* NEAR/6 carcinom*) OR hyperpigment* OR hypopigment* OR Café-au-lait OR albinis* OR vitiligo OR depigment* OR erythroderma* OR Hyperkeratosis OR Ecthyma OR Melanoma OR Panniculitis):ab,ti)
Google Scholar	"primary congenital genetic immunodeficiency immunodeficiencies" skin cutaneous dermal pigmentation (((((immun* OR antibod*) NEAR/3 (deficien* OR deficit* OR incompeten*)) OR immunodeficien* OR immunocomprom*):ab,ti) AND (((primar* OR congenital* OR hereditar* OR genetic* OR familial*) NEAR/6 (immunodeficien*)) OR ((primar* OR congenital* OR hereditar* OR genetic* OR familial*) NEAR/6 (immun* OR antibod*) NEAR/3 (deficien* OR deficit* OR incompeten*))):ab,ti)) OR (((severe* NEAR/3 combin* NEAR/3 immun*) OR (ataxi* NEAR/3 (telangiect* OR teleangiect*)) OR (Chediak NEAR/3 (Higashi OR anomal* OR syndrom*)) OR (common NEAR/3 variable NEAR/3 immun*) OR (complement* NEAR/3 deficienc*) OR (Congenital* NEAR/3 CD4 NEAR/3 deficien*) OR (Cyclic* NEAR/3 haematopoie*) OR DiGeorge OR (hyper NEXT/1 (IgE OR igm)) OR 'job syndrome' OR (hyperimmunoglobul* NEXT/1 (E or M)) OR (Idiopathic NEAR/3 CD4 NEAR/3 lymphocytopenia) OR ((immunoglobulin-A OR iga OR immunoglobulin-M OR igm OR irak OR IL-1-receptor-associated-kinase OR leukocyte-adhesion OR tap OR transporter-associated-with-antigen*) NEAR/3 deficien*) OR whim OR (Wart* NEAR/6 Hypogammaglobulin* NEAR/6 Immun NEAR/6 Myelokath*) OR ((linked OR bruton) NEXT/1 agammaglobulin*) OR (chronic NEAR/3 granulomat*) OR (chronic* NEAR/3 mucocutan* NEAR/3 candidias*) OR Griscelli OR (ocular NEAR/3 albin*) OR (Hermansky NEAR/3

	Pudlak) OR (Wiskott NEAR/3 Aldrich) OR Netherton OR IPEX-syndrome OR APECED OR PLAID OR ((DOCK8 OR Phosphoglucomutase-3 OR PGM3 OR Mammalian-sterile OR MST1 OR GATA2 OR CARD9 OR IL-17RA OR IL-17F OR ACT1) NEAR/3 deficient*) OR (mutation NEAR/3 (STAT1 OR Dectin-1))):ab,ti)) AND (((skin OR cutaneous OR dermal OR mucocutan* OR pigment*) NEAR/3 (disease* OR disorder* OR manifestat* OR infect* OR candid* OR abscess* OR granulom* OR lymphoma OR carcinoma* OR tumor* OR cancer* OR neoplas*)) OR eczem* OR verruca* OR Wart* OR dermatit* OR (lupus NEAR/3 erythemat*) OR (small-vessel* NEAR/3 vasculit*) OR Ulcer* OR decubit* OR pyoderm* OR atop* OR ((Staphylococcus OR s) NEXT/1 aureus) OR ((Basal OR squamous) NEXT/1 cell* NEAR/6 carcinom*) OR hyperpigment* OR hypopigment* OR Café-au-lait OR albinis* OR vitiligo OR depigment* OR erythroderma* OR Hyperkeratosis OR Ecthyma OR Melanoma OR Panniculitis):ab,ti)
--	---

1 **Table E2**

2 Study characteristics per study

3 Full reference details are provided in Appendix E4 of the Online Repository

	Country	Patients			IHE Quality Appraisal ^a	Primary immunodeficiency	<i>Staphylococcus aureus</i> ^b	Atopy ^c
		N	% Male	Mean age (y)				
Gernez 2018 ³⁵	USA	85	58.8	27.3 (med)	5.5/11	Autosomal dominant hyper IgE syndrome	-	47/82 eczema, 31/82 food allergy
Nguyen 2018 ⁹¹	USA	71	32.4	-	4.5/11	GATA2 deficiency	-	-
Sahin 2018 ⁸²	Turkey	8	75.0	16.9	7.5	Adenosine deaminase 2 deficiency	-	-
Zhou 2018 ³²	China	169	95.9	-	10.5/13	Chronic granulomatous disease	1/4 suppurative dermatitis	-
Azizi 2017 ⁶⁵	Iran	471	67.3	16.0 (med)	7.5/11	X-linked agammaglobulinemia	-	-
						Common variable immunodeficiency		
						Selective IgA deficiency		
Broides 2017 ³⁷	Israel	10	-	6.4	4.5/11	Autosomal recessive hyper IgE syndrome	-	10/10 eczema, 1/10 food allergy, 4/10 asthma
Deripapa 2017 ⁶³	Russia	35	45.7	13.4 (med)	8.5/13	Nijmegen breakage syndrome	-	-
Erkoçoğlu 2017 ⁴³	Turkey	81	-	10.4	7.5/13	Selective IgA deficiency	-	9/81 eczema, 1/81 food allergy, 28/81 asthma, 22/81 rhinoconjunctivitis
Magen 2017 ⁷²	Israel	347	54.2	24.1	8.5/11	Selective IgA deficiency	-	16/347 eczema
Muşabak 2017 ⁶⁹	Turkey	31	61.3	28 (med)	8/11	Common variable immunodeficiency	-	-
Wolach 2017 ⁸⁸	Israel	84	79.8	-	5.5/11	Chronic granulomatous disease	-	-
Wu 2017 ²⁹	China	17	47.1	11.8	7.5/11	Autosomal dominant hyper IgE syndrome	3/4 wounds	17/17 eczema, 2/17 food allergy
Wu 2017 ⁴⁵	China	48	91.7	5.8	5.5/11	Chronic granulomatous disease	-	8/48 eczema, 1/18 rhinoconjunctivitis
Zaidi 2017 ⁷⁷	India	22	-	14 (med)	7/11	Autoimmune polyendocrinopathy	-	-

						candidiasis ectodermal dystrophy		
Altun 2016 ³⁸	Turkey	258	58.5	59.3 mo	4.5/11	Hypogammaglobulinemia	-	0/12 eczema, 7/12 asthma, 8/12 rhinoconjunctivitis
						Selective IgA deficiency		4/12 eczema, 3/12 asthma, 8/12 rhinoconjunctivitis
						IgM deficiency		11/53 eczema, 19/53 asthma, 35/53 rhinoconjunctivitis
Baris 2016 ⁸⁰	Turkey	30	83.3	10.6	6.5/11	Immunodysregulation, polyendocrinopathy, enteropathy X-linked syndrome	-	2/5 eczema
Blazina 2016 ⁷¹	Slovenia	247	59.5	20	8/11	Selective IgA deficiency	-	-
						Autoimmune polyendocrinopathy		
						candidiasis ectodermal dystrophy		
Ferre 2016 ⁷⁶	USA	35	45.7	13.4 (med)	8.5/11	Chronic granulomatous disease		
						Autoimmune polyendocrinopathy		
						candidiasis ectodermal dystrophy		
Mehr 2016 ⁹⁴	Australia	18	66.7	12 (med)	7.5/11	Muckle-Wells syndrome	-	-
						Neonatal onset multisystem inflammatory disease		
Sobolewska 2016 ⁹²	Germany	37	43.2	-	4.5/11	Muckle-Wells syndrome	-	-
Szczawińska-Popłonyk 2016 ³⁹	Poland	78	53.8	17 mo	5/11	Hypogammaglobulinemia		46/78 eczema, 40/78 food allergy
Van Montfrans 2016 ⁸¹	The Netherlands, Belgium	9	77.8	17.3	10/13	Adenosine deaminase 2 deficiency	-	5/9 eczema
Aderibigbe 2015 ⁴⁶	USA	36	-	-	7/11	PLCG2 associated antibody deficiency and immune dysregulation	-	1/8 eczema, 3/8 food allergy, 3/8 asthma, 2/8 rhinoconjunctivitis ^d
Gualdi 2015 ⁶⁷	Italy	47 (CVID),	57.4 (CVID),	23.4 (CVID),	9/13	Common variable immunodeficiency	-	9/47 eczema

		102 (SIgAD)	54.9 (SIgAD)	8.6 (SIgAD)		Selective IgA deficiency		59/102 eczema
Aghamohammadi 2014 ⁵⁸	Iran	75	58.7	2 (med)	11.5/13	Hyper IgE syndrome unspecified	-	5/5 eczema
Celiksoy 2014 ⁶⁶	Turkey	160	61.3	1.2 (med)	11.5/13	Hypogammaglobulinemia	-	28/28 eczema
						Selective IgA deficiency		13/13 eczema
						IgM deficiency		11/11 eczema
						IgG deficiency		14/14 eczema
Greenberger 2013 ⁵⁴	Israel	32	59.4	11.8	7.5/11	Ataxia-telangiectasia	-	-
Uzel 2013 ⁷⁹	USA	5	80.0	9.9	10/13	Immunodysregulation, polyendocrinopathy, enteropathy X-linked syndrome	-	5/5 eczema
Chu 2012 ³⁶	USA	21	47.6	15.9	4.5/11	Autosomal recessive hyper IgE syndrome	-	19/21 eczema, 14/21 food allergy, 10/21 asthma
Olaiwan 2011 ²⁸	France	21	52.4	19	5/11	Autosomal dominant hyper IgE syndrome	2/5 papulopustular rash, 20/20 eczematous dermatitis, 20/20 cold abscesses	20/21 eczema, 5/21 asthma or rhinoconjunctivitis
Aghamohammadi 2009 ⁴²	Iran	23	65.2	8 (med)	8/13	Selective IgA deficiency	-	12/23 eczema, 7/23 food allergy, 12/23 asthma, 8/23 rhinoconjunctivitis
Gregorek 2009 ⁶¹	Poland	21	52.4	11.2 (med)	6/11	Nijmegen breakage syndrome	-	-
Halabi-Tawil 2009 ⁷⁸	France	10	100	-	5.5/11	Immunodysregulation, polyendocrinopathy, enteropathy X-linked syndrome	-	7/10 eczema
Renner 2009 ³¹	Germany	9	66.7	6 (med)	6/11	Comèl Netherton syndrome	8/9 recurrent or persistent skin infections	9/9 atopy: 8/9 eczema, 7/9 food allergy, 2/9 asthma, 5/9 rhinoconjunctivitis
Van den Berg 2009 ⁸⁴	The Netherlands	429	81.8	-	7/11	Chronic granulomatous disease	-	eczema
Gambineri 2008 ⁴⁴	Italy	14	-	range 0-19	7/11	Immunodysregulation, polyendocrinopathy, enteropathy X-linked syndrome	-	11/14 eczema, 1/14 food allergy, 1/14 asthma

Movahedi 2007 ⁸³	Iran	15	66.7	4 (med)	6.5/11	Leukocyte adhesion defect type I	-	-
Rezaei 2007 ⁸⁹	Iran	18	55.6	8.8	6/11	Severe congenital neutropenia	-	-
Collins 2006 ⁷⁴	Ireland	18	38.9	14.6	8.5/13	Autoimmune polyendocrinopathy candidiasis ectodermal dystrophy	-	-
El-Darouti 2006 ⁹³	Egypt	6	33.3	23.2	2/11	Muckle-Wells syndrome	-	-
Moin 2006 ⁶	Iran	210	58.1	-	5/11	Ataxia-telangiectasia	-	1/62 eczema
						Hyper IgE syndrome unspecified		5/6 eczema
						X-linked agammaglobulinemia		4/23 eczema
						Common variable immunodeficiency		6/28 eczema
						Selective IgA deficiency		5/17 eczema
						Leukocyte adhesion defect		-
						Chronic granulomatous disease		-
						Chronic mucocutaneous candidiasis		-
Eberting 2004 ³⁴	USA	43	37.2	23	5/11	Hyper IgE syndrome unspecified	-	28/43 eczema, 22/43 asthma, 14/43 rhinoconjunctivitis
Haas 2004 ⁹⁵	Germany	16	43.8	-	6.5/13	Muckle-Wells syndrome	-	-
Markert 2004 ⁶⁴	USA	5	100	-	9/13	DiGeorge syndrome	-	-
Renner 2004 ³⁰	Germany, USA	13	30.1	-	10/13	Autosomal recessive hyper IgE syndrome	skin abscesses (frequently)	13/13 eczema
Dupuis-Girod 2003 ⁵⁷	France	55	100	-	5/11	Wiskott-Aldrich syndrome	-	-
Ullbro 2003 ⁹⁰	Sweden	47	44.7	10 (med)	9/13	Papillon-Lefèvre syndrome	-	-
Chamlin 2002 ²⁷	USA	8	75.0	-	4/11	Hyper IgE syndrome unspecified	4/7 papulopustular eruption	5/8 eczema
Hiel 2001 ⁶²	The Netherlands	55	56.4	-	3/11	Nijmegen breakage syndrome	-	-
Berron-Ruiz 2000 ⁷	Mexico	130	-	range 0-17	3/11	Severe combined immunodeficiency	-	-

						Ataxia-telangiectasia		-
						Wiskott-Aldrich syndrome		5/5 eczema
						X-linked agammaglobulinemia		-
						Common variable immunodeficiency		-
						Selective IgA deficiency		-
Liese 2000 ⁸⁵	Germany	39	94.9	-	6/11	Chronic granulomatous disease	-	eczema
Lipsker 2000 ⁹⁶	France	47	-	-	5/11	C2 deficiency	-	-
Pruszkowski 2000 ⁵²	France	51	47.1	-	5/11	Omenn syndrome	-	-
						Comèl Netherton syndrome		-
Winkelstein 2000 ⁸⁶	USA	368	85.9	-	5/11	Chronic granulomatous disease	-	-
Grimbacher 1999 ⁵⁹	USA	30	33.3	26.5	5.5/11	Hyper IgE syndrome unspecified	-	30/30 eczema
Koskinen 1996 ⁴⁰	Finland	159	-	-	8/13	Selective IgA deficiency	-	12/159 eczema, 5/159 food allergy, 17/159 rhinoconjunctivitis
Sullivan 1994 ⁵⁶	USA	154	100	11	7/11	Wiskott-Aldrich syndrome	-	-
Porter 1993 ⁶⁸	England	15	86.7	range 3-14	4.5/11	Common variable immunodeficiency	-	4/15 eczema
Porter 1993 ⁷⁰	England	39	71.8	range 2-13	4.5/11	Selective IgA deficiency	-	2/39 eczema
Patrizi 1992 ⁴¹	Italy	142	57.0	6.0	6.5/13	Selective IgA deficiency	-	9/123 eczema
Paller 1991 ⁵⁵	USA	8	25.0	10.5	4.5/11	Ataxia-telangiectasia	-	-
Ahonen 1990 ⁷⁵	Finland	68	54.4	follow-up to age 10 mo-53 y at end of study	10/13	Autoimmune polyendocrinopathy candidiasis ectodermal dystrophy	-	-
Charon 1985 ⁶⁰	USA	27	44.4	20.3	3.5/11	Hyper E syndrome unspecified	-	-
						Chronic granulomatous disease		-
Cohen 1985 ⁸⁷	USA	6	66.6	range 17-32	4/11	Chronic granulomatous disease	-	-
Cohen 1984 ³³	USA	12	-	-	4.5/11	Ataxia-telangiectasia	-	0/12 atopy
Göring 1981 ⁷³	Germany	8	-	-	3.5/11	Selective IgA deficiency	-	-

Reed 1966 ⁵³	USA	22	63.6	13.4	6/11	Ataxia-telangiectasia	-	2/22 eczema
Abbreviations: N, number of patients; y, year; mo, months; med, median; CVID, common variable immunodeficiency; sIgAD, selective IgA deficiency; IHE, Institute of Health Economics ^a Prospective studies could reach a maximum score of 13 points and retrospective studies could reach a maximum score of 11 points according to the Institute of Health Economics Quality Appraisal Checklist for Case Series Studies ^b Number of patients with positive <i>Staphylococcus aureus</i> culture / number of patients with reported skin disorder ^c Number of patients with atopic disorder / number of patients included in the study ^d Number of patients with atopic disorder / number of patients with a history of cold urticarial (subgroup in original study)								

Figure E1

Skin disorders in patients with a primary immunodeficiency

- a. Exfoliative erythroderma in Omenn syndrome
- b. Dermatitis in Comèl Netherton syndrome
- c. Exfoliative erythroderma in Comèl Netherton syndrome
- d. Urticaria in common variable immunodeficiency
- e. Alopecia in immunodysregulation polyendocrinopathy enteropathy X-linked syndrome
- f. Onychodystrophy in immunodysregulation polyendocrinopathy enteropathy X-linked syndrome
- g. Discoid lupus erythematosus in chronic granulomatous disease
- h. Thrush in chronic mucocutaneous candidiasis

Appendix E1

Selected terms of primary immunodeficiency disease and skin disorder used for the electronic search

Primary immunodeficiency disease	Skin disorder
Agammaglobulinemia	Abscess
Autoimmune polyendocrinopathy candidiasis ectodermal dystrophy	Albinism
Ataxia-telangiectasia	Basal cell carcinoma
Chediak-Higashi syndrome	Café-au-lait
Chronic granulomatous disease	Candidiasis
Chronic mucocutaneous candidiasis	Carcinoma
Common variable immunodeficiency	Decubitus
DiGeorge syndrome	Depigmentation disorder
Griscelli syndrome	Dermatitis
Hermansky-Pudlak syndrome	Ecthyma
Hyper IgE syndrome	Eczema
Hyper IgM syndrome	Erythroderma
Hypogammaglobulinemia	Granuloma
Idiopathic CD4+ lymphocytopenia	Hyperkeratosis
IgA deficiency	Hyperpigmentation
IgM deficiency	Hypopigmentation
Interleukin-1 receptor-associated kinase-4 deficiency	Infection
Immunodysregulation polyendocrinopathy enteropathy X-linked syndrome	Lupus erythematosus
Leukocyte adhesion defect	Lymphoma
Comèl-Netherton syndrome	Melanoma
PLCG2 associated antibody deficiency and immune dysregulation	Panniculitis
Severe combined immunodeficiency	Pigment disorder
Transporter-associated-with-antigen deficiency	Pyoderma

Warts, hypogammaglobulinemia, immunodeficiency and myelokathexis syndrome	Small vessel vasculitis
Wiskott-Aldrich syndrome	Squamous cell carcinoma
X-linked agammaglobulinemia	Ulcer
	Verruca
	Vitiligo
	Wart

Appendix E2

Inclusion and exclusion criteria for selecting studies for this review

Types of studies

Inclusion criteria

- Original observational and experimental human studies which assess the presence of skin disorders in patients with a primary immunodeficiency disease.
- Original observational and experimental human studies which report a differential diagnosis of a specific skin disorder that includes a primary immunodeficiency disease.

Exclusion criteria

- Case reports (<5 patients per primary immunodeficiency disease), conference abstracts, letters, editorials and review articles.
- Studies reporting only a genetic mutation (suggestive for a primary immunodeficiency disease) instead of a primary immunodeficiency disease as clinical diagnosis.
- Studies reporting skin disorders in primary immunodeficiency diseases that developed after and/or during treatment or intervention.
- Studies in which the description of skin disorders in primary immunodeficiency diseases was not part of the results section.

Participants

Inclusion criteria

- Patients of all ages with a primary immunodeficiency disease according to Picard *et al.* both in hospital setting and the general population.¹⁶

Exclusion criteria

- Patients with an acquired immunodeficiency.

Controls

- No controls.

Outcome measures

- Primary: An overview of the presence of skin disorders described in patients with a PID, in order to compose a differential diagnosis of primary immunodeficiency diseases per skin disorder.
- Secondary: The prevalence of skin disorders in primary immunodeficiency diseases, *S. aureus* associated skin disorders in primary immunodeficiency diseases and primary immunodeficiency diseases associated with an atopic constitution (i.e. atopic dermatitis, food allergy, asthma, rhinoconjunctivitis).

Appendix E3

Quality assessment score

Institute of Health Economics (IHE) Quality Appraisal Checklist for Case Series Studies

Stars indicate the points allocated if the item criterion is met.¹⁹ A maximum score of 13 can be allocated in each article.

Criteria that are not applicable are excluded.

Study objective	
1.	Was the hypothesis/aim/objective of the study clearly stated? Yes: The hypothesis/aim/objective of the study was clearly reported (includes patients, intervention and outcome). ★ Partial: Only one or two components (patients, intervention, or outcome) were included. ★ No: The hypothesis/aim/objective was not reported.
Study design	
2.	Was the study conducted prospectively? Yes: It was clearly stated that the study was conducted prospectively. ★ Unclear: Unclear or no information was provided. No: The study clearly stated it was a retrospective study.
3.	Were the cases collected in more than one centre? Yes: Cases were collected in more than one centre (multicentre study). ★ Unclear: Unclear where the patients came from. No: Cases were collected from one centre.
4.	Were patients recruited consecutively? Yes: There was a clear statement or it was clear from the context that the patients were recruited consecutively; or the study stated that all eligible patients were recruited. ★ Unclear: No information was provided about the method used to recruit patients in the study. No: The study clearly stated that patients were not recruited consecutively; or the patients were recruited based on other criteria such as access to intervention determined by the distance or availability of resources.
Study population	
5.	Were the characteristics of the patients included in the study described? Yes: All of the most relevant characteristics of the patients were reported (for example, number, age,

	gender, ethnicity, severity of disease/condition, comorbidity, or etiology). ★ Partial: Some, but not all, of the most relevant characteristics were reported. ½★ No: Only the number of patients was reported. <i>Note: Assessor(s) should decide which aspects are important before using the checklist.</i>
6.	Were the eligibility criteria (i.e. inclusion and exclusion criteria) for entry into the study clearly stated? Yes: Both inclusion and exclusion criteria were reported. ★ Partial: Either the inclusion or exclusion criteria were reported. ½★ No: Neither inclusion nor exclusion criteria were reported. <i>Note: Assessor(s) should decide which aspects are important before using the checklist.</i>
7.	Did patients enter the study at a similar point in the disease? Yes: It was clear from the baseline data presented in the study (for example, tables of patients' characteristics) that the majority (at least 80%) of patients entered the study at a similar point in terms of the duration and severity of the disease/condition and the presence of co-morbidities/complications. ★ Unclear: There was no baseline information on patients' characteristics to make a judgment. No: There was a wide range in the severity of the disease/condition and co-morbidities/complications in patients at baseline. <i>Note: Assessor(s) should decide which aspects are important before using the checklist. It might be useful to discuss with specialists to determine the most important aspects that should be considered.</i>
Outcome measures	
10.	Were relevant outcome measures established a priori? Yes: All relevant outcome measures were stated in the introduction or methods section. ★ Partial: Some, but not all, of the relevant outcome measures were stated in the introduction or method section. ½★ No: None of the relevant outcome measures were stated in the introduction or method section.
12.	Were the relevant outcomes measured using appropriate objective/subjective methods? Yes: All relevant outcomes were measured with appropriate methods. These measures can be objective (for example, gold standard tests or standardized clinical tests), subjective (for example, self-administered questionnaires, standardized forms, or patient symptoms interview forms), or both. ★ Partial: Some, but not all, relevant outcomes were measured with appropriate methods. ½★ No: The methods used to measure the relevant outcomes were inappropriate. <i>Note: Assessor(s) should decide which methods are appropriate before using the checklist.</i>
Results and conclusions	
15.	Was follow-up long enough for important events and outcomes to occur?

	Yes: It was clear from the information provided that the follow-up period was long enough for the majority (at least 80%) of patients, to allow for important events and outcomes (for example, changes in clinical status, adverse events) to occur. ★ Unclear: The length of follow-up was not clearly reported. No: It is clear from the information provided that the follow-up period was not long enough to allow for important events and outcomes to occur. <i>Note: Assessor(s) should define the appropriate duration of follow-up for each outcome of interest (for example, short-term and long-term adverse events).</i>
16.	Were losses to follow-up reported? Yes: The number or proportion of patients lost to follow-up was clearly reported; the authors reported outcome results on all patients initially included; or the number lost to follow-up can be subtracted from the number of patients enrolled and the number of patients included in the final analysis. ★ Unclear: There was a discrepancy between the number or proportion of patients reported in tables, figures, and text. No: The number or proportion of patients lost to follow-up was not reported.
19.	Were the conclusions of the study supported by the results? Yes: The conclusions of the study were supported by the evidence presented in the results and discussion sections. ★ Unclear: Unclear conclusion statement that makes it difficult to link the presented evidence to conclusions. No: The conclusions were not supported by the evidence presented in the results and discussion sections.
Competing interests and sources of support	
20.	Were both competing interests and sources of support for the study reported? Yes: Both competing interests and sources of support (financial or other) received for the study were reported; or the absence of any competing interest and source of support was acknowledged. ★ Partial: Either the competing interest or source of support was reported. ½ ★ No: Neither competing interests nor sources of support were reported.

Appendix E4

Supplementary references

6. Moin A, Farhoudi A, Moin M, Pourpak Z, Bazargan N. Cutaneous manifestations of primary immunodeficiency diseases in children. *Iran J Allergy Asthma Immunol* 2006;**5**(3):121-126.
7. Berron-Ruiz A, Berron-Perez R, Ruiz-Maldonado R. Cutaneous markers of primary immunodeficiency diseases in children. *Pediatr Dermatol* 2000;**17**(2):91-96.
27. Chamlin SL, McCalmont TH, Cunningham BB, Esterly NB, Lai CH, Mallory SB, et al. Cutaneous manifestations of hyper-IgE syndrome in infants and children. *J Pediatr* 2002;**141**(4):572-575.
28. Olaiwan A, Chandesris MO, Fraitag S, Lortholary O, Hermine O, Fischer A, et al. Cutaneous findings in sporadic and familial autosomal dominant hyper-IgE syndrome: a retrospective, single-center study of 21 patients diagnosed using molecular analysis. *J Am Acad Dermatol* 2011;**65**(6):1167-1172.
29. Wu J, Chen J, Tian ZQ, Zhang H, Gong RL, Chen TX, et al. Clinical Manifestations and Genetic Analysis of 17 Patients with Autosomal Dominant Hyper-IgE Syndrome in Mainland China: New Reports and a Literature Review. *J Clin Immunol* 2017;**37**(2):166-179.
30. Renner ED, Puck JM, Holland SM, Schmitt M, Weiss M, Frosch M, et al. Autosomal recessive hyperimmunoglobulin E syndrome: a distinct disease entity. *J Pediatr* 2004;**144**(1):93-99.
31. Renner ED, Hartl D, Rylaarsdam S, Young ML, Monaco-Shawver L, Kleiner G, et al. Comel-Netherton syndrome defined as primary immunodeficiency. *J Allergy Clin Immunol* 2009;**124**(3):536-543.
32. Zhou Q, Hui X, Ying W, Hou J, Wang W, Liu D, et al. A Cohort of 169 Chronic Granulomatous Disease Patients Exposed to BCG Vaccination: a Retrospective Study from a Single Center in Shanghai, China (2004–2017). *J Clin Immunol* 2018:1-13.
33. Cohen LE, Tanner DJ, Schaefer HG, Levis WR. Common and uncommon cutaneous findings in patients with ataxia-telangiectasia. *J Am Acad Dermatol* 1984;**10**(3):431-438.
34. Eberting CL, Davis J, Puck JM, Holland SM, Turner ML. Dermatitis and the newborn rash of hyper-IgE syndrome. *Arch Dermatol* 2004;**140**(9):1119-1125.
35. Gernez Y, Freeman AF, Holland SM, Garabedian E, Patel NC, Puck JM, et al. Autosomal Dominant Hyper-IgE Syndrome in the USIDNET Registry. *J Allergy Clin Immunol Pract* 2018;**6**(3):996-1001.
36. Chu EY, Freeman AF, Jing H, Cowen EW, Davis J, Su HC, et al. Cutaneous manifestations of DOCK8 deficiency syndrome. *Arch Dermatol* 2012;**148**(1):79-84.
37. Broides A, Mandola AB, Levy J, Yerushalmi B, Pinsk V, Eldan M, et al. The clinical and laboratory spectrum of dedicator of cytokinesis 8 immunodeficiency syndrome in patients with a unique mutation. *Immunol Res* 2017;**65**(3):651-657.
38. Altun D, Akpinar M, Haskoloğlu ZS, Köste Bal S, Kavgacı A, Doğu EF, et al. Immunoglobulin isotype deficiency together with allergic diseases. *Asim Allerji Immunoloji* 2016;**14**(3):164-169.

39. Szczawińska-Popłonyk A, Komasińska P, Bręborowicz A. IgA deficiency: A risk factor for food allergy-related atopic dermatitis in infants and young children. *Postepy Dermatol Alergol* 2016;**33**(5):369-374.
40. Koskinen S. Long-term follow-up of health in blood donors with primary selective IgA deficiency. *J Clin Immunol* 1996;**16**(3):165-170.
41. Patrizi A, Ricci G, Cassoli C, Specchia F, Neri I, Masi M. [Dermatologic diseases associated with IgA deficiency]. *G Ital Dermatol Venereol* 1992;**127**(7-8):325-329.
42. Aghamohammadi A, Cheraghi T, Gharagozlou M, Movahedi M, Rezaei N, Yeganeh M, et al. IgA deficiency: correlation between clinical and immunological phenotypes. *J Clin Immunol* 2009;**29**(1):130-136.
43. Erkoçoğlu M, Metin A, Kaya A, Özcan C, Akan A, Civelek E, et al. Allergic and autoimmune disorders in families with selective IgA deficiency. *Turk J Med Sci* 2017;**47**(2):592-598.
44. Gambineri E, Perroni L, Passerini L, Bianchi L, Doglioni C, Meschi F, et al. Clinical and molecular profile of a new series of patients with immune dysregulation, polyendocrinopathy, enteropathy, X-linked syndrome: inconsistent correlation between forkhead box protein 3 expression and disease severity. *J Allergy Clin Immunol* 2008;**122**(6):1105-1112 e1101.
45. Wu J, Wang WF, Zhang YD, Chen TX. Clinical Features and Genetic Analysis of 48 Patients with Chronic Granulomatous Disease in a Single Center Study from Shanghai, China (2005-2015): New Studies and a Literature Review. *J Immunol Res* 2017;**2017**.
46. Aderibigbe OM, Priel DL, Lee CC, Ombrello MJ, Prajapati VH, Liang MG, et al. Distinct Cutaneous Manifestations and Cold-Induced Leukocyte Activation Associated With PLCG2 Mutations. *JAMA Dermatol* 2015;**151**(6):627-634.
52. Pruszkowski A, Bodemer C, Fraitag S, Teillac-Hamel D, Amoric JC, de Prost Y. Neonatal and infantile erythrodermas: a retrospective study of 51 patients. *Arch Dermatol* 2000;**136**(7):875-880.
53. Reed WB, Epstein WL, Boder E, Sedgwick R. Cutaneous manifestations of ataxia-telangiectasia. *JAMA* 1966;**195**(9):746-753.
54. Greenberger S, Berkun Y, Ben-Zeev B, Levi YB, Barzilai A, Nissenkorn A. Dermatologic manifestations of ataxia-telangiectasia syndrome. *J Am Acad Dermatol* 2013;**68**(6):932-936.
55. Paller AS, Massey RB, Curtis MA, Pelachyk JM, Dombrowski HC, Leickly FE, et al. Cutaneous granulomatous lesions in patients with ataxia-telangiectasia. *J Pediatr* 1991;**119**(6):917-922.
56. Sullivan KE, Mullen CA, Blaese RM, Winkelstein JA. A multiinstitutional survey of the Wiskott-Aldrich syndrome. *J Pediatr* 1994;**125**(6 Pt 1):876-885.
57. Dupuis-Girod S, Medioni J, Haddad E, Quartier P, Cavazzana-Calvo M, Le Deist F, et al. Autoimmunity in Wiskott-Aldrich syndrome: risk factors, clinical features, and outcome in a single-center cohort of 55 patients. *Pediatrics* 2003;**111**(5 Pt 1):e622-627.
58. Aghamohammadi A, Moghaddam ZG, Abolhassani H, Hallaji Z, Mortazavi H, Pourhamdi S, et al. Investigation of underlying primary immunodeficiencies in patients with severe atopic dermatitis. *Allergol Immunopathol (Madr)* 2014;**42**(4):336-341.
59. Grimbacher B, Holland SM, Gallin JI, Greenberg F, Hill SC, Malech HL, et al. Hyper-IgE syndrome with recurrent infections--an autosomal dominant multisystem disorder. *N Engl J Med* 1999;**340**(9):692-702.

60. Charon JA, Mergenhagen SE, Gallin JI. Gingivitis and oral ulceration in patients with neutrophil dysfunction. *J Oral Pathol* 1985;**14**(2):150-155.
61. Gregorek H, Olczak-Kowalczyk D, Dembowska-Baginska B, Pietrucha B, Wakulinska A, Gozdowski D, et al. Oral findings in patients with Nijmegen breakage syndrome: a preliminary study. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2009;**108**(5):e39-45.
62. Hiel JA, Weemaes CM, van Engelen BG, Smeets D, Ligtenberg M, van Der Burgt I, et al. Nijmegen breakage syndrome in a Dutch patient not resulting from a defect in NBS1. *J Med Genet* 2001;**38**(6):E19.
63. Deripapa E, Balashov D, Rodina Y, Laberko A, Myakova N, Davydova NV, et al. Prospective study of a cohort of Russian Nijmegen breakage syndrome patients demonstrating predictive value of low kappa-deleting recombination excision circle (KREC) numbers and beneficial effect of hematopoietic stem cell transplantation (HSCT). *Front Immunol* 2017;**8**(JUL).
64. Markert ML, Alexieff MJ, Li J, Sarzotti M, Ozaki DA, Devlin BH, et al. Complete DiGeorge syndrome: development of rash, lymphadenopathy, and oligoclonal T cells in 5 cases. *J Allergy Clin Immunol* 2004;**113**(4):734-741.
65. Azizi G, Tavakol M, Rafiemanesh H, Kiaee F, Yazdani R, Heydari A, et al. Autoimmunity in a cohort of 471 patients with primary antibody deficiencies. *Expert Rev Clin Immunol* 2017;**13**(11):1099-1106.
66. Celiksoy MH, Topal E, Sancak R, Catal F, Sogut A. Relationship between hypogammaglobulinemia and severity of atopic dermatitis. *Ann Allergy Asthma Immunol* 2014;**113**(4):467-469.
67. Gualdi G, Lougaris V, Baronio M, Vitali M, Tampella G, Moratto D, et al. Burden of Skin Disease in Selective IgA Deficiency and Common Variable Immunodeficiency. *J Investig Allergol Clin Immunol* 2015;**25**(5):369-371.
68. Porter SR, Scully C. Orofacial manifestations in primary immunodeficiencies: common variable immunodeficiencies. *J Oral Pathol Med* 1993;**22**(4):157-158.
69. Muşabak UH, Demirel F, Yeşillik S, Baysan A, Selçuk A, Kartal Ö, et al. Adults with common variable immunodeficiency: A single-center experience. *Turk J Med Sci* 2017;**47**(1):1-12.
70. Porter SR, Scully C. Orofacial manifestations in primary immunodeficiencies involving IgA deficiency. *J Oral Pathol Med* 1993;**22**(3):117-119.
71. Blazina Š, Markelj G, Jeverica AK, Toplak N, Bratanič N, Jazbec J, et al. Autoimmune and Inflammatory Manifestations in 247 Patients with Primary Immunodeficiency—a Report from the Slovenian National Registry. *J Clin Immunol* 2016;**36**(8):764-773.
72. Magen E, Masalha A, Waitman DA, Kahan N, Viner I, Klassov L, et al. Prevalence of dermatologic diseases among patients with selective immunoglobulin A deficiency. *Allergy Asthma Proc* 2017;**38**(1):70-77.
73. Goring HD. [IgA deficiency in the dermatologic clinic. Frequency, clinical relevance, diagnosis and therapy]. *Hautarzt* 1981;**32**(10):505-511.
74. Collins SM, Dominguez M, Ilmarinen T, Costigan C, Irvine AD. Dermatological manifestations of autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy syndrome. *Br J Dermatol* 2006;**154**(6):1088-1093.

75. Ahonen P, Myllarniemi S, Sipila I, Perheentupa J. Clinical variation of autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED) in a series of 68 patients. *N Engl J Med* 1990;**322**(26):1829-1836.
76. Ferre EM, Rose SR, Rosenzweig SD, Burbelo PD, Romito KR, Niemela JE, et al. Redefined clinical features and diagnostic criteria in autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy. 2016.
77. Zaidi G, Bhatia V, Sahoo SK, Sarangi AN, Bharti N, Zhang L, et al. Autoimmune polyendocrine syndrome type 1 in an Indian cohort: a longitudinal study. *Endocr. Connect.* 2017;**6**(5):289-296.
78. Halabi-Tawil M, Ruemmele FM, Fraitag S, Rieux-Laucat F, Neven B, Brousse N, et al. Cutaneous manifestations of immune dysregulation, polyendocrinopathy, enteropathy, X-linked (IPEX) syndrome. *Br J Dermatol* 2009;**160**(3):645-651.
79. Uzel G, Sampaio EP, Lawrence MG, Hsu AP, Hackett M, Dorsey MJ, et al. Dominant gain-of-function STAT1 mutations in FOXP3 wild-type immune dysregulation-polyendocrinopathy-enteropathy-X-linked-like syndrome. *J Allergy Clin Immunol* 2013;**131**(6):1611-1623.
80. Barış HE, Kiyıkım A, Nain E, Özen AO, Karakoç-Aydiner E, Barış S. The plethora, clinical manifestations and treatment options of autoimmunity in patients with primary immunodeficiency. *Turk Pediatr Ars* 2016;**51**(4):186-192.
81. Van Montfrans JM, Hartman EAR, Braun KPJ, Hennekam EAM, Hak EA, Nederkoorn PJ, et al. Phenotypic variability in patients with ADA2 deficiency due to identical homozygous R169Q mutations. *Rheumatology* 2016;**55**(5):902-910.
82. Sahin S, Adrovic A, Barut K, Ugurlu S, Turanli ET, Ozdogan H, et al. Clinical, imaging and genotypical features of three deceased and five surviving cases with ADA2 deficiency. *Rheumatol Int* 2018;**38**(1):129-136.
83. Movahedi M, Entezari N, Pourpak Z, Mamishi S, Chavoshzadeh Z, Gharagozlou M, et al. Clinical and laboratory findings in Iranian patients with leukocyte adhesion deficiency (study of 15 cases). *J Clin Immunol* 2007;**27**(3):302-307.
84. van den Berg JM, van Koppen E, Ahlin A, Belohradsky BH, Bernatowska E, Corbeel L, et al. Chronic granulomatous disease: the European experience. *PLoS One* 2009;**4**(4):e5234.
85. Liese J, Kloos S, Jendrosseck V, Petropoulou T, Wintergerst U, Notheis G, et al. Long-term follow-up and outcome of 39 patients with chronic granulomatous disease. *J Pediatr* 2000;**137**(5):687-693.
86. Winkelstein JA, Marino MC, Johnston RB, Jr., Boyle J, Curnutte J, Gallin JJ, et al. Chronic granulomatous disease. Report on a national registry of 368 patients. *Medicine (Baltimore)* 2000;**79**(3):155-169.
87. Cohen MS, Leong PA, Simpson DM. Phagocytic cells in periodontal defense. Periodontal status of patients with chronic granulomatous disease of childhood. *J Periodontol* 1985;**56**(10):611-617.
88. Wolach B, Gavrieli R, de Boer M, van Leeuwen K, Berger-Achituv S, Stauber T, et al. Chronic granulomatous disease: Clinical, functional, molecular, and genetic studies. The Israeli experience with 84 patients. *Am J Hematol* 2017;**92**(1):28-36.
89. Rezaei N, Moin M, Pourpak Z, Ramyar A, Izadyar M, Chavoshzadeh Z, et al. The clinical, immunohematological, and molecular study of Iranian patients with severe congenital neutropenia. *J Clin Immunol* 2007;**27**(5):525-533.

90. Ullbro C, Crossner CG, Nederfors T, Alfadley A, Thestrup-Pedersen K. Dermatologic and oral findings in a cohort of 47 patients with Papillon-Lefevre syndrome. *J Am Acad Dermatol* 2003;**48**(3):345-351.
91. Nguyen J, Alexander T, Jiang H, Hill N, Abdullaev Z, Pack SD, et al. Melanoma in patients with GATA2 deficiency. *Pigm Cell Melanoma Res* 2018;**31**(2):337-340.
92. Sobolewska B, Angermair E, Deuter C, Doycheva D, Kuemmerle-Deschner J, Zierhut M. NLRP3 A439V Mutation in a Large Family with Cryopyrin-associated Periodic Syndrome: Description of Ophthalmologic Symptoms in Correlation with Other Organ Symptoms. Erratum appears in *J Rheumatol*. 2016 Jul;**43**(7):1451; PMID: 27371654. *J Rheumatol* 2016;**43**(6):1101-1106.
93. El-Darouti MA, Marzouk SA, Abdel-Halim MR. Muckle-Wells syndrome: report of six cases with hyperpigmented sclerodermoid skin lesions. *Int J Dermatol* 2006;**45**(3):239-244.
94. Mehr S, Allen R, Boros C, Adib N, Kakakios A, Turner PJ, et al. Cryopyrin-associated periodic syndrome in Australian children and adults: Epidemiological, clinical and treatment characteristics. *J Paediatr Child Health* 2016;**52**(9):889-895.
95. Haas N, Kuster W, Zuberbier T, Henz BM. Muckle-Wells syndrome: clinical and histological skin findings compatible with cold air urticaria in a large kindred. *Br J Dermatol* 2004;**151**(1):99-104.
96. Lipsker DM, Schreckenberg-Gilliot C, Uring-Lambert B, Meyer A, Hartmann D, Grosshans EM, et al. Lupus erythematosus associated with genetically determined deficiency of the second component of the complement. *Arch Dermatol* 2000;**136**(12):1508-1514.