

Received Date : 14-Jul-2016

Revised Date : 12-Nov-2016

Accepted Date : 25-Nov-2016

Article type : Original

**The association between parental consanguinity and primary immunodeficiency diseases:
a systematic review and meta-analysis**

Hasti Hadizadeh*, M.D.^{1, 2}

M.Masoud Salehi*, M.D.^{1, 2}

Shabnam Khoramnejad, M.D.¹

Kia Vosoughi, M.D.^{1, 2}

Nima Rezaei, M.D. PhD.^{2, 3}

1. Doctor of Medicine (MD), Student Research Committee, Faculty of Medicine, Iran University of Medical Sciences, Tehran, Iran
2. Systematic Review and Meta-Analysis Expert Group (SRMEG), Universal Scientific Education and Research Network (USERN), Tehran, Iran
3. Research Center for Immunodeficiencies, Children's Medical Center, Tehran University of Medical Sciences, Tehran, Iran

Parental consanguinity and primary immunodeficiency diseases

Corresponding Author: M. Masoud Salehi, Medical Student Research Committee, School of Medicine, Iran University of Medical Sciences, Hemmat Expressway, Tehran, Iran. Tel and Fax: +982186703072;
mmasood.salehi@gmail.com

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/pai.12685

This article is protected by copyright. All rights reserved.

Hadizadeh (HH), Salehi (MS), Khoramnejad (SK), Kia Vosoughi (KV), Rezaei (NR)

The association between parental consanguinity and primary immunodeficiency diseases: a systematic review and meta-analysis

Pediatr Allergy Immunol

Abstract

Background: We aimed to establish the prevalence of parental consanguinity among patients with primary immunodeficiency diseases (PID), and compare the prevalence with the general population.

Method: We searched PubMed, EMBASE, and Scopus for studies mentioning parental consanguinity prevalence in PID patients and calculated the prevalence odds ratio (POR) of parental consanguinity in each study, compared to a matched healthy population.

Results: We identified 21 eligible studies with a total population of 18091 accounting for sample overlap. The POR among studies on a sample of mixed PID patients ranged from 0.6 to 21.9 with the pooled POR of 3.0 ($P < 0.001$; $I^2 = 89\%$, 95% CI 2.5-3.7).

Conclusion: PIDs with an autosomal recessive pattern of inheritance had significant odds of parental consanguinity compared to the healthy population; a phenomenon not observed in other inheritance patterns. Determining the extent of the impact that consanguinity impose upon the progeny paves the way for convincing healthcare policy makers in highly consanguineous communities to act more diligently in informing the masses about the consequences of practicing inbreeding.

Keywords: Consanguinity; epidemiology; meta-analysis; prevalence; primary immunodeficiencies.

M. Masoud Salehi, Medical Student Research Committee, School of Medicine, Iran University of Medical Sciences, Hemmat Expressway, Tehran, Iran.

INTRODUCTION

Primary immunodeficiencies (PIDs) are a diverse group of rare, chronic disorders (1) which causes immune dysregulation and imbalance, leading to increased susceptibility to microorganisms and infections, the frequent occurrence of allergy, and predisposition to the development of immunological disorders, autoimmune diseases, and malignancies (2).

Most primary immunodeficiency diseases are inherited. Advances in molecular biology have shown various primary biologic defects in most of these disorders. The mode of inheritance in most primary immunodeficiency diseases is in one of these ways: X-linked recessive, autosomal recessive (AR) or autosomal dominant (AD) (3).

Consanguineous marriage –defined as a union between two persons related as second cousins or closer- increases the risk of AR disorders (4,5). Since the probability that the offspring of consanguineous unions will inherit mutated AR gene increases, the risk for recessive disorders is probably higher in the offspring of consanguineous parents (6).

Knowing more about the impacts of consanguineous marriage helps to implement appropriate community programs to improve family health. To provide more evidence on the contribution of consanguineous marriage in primary immunodeficiency diseases (PID), this systematic review, and meta-analysis, aims to summarize (1) the worldwide prevalence of parental consanguinity among patients with PIDs (2) the prevalence odds ratio (POR) for parental consanguinity in PID patients compared to general population.

MATERIALS AND METHODS

We conducted this systematic review and meta-analysis according to PRISMA statement guidelines (7).

Data sources and search strategies

We constructed an electronic search in PubMed, EMBASE, and Scopus databases to gather articles published up to May 2016. Our target population was patients diagnosed with any diseases categorized under the term “Primary Immunologic Deficiency Syndrome”.

This study’s primary outcome was the consanguinity prevalence among patients’ parents. We combined the keywords and MeSH terms for “Primary Immunologic Deficiency Syndrome” and “consanguinity” using the term “AND”.

Study selection

Inclusion criteria were: (1) English-language studies (2) study designed as cross-sectional, prospective or retrospective cohort (3) data collected from national registries, national referral centers for immunodeficiency diseases, national surveys addressing clinical immunologists throughout the countries, or multiple centers (4) consanguinity prevalence

measured by any definition (5) obtaining data on patients with PIDs (6) PID diagnosed by a specialist (7) PID diagnosed by any diagnostic criteria

Exclusion criteria: (1) the presence of selection bias (2) overlap of the database or patients included in studies performed in a single country (3) not addressing PID patients, separately

A team of two medical doctors performed the screening process.

Data extraction and methodological quality

The data for each study were extracted using a data extraction sheet which included:

- Article's ID
- Original country of the study
- Study design
- Study's time span
- The population characteristics
- Parental consanguinity prevalence
- Possible biases

The extracted articles went under critical analysis using Munn et al. Critical Appraisal tool for systematic reviews addressing questions of prevalence. (8) The authors resolved disagreements by consensus. Eventually, studies were graded as "High", "Moderate", or "Low" quality, and then the low-quality articles were omitted.

Statistical analysis

We analyzed data using STATA 13.0 software (Stata-Crop. 2013. Stata Statistical Software: Release 13, College Station, TX). We used "metan" command to generate pooled prevalence and RevMan 5.3 software (9) to generate forest plots.

The included studies were stratified first according to:

1. Geographic region based on world health organization (WHO) classification (10):
 - The Americas
 - Europe
 - the Eastern Mediterranean
 - the Western Pacific
2. The most common types of PIDs in the included articles:
 - Common variable immunodeficiency (CVID)
 - Selective IgA Deficiency (SIgAD)
 - severe combined immunodeficiency (SCID)
 - agammaglobulinemia
 - IgG subclass deficiency (IgGSD)
 - chronic granulomatous disease (CGD)

- ataxia telangiectasia (AT)
- DiGeorge syndrome (DGS)
- Wiskott-Aldrich syndrome (WAS)
- hyper-IgE syndrome (HIES)
- leukocyte adhesion deficiency (LAD)
- Chediak-Higashi syndrome (CHS)

Consanguinity prevalence in population

We obtained the information about the prevalence of consanguinity in the general population (the denominator) from national data that most closely matched the community of the study on PID patients.

There were times when the study of PID patients was performed in a large population/country like China where the practicing of consanguinity varies in different provinces. In such cases, we weighted consanguinity prevalence in the population based on the same composition as PID patients.

Same as above, for one study consisting of data mainly from 8 countries in Europe (11), we weighted the consanguinity prevalence in each country.

Exploring heterogeneity

We estimated the summary of odds ratios using the Mantel-Haenszel method. To assess heterogeneity between studies, we used I^2 statistics (12) as well as the p-value from Cochran's Q (13). In cases of low heterogeneity, we performed the meta-analysis using fixed effect model (14). When there was moderate, substantial, or considerable heterogeneity (15), we used random effects model was used.

RESULTS

Figure 1 summarizes the literature search process. Figure 2 depicts the statistics of the 21 studies that passed the quality appraisal. The final 21 studies were published from

December 1985 to January 2016; data were collected since 1963 up until 2014. The sum of studies' sample sizes reached 24968; but, since there were times that we included more than one study from a country, we omitted the potential duplicate patients, so the number of individuals reached a total of 18091 patients diagnosed with different types of PIDs.

Eleven studies' data were extracted from national registries, six were done on patients from

referral centers for immunodeficiency diseases, three were conducted using national surveys addressing clinical immunologists throughout the countries, and one was done based on data from multiple centers.

The most common diagnostic criteria was based on The European Society for Immunodeficiencies (ESID) and The International Union of Immunological Societies (IUIS) (11,16–27).

Among all, only three studies were conducted on a particular type of PID (28–30). Overall male to female ratio was 1.44:1. Based on twelve studies, the mean age at diagnosis ranged from 12.0 to 74.5 months with considerable heterogeneity throughout the studies ($P<0.001$; $I^2=96.5\%$). The random-effects pooled mean age of patients at the time of diagnosis was 43.6 months (95% CI 36.9-50.4).

Table 1 shows the studies with their characteristics in details. The study with the lowest prevalence of parental consanguinity was from China with 2% (31), and the highest rated consanguinity was from a study from Oman, which was 81% (22).

To speculate the overall consanguinity prevalence in PID patients of each geographic region we meta-analyzed the consanguinity prevalence from articles that included all PID patients regardless of their subtype (we omitted three studies (28–30) which were on a particular PID subtype and four studies (18,23,26,27) that were from the same database as a larger study (11) to exclude the potential duplicates). Considering the high heterogeneity, the pooled consanguinity prevalence from the random-effect analysis was 65.7% ($P<0.001$; $I^2=87.6$, 95% CI 57.8-73.6) in PID patients of Eastern-Mediterranean region, 5.6% ($P<0.001$; $I^2=98.3$, 95% CI -0.8%-12%) in Europe, and 2.3% ($P: 0.712$; $I^2=0\%$, 95% CI 0.9%-3.7%) in the Western Pacific.

Our findings revealed that Eastern-Mediterranean communities tended to engage in consanguineous relationships widespread. Particularly, people of middle-eastern countries were on top of the list, with Oman and Saudi Arabia being the first (56.3% and 56.0%, respectively).

Overall, the range of odds ratios stretched from 0.6 (Turkey) -with a sample of mixed subtypes of PIDs, through 117 (Australia) -which shows the ratio in SCID patients. Among the studies with a sample of mixed subtypes of PID patients, the French study sat at highest with an odds ratio of 21.9.

Figure 3 illustrates the output of the meta-analysis of overall POR estimation. The pooled odds ratio (random-effects) was 3.02 ($P < 0.001$; $I^2 = 89.1\%$, 95% CI 2.46 - 3.70).

By going through the articles and looking for the prevalent subtypes in each study, the analysis team agreed on 12 most common diseases to be included in further investigations.

These specified diseases constitute 50% of all of our study patients. Figure S1 portrays the distribution of these subtypes and indicates their proportion in relevance to the whole.

Worth mentioning that CHD (POR=16.4, $P < 0.001$) was the disease with the highest odds ratio between all. Figure S2 depicts the disease categorized meta-analysis in detail.

The studies result had considerable heterogeneity in CVID, SCID, and IgGSD subgroup analysis, substantial heterogeneity in SigAD and agammaglobulinemia, moderate heterogeneity in CGD, DGS, WAS, and LAD, and finally no heterogeneity in AT, HIES, and CHS.

DISCUSSION

We reported the parental consanguinity prevalence in format of 3 WHO regions based on 18091 individuals. As expected, Parental consanguinity in PID patients was dramatically high in Eastern-Mediterranean region (65.7%), mostly due to high inbreeding rate in Middle East. When comparing Eastern-Mediterranean region with Europe and western pacific, we observe a drastic decrease in consanguinity prevalence (5.6%, 2.3% respectively). Regarding the similarities in ethnicity and social beliefs we did not expect a considerable difference in consanguinity prevalence in the two groups. However, the prevalence of parental consanguinity in the North America is based on the data from one study arising from a single national referral center in Mexico, and cannot be representative for the Americas.

We found that offspring of consanguineous marriages were more likely to suffer from a PID disease throughout life than the children with no parental consanguinity (Pooled POR: 3.0, 95% CI, 2.5-3.7). Although, by reporting odds ratio we tried to match variables that could contribute to the prevalence of parental consanguinity in PID patients (such as culture or geographic region)(32), we noticed a high degree of heterogeneity even in the pooled “odds ratio.”

After pooling the data at the level of each disease, our findings followed a notable scheme. All of the diseases with an AR pattern of inheritance had the highest prevalence odds ratios among all; and plus CGD, were the only conditions that reached a significant result. The other six diseases -although all with odds ratios greater than one- didn't approve as statistically significant. The latter PID diseases fall into various inheritance patterns: WAS (X-linked); DGS (autosomal dominant); SIgAD, IgGSD, Agammaglobulinemia and CVID (Variable) (33).

There is a prevailing mentality that inbreeding increases the incidence of autosomal recessive disorders (ARDs). The rationale behind such hypothesis is that the chance of homozygosity boosts in the offspring of inbred couples, thus expressing the deleterious alleles (16). Several studies reported that ARDs are more common in communities with high consanguinity prevalence (34–36). The majority of these studies are geographically limited to specific regions of the world with high inbreeding rates, and also cease at the level of reporting a difference in inbreeding rate rather than odds ratio (37).

We may declare with considerable confidence that there is a definitive high correlation between inbreeding and the likelihood of autosomal recessive PIDs. The odds ratios for AR PIDs were considerably higher than the mean pooled odds ratio for PIDs in general: SCID and AT (two-folds); LAD (three-folds), CHS (more than 5-folds). The only exception here is HIES. The pooled POR for this disease was 2.9 -close to the mean POR for PIDs. This observation may be understandable; as this particular disease has mainly a dual mechanism of inheritance based on the genotype: HIES may both follow an AR or an AD pattern (associated with DOCK-8 and STAT-3; respectively) (38,39). We may assume that our sample of HIES cases included a combination of both genotypes of the disease.

Among diseases with other inheritance patterns, CGD was the only one which resulted in a significantly high odds ratio. CGD is mostly inherited as an x-linked trait (CYBB), and only about one-third of cases would be the autosomal recessive variants (associated with CYBA, NCF1, NCF2, and NCF4) (40); however, these epidemiologic statistics come from western countries with low consanguinity prevalence. The data from high consanguine countries demonstrates the dominance of autosomal recessive cases regarding prevalence rates (41,42). Although the odds ratio reached statistical significance, POR of 2.7 was relatively

low compared to the other significant odds ratios in ARDs which can be explained by the duality of CGD inheritance pattern.

Our results also confirm the evidence that suggests that impact of consanguinity on monogenic disorders is disproportionate to the prevalence of the disorder in the population (43). As depicted in S2, the highest odds ratio belongs to CHS -the rarest disease in our study- followed by LAD.

There was significant heterogeneity in the prevalence of consanguinity, and also in the prevalence odds ratios (POR). Apart from geographic and ethnic diversities, the use of different definitions of consanguinity by the studies and variations among registration systems are possible explanations. We also noted variations in the PORs among PIDs with AR inheritance pattern (such as CHD, LAD or SCID). Different genetic background in different ethnicities explains a part of these variations. For example, it seems that there are common ancestors with same mutations for RAG1 and RAG2 in some regions of Middle East. Al-Herz et al (44) demonstrated that deficiencies in RAG1, RAG2, and MHC II (AR traits) are the main origin of CID in Middle Eastern nations like Kuwait; whereas in western countries, the x-linked SCID is more prevalent due to common gamma chain (γ c) deficiency. It is competent to put this finding down to consanguinity being more common in the Middle East. We should also mention the fact that not all the PID cases have been registered in the studies. The lack of perfect registration systems especially leads to discrepancies among the ORs of the rarest PIDs - such as the AR inherited immunodeficiencies.

The estimated pooled mean of age at onset in this study was 43.6 months (95% CI 36.9-50.4). Although data of age at diagnosis were incomplete, it did not show any trend among different geographic regions. Previously Gathmann et al (11) reported that diagnostic delay

(time period between the onset of disease and the diagnosis) has been decreased for IgG subclass deficiency and agammaglobulinemia from 1987 to 2010. Longitudinal studies with large sample sizes are needed to investigate the trend of diagnostic delay through time and geographic regions.

To the authors' knowledge, this is the first systematic review and meta-analysis assessing parental consanguinity in PID patients, and in the matched general population. We hope that Genetic counselors, health educators, and consanguineous couples seeking preconception counseling, all over the world, benefit from the results of this systematic review and meta-analysis.

A limitation of the present meta-analysis was that most of the included articles did not define "consanguinity", and even those who defined were not using a unique definition. So, the prevalence of parental consanguinity and the odds ratios may have been underestimated or overestimated. Probable imperfections of data registry systems can be the other important limitation of this review.

With this systematic review we manifested that odds of consanguinity are three folds in parents of PID patients compared to the normal healthy population. We also benefited from meta-analyzing the available data and displayed the magnitude of inbreeding influence on the 12 globally most common PID diseases. The analysis revealed that impact of consanguinity only kicks in when dealing with autosomal recessive PID diseases -or those with a dominant autosomal recessive side.

Acknowledgements

We would like to thank Seyed-Mohammad Fereshtehnejad and Somayye Barati for comments and statistical advice that significantly improved the manuscript.

References

1. de Vries E. Patient-centred screening for primary immunodeficiency: a multi-stage diagnostic protocol designed for non-immunologists. *Clin Exp Immunol* 2006; **145**:204–214.
2. Arkwright PD, Gennery AR. Ten warning signs of primary immunodeficiency: a new paradigm is needed for the 21st century. *Ann N Y Acad Sci* 2011; **1238**:7–14.
3. Picard C, Al-Herz W, Bousfiha A et al. Primary Immunodeficiency Diseases: an Update on the Classification from the International Union of Immunological Societies Expert Committee for Primary Immunodeficiency 2015. *J Clin Immunol* 2015; **35**:696–726.
4. Port K, Bittles AH. A population-based estimate of the prevalence of consanguineous marriage in Western Australia. *Public Health Genomics* 2001; **4**:97–101.
5. Bittles AH, Mason WM, Greene J et al. Reproductive behavior and health in consanguineous marriages. *Science* 1991; **252**:789–794.
6. Bennett RL, Motulsky AG, Bittles A et al. Genetic Counseling and Screening of Consanguineous Couples and Their Offspring: Recommendations of the National Society of Genetic Counselors. *J Genet Couns* 2002; **11**:97–119.
7. Moher D, Liberati A, Tetzlaff J et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med* 2009; **151**:264–269.
8. Munn Z, Moola S, Riitano D et al. The development of a critical appraisal tool for use in systematic reviews addressing questions of prevalence. *Int J Heal policy Manag* 2014; **3**:123–128.
9. Review Manager (RevMan). 2014.
10. Page. WHO regional offices, 2016. <http://www.who.int/about/regions/en/> (accessed 7 May2016).
11. Gathmann B, Binder N, Ehl S et al. The European internet-based patient and research database for primary immunodeficiencies: update 2011. *Clin Exp Immunol* 2012; **167**:479–491.
12. Higgins JPT, Thompson SG, Deeks JJ et al. Measuring inconsistency in meta-analyses. *Bmj* 2003; **327**:557–560.
13. Higgins J, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002; **21**:1539–1558.
14. Hedges L V, Vevea JL. Fixed-and random-effects models in meta-analysis. *Psychol Methods* 1998; **3**:486.
15. Page. 9.5.2 Identifying and measuring heterogeneity. http://handbook.cochrane.org/chapter_9/9_5_2_identifying_and_measuring_heterogeneity.htm (accessed 26 May2016).

- Accepted Article
16. Lugo Reyes SO, Ramirez-Vazquez G, Cruz Hernández A et al. Clinical Features, Non-Infectious Manifestations and Survival Analysis of 161 Children with Primary Immunodeficiency in Mexico: A Single Center Experience Over two Decades. *J Clin Immunol* 2016; **36**:56–65.
 17. Ehlayel M, Bener A, Laban MA. Effects of family history and consanguinity in primary immunodeficiency diseases in children in Qatar. *Open J Immunol* 2013; **3**:47–53.
 18. Gathmann B, Goldacker S, Klima M et al. The German national registry for primary immunodeficiencies (PID). *Clin Exp Immunol* 2013; **173**:372–380.
 19. Marschall K, Hoernes M, Bitzenhofer-Grüber M et al. The Swiss National Registry for Primary Immunodeficiencies: report on the first 6 years' activity from 2008 to 2014. *Clin Exp Immunol* 2015; **182**:45–50.
 20. Mellouli F, Mustapha I Ben, Khaled M Ben et al. Report of the Tunisian Registry of Primary Immunodeficiencies: 25-Years of Experience (1988-2012). *J Clin Immunol* 2015; **35**:745–753.
 21. Bousfiha AA, Jeddane L, El Hafidi N et al. First report on the Moroccan registry of primary immunodeficiencies: 15 years of experience (1998-2012). *J Clin Immunol* 2014; **34**:459–468.
 22. Al-Tamemi S, Elnour I, Dennison D. Primary immunodeficiency diseases in oman: five years' experience at sultan qaboos university hospital. *World Allergy Organ J* 2012; **5**:52–56.
 23. CEREDIH. The French national registry of primary immunodeficiency diseases. *Clin Immunol* 2010; **135**:264–272.
 24. Al-Herz W, Moussa MAA. Survival and predictors of death among primary immunodeficient patients: a registry-based study. *J Clin Immunol* 2012; **32**:467–473.
 25. Aghamohammadi A, Mohammadinejad P, Abolhassani H et al. Primary immunodeficiency disorders in Iran: update and new insights from the third report of the national registry. *J Clin Immunol* 2014; **34**:478–490.
 26. Kilic SS, Ozel M, Hafizoglu D et al. The prevalences [correction] and patient characteristics of primary immunodeficiency diseases in Turkey--two centers study. *J Clin Immunol* 2013; **33**:74–83.
 27. Edgar JDM, Buckland M, Guzman D et al. The United Kingdom Primary Immune Deficiency (UKPID) Registry: report of the first 4 years' activity 2008-2012. *Clin Exp Immunol* 2014; **175**:68–78.
 28. Yee A, De Ravin SS, Elliott E et al. Severe combined immunodeficiency: a national surveillance study. *Pediatr Allergy Immunol* 2008; **19**:298–302.
 29. Hayakawa H, Kobayashi N, Yata J. Chronic Granulomatous Disease in Japan: A Summary of the Clinical Features of 84 Registered Patients. *Pediatr Int* 1985; **27**:501–510.
 30. Nagai K, Ochi F, Terui K et al. Clinical characteristics and outcomes of chédiak-Higashi syndrome: a nationwide survey of Japan. *Pediatr Blood Cancer* 2013; **60**:1582–1586.

31. Wang L-L, Jin Y-Y, Hao Y-Q et al. Distribution and clinical features of primary immunodeficiency diseases in Chinese children (2004-2009). *J Clin Immunol* 2011; **31**:297–308.
32. Bittles AH. A community genetics perspective on consanguineous marriage. *Community Genet* 2008; **11**:324–330.
33. Al-Herz W, Bousfiha A, Casanova J-L et al. Primary immunodeficiency diseases: an update on the classification from the international union of immunological societies expert committee for primary immunodeficiency. *Front Immunol* 2011; **2**:54.
34. Jaouad IC, Elalaoui SC, Sbiti A et al. Consanguineous marriages in Morocco and the consequence for the incidence of autosomal recessive disorders. *J Biosoc Sci* 2009; **41**:575–581.
35. Bittles AH, Hamamy HA. Genetic Disorders Among Arab Populations. In: Teebi SA, ed. . Berlin, Heidelberg: Springer Berlin Heidelberg 2010: 85–108.
36. Romdhane L, Abdelhak S. Genetic diseases in the Tunisian population. *Am J Med Genet A* 2011; **155A**:238–267.
37. Kelmemei W, Chelly I, Kharrat M et al. Consanguinity and homozygosity among Tunisian patients with an autosomal recessive disorder. *J Biosoc Sci* 2015; **47**:718–726.
38. Ben Arab S, Masmoudi S, Beltaief N et al. Consanguinity and endogamy in Northern Tunisia and its impact on non-syndromic deafness. *Genet Epidemiol* 2004; **27**:74–79.
39. Kumaramanickavel G, Joseph B, Vidhya A et al. Consanguinity and ocular genetic diseases in South India: analysis of a five-year study. *Community Genet* 2002; **5**:182–185.
40. Barbouche M-R, Galal N, Ben-Mustapha I et al. Primary immunodeficiencies in highly consanguineous North African populations. *Ann N Y Acad Sci* 2011; **1238**:42–52.
41. Hamamy HA, Masri AT, Al-Hadidy AM et al. Consanguinity and genetic disorders. Profile from Jordan. *Saudi Med J* 2007; **28**:1015–1017.
42. Zlotogora J. Genetic disorders among Palestinian Arabs: 1. Effects of consanguinity. *Am J Med Genet* 1997; **68**:472–475.
43. Freire-Maia N. Inbreeding levels in American and Canadian populations: a comparison with Latin America. *Eugen Q* 1968; **15**:22–33.
44. Al-Herz W, Notarangelo LD, Sadek A et al. Combined immunodeficiency in the United States and Kuwait: Comparison of patients' characteristics and molecular diagnosis. *Clin Immunol* 2015; **161**:170–173.
45. Sutter J, Goux J-M. Decline of consanguineous marriages in France from 1926 to 1958. *Eugen Q* 1964; **11**:127–140.
46. Koc I. Prevalence and sociodemographic correlates of consanguineous marriages in Turkey. *J*

Biosoc Sci 2008: **40**:137–148.

47. Fraccaro M. Consanguineous marriages in Italy: a note. *Eugen Q* 1957: **4**:36–39.
48. Freire-Maia N. Inbreeding levels in different countries. *Eugen Q* 1957: **4**:127–138.
49. Masterson JG. Cousin marriage in Ireland. *Genet Var Britain* 1973: :59–66.
50. Pinto-Cisternas J, Zei G, Moroni A. Consanguinity in Spain, 1911-1943: general methodology, behavior of demographic variables, and regional differences. *Soc Biol* 1979: **26**:55–71.
51. Coleman DA. A Note on the Frequency of Consanguineous Marriages in Reading, England in 1972/1973. *Hum Hered* 1980: **30**:278–285.
52. Reda SM, Afifi HM, Amine MM. Primary immunodeficiency diseases in Egyptian children: a single-center study. *J Clin Immunol* 2009: **29**:343–351.
53. Hafez M, El-Tahan H, Awadalla M et al. Consanguineous matings in the Egyptian population. *J Med Genet* 1983: **20**:58–60.
54. Rezaei N, Pourpak Z, Aghamohammadi A et al. Consanguinity in primary immunodeficiency disorders; the report from Iranian Primary Immunodeficiency Registry. *Am J Reprod Immunol* 2006: **56**:145–151.
55. Saadat M, Ansari-Lari M, Farhud DD. Consanguineous marriage in Iran. *Ann Hum Biol*: **31**:263–269.
56. Al-Awadi SA, Naguib KK, Moussa MA et al. The effect of consanguineous marriages on reproductive wastage. *Clin Genet* 1986: **29**:384–388.
57. Lamdouar Bouazzaoui L, MOZZICONACCI P, BLANCHER G et al. Consanguinit{é} et sant{é} publique au Maroc. Discussion. *Bull l'Acad{é}mie Natl m{é}decine* 1994: **178**:1013–1027.
58. Rajab A, Patton MA. A study of consanguinity in the Sultanate of Oman. *Ann Hum Biol*: **27**:321–326.
59. Bener A, Alali KA. Consanguineous marriage in a newly developed country: the Qatari population. *J Biosoc Sci* 2006: **38**:239–246.
60. Al-Saud B, Al-Mousa H, Al Gazlan S et al. Primary Immunodeficiency Diseases in Saudi Arabia: a Tertiary Care Hospital Experience over a Period of Three Years (2010-2013). *J Clin Immunol* 2015: **35**:651–660.
61. El-Mouzan MI, Al-Salloum AA, Al-Herbish AS et al. Regional variations in the prevalence of consanguinity in Saudi Arabia. *Saudi Med J* 2007: **28**:1881–1884.
62. Riou S, el Younsi C, Chaabouni H. [Consanguinity in the population of northern Tunisia]. *La Tunisie médicale* 1989: **67**:167–172.
63. Wang L-L, Jin Y-Y, Hao Y-Q et al. Distribution and clinical features of primary immunodeficiency diseases in Chinese children (2004-2009). *J Clin Immunol* 2011: **31**:297–

308.

64. Wu L. Investigation of consanguineous marriages among 30 Chinese ethnic groups. *Hered Dis* 1987; **4**:163–166.
65. Zhan J, Qin W, Zhou Y et al. Effects of consanguineous marriages on hereditary diseases: a study of the Han ethnic group in different geographic districts of Zejiang province. *Nat Med J China* 1992; **172**:674–676.
66. Lee PP-W, Lau Y-L. Improving care, education, and research: the Asian primary immunodeficiency network. *Ann N Y Acad Sci* 2011; **1238**:33–41.
67. Imaizumi Y. A recent survey of consanguineous marriages in Japan. *Clin Genet* 1986; **30**:230–233.




Figure 1. Flow diagram of the review process



Figure 2. Results of the methodological quality item assessment presented as percentage among all included studies




Figure 3. Estimated prevalence odds ratio (POR) of parental consanguinity in patients with primary immunodeficiency diseases

Table 1: Detailed characteristics of the studies included in the systematic review

Study	Country	Year	N ^a	Type	Quality	Study CP ^b	Population CP	POR ^c (95% CI)
the Americas								
Reyes et al,(16) 2015	Mexico	1991-2012	161	Mixed	High	11.2%	1.27% (43)	9.79 (5.93-16.17)
Europe								
Marschall et al,(19) 2015	Swiss	2008-2014	348	Mixed	High	2.3%	N/A	N/A
CEREDIH, (23) 2010	France	2005-2009	3083	Mixed	High	15%	0.8% (45)	21.86 (19.71-24.24)
Gathmann et al, (18) 2013	Germany	1963-2010	917	Mixed	High	8.6%	N/A	N/A
Kilic et al, (26) 2013	Turkey	2004-2010	1130	Mixed	High	14.3%	22% (46)	0.59 (0.49-0.70)
Gathmann et al, (11) 2012	Europe	Up to 2011	13708	Mixed	High	8.8%	4.8% (45–50)	1.92 (1.72-2.16)
Edgar et al, (27) 2014	The UK ^d	2008- 2012	924	Mixed	High	2.9%	0.4% (51)	6.91 (2.43-19.67)
Eastern Mediterranean								
Reda et al, (52) 2009	Egypt	2004-2008	64	Mixed	Mod ^e	62.5%	28.96% (53)	4.09 (2.46-6.78)
Rezaei et al, (54) 2006	Iran	1980-2006	930	Mixed	High	68.5%	38.6% (55)	3.46 (3.01-3.97)

Aghamohammadi et al, (25) 2014	Iran	2006-2013	731	Mixed	High	63.1%	38.6% (55)	2.72 (2.34-3.16)
Al-Herz et al, (24) 2012	Kuwait	2004-2011	176	Mixed	High	78%	54.3% (56)	3.06 (2.12-4.39)
Bousfiha et al, (21) 2014	Morocco	1998-2012	421	Mixed	Mod	43.2%	19.9% (57)	3.06 (2.50-3.76)
Al-Tamemi et al, (22) 2012	Oman	2005-2010	90	Mixed	Mod	81%	56.3% (58)	3.33 (2.50-3.76)
Ehlayel et al, (17) 2013	Qatar	1998-2012	131	Mixed	Mod	61.1%	54% (59)	2.07 (1.36-3.14)
Al-saud et al, (60) 2015	KSA ^f	2010-2013	502	Mixed	High	75%	56% (61)	2.34 (1.91-2.88)
Mellouli et al, (20) 2015	Tunisia	2002-2003	710	Mixed	High	58.2%	26.9% (62)	3.78 (3.22-4.44)
Western Pacific								
Yee et al, (28) 2008	Australia	1995-2001	33	SCID ^g	High	22%	0.23% (4)	116.68 (49.85-273.10)
Wang et al, (63) 2011	China	2004-2009	195	Mixed	Mod	2%	1.9% (64,65)	1.06 (0.39-2.88)
Lee et al, (66) 2011	China	2001-2011	272	Mixed	Mod	<3%	1.9% (64,65)	1.37 (0.64-2.92)
Hayakawa et al, (29) 1985	Japan	Up to 1984	84	CGD ^h	Mod	3.6%	3.9% (67)	0.91 (0.29-2.90)
Nagai et al, (30) 2013	Japan	2000-2010	15	CHS ⁱ	High	13%	3.9% (67)	3.79 (0.85-16.85)

^a Sample Size ^b Consanguinity prevalence ^c prevalence odds ratio ^d The United Kingdom ^e Moderate ^f Kingdom of Saudi Arabia ^g Severe Combined Immunodeficiency ^h Chronic Granulomatous Disease ⁱ Chediak Higashi Syndrome




Figure S1. Distribution of the 12 most prevalent primary immunodeficiency subtypes included in the additional meta-analysis, (CVID: Common Variable immunodeficiency; SIgAD: Selective IgA Deficiency; SCID: Severe Combined Immunodeficiency; IgGSD: IgG Subclass Deficiency; CGD: Chronic Granulomatous Disease; AT: Ataxia Telangectasia; DGS: DiGeorge Syndrome; WAS: Wiskott-Aldrich Syndrome; HIES: Hyper-IgE Syndrome; LAD: Leukocyte Adhesion Deficiency; CHS: Chediak-Higashi Syndrome)