

Biologics for Treatment of Atopic Dermatitis: Current Status and Future Prospect



Thanaporn Ratchataswan, MD^a, Tina M. Banzon, MD^{a,b}, Jacob P. Thyssen, MD, PhD, DmSci^c,
Stephan Weidinger, MD, PhD, MaHM^d, Emma Guttman-Yassky, MD, PhD^e, and Wanda Phipatanakul, MD, MS^{a,b} *Boston, Mass, Copenhagen, Denmark; Kiel, Germany; and New York, NY*

INFORMATION FOR CATEGORY 1 CME CREDIT

Credit can now be obtained, free for a limited time, by reading the review articles in this issue. Please note the following instructions.

Method of Physician Participation in Learning Process: The core material for these activities can be read in this issue of the Journal or online at the *JACI: In Practice* Web site: www.jaci-inpractice.org/. The accompanying tests may only be submitted online at www.jaci-inpractice.org/. Fax or other copies will not be accepted.

Date of Original Release: March 1, 2021. Credit may be obtained for these courses until February 28, 2022.

Copyright Statement: Copyright © 2021-2023. All rights reserved.

Overall Purpose/Goal: To provide excellent reviews on key aspects of allergic disease to those who research, treat, or manage allergic disease.

Target Audience: Physicians and researchers within the field of allergic disease.

Accreditation/Provider Statements and Credit Designation: The American Academy of Allergy, Asthma & Immunology (AAAAI) is accredited by the Accreditation Council for Continuing Medical Education (ACCME) to provide continuing medical education for physicians. The AAAAI designates this journal-based CME activity for 1.00 *AMA PRA Category 1 Credit*TM. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

List of Design Committee Members: Thanaporn Ratchataswan, MD, Tina M. Banzon, MD, Jacob P. Thyssen, MD, PhD, DmSci, Stephan Weidinger, MD, PhD, MaHM, Emma Guttman-Yassky, MD, PhD, and Wanda Phipatanakul, MD, MS (authors); Scott H. Sicherer, MD (editor)

Learning objectives:

1. To discuss clinical efficacy and limitations of approved and new biologics, including biomarkers for atopic dermatitis (AD) from recent clinical trials results.

2. To identify the mechanism of action of approved and new biologic drugs for AD.

Recognition of Commercial Support: This CME has not received external commercial support.

Disclosure of Relevant Financial Relationships with Commercial Interests: T. Ratchataswan declares no relevant conflicts of interest. T. M. Banzon is supported by T32 AI 007512 from the National Institutes of Health. J. P. Thyssen has been an advisor/investigator/speaker for Regeneron, Sanofi-Genzyme, AbbVie, LEO Pharma, Eli Lilly & Co, and Pfizer. S. Weidinger has received institutional research grants from Sanofi Deutschland GmbH, Leo Pharma, and La Roche Posay; has performed consultancies for Sanofi-Genzyme, Regeneron, LEO Pharma, AbbVie, Pfizer, Eli Lilly, Kymab, and Novartis; has lectured at educational events sponsored by Sanofi-Genzyme, Regeneron, LEO Pharma, AbbVie, Novartis, and Galderma; and is involved in performing clinical trials with many pharmaceutical industries that manufacture drugs used for the treatment of psoriasis and atopic dermatitis. E. Guttman-Yassky is an employee of Mount Sinai and has received research funds (grants paid to the institution) from AbbVie, Almirall, Amgen, AnaptysBio, Asana Biosciences, Boehringer-Ingelheim, Celgene, Dermavant, DS Biopharma, Eli Lilly, Galderma, Ichnos Sciences, Innovaderm, Janssen, Kiniska, Kyowa Kirin, Leo Pharma, Novan, Pfizer, Ralexar, Regeneron Pharmaceuticals, Inc, Sienna Biopharma, UCB, and Union Therapeutics; and is a consultant for AbbVie, Aditum Bio, Almirall, Amgen, Asana Biosciences, AstraZeneca, Boehringer-Ingelheim, Cara Therapeutics, Celgene, Concert, DBV, Dermira, DS Biopharma, Eli Lilly, EMD Serono, Escalier, Galderma, Ichnos Sciences, Incyte Kyowa Kirin, Leo Pharma, Mitsubishi Tanabe, Pandion Therapeutics, Pfizer, RAPT Therapeutics, Regeneron Pharmaceuticals, Inc, Sanofi, Sienna Biopharma, Target PharmaSolutions, and Union Therapeutics. W. Phipatanakul is supported by the grants U01 AI 152033 and K24 AI 106822 from the National Institutes of Health and Allergy Asthma Awareness Initiative, Inc; and reports consultancy fees from Genentech, Novartis, Regeneron, Sanofi, GSK, and Astra Zeneca for therapeutics related to Asthma, outside the submitted work. S.H. Sicherer declares no relevant conflicts of interest.

^aDepartment of Allergy and Immunology, Boston Children's Hospital, Boston, Mass

^bDepartment of Pediatrics, Harvard Medical School, Boston, Mass

^cDepartment of Dermatology and Allergy, Herlev and Gentofte Hospital, University of Copenhagen, Copenhagen, Denmark

^dDepartment of Dermatology and Allergy, University Hospital Schleswig-Holstein, Campus Kiel, Kiel, Germany

^eDepartment of Dermatology and Laboratory of Inflammatory Skin Diseases, Icahn School of Medicine at Mount Sinai, New York, NY

No funding was received for this work.

Conflicts of interest: T. Ratchataswan declares no relevant conflicts of interest. T. M. Banzon is supported by T32 AI 007512 from the National Institutes of Health. J.

P. Thyssen has been an advisor/investigator/speaker for Regeneron, Sanofi-Genzyme, AbbVie, LEO Pharma, Eli Lilly & Co, and Pfizer. S. Weidinger has received institutional research grants from Sanofi Deutschland GmbH, Leo Pharma, and La Roche Posay; has performed consultancies for Sanofi-Genzyme, Regeneron, LEO Pharma, AbbVie, Pfizer, Eli Lilly, Kymab, and Novartis; has lectured at educational events sponsored by Sanofi Genzyme, Regeneron, LEO Pharma, AbbVie, Novartis, and Galderma; and is involved in performing clinical trials with many pharmaceutical industries that manufacture drugs used for the treatment of psoriasis and atopic dermatitis. E. Guttman-Yassky is an employee of Mount Sinai and has received research funds (grants paid to the institution) from AbbVie, Almirall, Amgen, AnaptysBio, Asana Biosciences, Boehringer

Abbreviations used

ACD- Allergic contact dermatitis
 AD- Atopic dermatitis
 AE- Adverse event
 DLQI- Dermatology Life Quality Index
 DPP-4- Dipeptidyl peptidase-4
 EASI- Eczema Area and Severity Index
 EASI-50- Greater than 50% improvement in EASI score
 EASI-75- Greater than 75% improvement in EASI score
 EASI-90- Greater than 90% improvement in EASI score
 EMA- European Medicines Agency
 FDA- Food and Drug Administration
 IGA- Investigator's Global Assessment
 ITT- Intended to treat
 IV- Intravenous
 mAb- Monoclonal antibody
 NRS- Numeric rating scale
 SC- Subcutaneously
 SCORAD- Scoring of Atopic Dermatitis
 SCORAD-50- Scoring of Atopic Dermatitis improvements of $\geq 50\%$
 TARC- Thymus- and activation-regulated chemokine
 TCS- Topical corticosteroid
 TEAE- Treatment emergent adverse event
 Th- T-helper
 TSLP- Thymic stromal lymphopoietin
 VAS- Visual analog scale

Atopic dermatitis (AD) is a common inflammatory skin disease characterized by intense pruritus and recurrent eczematous lesions that significantly impair quality of life. It is a heterogeneous disease affecting both children and adults. The treatment of moderate-to-severe forms of AD is challenging, as topical corticosteroids are often insufficient to achieve disease control or inappropriate and off-label use of immunosuppressants may have significant undesirable side effects. The development of targeted biologic therapies specifically for AD is thus highly desirable. Dupilumab is the only biologic therapy that is Food and Drug Administration approved for the treatment of moderate-to-severe AD in patients 6 years and older, with consistent long-term efficacy and safety trial data. In this article, we review the mechanisms, safety, and efficacy of dupilumab from recent clinical trials, and we review the current data, mechanism of action, clinical efficacy, and limitations of new biologics currently in phase 2 and 3 clinical trials (lebrikizumab, tralokinumab, nemolizumab, tezepelumab, and ISB 830). © 2020 American Academy of Allergy, Asthma & Immunology (J Allergy Clin Immunol Pract 2021;9:1053-65)

Key words: Atopic dermatitis; Biologic therapy; Dupilumab; Eczema; GBR 830; ISB 830; Lebrikizumab; Nemolizumab; Tezepelumab; Tralokinumab

Atopic dermatitis (AD) is one of the most common chronic inflammatory skin diseases. Describing its incidence and prevalence is challenging due to substantial variation in diagnostic criteria used for its identification, fluctuating course, and geographical differences. The majority of cases start in early childhood, but AD can have heterogeneous trajectory patterns, ranging from transient disease in early life, to relapsing remitting AD, chronic persistent AD, and long periods of remission followed by recurrence, or adult-onset AD.¹⁻³ Although cumulative lifetime prevalence up to 30% has been reported, period prevalences are estimated to be 7% to 14% in children and 5% to 12% in adults.¹⁻³

The clinical hallmarks of AD are eczematous lesions and intense itch. Many patients suffer from dry scaly skin and IgE-mediated sensitizations. Lesions can affect any part of the body but typically show age-related morphology and distribution, with face, trunk, and extensor limb inflammatory involvement in infants and young children, and lichenified, chronic, dry flexural distribution in adults.⁴ However, compared with childhood-onset AD, presentation of adult-onset AD is more heterogeneous, with more variation in lesional morphology and distribution, and greater predilection for the head, neck, hands, and feet.⁵

The pathophysiology of AD is complex and multifactorial, with strong genetic susceptibility, important contributions of immune dysregulation due to excessive T-helper (Th)2 and Th22 activity, with variable contributions of Th1/Th17, skin barrier dysfunction, and cutaneous dysbiosis (Figure 1).⁶⁻¹⁰ Skin barrier dysfunction is characterized by downregulation of epidermal barrier proteins, including filaggrin, keratins, locrin, involucrin, and cell adhesion molecules, a disturbed intercellular lipid composition, and an altered composition of the skin microbiome, which together lead to increased permeability and proinflammatory signaling. Cutaneous inflammation is characterized by upregulated type 2 cytokines, such as IL-4, IL-13, and IL-31, heightening sensitization to allergens, risk of food allergy, airway hyperreactivity, and contributing to skin barrier dysfunction.¹¹⁻¹⁴

"The Th22 pathway has also been shown to be involved in AD, with IL-22 participating in epidermal disorders. By attenuating keratinocyte terminal differentiation and inhibiting tight-junction formation, IL-22 is thought to contribute to barrier dysfunction."^{15,16} However, there is increasing evidence that AD is not dominated by 1 disease spectrum, but rather involves multiple alternating immune pathways comprised by several endotypes.^{9,17-19} Several subtypes, such as intrinsic, Asian,

Ingelheim, Celgene, Dermavant, DS Biopharma, Eli Lilly, Galderma, Ichnos Sciences, Innovaderm, Janssen, Kiniska, Kyowa Kirin, Leo Pharma, Novan, Pfizer, Ralexar, Regeneron Pharmaceuticals, Inc, Sienna Biopharma, UCB, and Union Therapeutics; and is a consultant for AbbVie, Aditum Bio, Almirall, Amgen, Asana Biosciences, AstraZeneca, Boehringer-Ingelheim, Cara Therapeutics, Celgene, Concert, DBV, Dermira, DS Biopharma, Eli Lilly, EMD Serono, Escalier, Galderma, Ichnos Sciences, Incyte Kyowa Kirin, Leo Pharma, Mitsubishi Tanabe, Pandion Therapeutics, Pfizer, RAPT Therapeutics, Regeneron Pharmaceuticals, Inc, Sanofi, Sienna Biopharma, Target PharmaSolutions, and Union Therapeutics. W. Phipatanakul is supported by the grants U01 AI 152033 and K24 AI 106822 from the National Institutes of Health and Allergy Asthma Awareness Initiative, Inc; and reports consultancy fees from Genentech, Novartis,

Regeneron, Sanofi, GSK, and Astra Zeneca for therapeutics related to Asthma, outside the submitted work.

Received for publication September 14, 2020; revised November 18, 2020; accepted for publication November 24, 2020.

Corresponding author: Wanda Phipatanakul, MD, MS, Department of Allergy and Immunology, Boston Children's Hospital, 300 Longwood Avenue, Fegan, 6th Floor, Boston, MA 02115. E-mail: Wanda.Phipatanakul@childrens.harvard.edu. 2213-2198

© 2020 American Academy of Allergy, Asthma & Immunology

<https://doi.org/10.1016/j.jaip.2020.11.034>




FIGURE 1. Atopic dermatitis pathogenesis and targets of biologics approved and in clinical development for atopic dermatitis. DC, Dendritic cell; IFN, interferon; IgE, immunoglobulin E; IL, interleukin; ILC, innate lymphoid cell; LC, langerhans cell; Th, T-helper; TNF, tumour necrosis factor; TSLP, thymic stromal lymphopoietin.

pediatric, and filaggrin-positive AD subcategories, were shown to have differential upregulation in Th17/Th22 or Th1 axes. Thus, Th1 and Th17/Th22 modulation in addition to Th2 might provide broader and/or more sustained therapeutic benefit.¹⁰

AD is associated with numerous atopic comorbidities, including asthma, allergic rhinitis, food allergy, and eosinophilic esophagitis.²⁰ Increasing evidence supports the association of AD with other systemic inflammatory diseases, including alopecia areata, vitiligo, rheumatoid arthritis, and inflammatory bowel disease,²¹⁻²⁵ although well-powered longitudinal studies are needed to confirm causal links. A systematic review of 74 studies found no significant overall association between AD and allergic contact dermatitis (ACD), and in children referred for patch testing, ACD was more common in those without AD.^{26,27} It was also shown that ACD in the setting of AD has attenuated responses.²⁸ Furthermore, patients with AD have increased risk of psychiatric conditions, including attention-deficit hyperactivity disorder, anxiety, and depression.^{20,29}

The aim of AD management is to improve symptoms and achieve long-term eczema control with a multistep approach tailored according to disease severity (Figure 2). For all patients, basic management consists of continuous epidermal barrier repair with emollients and avoidance of individual triggers. Oral antihistamines are not recommended, as little evidence supports that they are effective for AD signs and symptoms, including pruritus.⁴

In moderate-to-severe AD, topical corticosteroids (TCSs) are the mainstay of treatment. However, patients with more severe AD often require treatment with systemic anti-inflammatory drugs, which may lead to significant side effects and hence treatment cessation, and efficacy is also moderate. Therefore, targeted

biologics, large proteins that are injected and do not penetrate the lipid bilayer cell membrane, involving pathways directly responsible for AD are an attractive treatment. Dupilumab is the only biologic approved by the United States Food and Drug Administration (FDA) and European Medicines Agency (EMA) for moderate-to-severe AD.^{30,31} Although dupilumab shows good efficacy, only approximately one-third of patients have complete clearance.³² There is thus a large need for further innovative therapeutics, including other biologics selectively targeting cytokines involved in the inflammatory pathway of AD.⁹

The purpose of this review is to discuss the mechanisms, safety, and efficacy of biologics recently approved and currently in phase 2 and 3 clinical trials for treatment of moderate-to-severe AD (Table I). In addition to biologics, other various targeted therapies, including JAK inhibitors and phosphodiesterase 4 inhibitors, are also in the pipeline for AD treatment at different stages of clinical trial.

DUPILUMAB

Dupilumab is a human IgG4K monoclonal antibody (mAb) blocking IL-4 receptor (IL-4Rα), a shared receptor for IL-4 and IL-13,³³ key cytokines in Th2-mediated inflammation playing major roles in AD pathogenesis. Dupilumab was first approved by the FDA in 2017 and EMA in 2019 for use in moderate-to-severe AD in adults and adolescents (≥12 years) and recently approved in pediatric patients aged 6 to 11 years.^{30,31,34} In 2018, dupilumab was approved in the United States and Europe as add-on maintenance treatment in patients with moderate-to-severe asthma with eosinophilic phenotype or oral corticosteroid-dependent asthma,^{31,35} and was approved in 2019 for chronic rhinosinusitis with nasal




FIGURE 2. Step-care management of atopic dermatitis (AD). Acute and maintenance treatments for AD across the spectrum of disease severity. ^aFor patients aged ≥ 2 years with mild-to-moderate AD. ^bFor patients aged ≥ 6 years with moderate-to-severe AD. ^cNot approved by the Food and Drug Administration to treat AD. ^dNot recommended for long-term maintenance. TCI, Topical calcineurin inhibitor; TCS, topical corticosteroid. Adapted from Fishbein et al.⁸¹

polyposis.³⁶ Beneficial effects of dupilumab were also reported in patients with eosinophilic esophagitis.³⁷

Efficacy and safety of dupilumab for treatment of AD has been studied in several phase 3 clinical trials, including SOLO-1, SOLO-2,³² LIBERTY AD CHRONOS,³⁸ and AD-1526.³⁹ In SOLO-1 and SOLO-2, dupilumab was used as monotherapy in adults with moderate-to-severe AD inadequately controlled on topical therapies. Patients were randomized to 2 dupilumab arms and placebo, with a loading dose of 600 mg subcutaneously (SC) followed by 300 mg SC weekly, or a loading dose of 600 mg SC followed by 300 mg SC every other week, or placebo once weekly.³² Dupilumab achieved significant improvement in the primary endpoints, which was Investigator's Global Assessment (IGA) score of 0/1 and improvement of ≥ 2 points in IGA from baseline score at week 16. In SOLO-1, the primary outcome was achieved in 37% receiving weekly dupilumab and 38% receiving dupilumab every other week, versus 10% on placebo (Figure 3, A). Results from SOLO-2 were similar. Both SOLO-1 and SOLO-2 demonstrated a higher proportion of patients receiving dupilumab achieving EASI-75 (Eczema Area and Severity Index) versus placebo (44%-52% dupilumab; 12%-15% placebo, Figure 3, B). Reductions in pruritus and depression, as well as improved quality of life, also showed statistically significant differences versus placebo. In a pooled analysis, dupilumab significantly improved pruritus within 1 to 3 days of treatment initiation.⁴⁰ Dupilumab was overall well tolerated, with conjunctivitis and injection site reactions reported at a higher rate in treatment groups versus placebo.

The LIBERTY AD CHRONOS trial studied long-term safety and efficacy of dupilumab using similar dosing regimens to the SOLO trials, but on background TCS and for a period of 52 weeks.³⁸ Similar to SOLO trials, the most frequent adverse events (AEs) were conjunctivitis and injection site reactions. The trial had 2 primary endpoints: IGA 0/1 and ≥ 2 points improvement from baseline, and EASI-75 improvement. At week 16, a higher proportion of patients receiving both dupilumab regimens significantly achieved primary outcomes versus placebo (Figure 3, A and B), with these efficacies maintained at week 52.

In the LIBERTY AD SOLO-CONTINUE trial, high-responding patients treated with dupilumab in SOLO were re-randomized to continue their original dupilumab regimen, 300 mg SC weekly or every 2 weeks (q2w), 300 mg SC every 4 weeks (q4w) or 8 weeks (q8w), or placebo for 36 weeks.⁴¹ Co-primary endpoints were the percentage change in EASI score from baseline and the proportion of patients with EASI-75 at week 36. Patients on dupilumab weekly or q2w showed continued response over time, with an insignificant change in percent EASI improvement from SOLO baseline versus placebo (Table I). In addition, the percent change with the other regimens dose dependently worsened. A significantly higher proportion of patients on dupilumab weekly or q2w maintained EASI-75 versus patients on placebo (71.6% vs 30.4%). Longer dosage intervals of dupilumab resulted in diminution of EASI-75 response (58.3% on q4w, 54.9% on q8w). Consistent with the continuous co-primary EASI endpoint, there was no overall loss of

TABLE I. Biologics currently approved or in clinical trials for atopic dermatitis

Drug	Mechanism	Phase trial	N	Age group (y)	Duration (wk)	Concomitant TCS/rescue therapy	Primary endpoint	Adverse events	Current phase of clinical trials underway	Strength of recommendation, grades of evidence ^{*79,80}
Dupilumab	Anti-IL-4R α mAb	Phase 3 (SOLO-CONTINUE)	422	≥ 18	36	No	Percent change in EASI score (difference between SOLO-CONTINUE baseline and week 36) Treatment group weekly or q2w: no significant change from SOLO-CONTINUE baseline (−0.06%) q4w: −3.8% q8w: −6.8% Placebo: −21.7% EASI-75 Weekly or q2w: 71.6% [†] q4w: 58.3% [‡] q8w: 54.9% [‡] Placebo: 30.4%	Conjunctivitis (5.4% weekly or q2w, 4.6% q4w 3.6% q8w vs 4.9% placebo), injection-site reactions (10.8% weekly or q2w, 6.9% q4w 7.1% q8w vs 8.5% placebo)	Phase 2: NCT03861455 NCT03346434 Phase 3: NCT02612454 NCT01949311 Phase 4: NCT03667014 NCT03389893 NCT03293030 NCT04447417 NCT04033367	A, 1a ⁴⁶
		Phase 3 (AD-1526)	251	12-18	16	Yes	Percentage of participants achieving an IGA response 200/300 mg q2w: 24.4% [†] 300 mg q4w: 17.9% [†] vs placebo: 2.4% EASI-75 200/300 mg q2w: 41.5% [†] 300 mg q4w: 38.1% [†] vs placebo: 8.2%	Conjunctivitis (9.8% q2w, 10.8% q4w vs 4.7% placebo), injection-site reactions (8.5% q2w, 6.0% q4w vs 3.5% placebo)		
Lebrikizumab	Anti-IL-13 mAb	Phase 2b	280	≥ 18	16	Yes	Percent change in EASI score 125 mg q4w: −62.3% 250 mg q4w: −69.2% [†] 250 mg q2w: −72.1% [†] vs placebo: −41.1%	Upper respiratory tract infection (7.5% all lebrikizumab groups, placebo 5.8%), nasopharyngitis (6.6% all lebrikizumab groups, placebo 3.8%)	Phase 3 NCT04146363 NCT04178967 NCT04392154 NCT04250350 NCT04250337	B, 2b
Tralokinumab	Anti-IL-13 mAb	Phase 2b	204	18-75	12	Yes	Change from baseline in EASI score 150 mg: −4.36 [‡] 300 mg: −4.94 [‡]	Upper respiratory tract infection (3.9% all tralokinumab groups vs 3.9% placebo)	Phase 3 NCT03587805 NCT03761537 NCT03526861	B, 2b

(continued)

TABLE I. (Continued)

Drug	Mechanism	Phase trial	N	Age group (y)	Duration (wk)	Concomitant TCS/rescue therapy	Primary endpoint	Adverse events	Current phase of clinical trials underway	Strength of recommendation, grades of evidence* ^{79,80}
Nemolizumab	Anti-IL-31R α mAb	Phase 2b	226	≥ 18	24	Yes	<i>Percent change in EASI score</i> At week 24: Greatest difference in 30 mg dose: $-68.8\%^{\dagger}$ vs placebo: -52.1%	Dose-dependent increase in asthma exacerbations in treatment group (3.6% 10 mg, 12.3% 30 mg, 17.5% 90 mg vs 1.8% placebo)	Phase 2 NCT03921411 Phase 3 NCT04365387 NCT03985943 NCT03989349 NCT03989206	A, 1b
Tezepelumab	Anti-TSLP mAb	Phase 2a	113	18-75	12	Yes	<i>EASI-50</i> Failed to reach statistical significance 280 mg: 64.7% vs placebo: 48.2%	Injection-site erythema (5.4% vs 0% placebo)	Phase 2b NCT03809663	B, 2b
ISB 830	Anti-OX40 mAb	Phase 2a	64	≥ 18	12	No	<i>Incidence of TEAEs</i> Similar across treatment groups ISB 830: 63.0% (29/46) vs placebo: 63.0% (10/16) <i>Changes from baseline in biomarkers (epidermal hyperplasia and cytokine) at day 71</i> Significant reductions in: - hyperplasia measures [†] - mRNA signature for Th1, Th2, and Th17/Th22 axes [†] - OX40 ⁺ T cells and OX40L ⁺ dendritic cells [†]	Headache (16%), atopic dermatitis (13%) nasopharyngitis (10%)	Phase 2b NCT03568162	B, 2b

EASI, Eczema Area and Severity Index; *EASI-50*, greater than 50% improvement in EASI score; *EASI-75*, greater than 75% improvement in EASI score; *IGA*, Investigator's Global Assessment; *mAb*, monoclonal antibody; *q2w*, every 2 weeks; *q4w*, every 4 weeks; *q8w*, every 8 weeks; *TCS*, topical corticosteroid; *TEAE*, treatment-emergent adverse event.

*(Strength of recommendation, grades of evidence) = (A, 1a/1b), (B, 2a/2b/3a/3b), (C, 4), (D, Expert opinion).⁷⁹

[†]Statistical significance of at least $P < .001$.

[‡]Statistical significance of at least $P < .05$.




FIGURE 3. Primary endpoint (**A**) and key efficacy endpoints (**B, C**) of phase 3 trials of dupilumab for the treatment of moderate-to-severe atopic dermatitis. **A**, The proportions of patients who achieved IGA 0/1 with ≥ 2 reductions from baseline at week 16 among patients who received dupilumab q1w, q2w, or placebo in SOLO 1 and SOLO 2. Proportions of patients who achieved IGA 0/1 at week 16 among were a primary endpoint for AD-1526. **B**, The proportions of patients who achieved EASI-75 at week 16 among patients who received dupilumab and placebo. EASI-75 was a co-primary endpoint in CHRONOS. **C**, The proportions of patients with ≥ 4 -point improvement in peak pruritus NRS score from baseline at week 16 among patients who received dupilumab and placebo. $P < .001$ for all comparisons with placebo in SOLO1/SOLO2. $P < .0001$ for all comparisons with placebo in CHRONOS. $P < .001$ for all comparisons with placebo in AD-1526. EASI-75, Greater than 75% improvement in Eczema Area and Severity Index score; IGA, Investigator's global assessment; NRS, numeric rating scale; q1w, every week; q2w, every other week. *In CHRONOS, patients also received concomitant topical corticosteroid in every treatment arm. **In AD-1526, patients received dupilumab every 2 weeks (green), every 4 weeks (blue), or placebo.

efficacy with dupilumab weekly or q2w (baseline vs week 35) in percentage change in peak pruritus numeric rating scale (NRS), whereas there was a dose-dependent return of pruritus for the q4w, q8w, and placebo groups, particularly after week 12. No new safety signals were identified. These findings confirm dupilumab's efficacy and safety and support the regimen of 300 mg SC q2w for long-term treatment.

In a phase 3 trial in adolescents aged 12 to 18 years (AD-1526), patients with moderate-to-severe AD were randomized to dupilumab 300 mg SC q4w, 200/300 mg SC q2w (based on weight < 60 kg or ≥ 60 kg, respectively), or placebo.³⁹ Primary outcomes were the percentage of patients reaching the IGA score of 0/1 and the percentage of patients achieving EASI-75 at week 16. A greater percentage of adolescents in the treatment arms achieved primary outcomes versus placebo (Table I, Figure 3, A and B). Safety profiles in adolescents were similar to adults, with

conjunctivitis and injection site reactions the most common AEs. Furthermore, pooled analysis from laboratory findings from 3 randomized controlled trials (SOLO-1, SOLO-2, and LIBERTY AD CHRONOS) assessing the need for routine safety testing of dupilumab concluded that patients using dupilumab for moderate-to-severe AD do not need routine laboratory testing, as there was no clinically major change in routine laboratory parameters attributed to dupilumab.⁴²

Moreover, the effectiveness and safety of dupilumab treatment in real-life clinical setting are comparable with that of clinical trials. Twenty-two unique studies encompassing 3303 patients with AD showed that after 16 weeks of dupilumab therapy, the pooled proportion of patients achieving EASI-75 was 59.8%. Conjunctivitis was the most common AE, reported in a pooled proportion of 26.1% compared with 8% in pooled data from clinical trials.⁴³ This AE seems to be specific to AD, as it was not observed in

patients involved in trials assessing dupilumab in conjunction with asthma and nasal polyposis.⁴⁴ Although it is currently not possible to predict who will develop conjunctivitis during treatment with dupilumab, patients with a history of allergic conjunctivitis seem to have a higher risk. However, the pathophysiology of dupilumab-induced eye disorders remains unknown.⁴³ Furthermore, biomarker analysis of a cohort study of daily practice dupilumab treatment up to 16 weeks found that treatment with dupilumab also significantly suppressed disease severity-related serum biomarkers thymus- and activation-regulated chemokine (TARC), pulmonary and activation-regulated chemokine, periostin, and IL-22, and eosinophil-related markers eotaxin-1 and eotaxin-3.⁴⁵

In adult patients with moderate-to-severe AD, dupilumab improved their health outcomes compared with best supportive care, and additional costs led to an incremental cost-effectiveness ratio ranging from £28,500 (low certainty) to \$124,541 (US dollars; moderate certainty), which are still likely to result in overall cost-effectiveness. However, all economic analyses were performed in high-income countries in line with their health system perspectives. Thus, their results may not be applicable to other countries.⁴⁶

Phase 1 and 2 mechanistic studies involving skin biopsies and blood from patients treated with dupilumab versus placebo showed highly significant changes in the molecular signatures only in patients treated with dupilumab. These included as expected Th2-related products, but also extended to markers related to other immune axes, such as Th17/Th22, but not Th1.^{47,48} A recent meta-analysis approach showed that the molecular improvements with dupilumab in certain immune pathways (such as Th2) are on par to those induced by broad systemic treatments, such as cyclosporine A.^{49,50} Furthermore, a systematic literature review comparing efficacies of systemic therapies for the treatment of AD showed that the strongest evidence currently exists for dupilumab and cyclosporine at improving clinical disease severity and quality of life.⁵¹

Several phase 3 clinical trials indicate that dupilumab is effective in AD, with the response maintained for at least 1 year of continuous treatment in the majority of patients and with a tolerable safety profile, but showed a higher incidence of conjunctivitis. Approximately one-third of all treated adult patients are clear in IGA from their AD. Up to 70% of patients achieve an EASI-75, and it takes approximately 4 weeks to reach the full clinical outcome.

LEBRIKIZUMAB

Lebrikizumab is a humanized IgG4K mAb that inhibits IL-13 signaling by binding free IL-13 with very high affinity, blocking ability of IL-13 to bind IL-4R α . This prevents heterodimerization of IL-4R α and IL-13 receptor alpha 1 chain (IL-13R α 1) subunits.⁵² IL-13 also downregulates filaggrin expression, an integral protein in formation and maintenance of the epidermal barrier.⁵³

A phase 2 study assessed the safety and efficacy of lebrikizumab administered subcutaneously at 3 different doses in adults with moderate-to-severe AD inadequately controlled with TCS and regular emollients for ≥ 1 month (125 mg single dose, 250 mg single dose, 125 mg q4w, or placebo q4w for 12 weeks after a 2-week period of TCS run-in). At the primary endpoint at week 12, EASI-50 was achieved in patients on 125 mg q4w (82.4%; $P = .026$), whereas a 62.3% response was observed in

the placebo group treated with TCS. Furthermore, 125 mg q4w treatment showed significantly more patients achieving EASI-75 (54.9% vs 34.0% placebo; $P = .036$) and Scoring of Atopic Dermatitis improvements of $\geq 50\%$ (SCORAD-50) (51.0% vs 26.4% placebo; $P = .012$), whereas no statistically significant response was seen in the single-dose groups. None of the 3 different doses of lebrikizumab achieved statistically significant improvement in pruritus visual analog scale (VAS), Atopic Dermatitis Impact Questionnaire, or Dermatology Life Quality Index (DLQI). Lebrikizumab was well tolerated, with only mild-to-moderate AEs. No significant differences in AEs among treatment and placebo groups were seen (66.7%; placebo 66.0%). The most frequent treatment associated AEs included conjunctivitis (9.6%; placebo 8.0%) and herpetic infections (7.7%; placebo 0%).⁵⁴

A 16-week phase 2b study investigated the safety and efficacy of lebrikizumab monotherapy 125 mg SC q4w, 250 mg SC q4w, or 250 SC mg q2w (following a loading dose) for the treatment of patients with moderate-to-severe AD with chronic AD ≥ 1 year and uncontrolled by topical treatments. Patients requiring rescue therapy could use TCS for as brief a period as possible. The study showed significant dose-dependent improvement in the average percentage of change in EASI scores and pruritus NRS versus placebo. Improvements in EASI score were achieved in 62.3% of patients in the 125 mg q4w group ($P = .165$), 69.2% of patients in the 250 mg q4w group ($P = .0022$), and 72.1% in the 250 mg q2w group ($P = .0005$), versus 41.1% in the placebo group. Furthermore, the 250 mg groups showed a significant difference versus placebo in secondary endpoints, including percent with score of 0/1 on IGA, EASI-50, EASI-75, and EASI-90. In addition, no dose-response relationship in treatment emergent adverse events (TEAEs) was observed. Rescue medication was used less in lebrikizumab-treated patients versus placebo (125 mg q4w: 12.3%; 250 mg q4w: 12.5%; 250 mg q2w: 13.3%; placebo: 34.6%), and lebrikizumab-treated patients had a shorter duration of topical medication use. AEs were mild to moderate, with the most common AEs upper respiratory tract infection (7.5% all lebrikizumab groups, placebo 5.8%) and nasopharyngitis (6.6%; placebo 3.8%).⁵⁵

Although phase 2 trials of lebrikizumab demonstrated that blocking IL-13 alone may be sufficient to see the efficacy signals in AD, head-to-head active comparator studies are needed to provide the comparative efficacy of anti-IL-13 therapy alone versus anti-IL-4/IL-13. Phase 3 trials investigating the long-term safety and efficacy of lebrikizumab as monotherapy of 52 weeks' duration for the treatment of moderate-to-severe AD are also currently underway (NCT04146363, NCT04178967, NCT04392154, NCT04250350).

TRALOKINUMAB

Tralokinumab is a fully human IgG4K mAb binding to unbound IL-13 cytokine with high affinity, similar to lebrikizumab but at a different epitope, preventing IL-13 from binding to both IL-13R α 1 and IL-13R α 2.⁵⁶ It was initially developed for the treatment of severe asthma. However, in a phase 2b study, tralokinumab did not significantly reduce asthma exacerbation rates in patients with severe uncontrolled asthma.⁵⁷

In a 12-week phase 2b trial, tralokinumab was studied in adults with moderate-to-severe AD. Participants were randomized to receive placebo or tralokinumab SC q2w (45, 150, or 300

mg) with concomitant TCS, after a 2-week run-in period with TCS. At baseline, the mean EASI score ranged from 24.8 to 27.3 for all tralokinumab groups and 26.4 for placebo. The 150 or 300 mg group achieved clinically significant improvements in EASI scores starting at week 4 versus placebo. At week 12, the adjusted mean difference from baseline EASI score was significantly different from placebo in patients treated with 150 mg (-4.36 ; $P = .03$) and 300 mg (-4.94 ; $P = .01$). The greatest improvements in IGA score were observed in the 300 mg group but did not achieve statistical significance (26.7% of tralokinumab-treated patients vs 11.8% placebo; $P = .06$). The 300 mg group also had a statistically significant difference in secondary endpoints (DLQI, pruritus NRS score, and SCORAD score) versus placebo.⁵⁸ Furthermore, the exploratory analyses stratified the patients by baseline serum biomarkers for IL-13 signaling, including dipeptidyl peptidase-4 (DPP-4) and periostin.^{58,59} Tralokinumab-treated groups in the DPP-4 high and periostin high subgroups had greater improvements in EASI scores versus the intended to treat (ITT) population. Differences in IGA response for the treatment arm were also improved in the same subgroups versus ITT population. This presents the concept that the biomarkers DPP-4 and periostin may help identify patients who may have better targeted treatment response. AEs were mild to moderate, with the most frequent event upper respiratory tract infection (3.9% all tralokinumab groups; 3.9% placebo). Conjunctivitis was also reported (2.0% 45 mg, 5.9% 150 mg vs 3.9% placebo). In terms of interpretation of this trial, because of the run-in period and concomitant use of TCS, the real efficacy of this biologic may be confounded.⁵² The observed improvements in placebo-treated participants indicate that TCS provided partial benefit, despite all participants having inadequate disease control with such therapy at enrollment. Although a study of tralokinumab monotherapy would provide a more definitive measurement of efficacy than in combination with topical glucocorticoids, in clinical practice, it is expected that biologics will be prescribed concomitantly to topical therapies.⁵² Phase 3 trials of tralokinumab in adults and adolescents with moderate-to-severe AD have recently been completed with results pending (NCT03131648, NCT03160885, NCT03363854). A phase 3 trial investigating the long-term safety and efficacy of tralokinumab in patients who participated in previous trials is also currently underway (NCT03587805). In addition, there is an active phase 3 trial studying the efficacy of tralokinumab used as monotherapy for adolescents with moderate-to-severe AD (NCT03526861).

NEMOLIZUMAB

Nemolizumab is a humanized IgG2 κ anti-IL-31 receptor α (IL-31R α) mAb.⁶⁰ IL-31, a proinflammatory cytokine playing an important role in mediating pruritus via overexpression of IL-31 receptors on sensory nerves,⁶¹ is known as a perpetuator of the itch-scratch cycle that results in disruption of the skin barrier in AD.⁶² A phase 2b study randomized patients with moderate-to-severe AD and severe AD-associated pruritus uncontrolled by topical treatments to 10, 30, or 90 mg of subcutaneous nemolizumab q4w or placebo until week 20, with a 12-week follow-up period until week 32.⁶⁰ Both treatment and placebo groups applied TCS and moisturizers in addition to treatment. At

baseline, the mean EASI score ranged from 24.2 to 25.9 for all nemolizumab groups and 27.0 for placebo. All groups showed improvement in the primary endpoint, the mean percentage change in baseline EASI at 24 weeks versus placebo, with the 30 mg group the most effective (-68.8% vs -52.1% ; $P = .016$). All doses of nemolizumab also showed statistically significant improvements in peak pruritus NRS as early as week 1 ($P < .05$), with the most efficacious response seen in the 30 mg treatment arm (67.3% vs 35.8% placebo at week 24, $P < .001$). Furthermore, the 30 mg group had a significant difference from placebo in secondary endpoints, including the percentage of patients achieving EASI-50, EASI-75, and EASI-90 ($P < .05$). Nemolizumab was well tolerated and had a good safety profile, with the most common AEs nasopharyngitis, upper respiratory tract infection, and exacerbation of AD. In addition, a dose-dependent increase in asthma-related events was observed in patients with a history of asthma. Some of these events might have occurred because effective treatment with nemolizumab led to improved overall well-being and increased activity levels that, in turn, triggered asthma symptoms. All asthma-related events were mild and manageable.

In a 16-week phase 3 trial, Japanese patients with AD and moderate-to-severe pruritus and inadequate response to topical agents and antihistamines were randomized to receive SC nemolizumab (60 mg) or placebo q4w with concomitant topical agents. The primary endpoint was the mean percent change in the VAS score for pruritus.⁶³ At baseline, the median VAS score for pruritus was 75 and the median EASI score ranged from 22.7 (placebo) to 24.2 (nemolizumab). At week 16, the mean percent change in the VAS score was -42.8% in the nemolizumab group and -21.4% in the placebo group ($P < .001$). However, for the secondary efficacy endpoints, including the change in EASI score and the time course of change in the VAS score for pruritus up to 4 weeks, no adjustments were made for multiple comparisons, from which no clinical inferences can be made. The mean percent change in EASI score was -45.9% with nemolizumab and -33.2% with placebo. At day 15, the percent change in the daily mean VAS score for pruritus was reported as early as day 2 (-10.3% nemolizumab vs -4.4% placebo). Overall, nemolizumab was well tolerated, with only mild-to-moderate AEs. Incidence of injection-site reactions was greater with nemolizumab versus placebo (8% vs 3%).⁶³

A phase 2 study testing the safety and pharmacokinetics of nemolizumab in adolescent patients with AD is ongoing (NCT03921411). Two phase 3 studies are currently recruiting to further test the efficacy and safety profile of nemolizumab in adults with moderate-to-severe AD (NCT03985943, NCT03989349).

TEZEPELUMAB

Tezepelumab is a fully human IgG2 λ mAb binding thymic stromal lymphopoietin (TSLP), an epidermal keratinocyte-derived cytokine that activates dendritic cells to induce the production of type 2 cytokines, including IL-4, IL-5, IL-13, and TNF- α , and contributes to pruritus in AD by activating cutaneous sensory neurons.^{64,65} Overexpression of TSLP in keratinocytes has been found in patients with acute or chronic AD.⁶⁶

High levels of TSLP in the serum have also been observed in children with AD.⁶⁷

In a phase 2a trial, patients were randomized to receive 280 mg tezepelumab SC or placebo q2w with concomitant TCS. However, the study failed to reach statistical significance in the primary endpoint, EASI-50 versus placebo, at week 12 (64.7% vs 48.2%; $P = .091$).⁶⁵ Similarly, only a numerical difference between the 2 groups in IGA response rate was observed at week 12 (19.3% vs 12.8%; $P = .36$). At other secondary endpoints, including EASI-75, EASI-90, SCORAD, and NRS scores at week 12, patients treated with tezepelumab did not achieve statistically significant improvements versus placebo. In addition, in an exploratory biomarker subgroup analysis, tezepelumab-treated patients who were DPP-4 high, periostin low, CCL17/TARC low, or IgE high were found to have only numerically greater response at week 12 compared with patients in opposing biomarker subgroups. AEs were comparable between groups and were mild to moderate. Most AEs were not considered treatment related, except for injection site erythema. Additional common AEs were nasopharyngitis, diarrhea, upper respiratory tract infection, and headache.⁶⁵ Currently, a phase 2b trial is recruiting to assess the safety and efficacy of tezepelumab as monotherapy and adjunct therapy with TCS (NCT03809663).

ISB 830 (PREVIOUSLY GBR 830)

ISB 830 is a humanized IgG1 anti-OX40 mAb. Engagement of OX40 with its ligand (OX40L) expressed on antigen-presenting cells potentiates effector T-cell responses.^{10,68} In a phase 2a trial, the safety and efficacy of ISB 830 was assessed in adults with moderate-to-severe AD and inadequate response to topical treatments. Patients were randomized 3:1 to 10 mg/kg intravenous (IV) ISB 830 or placebo on day 1 (baseline) and day 29. Biopsies were collected on days 1, 29, and 71. Primary endpoints were TEAEs and changes in baseline epidermal hyperplasia and gene expression of lesional biomarkers. Secondary endpoints included percent improvement from baseline in SCORAD, IGA, body surface area, and EASI scores. However, the study analyzed statistical differences between treatment groups only in primary endpoints and not in secondary endpoints. ISB 830 was well tolerated with similar TEAEs between treatment groups (63.0% vs 63.0%). Nasopharyngitis was the only AE considered treatment related (8.7%).⁴³ Biomarker analysis demonstrated that ISB 830 post-treatment lesional skin had a significant decline in levels of OX40⁺ T cells and OX40L⁺ dendritic cells ($P < .001$). Furthermore, significant reductions in hyperplasia, epidermal thickness ($P < .001$), Keratin 16 (K16) mRNA expression ($P < .01$), and Ki67⁺ cells ($P < .001$) were noted in ISB 830-treated groups by day 71. ISB 830 significantly decreased the mRNA expression of cytokines along the Th1 axis—IFN- γ ($P < .01$) and CXCL10 ($P < .001$); Th2 axis—IL-31 ($P < .05$), CCL11 ($P < .001$), CCL17 ($P < .001$), and TSLP receptor ($P < .001$); and Th17/Th22 axis—IL-23p19 ($P < .001$), IL-8 ($P < .01$), and S100A12 ($P < .001$). However, other main Th2 (IL-4, IL-13) and Th17/Th22 (IL-17A, IL-22) cytokines were not significantly reduced with ISB 830. In terms of secondary outcome measures, the ISB 830 group had a greater proportion of patients achieving EASI-50 versus placebo (76.9% vs 37.5%). Similarly, an IGA response was reached by 23.1% of ISB 830-treated groups versus 12.5% of placebo-treated patients at day 71. Of note, 2 IV doses of the drug administered 4

weeks apart induced significant improvement of tissue and clinical measurements even 42 days after the last dose, suggesting that ISB 830 may provide a novel therapeutic paradigm for patients with moderate-to-severe AD. ISB 830 is currently in a phase 2b clinical trial in AD (NCT03568162).

BIOLOGIC THERAPIES CURRENTLY NOT UNDERGOING TRIALS IN AD

Fezakinumab

Fezakinumab is a human IgG1 λ anti-IL-22 mAb.⁶⁹ IL-22, produced by Th22 cells, is involved in epidermal hyperplasia and barrier defects in patients with AD. IL-22 levels have been reported to correlate with AD disease severity and response to AD treatment.^{15,70} A small study of 60 patients with moderate-to-severe AD treated for 10 weeks with fezakinumab SC q2w, with primary endpoint at week 12 and a secondary endpoint at week 20, showed clinical benefit in a portion of patients with severe disease, but not in moderate patients.⁶⁹ A follow-up mechanistic study showed that responders were those with high levels of IL-22, whereas patients with low IL-22 levels did not respond, and perhaps even showed an exacerbation.⁷¹ This study suggests the potential for a personalized medicine approach based on distinct molecular mechanisms.

Anti-IL-17 therapy

MOR106, a mouse and human IgG1 anti-IL-17C mAb, showed promising results from a phase 1 study. Eighty-three percent of patients with AD treated with MOR106 achieved EASI-50 versus less than 20% in the placebo at week 4, with continued improvements in the treatment group over the 10-week follow-up period.⁷² However, 2 phase 2 trials testing the safety, efficacy, and tolerability of MOR106 in patients with moderate-to-severe AD were terminated, as they did not meet primary endpoints (NCT03568071, NCT03864627).⁷³ Secukinumab, a human IgG1K anti-IL-17A mAb, has recently completed the phase 2 trial (NCT02594098), with no significant changes in EASI or IGA scores. Another phase 2 trial investigating secukinumab for moderate-to-severe AD recently completed with pending results (NCT03568136).

Omalizumab and ligelizumab

Omalizumab, a recombinant IgG1K anti-IgE mAb, has been used in AD with variable results. A systematic review and meta-analysis of omalizumab in AD found that fewer than 50% of the patients treated with this biologic achieved a significant clinical improvement (defined as SCORAD-50, EASI-75, or IGA 0 or reduction by 2 points).⁷⁴ Recently, a phase 4 trial in children with severe AD found that omalizumab significantly reduced disease severity and topical steroid use.⁷⁵ A phase 2 trial of ligelizumab (QGE031), an mAb with greater affinity for IgE than omalizumab, was completed in 2013 with no results posted, suggesting lack of efficacy in AD (NCT01552629).

CONCLUSION

Targeted biologic agents play an increasing role in the treatment of patients with AD refractory to topical treatments. Compared with the off-label use of immunosuppressant medications that are still a mainstay of treatment for severe uncontrolled AD in some countries, these novel biologic therapies appear to demonstrate considerably better risk-benefit ratios, although there is little long-term data to date.⁷⁶ So far, only dupilumab is FDA approved, for which trials and real-life studies

have demonstrated considerable short- and mid-term efficacy and safety.^{32,33,38,39,41-44} There is a large unmet need for the development of additional mAbs targeting distinct immune pathways necessary to increase probability of achieving disease control in patients with AD, and combination therapy such as dupilumab plus another systemic immunomodulatory agent and/or phototherapy has not been adequately studied.⁷⁷

Apart from dupilumab, several novel biologic agents have demonstrated promising results in clinical trial. Lebrikizumab, tralokinumab, nemolizumab, and ISB 830 have shown favorable results in achieving improvement in disease severity and multiple endpoint outcomes.^{26,54,55,58,60} Compared with other treatments, nemolizumab appears to be superior in improving pruritus, because it has a rapid onset of pruritus reduction as early as dupilumab, which was within 2 days of treatment initiation,^{40,60} and also maintained good long-term effectiveness.⁷⁸ Tezepelumab, however, did not show statistically significant changes in primary outcome endpoints.⁶⁵

There are several limitations in analyzing clinical trials in AD. Interpretation of the efficacy of biologics can be confounded by concomitant TCS use, and the clinical application of these drugs is limited, as children were excluded from most AD trials. Therefore, it is imperative that future large-scale studies investigate the efficacy of biologics as monotherapy in children with moderate-to-severe AD. Furthermore, long-term follow-up studies assessing safety and persistence of efficacy of each biologic agent are critical for patients with chronic AD. To further guide treatment decisions and guidelines, network meta-analysis is also necessary to provide the best comparative effectiveness of biologic therapies.

REFERENCES

- Thyssen JP, Andersen Y, Halling AS, Williams HC, Egeberg A. Strengths and limitations of the United Kingdom Working Party criteria for atopic dermatitis in adults. *J Eur Acad Dermatol Venereol* 2020;34:1764-72.
- Abuabara K, Ye M, McCulloch CE, Sullivan A, Margolis DJ, Strachan DP, et al. Clinical onset of atopic eczema: results from 2 nationally representative British birth cohorts followed through midlife. *J Allergy Clin Immunol* 2019;144:710-9.
- Barbarot S, Auziere S, Gadkari A, Girolomoni G, Puig L, Simpson EL, et al. Epidemiology of atopic dermatitis in adults: results from an international survey. *Allergy* 2018;73:1284-93.
- Weidinger S, Novak N. Atopic dermatitis. *Lancet* 2016;387:1109-22.
- Silverberg JI. Adult-onset atopic dermatitis. *J Allergy Clin Immunol Pract* 2019;7:28-33.
- Guttman-Yassky E, Krueger JG, Lebwohl MG. Systemic immune mechanisms in atopic dermatitis and psoriasis with implications for treatment. *Exp Dermatol* 2018;27:409-17.
- Langan SM, Irvine AD, Weidinger S. Atopic dermatitis. *Lancet* 2020;396:345-60.
- Brunner PM, Guttman-Yassky E, Leung DY. The immunology of atopic dermatitis and its reversibility with broad-spectrum and targeted therapies. *J Allergy Clin Immunol* 2017;139:S65-76.
- Czarnowicki T, He H, Krueger JG, Guttman-Yassky E. Atopic dermatitis endotypes and implications for targeted therapeutics. *J Allergy Clin Immunol* 2019;143:1-11.
- Guttman-Yassky E, Pavel AB, Zhou L, Estrada YD, Zhang N, Xu H, et al. GBR 830, an anti-OX40, improves skin gene signatures and clinical scores in patients with atopic dermatitis. *J Allergy Clin Immunol* 2019;144:482-493.e7.
- Hirasawa Y, Takai T, Nakamura T, Mitsuishi K, Gunawan H, Suto H, et al. *Staphylococcus aureus* extracellular protease causes epidermal barrier dysfunction. *J Invest Dermatol* 2010;130:614-7.
- Sonesson A, Bartosik J, Christiansen J, Roscher I, Nilsson F, Schmidtchen A, et al. Sensitization to skin-associated microorganisms in adult patients with atopic dermatitis is of importance for disease severity. *Acta Derm Venereol* 2013;93:340-5.
- van Drongelen V, Haisma EM, Out-Luiting JJ, Nibbering PH, El Ghalbzouri A. Reduced filaggrin expression is accompanied by increased *Staphylococcus aureus* colonization of epidermal skin models. *Clin Exp Allergy* 2014;44:1515-24.
- Kopfnael V, Harder J, Werfel T. Expression of antimicrobial peptides in atopic dermatitis and possible immunoregulatory functions. *Curr Opin Allergy Clin Immunol* 2013;13:531-6.
- Gittler JK, Shemer A, Suárez-Fariñas M, Fuentes-Duculan J, Gulewicz KJ, Wang CQ, et al. Progressive activation of T(H)2/T(H)22 cytokines and selective epidermal proteins characterizes acute and chronic atopic dermatitis. *J Allergy Clin Immunol* 2012;130:1344-54.
- Ahn K, Kim BE, Kim J, Leung DY. Recent advances in atopic dermatitis. *Curr Opin Immunol* 2020;66:14-21.
- Noda S, Suárez-Fariñas M, Ungar B, Kim SJ, de Guzman Strong C, Xu H, et al. The Asian atopic dermatitis phenotype combines features of atopic dermatitis and psoriasis with increased TH17 polarization. *J Allergy Clin Immunol* 2015;136:1254-64.
- Czarnowicki T, Esaki H, Gonzalez J, Malajian D, Shemer A, Noda S, et al. Early pediatric atopic dermatitis shows only a cutaneous lymphocyte antigen (CLA)(+) TH2/TH1 cell imbalance, whereas adults acquire CLA(+) TH22/TC22 cell subsets. *J Allergy Clin Immunol* 2015;136:941-951.e3.
- Sanyal RD, Pavel AB, Glickman J, Chan TC, Zheng X, Zhang N, et al. Atopic dermatitis in African American patients is T(H)2/T(H)22-skewed with T(H)1/T(H)17 attenuation. *Ann Allergy Asthma Immunol* 2019;122:99-110.e6.
- Silverberg JI. Comorbidities and the impact of atopic dermatitis. *Ann Allergy Asthma Immunol* 2019;123:144-51.
- Schmitt J, Schwarz K, Baurecht H, Hotze M, Folster-Holst R, Rodriguez E, et al. Atopic dermatitis is associated with an increased risk for rheumatoid arthritis and inflammatory bowel disease, and a decreased risk for type 1 diabetes. *J Allergy Clin Immunol* 2016;137:130-6.
- Mohan GC, Silverberg JI. Association of vitiligo and alopecia areata with atopic dermatitis: a systematic review and meta-analysis. *JAMA Dermatol* 2015;151:522-8.
- Drucker AM, Thompson JM, Li WQ, Cho E, Li T, Guttman-Yassky E, et al. Incident alopecia areata and vitiligo in adult women with atopic dermatitis: Nurses' Health Study 2. *Allergy* 2017;72:831-4.
- Kridin K, Renert-Yuval Y, Guttman-Yassky E, Cohen AD. Alopecia areata is associated with atopic diathesis: results from a population-based study of 51,561 patients. *J Allergy Clin Immunol Pract* 2020;8:1323-1328.e1.
- Glickman JW, Dubin C, Renert-Yuval Y, Dahabreh D, Kimmel GW, Auyeung K, et al. Cross-sectional study of blood biomarkers of patients with moderate to severe alopecia areata reveals systemic immune and cardiovascular biomarker dysregulation. *J Am Acad Dermatol* 2021;84:370-80.
- Hamann CR, Hamann D, Egeberg A, Johansen JD, Silverberg J, Thyssen JP. Association between atopic dermatitis and contact sensitization: a systematic review and meta-analysis. *J Am Acad Dermatol* 2017;77:70-8.
- Simonsen AB, Johansen JD, Deleuran M, Mortz CG, Sommerlund M. Contact allergy in children with atopic dermatitis: a systematic review. *Br J Dermatol* 2017;177:395-405.
- Correa da Rosa J, Malajian D, Shemer A, Rozenblit M, Dhingra N, Czarnowicki T, et al. Patients with atopic dermatitis have attenuated and distinct contact hypersensitivity responses to common allergens in skin. *J Allergy Clin Immunol* 2015;135:712-20.
- Hale G, Davies E, Grindlay DJC, Rogers NK, Harman KE. What's new in atopic eczema? An analysis of systematic reviews published in 2017. Part 2: epidemiology, aetiology and risk factors. *Clin Exp Dermatol* 2019;44:868-73.
- Silverberg JI. Atopic dermatitis treatment: current state of the art and emerging therapies. *Allergy Asthma Proc* 2017;38:243-9.
- Agency EM. Dupixent (dupilumab): an overview of dupixent and why it is authorised in the EU: European Medicines Agency. Updated October 3, 2019. Available from: <https://www.ema.europa.eu/en/medicines/human/EPAR/dupixent>. Accessed August 15, 2020.
- Simpson EL, Bieber T, Guttman-Yassky E, Beck LA, Blauvelt A, Cork MJ, et al. Two phase 3 trials of dupilumab versus placebo in atopic dermatitis. *N Engl J Med* 2016;375:2335-48.
- Thaçi D, Simpson EL, Beck LA, Bieber T, Blauvelt A, Papp K, et al. Efficacy and safety of dupilumab in adults with moderate-to-severe atopic dermatitis inadequately controlled by topical treatments: a randomised, placebo-controlled, dose-ranging phase 2b trial. *Lancet* 2016;387:40-52.
- Sanofi S. FDA approves Dupixent® (dupilumab) as first biologic medicine for children aged 6 to 11 years with moderate-to-severe atopic dermatitis. Available from: <https://www.sanofi.com/en/media-room/press-releases/2020/2020-05-26-17-40-00#:~:text=PARIS%20and%20TARRYTOWN%2C%20N.Y.,those%20therapies%20are%20not%20advisable>. Accessed July 24, 2020.

35. Regeneron. FDA approves asthma indication for Dupixent® (dupilumab); 2018. Updated October 19, 2018. Available from: <https://investor.regeneron.com/news-releases/news-release-details/fda-approves-asthma-indication-dupixent-dupilumab>. Accessed August 15, 2020.
36. US Food and Drug Administration. FDA approves first treatment for chronic rhinosinusitis with nasal polyps; 2019. Available from: <https://www.fda.gov/news-events/press-announcements/fda-approves-first-treatment-chronic-rhinosinusitis-nasal-polyps>. Accessed August 16, 2020.
37. Hirano I, Dellon ES, Hamilton JD, Collins MH, Peterson K, Chehade M, et al. Efficacy of dupilumab in a phase 2 randomized trial of adults with active eosinophilic esophagitis. *Gastroenterology* 2020;158:111-122.e10.
38. Blauvelt A, de Bruin-Weller M, Gooderham M, Cather JC, Weisman J, Pariser D, et al. Long-term management of moderate-to-severe atopic dermatitis with dupilumab and concomitant topical corticosteroids (LIBERTY AD CHRONOS): a 1-year, randomised, double-blinded, placebo-controlled, phase 3 trial. *Lancet* 2017;389:2287-303.
39. Simpson EL, Paller AS, Siegfried EC, Boguniewicz M, Sher L, Gooderham MJ, et al. Efficacy and safety of dupilumab in adolescents with uncontrolled moderate to severe atopic dermatitis: a phase 3 randomized clinical trial. *JAMA Dermatol* 2020;156:44-56.
40. Thaçi D, Simpson EL, Deleuran M, Kataoka Y, Chen Z, Gadkari A, et al. Efficacy and safety of dupilumab monotherapy in adults with moderate-to-severe atopic dermatitis: a pooled analysis of two phase 3 randomized trials (LIBERTY AD SOLO 1 and LIBERTY AD SOLO 2). *J Dermatol Sci* 2019;94:266-75.
41. Worm M, Simpson EL, Thaçi D, Bissonnette R, Lacour JP, Beissert S, et al. Efficacy and safety of multiple dupilumab dose regimens after initial successful treatment in patients with atopic dermatitis: a randomized clinical trial. *JAMA Dermatol* 2020;156:131-43.
42. Wollenberg A, Beck LA, Blauvelt A, Simpson EL, Chen Z, Chen Q, et al. Laboratory safety of dupilumab in moderate-to-severe atopic dermatitis: results from three phase III trials (LIBERTY AD SOLO 1, LIBERTY AD SOLO 2, LIBERTY AD CHRONOS). *Br J Dermatol* 2020;182:1120-35.
43. Halling A-S, Loft ND, Silverberg JI, Guttman-Yassky E, Thyssen JP. Real-world evidence of dupilumab efficacy and risk of adverse events: a systematic review and meta-analysis. *J Am Acad Dermatol* 2021;84:139-47.
44. Faiz S, Giovannelli J, Podevin C, Jachiet M, Bouaziz JD, Reguiai Z, et al. Effectiveness and safety of dupilumab for the treatment of atopic dermatitis in a real-life French multicenter adult cohort. *J Am Acad Dermatol* 2019;81:143-51.
45. Ariens LFM, van der Schaft J, Bakker DS, Balak D, Romeijn MLE, Kouwenhoven T, et al. Dupilumab is very effective in a large cohort of difficult-to-treat adult atopic dermatitis patients: first clinical and biomarker results from the BioDay registry. *Allergy* 2020;75:116-26.
46. Agache I, Song Y, Posso M, Alonso-Coello P, Rocha C, Sola I, et al. Efficacy and safety of dupilumab for moderate-to-severe atopic dermatitis: a systematic review for the EAACI biologicals guidelines. *Allergy* 2021;76:45-58.
47. Hamilton JD, Suárez-Fariñas M, Dhingra N, Cardinale I, Li X, Kostic A, et al. Dupilumab improves the molecular signature in skin of patients with moderate-to-severe atopic dermatitis. *J Allergy Clin Immunol* 2014;134:1293-300.
48. Guttman-Yassky E, Bissonnette R, Ungar B, Suárez-Fariñas M, Ardeleanu M, Esaki H, et al. Dupilumab progressively improves systemic and cutaneous abnormalities in patients with atopic dermatitis. *J Allergy Clin Immunol* 2019;143:155-72.
49. Glickman JW, Dubin C, Han J, Dahabreh D, Garcet S, Krueger JG, et al. Comparing cutaneous molecular improvement with different treatments in atopic dermatitis patients. *J Allergy Clin Immunol* 2020;145:1285-8.
50. Glickman JW, Han J, Garcet S, Krueger JG, Pavel AB, Guttman-Yassky E. Improving evaluation of drugs in atopic dermatitis by combining clinical and molecular measures. *J Allergy Clin Immunol Pract* 2020;8:3622-3625.e19.
51. Seger EW, Wechter T, Strowd L, Feldman SR. Relative efficacy of systemic treatments for atopic dermatitis. *J Am Acad Dermatol* 2019;80:411-416.e4.
52. Bieber T. Interleukin-13: targeting an underestimated cytokine in atopic dermatitis. *Allergy* 2020;75:54-62.
53. Elias PM, Steinhoff M. "Outside-to-inside" (and now back to "outside") pathogenic mechanisms in atopic dermatitis. *J Invest Dermatol* 2008;128:1067-70.
54. Simpson EL, Flohr C, Eichenfield LF, Bieber T, Sofen H, Taieb A, et al. Efficacy and safety of lebrikizumab (an anti-IL-13 monoclonal antibody) in adults with moderate-to-severe atopic dermatitis inadequately controlled by topical corticosteroids: a randomized, placebo-controlled phase II trial (TREBLE). *J Am Acad Dermatol* 2018;78:863-871.e11.
55. Guttman-Yassky E, Blauvelt A, Eichenfield LF, Paller AS, Armstrong AW, Drew J, et al. Efficacy and safety of lebrikizumab, a high-affinity interleukin 13 inhibitor, in adults with moderate to severe atopic dermatitis: a phase 2b randomized clinical trial. *JAMA Dermatol* 2020;156:411-20.
56. Popovic B, Breed J, Rees DG, Gardener MJ, Vinall LM, Kemp B, et al. Structural characterisation reveals mechanism of IL-13-neutralising monoclonal antibody tralokinumab as inhibition of binding to IL-13Ralpha1 and IL-13Ralpha2. *J Mol Biol* 2017;429:208-19.
57. Brightling CE, Chaney P, Leigh R, O'Byrne PM, Korn S, She D, et al. Efficacy and safety of tralokinumab in patients with severe uncontrolled asthma: a randomised, double-blind, placebo-controlled, phase 2b trial. *Lancet Respir Med* 2015;3:692-701.
58. Wollenberg A, Howell MD, Guttman-Yassky E, Silverberg JI, Kell C, Ranade K, et al. Treatment of atopic dermatitis with tralokinumab, an anti-IL-13 mAb. *J Allergy Clin Immunol* 2019;143:135-41.
59. Izuhara K, Arima K, Ohta S, Suzuki S, Inamitsu M, Yamamoto K-i. Periostin in allergic inflammation. *Allergol Int* 2014;63:143-51.
60. Silverberg JI, Pinter A, Pulka G, Poulin Y, Bouaziz JD, Wollenberg A, et al. Phase 2B randomized study of nemolizumab in adults with moderate-to-severe atopic dermatitis and severe pruritus. *J Allergy Clin Immunol* 2020;145:173-82.
61. Miake S, Tsuji G, Takemura M, Hashimoto-Hachiya A, Vu YH, Furue M, et al. IL-4 augments IL-31/IL-31 receptor alpha interaction leading to enhanced Ccl 17 and Ccl 22 production in dendritic cells: implications for atopic dermatitis. *Int J Mol Sci* 2019;20:4053.
62. Feld M, Garcia R, Buddenkotte J, Katayama S, Lewis K, Muirhead G, et al. The pruritus- and TH2-associated cytokine IL-31 promotes growth of sensory nerves. *J Allergy Clin Immunol* 2016;138:500-508.e24.
63. Kabashima K, Matsumura T, Komazaki H, Kawashima M, Nemolizumab JPSG. Trial of nemolizumab and topical agents for atopic dermatitis with pruritus. *N Engl J Med* 2020;383:141-50.
64. Wilson SR, The L, Batia LM, Beattie K, Katibah GE, McClain SP, et al. The epithelial cell-derived atopic dermatitis cytokine TSLP activates neurons to induce itch. *Cell* 2013;155:285-95.
65. Simpson EL, Parnes JR, She D, Crouch S, Rees W, Mo M, et al. Tezepelumab, an anti-thymic stromal lymphopoietin monoclonal antibody, in the treatment of moderate to severe atopic dermatitis: a randomized phase 2a clinical trial. *J Am Acad Dermatol* 2019;80:1013-21.
66. Soumelis V, Reche PA, Kanzler H, Yuan W, Edward G, Homey B, et al. Human epithelial cells trigger dendritic cell mediated allergic inflammation by producing TSLP. *Nat Immunol* 2002;3:673-80.
67. Indra AK. Epidermal TSLP: a trigger factor for pathogenesis of atopic dermatitis. *Expert Rev Proteomics* 2013;10:309-11.
68. Webb GJ, Hirschfield GM, Lane PJ. OX40, OX40L and autoimmunity: a comprehensive review. *Clin Rev Allergy Immunol* 2016;50:312-32.
69. Guttman-Yassky E, Brunner PM, Neumann AU, Khattri S, Pavel AB, Malik K, et al. Efficacy and safety of fezakinumab (an IL-22 monoclonal antibody) in adults with moderate-to-severe atopic dermatitis inadequately controlled by conventional treatments: a randomized, double-blind, phase 2a trial. *J Am Acad Dermatol* 2018;78:872-881.e6.
70. Nogralas KE, Zaba LC, Shemer A, Fuentes-Duculan J, Cardinale I, Kikuchi T, et al. IL-22-producing "T22" T cells account for upregulated IL-22 in atopic dermatitis despite reduced IL-17-producing TH17 T cells. *J Allergy Clin Immunol* 2009;123:1244-1252.e2.
71. Brunner PM, Pavel AB, Khattri S, Leonard A, Malik K, Rose S, et al. Baseline IL-22 expression in patients with atopic dermatitis stratifies tissue responses to fezakinumab. *J Allergy Clin Immunol* 2019;143:142-54.
72. Morphosys. MorphoSys and Galapagos report first promising signs of clinical activity in a phase 1 study with IL-17C-antibody MOR106 in atopic dermatitis patients; 2017. Available from: <https://www.morphosys.com/media-investors/media-center/morphosys-and-galapagos-report-first-promising-signs-of-clinical>. Accessed August 17, 2020.
73. Morphosys. MorphoSys AG: MOR106 clinical development in atopic dermatitis stopped; 2019. Updated October 28, 2019. Available from: <https://www.morphosys.com/media-investors/media-center/morphosys-ag-mor106-clinical-development-in-atopic-dermatitis-stopped>. Accessed August 17, 2020.
74. Wang HH, Li YC, Huang YC. Efficacy of omalizumab in patients with atopic dermatitis: a systematic review and meta-analysis. *J Allergy Clin Immunol* 2016;138:1719-1722.e1.
75. Chan S, Cornelius V, Cro S, Harper JJ, Lack G. Treatment effect of omalizumab on severe pediatric atopic dermatitis: the ADAPT randomized clinical trial. *JAMA Pediatr* 2019;174:29-37.

76. Chun PIF, Lehman H. Current and future monoclonal antibodies in the treatment of atopic dermatitis. *Clin Rev Allergy Immunol* 2020;59:208-19.
77. Johnson BB, Franco AI, Beck LA, Prezzano JC. Treatment-resistant atopic dermatitis: challenges and solutions. *Clin Cosmet Investig Dermatol* 2019;12:181-92.
78. Alexander H, Patton T, Jabbar-Lopez ZK, Manca A, Flohr C. Novel systemic therapies in atopic dermatitis: what do we need to fulfil the promise of a treatment revolution? *F1000Res* 8:F1000.
79. Wollenberg A, Barbarot S, Bieber T, Christen-Zaech S, Deleuran M, Fink-Wagner A, et al. Consensus-based European guidelines for treatment of atopic eczema (atopic dermatitis) in adults and children: part II. *J Eur Acad Dermatol Venereol* 2018;32:850-78.
80. Ebell MH, Siwek J, Weiss BD, Woolf SH, Susman J, Ewigman B, et al. Strength of recommendation taxonomy (SORT): a patient-centered approach to grading evidence in the medical literature. *Am Fam Physician* 2004;69:548-56.
81. Fishbein AB, Silverberg JI, Wilson EJ, Ong PY. Update on atopic dermatitis: diagnosis, severity assessment, and treatment selection. *J Allergy Clin Immunol Pract* 2020;8:91-101.