



Global diversity in the human salivary microbiome

Ivan Nasidze, Jing Li, Dominique Quinque, et al.

Genome Res. 2009 19: 636-643 originally published online February 27, 2009
Access the most recent version at doi:[10.1101/gr.084616.108](https://doi.org/10.1101/gr.084616.108)

**Supplemental
Material**

<http://genome.cshlp.org/content/suppl/2009/02/27/gr.084616.108.DC1.html>

References

This article cites 45 articles, 23 of which can be accessed free at:
<http://genome.cshlp.org/content/19/4/636.full.html#ref-list-1>

Article cited in:

<http://genome.cshlp.org/content/19/4/636.full.html#related-urls>

**Email alerting
service**

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#)

To subscribe to *Genome Research* go to:
<http://genome.cshlp.org/subscriptions>

Global diversity in the human salivary microbiome

Ivan Nasidze,¹ Jing Li,^{2,3} Dominique Quinque,¹ Kun Tang,^{1,2} and Mark Stoneking^{1,4}

¹Max Planck Institute for Evolutionary Anthropology, D-04103 Leipzig, Germany; ²Institute for Computational Biology, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai 20031, China; ³National Drug Screening Laboratory, China Pharmaceutical University, Nanjing City 21009, China

The human salivary microbiome may play a role in diseases of the oral cavity and interact with microbiomes from other parts of the human body (in particular, the intestinal tract), but little is known about normal variation in the salivary microbiome. We analyzed 14,115 partial (~500 bp) 16S ribosomal RNA (rRNA) sequences from saliva samples from 120 healthy individuals (10 individuals from each of 12 worldwide locations). These sequences could be assigned to 101 known bacterial genera, of which 39 were not previously reported from the human oral cavity; phylogenetic analysis suggests that an additional 64 unknown genera are present. There is high diversity in the salivary microbiome within and between individuals, but little geographic structure. Overall, ~13.5% of the total variance in the composition of genera is due to differences among individuals, which is remarkably similar to the fraction of the total variance in neutral genetic markers that can be attributed to differences among human populations. Investigation of some environmental variables revealed a significant association between the genetic distances among locations and the distance of each location from the equator. Further characterization of the enormous diversity revealed here in the human salivary microbiome will aid in elucidating the role it plays in human health and disease, and in the identification of potentially informative species for studies of human population history.

[Supplemental material is available online at www.genome.org. The sequence data from this study have been submitted to GenBank (<http://www.ncbi.nlm.nih.gov/Genbank/>) under accession nos. EU984515–EU998629.]

Molecular analyses have revealed enormous variation in the microbiome of diverse environments, including the human body (Eckburg et al. 2005; Dethlefsen et al. 2007; Grice et al. 2008; Oakley et al. 2008). These studies have revealed a wealth of not-yet-cultivated bacterial species, and several projects to study the human microbiome have been proposed and/or initiated (Turnbaugh et al. 2007; Mullard 2008). However, comparatively little attention has been paid to the human salivary microbiome, as most studies of the human oral cavity have focused on identifying bacteria that might be associated with diseases (Becker et al. 2002; Kumar et al. 2003; Diaz et al. 2006; Kilian et al. 2006; Machado de Oliveira et al. 2007; Faveri et al. 2008). More than 600 different species have been described from the human oral cavity (Fabian et al. 2008), but only limited information is available on the normal microflora of healthy individuals (Aas et al. 2005; Kang et al. 2006). A comprehensive understanding of the breadth of diversity in the human oral cavity microbiome is necessary to fully gauge the role of bacteria in diseases of the oral cavity (Aas et al. 2005; Paster et al. 2006; Fabian et al. 2008). Moreover, as the oral cavity is often the entry point of bacteria into the body, there may be important interactions between the saliva microbiome and other microbiomes in the human body, in particular, that of the intestinal tract (Schipper et al. 2007; Fabian et al. 2008).

A further reason for analyzing the human salivary microbiome is that since saliva is increasingly preferred in sampling humans as a source of DNA for epidemiologic and population genetic studies (Quinque et al. 2006; Hansen et al. 2007), it would be useful to identify bacterial taxa in saliva that may be able to provide insights into human population structure and migrations.

When humans migrate, they take their bacteria with them, and it has previously been shown that patterns of variation in the stomach bacteria *Helicobacter pylori* correspond closely to migrations inferred from analyses of human DNA variation (Falush et al. 2003; Linz et al. 2007). Moreover, it was shown that two human ethnic groups, which could not be distinguished on the basis of human DNA markers, could be distinguished based on their patterns of *H. pylori* variation (Wirth et al. 2004). This result could reflect either genetic isolation of these two communities that is too recent to be reflected in the human DNA markers, or some cultural factor that influences *H. pylori* variation (such as diet) that differs between these communities; in either event, the *H. pylori* variation does provide insights into human populations that are not revealed by analysis of human DNA markers.

However, sampling *H. pylori* requires stomach biopsy material, so there would be obvious advantages if similarly informative bacterial species could be identified in saliva. The most informative bacterial species for this purpose should exhibit high levels of vertical versus horizontal transmission. Although this information remains largely unknown for bacteria in the oral cavity, there is some preliminary evidence from mother–child (Li et al. 2007) and twin (Corby et al. 2007) studies to suggest a significant role for vertical transmission. And even bacterial species that exhibit high levels of horizontal transfer may nonetheless provide evidence of recent contact between populations that would not be revealed by human DNA markers. Characterization of patterns of diversity in the human salivary microbiome thus may provide useful insights into human population structure and migration.

Here, we present the first description of patterns of global diversity in any human microbiome, based on analyses of partial 16S ribosomal RNA (rRNA) sequences from 120 individuals from diverse locations around the world. Our results provide a framework for further investigations of the ecological and cultural factors that may influence the composition of the human salivary

⁴Corresponding author.

E-mail stoneking@eva.mpg.de; fax 49-341-3550-555.

Article published online before print. Article and publication date are at <http://www.genome.org/cgi/doi/10.1101/gr.084616.108>.




Figure 1. Map of the sampling locations. Ten saliva samples were obtained from each of the 12 sampling locations, for a total of 120 saliva samples analyzed.

microbiome and identify promising taxa for further characterization that may provide novel insights into human population structure, migration, and contact.

Results

Saliva samples were obtained from 120 individuals, consisting of 10 individuals from each of two locations from each of six geographic regions (Fig. 1). A highly variable portion of the 16S rRNA gene of ~500 bp was amplified and cloned, and approximately 120 clones were sequenced from each individual; this number of clones is sufficient to capture most of the variation at the level of bacterial genera inferred from the partial 16S rRNA sequences (Supplemental Fig. S1). A total of 14,691 sequences were obtained, which were then analyzed for possible chimeras or other artifacts (Ashelford et al. 2006). We identified 576 potentially chimeric sequences, which represents ~3.9% of the sequences. This is less than the average of 9% potentially chimeric sequences reported previously for full-length 16S rRNA clone libraries (Ashelford et al. 2006), probably because our partial 16S rRNA sequences provide a smaller target for jumping PCR or other processes that produce chimeric sequences. Excluding these 576 potentially chimeric sequences leaves 14,115 sequences for further analysis (Table 1); these sequences have been deposited in GenBank under accession numbers EU984515–EU998629. The sequences were analyzed in two ways: First, a genus was assigned to each sequence by comparison to the RDPII database (Cole et al. 2007), and the compo-

sition of genera across individuals was subsequently analyzed. Second, all sequences were used to generate a phylogenetic tree, which was subsequently analyzed.

The distribution of bacterial genera assigned to the sequences from each individual (Supplemental Tables S1–S12) varied greatly. Overall, 196 sequences (1.4%) did not match any sequence in RDPII above the cutoff value of 90% similarity, while 57 sequences (0.4%) matched a sequence in RDPII for which the genus was unclassified. To estimate the number of genera that might be present among these unknown sequences, a phylogenetic tree was constructed (Supplemental Fig. S2). There are 64 clusters of sequences in this tree at the cutoff value of 90% similarity, indicating that there are an estimated 64 new genera among these sequences.

The number of genera identified in each individual ranged from six to 30 (Supplemental Tables S1–S12), while the number of genera per location ranged from 39 to 55 (Table 1), which is comparable to previous studies of the oral cavity microbiome (Aas et al. 2005; Paster et al. 2006). In total, 101 different genera were identified, 39 of which have not been previously reported from the human oral cavity (Supplemental Table S13). In combination with the unknown genera, we have therefore identified an estimated 103 previously undescribed bacterial genera from human saliva.

A few genera are quite frequent, while most are rare (Supplemental Fig. S3); 16 genera were observed only once, while the most frequent genus (*Streptococcus*) accounted for 22.7% of the sequences (Supplemental Table S13). In general, we identified 85%–95% of the genera that have previously been found in molecular surveys of the human oral cavity (Kumar et al. 2003; Aas et al. 2005; Diaz et al. 2006; Kang et al. 2006; Paster et al. 2006); the previously reported genera not found in this study include *Eikenella*, *Lautropia*, *Synergistes*, and *Bacteroides*.

The relative abundance of each genus in each individual appears qualitatively to be fairly uniform (Fig. 2). To quantitate differences in the distribution of genera, we apportioned the total variance within each location into between-individual and within-individual components (Table 1). California and the Congo showed the biggest differences between individuals, while Georgia and Turkey showed the smallest differences between individuals. Overall, ~13.5% of the variation in the distribution of genera can be attributed to differences between individuals (i.e., any one individual has ~86.5% of the variation that is observed when all locations are sampled).

Table 1. Summary statistics for the 12 sampling locations, based on the composition of bacterial genera identified from the partial 16S rRNA sequences

	No. of sequences	Unclassified (%)	Unknown (%)	No. of genera	Variance between individuals (%)	Variance within individuals (%)
Argentina (Arg)	1188	0.51	0.76	46	6.8	93.2
Bolivia (Bol)	1221	0.08	0.90	43	19.1	80.9
California (Cal)	1182	0.34	0.68	39	20.4	79.6
Louisiana (Lou)	1115	0.54	2.60	41	7.9	92.1
Germany (Ger)	1155	0.26	1.13	43	5.9	94.1
Poland (Pol)	1167	0.60	1.37	46	13.9	86.1
Georgia (Geo)	1186	0.84	1.69	55	4.6	95.4
Turkey (Tur)	1145	0.17	2.45	45	4.6	95.4
Congo (Con)	1178	0.42	1.19	42	25.4	74.6
South Africa (Saf)	1247	0.24	1.52	41	15.1	84.9
China (Chi)	1149	0.35	1.91	49	7.0	93.0
Philippines (Phi)	1182	0.51	0.59	47	11.2	88.8
Total	14,115	0.40	1.39	101	13.5	86.5




Figure 2. Heat plot of the abundance of each bacterial genus in each individual, based on the partial 16S rRNA sequences. Each horizontal row corresponds to an individual saliva sample, with different colored shadings indicating the 10 individuals from each sampling location, and sampling locations from the same broad geographic region indicated by similar colors. Each column is a genus, with the numbers corresponding to the genus, as in Supplemental Table S13. The abundance of each genus (columns) is indicated by the grayscale value, according to the scale at the bottom of the plot.

The between-individual component of the total variation can be further subdivided into the amount of variation among individuals from the same location, and the amount of variation among individuals from different locations. The former estimate is 11.7%, and the latter is just 1.8%. Although both estimates are significantly different from zero ($P < 0.001$) based on a permutation analysis (Excoffier et al. 2005), this analysis indicates that differences in the composition of genera are larger among individuals from the same location than among individuals from

different locations, and thus that differences in the composition of genera among individuals are not structured by geography.

Another way to examine this question is to determine the probability that two sequences chosen at random will come from different genera, when the sequences are chosen from the same individual, from different individuals from the same location, or from different individuals from different locations (Fig. 3). The average probability that two sequences are from different genera when chosen at random from the same individual is 0.772, while for two sequences chosen from two different individuals from the same location, the corresponding probability is 0.875, which is significantly higher (Mann-Whitney test, $P < 0.001$). However, the average probability that two sequences are from different genera when chosen from two individuals from different locations is 0.892, which is not significantly different from the above probability of 0.875 for two sequences chosen from two individuals from the same location (Mann-Whitney test, $P > 0.05$). Thus, there is significantly more variation among sequences from different individuals than among sequences from the same individual, but there is not significantly more variation among individuals from different locations than among individuals from the same location.

The genetic structure of human populations is strongly influenced by geography (Ramachandran et al. 2005; Li et al. 2008). However, a multidimensional-scaling (MDS) plot based on the differences in the composition of genera among locations does not indicate any strong influence of geography (Fig. 4A). The first dimension separates the Congo from the other locations, while the second dimension separates Louisiana from the other locations, and overall there is no grouping by geography. In agreement with the MDS plot, the distances among locations based on the composition of genera (Supplemental Table S14) are not correlated with the geographic distances among locations (Mantel test, $r = -0.07$,

$P = 0.65$). Moreover, an MDS plot based on differences in the composition of genera among individuals, rather than locations, also does not reveal any geographic clustering (Supplemental Fig. S4).

The above analyses are all based on classifying the genus of each sequence; to assess independently the geographic structure in the data without first classifying the sequences, we also carried out a UniFrac analysis (Lozupone and Knight 2005). We constructed a phylogenetic tree for the entire set of 14,115 sequences, which was then used to estimate the UniFrac distance (the fraction




Figure 3. Frequency distribution of the probability that two sequences will be assigned to different genera when drawn from the same individual (values are for each of the 120 individuals); different individuals from the same location (values are for all pairwise comparisons of the 10 individuals from the same location, for all 12 locations); and different individuals from different locations (values are for all pairwise comparisons of the 12 locations).

of the tree that is not shared in common) between each pair of locations. The PCA plot of the UniFrac distances (Fig. 4B) is quite similar to the MDS plot for the distances based on genus composition (Fig. 4A), and the UniFrac distances (Supplemental Table S14) are similarly not correlated with the geographic distances among locations (Mantel test, $r = -0.14$, $P = 0.79$). In agreement with the close similarity between the plots (Fig. 4), the UniFrac distances are highly and significantly correlated with the genus composition distances (Mantel test, $r = 0.94$, $P < 0.001$). Thus, the lack of any significant geographic structure to the human saliva microbiome is not a consequence of the procedure used to assign the genus of each sequence, but is an inherent property of the sequences retrieved from the saliva samples.

Although overall there is no significant geographic patterning with respect to the human salivary microbiome, specific genera do vary significantly in frequency among locations (Fig. 2; Supplemental Table S13). For example, the most extreme variation in frequency across locations is exhibited by *Enterobacter*, which accounts for 28% of the sequences obtained from the Congo (Supplemental Table S13) but was completely absent from California, China, Germany, Poland, and Turkey. Several other genera were observed at higher frequency in the Congo than elsewhere (Fig. 2; Supplemental Table S13), reinforcing the uniqueness of the salivary microbiome from the Congo (Fig. 4; Table 1). Another genus that occurs at relatively high frequency in several individuals from one particular location is *Serratia* (Bolivia). However, several genera that vary significantly in frequency (Supplemental Table S13) occur at high frequency only in one individual from a particular location (Fig. 2); more intensive sampling would be required to determine if these genera are truly characteristic of these particular locations.

A prime question remains the identification of environmental and cultural factors that influence the human salivary microbiome. As a first step in this direction, we carried out a multivariate regression analysis (Anderson 2001, 2004) of associations between four factors (population size, distance from the equator, average annual temperature, and average annual rainfall) (Supplemental Table S15) and the UniFrac distances among the 12 locations. We also tested for associations between age and gender of the donor and the F_{ST}

distances based on the composition of bacterial genera for each donor. The only significant association involved the distance of each location from the equator, which explained $\sim 24\%$ of the variation in the UniFrac distances (Fig. 5). However, this apparently significant association should be viewed with caution. Many factors vary with distance from the equator (such as UV index), so further work will be required to ascertain what aspect(s) of the distance from the equator is responsible for this association. Moreover, the association is primarily driven by the Congo, which has high distances to other locations and is also closest to the equator; investigation of additional populations that are similarly close to the equator would be necessary to verify this association.

Discussion

The analyses of the human salivary microbiome presented here provide the first detailed investigation of global patterns of diversity in any human microbiome. There is high diversity in the human salivary microbiome; assignment of the partial 16S rRNA sequences generated in this study to bacterial genera via comparison to the RDP-II database identified a total of 101 bacterial genera, including 39 that had not been previously described from the human oral cavity. Moreover, we estimate that an additional 64 unknown genera are represented in the sequences. Understanding the role of these known and unknown genera in the health of the human oral cavity and how they interact with the microbiomes of other parts of the human body (in particular, the intestinal tract) would be of considerable interest.

A potential criticism of this study is that there are many environmental and cultural factors (in particular, diet and hygiene) that were not controlled for during the sampling. Indeed, a limited investigation of some environmental factors revealed a significant association between the genetic distances among locations and the distance of the location from the equator (Fig. 5). However, we emphasize that a main conclusion of our study (discussed below) is that the human salivary microbiome does not vary greatly between different geographic locations. Therefore, if these uncontrolled environmental and/or cultural factors are having a significant impact on our results, then controlling for them would presumably result in even less geographic differentiation in the patterns of variation. That is, if (for example) differences in diet between locations are responsible for differences in the salivary microbiome composition, then controlling for diet would further decrease the already small differences between locations.

Another potential weakness is that possible temporal change in the salivary microbiome was not studied. In this respect, the sampling in this study should be regarded as “environmental,” that is, a single snapshot in time of the salivary microbiome diversity of an individual was captured, much as sampling the microbiome of hot springs or other environmental locations often use a single time point. To be sure, it would be of considerable interest to determine the extent of change in an individual’s salivary microbiome over time and how it might be influenced by other factors. For example, what happens to the salivary microbiome if individuals change their diet? And what happens if an individual moves to a new




Figure 4. Graphical depictions of the relationships among sampling locations based on the partial 16S rRNA sequences. (A) MDS plot, based on linearized F_{ST} distances (Slatkin 1995) between each pair of locations, calculated from the genus composition. The location abbreviations are as in Table 1. The stress value is 0.03. (B) PCA plot, based on the UniFrac distances (Lozupone and Knight 2005); same abbreviations as in A. The percentage of the total variance explained by each PC is indicated in parentheses.

location? Although these questions were not addressed in this study, the results of this study do provide a framework for further studies designed to answer such questions about the temporal stability of the human salivary microbiome.

This high diversity in the human salivary microbiome does not appear to be geographically structured (Fig. 2). While there is significantly more diversity in bacterial genera compared from different individuals than from the same individual, the diversity among individuals from the same location is nearly the same as the diversity among individuals from different locations (Fig. 3). Overall, the between-population component of the variance in the bacterial genera composition is ~13.5% (Table 1), which means that any one individual contains on average 86.5% of the total variance observed when all individuals are sampled. This is remarkably similar to the amount of between-population variation typically observed among human populations for neutral genetic markers (Romualdi et al. 2002; Li et al. 2008). Thus, in this sense, the distribution of genera among individuals is behaving as a neutral genetic marker, if one equates the bacterial composition of individuals to the genetic composition of human populations.

It should be emphasized that this conclusion of an overall lack of geographic structure extends only to the pool of 16S rRNA

sequences and the bacterial genera identified from them. Sequence variation within particular bacterial taxa may very well exhibit geographic structure that would provide novel insights into human population structure, relationships, and migrations, as has been previously observed for *H. pylori* (Falush et al. 2003; Linz et al. 2007). Multi-locus sequence typing (MLST) is a particularly promising approach (Urwin and Maiden 2003; Maiden 2006) that has shown geographic structure in some bacteria that was not evident in 16S rRNA comparisons (Margos et al. 2008). Additional characterization of coding sequence variation in particular taxa by MLST or other means would be highly desirable, and may very well reveal geographic structure that is not apparent in the overall composition of the salivary microbiome. For tracing human migrations, it would be most useful to focus on bacteria that are likely to be transferred primarily vertically (i.e., within families or households). However, even bacteria that are primarily transferred horizontally may provide useful markers of population contact that is too recent to be detected by analysis of human DNA polymorphisms. Perhaps the most useful taxa to start with would be those that are both widespread (i.e., found in many different locations) and frequent (i.e., found in multiple individuals from a location); these would include *Streptococcus*, *Prevotella*, *Veillonella*, *Neisseria*, *Haemophilus*, *Rothia*, *Porphyromonas*, and *Fusobacterium* (Fig. 2). Together, these eight genera account for >70% of the total number of sequences

(Supplemental Table S13).

In conclusion, our results provide a framework for further studies of human microbiome diversity and for investigations of the environmental, dietary, and genetic factors that influence an individual's salivary microbiome. These, in turn, will lead to a better understanding of the role of the salivary microbiome in diseases of the oral cavity and how the salivary microbiome interacts with the intestinal and other microflora of the human body. In addition, further characterization of some of the taxa identified in this study by MLST or other means should provide useful markers for studies of human population history and/or recent population contact.

Methods

Samples and DNA extraction

Saliva samples from 10 individuals from each of 12 locations (two locations from each of six geographic regions) (Fig. 1) were collected with informed consent. The criteria for selection of individuals were that individuals should not be from the same household, and they should not have been away from the sampled




Figure 5. Amount of variation explained (in percent) in the F_{ST} distances among individuals (for the age and gender of the donors), and in the UniFrac distances among locations (for the variables population size, distance from the equator, average annual temperature, and average annual rainfall) (Supplemental Table S15). There is a significant association between the UniFrac distances and the variable "Distance from Equator" ($P = 0.006$).

location within the past two months. Although dental health was not examined in detail, all individuals were in good health, and no individual was suffering from any obvious dental disease at the time of sampling.

Two milliliters of saliva was collected from each individual into a tube containing an equal volume of lysis buffer (Quinque et al. 2006). Samples were stored at ambient temperature and transported to the laboratory (MPI-EVA, Leipzig, Germany), and DNA was extracted as described previously (Quinque et al. 2006).

PCR amplification of the microbial 16S rRNA gene

We designed primers to amplify an informative segment of the microbial 16S rRNA gene that would be of optimal size for cloning and DNA sequence analysis. We downloaded and aligned ~100,000 complete microbial 16S rRNA sequences from the Ribosomal Database Project II (RDPII) database (Cole et al. 2007) and searched for highly conserved segments that flank highly variable segments of the gene. The gene fragment we amplified corresponds to nucleotide positions 2,769,690 to 2,770,259 in the *Escherichia coli* genome (Blattner et al. 1997), with an amplicon length of 570 bp. The primer sequences used in this study are: forward primer, 5'-CTACGTGCCAGCAGCCGCGG-3'; and reverse primer, 5'-CTCACGACACGAGCTGACGA-3'. PCR amplification was carried out in a 50- μ L total volume, including 38 μ L of ddH₂O, 5 μ L of 10 $\times$ PCR buffer (500 mM KCl, 100 mM Tris-HCl at pH 9.0, 1.0% Triton X-100), 3 μ L of 25 mM MgCl₂, 1 μ L of 10 mM dNTPs (10 mM each dATP, dTTP, dGTP, dCTP), 1 μ L of 20 μ M forward primer, 1 μ L of 20 μ M reverse primer, 0.5 μ L of *Taq* polymerase (Applied Biosystems), and 20 ng of template DNA, for 35 cycles using the following conditions: denaturation step for 1 min at 95°C, annealing step for 1 min at 62°C, and extension step for 1 min at 72°C.

Cloning and sequencing of amplicons

Amplicons were cloned using the TOPO TA cloning kit (Invitrogen), following the protocol recommended by the supplier. Colony PCR was carried out using the M13 forward and reverse primers supplied with the cloning kit in a 50- μ L total volume that included the same reagents described above, to which a small amount of cells were added with a sterile pipette tip, and the following thermocycling conditions: an initial incubation for 5 min at 95°C to lyse cells and denature the DNA, followed by 30 cycles of 1 min at 95°C, 1.5 min at 54°C, and 1 min at 72°C, followed by

a final incubation for 5 min at 72°C. The forward stands of colony PCR products with appropriate inserts were sequenced using the 16S rRNA PCR primers and the Big Dye DNA Sequencing Kit (Perkin-Elmer), following the protocol recommended by the supplier, and an ABI 3730 automated DNA sequencer. Sequence quality was assessed via the software package DNASTAR (DNASTAR Inc.).

Data analysis

Chimeric or other potentially artifactual sequences were identified with the Malard program (Ashelford et al. 2006), using an iterative procedure and a quantile value of 95% as the cutoff line for determining outliers. This procedure identified 576 potentially chimeric sequences, which were excluded from further analyses, leaving 14,115 sequences.

These 14,115 sequences were then searched against the RDPII database (Cole et al. 2007), using the online program SeqMatch (Wang et al. 2007) and a threshold setting for the similarity score of 90%, to assign a genus to each sequence. The appropriate similarity score for defining bacterial genera is a matter of debate, generally ranging from 90% to 95%; we chose a value at the low end of the range, in accordance with some previous studies (Pei et al. 2004; Gao et al. 2007), in order to minimize falsely designating new taxa (at the possible expense of mis-assigning some taxa). In the event of multiple matches to sequences in the database, the genus was assigned according to the highest similarity score. Comparing the sequences in a similar fashion to GenBank resulted in a highly similar assignment of genera, except that more sequences were classified as unknown (data not shown). Supplemental Tables S1–S12 give the frequency of each genus for each individual, as well as the frequency of unclassified sequences (those that matched a sequence in RDPII, but the genus of the matching sequence has not been determined) and unknown sequences (those that did not match any sequence in RDPII above the 90% threshold). The apportionment of variation based on the frequency distribution of genera within and between individuals and locations, and linearized F_{ST} distances (Slatkin 1995) were calculated with Arlequin 3.1 (Excoffier et al. 2005); for these analyses, each genus was considered equivalent to an allele at a single haploid locus. Multidimensional scaling (MDS) analysis was carried out using STATISTICA 7.1 (StatSoft, Inc.).

We also compared the differences among sequences from different individuals and locations, without first assigning sequences to genera, using the UniFrac method (Lozupone and Knight 2005; Lozupone et al. 2006). To implement this method, we first carried out a multiple alignment of all of the sequences using ClustalW2 (Larkin et al. 2007). A matrix of the sequence differences between each pair of sequences was then determined using the F84 model (Kishino and Hasegawa 1989) as implemented in the DNADIST program in PHYLIP 3.67 (Felsenstein 1989). A UPGMA tree was then constructed from the pairwise distance matrix, using the NEIGHBOR program in PHYLIP, and the UniFrac distance metric (Lozupone et al. 2006) was calculated between each pair of locations with the online resource (<http://bmf2.colorado.edu/unifrac/>). These distances were used as input for principal coordinates analysis; a neighbor-joining tree resulted in UniFrac distances that were nearly identical to the distances based on the UPGMA tree (Mantel test, $r = 0.97$, $P < 0.001$), and the

PC plot based on the neighbor-joining distances was virtually identical to that based on UPGMA distances (data not shown). Multivariate regression analysis (Anderson 2001; McArdle and Anderson 2001) for associations between the genetic distance matrices and potential predictor variables of interest was carried out with the DISTLM program (Anderson 2004); this method treats the entire distance matrix as the response variable and uses a permutation method to test for associations with potential explanatory variables, in a regression analysis framework.

Acknowledgments

We thank all individuals who kindly donated a sample for this study; Rebecca Atencia, Jarek Bryk, Heather Buckner, Bryan Buckner, Daniel Corach, Anne Fischer, Michel Halbwax, Janet Kelso, Marina Nagveradze, Samra Sardas, Beth Trachtenberg, and Guido Valverde for assistance with sample collections; Susanne Gutsmuths and Anna Schönberg for technical assistance; and Micah Hamady for assistance with the UniFrac distance calculations. Funding was provided by the Max Planck Society.

References

- Aas, J., Paster, B.J., Stokes, L.N., Olsen, I., and Dewhirst, F.E. 2005. Defining the normal bacterial flora of the oral cavity. *J. Clin. Microbiol.* **43**: 5721–5732.
- Anderson, M.J. 2001. A new method for non-parametric multivariate analysis of variance. *Austral. Ecol.* **26**: 32–46.
- Anderson, M.J. 2004. *DISTLM v.5: A FORTRAN computer program to calculate a distance-based multivariate analysis for a linear model*. Department of Statistics, University of Auckland, New Zealand.
- Ashelford, K.E., Chuzhanova, N.A., Fry, J.C., Jones, A.J., and Weightman, A.J. 2006. New screening software shows that most recent large 16S rRNA gene clone libraries contain chimeras. *Appl. Environ. Microbiol.* **72**: 5734–5741.
- Becker, M.R., Paster, B.J., Leys, E.J., Moeschberger, M.L., Kenyon, S.G., Galvin, J.L., Boches, S.K., Dewhirst, F.E., and Griffen, A.L. 2002. Molecular analysis of bacterial species associated with childhood caries. *J. Clin. Microbiol.* **40**: 1001–1009.
- Blattner, F.R., Plunkett III, G., Bloch, C.A., Perna, N.T., Burland, V., Riley, M., Collado-Vides, J., Glasner, J.D., Rode, C.K., Mayhew, G.F., et al. 1997. The complete genome sequence of *Escherichia coli* K-12. *Science* **277**: 1453–1474.
- Cole, J.R., Chai, B., Farris, R.J., Wang, Q., Kulam-Syed-Mohideen, A.S., McGarrell, D.M., Bandela, A.M., Cardenas, E., Garrity, G.M., and Tiedje, J.M. 2007. The ribosomal database project (RDP-II): Introducing myRDP space and quality controlled public data. *Nucleic Acids Res.* **35**: D169–D172.
- Corby, P.M., Bretz, W.A., Hart, T.C., Schork, N.J., Wessel, J., Lyons-Weiler, J., and Paster, B.J. 2007. Heritability of oral microbial species in caries-active and caries-free twins. *Twin Res. Hum. Genet.* **10**: 821–828.
- Dethlefsen, L., McFall-Ngai, M., and Relman, D.A. 2007. An ecological and evolutionary perspective on human-microbe mutualism and disease. *Nature* **449**: 811–818.
- Diaz, P.I., Chalmers, N.I., Rickard, A.H., Kong, C., Milburn, C.L., Palmer, R.J., and Kolenbrander, P.E. 2006. Molecular characterization of subject-specific oral microflora during initial colonization of enamel. *Appl. Environ. Microbiol.* **72**: 2837–2848.
- Eckburg, P.B., Bik, E.M., Bernstein, C.N., Purdom, E., Dethlefsen, L., Sargent, M., Gill, S.R., Nelson, K.E., and Relman, D.A. 2005. Diversity of the human intestinal microbial flora. *Science* **308**: 1635–1638.
- Excoffier, L., Laval, G., and Schneider, S. 2005. Arlequin (version 3.0): An integrated software package for population genetics data analysis. *Evol. Bioinform. Online* **1**: 47–50.
- Fabian, T.K., Fejerdy, P., and Csermely, P. 2008. Salivary genomics, transcriptomics, and proteomics: The emerging concept of the oral ecosystem and their use in the early diagnosis of cancer and other diseases. *Cur. Genomics* **9**: 11–21.
- Falush, D., Wirth, T., Linz, B., Pritchard, J.K., Stephens, M., Kidd, M., Blaser, M.J., Graham, D.Y., Vacher, S., Perez-Perez, G.I., et al. 2003. Traces of human migrations in *Helicobacter pylori* populations. *Science* **299**: 1582–1585.
- Faveri, M., Mayer, M.P.A., Feres, M., de Figueiredo, L.C., Dewhirst, F.E., and Paster, B.J. 2008. Microbiological diversity of generalized aggressive periodontitis by 16S rRNA clonal analysis. *Oral Microbiol. Immunol.* **23**: 112–118.
- Felsenstein, J. 1989. PHYLIP—Phylogeny Inference Package (version 3.2). *Cladistics* **5**: 164–166.
- Gao, Z., Tseng, C., Pei, Z., and Blaser, M.J. 2007. Molecular analysis of human forearm superficial skin bacterial biota. *Proc. Natl. Acad. Sci.* **104**: 2927–2932.
- Grice, E.A., Kong, H.H., Renaud, G., Young, A.C., Bouffard, G.G., Blakesley, R.W., Wolfsberg, T.G., Turner, M.L., and Segre, J.A. 2008. A diversity profile of the human skin microbiota. *Genome Res.* **18**: 1043–1050.
- Hansen, T.V., Simonsen, M.K., Nielsen, F.C., and Hundrup, Y.A. 2007. Collection of blood, saliva, and buccal cell samples in a pilot study on the Danish nurse cohort: Comparison of the response rate and quality of genomic DNA. *Cancer Epidemiol. Biomarkers Prev.* **16**: 2072–2076.
- Kang, J.G., Kim, S.H., and Ahn, T.Y. 2006. Bacterial diversity in the human saliva from different ages. *J. Microbiol.* **44**: 572–576.
- Kilian, M., Frandsen, E.V.G., Haubek, H., and Poulsen, K. 2006. The etiology of periodontal disease revisited by population genetic analysis. *Periodontol.* **2000** **42**: 158–179.
- Kishino, H. and Hasegawa, M. 1989. Evaluation of the maximum likelihood estimate of the evolutionary tree topologies from DNA sequence data, and the branching order in hominoidae. *J. Mol. Evol.* **29**: 170–179.
- Kumar, P.S., Griffen, A.L., Barton, J.A., Paster, B.J., Moeschberger, M.L., and Leys, E.J. 2003. New bacterial species associated with chronic periodontitis. *J. Dent. Res.* **82**: 338–344.
- Larkin, M.A., Blackshields, G., Brown, N.P., Chenna, R., McGettigan, P.A., McWilliam, H., Valentin, F., Wallace, I.M., Wilm, A., Lopez, R., et al. 2007. Clustal W and Clustal X version 2.0. *Bioinformatics* **23**: 2947–2948.
- Li, Y., Ismail, A.I., Ge, Y., Tellez, M., and Sohn, W. 2007. Similarity of bacterial populations in saliva from African-American mother-child dyads. *J. Clin. Microbiol.* **45**: 3082–3085.
- Li, J.Z., Absher, D.M., Tang, H., Southwick, A.M., Casto, A.M., Ramachandran, S., Cann, H.M., Barsh, G.S., Feldman, M., Cavalli-Sforza, L.L., et al. 2008. Worldwide human relationships inferred from genome-wide patterns of variation. *Science* **319**: 1100–1104.
- Linz, B., Balloux, F., Moodley, Y., Manica, A., Liu, H., Roumagnac, P., Falush, D., Stamer, C., Prugnolle, F., van der Merwe, S.W., et al. 2007. An African origin for the intimate association between humans and *Helicobacter pylori*. *Nature* **445**: 915–918.
- Lozupone, C. and Knight, R. 2005. UniFrac: A new phylogenetic method for comparing microbial communities. *Appl. Environ. Microbiol.* **71**: 8228–8235.
- Lozupone, C., Hamady, M., and Knight, R. 2006. UniFrac—An online tool for comparing microbial community diversity in a phylogenetic context. *BMC Bioinformatics* **7**: 371. doi: 10.1186/1471-2105-7-371.
- Machado de Oliveira, J.C., Siqueira, J.F., Roca, I.N., Baumgartner, J.C., Xia, T., Peixoto, R.S., and Rosado, A.S. 2007. Bacterial community profiles of endodontic abscesses from Brazilian and USA subjects as compared by denaturing gradient gel electrophoresis analysis. *Oral Microbiol. Immunol.* **22**: 14–18.
- Maiden, M.C.J. 2006. Multilocus sequence typing of bacteria. *Annu. Rev. Microbiol.* **60**: 561–588.
- Margos, G., Gatewood, A.G., Aanensen, D.M., Hanincova, K., Terekhova, D., Vollmer, S.A., Cornet, M., Piesman, J., Donaghy, M., Bormane, A., et al. 2008. MLST of housekeeping genes captures geographic population structure and suggests a European origin of *Borrelia burgdorferi*. *Proc. Natl. Acad. Sci.* **105**: 8730–8735.
- McArdle, B.H. and Anderson, M.J. 2001. Fitting multivariate models to community data: A comment on distance-based redundancy analysis. *Ecology* **82**: 290–297.
- Mullard, A. 2008. Microbiology: The inside story. *Nature* **453**: 578–580.
- Oakley, B.B., Fiedler, T.L., Marrazzo, J.M., and Fredricks, D.N. 2008. The diversity of human vaginal bacterial communities and their association with clinically defined bacterial vaginosis. *Appl. Environ. Microbiol.* **74**: 4898–4909.
- Paster, B.J., Olsen, I., Aas, J., and Dewhirst, F.E. 2006. The breadth of bacterial diversity in the human periodontal pocket and other oral sites. *Periodontol.* **2000** **42**: 80–87.
- Pei, Z., Bini, E.J., Yang, L., Zhou, M., Francois, F., and Blaser, M.J. 2004. Bacterial biota in the human distal esophagus. *Proc. Natl. Acad. Sci.* **101**: 4250–4255.
- Quinque, D., Kittler, R., Kayser, M., Stoneking, M., and Nasidze, I. 2006. Evaluation of saliva as a source of human DNA for population and association studies. *Anal. Biochem.* **353**: 272–277.
- Ramachandran, S., Deshpande, O., Rosemann, C.C., Rosenberg, N.A., Feldman, M.W., and Cavalli-Sforza, L.L. 2005. Support from the relationship of genetic and geographic distance in human populations for a serial founder effect originating in Africa. *Proc. Natl. Acad. Sci.* **102**: 15942–15947.

- Romualdi, C., Balding, D., Nasidze, I.S., Risch, G., Robichaux, M., Sherry, S.T., Stoneking, M., Batzer, M.A., and Barbujani, G. 2002. Patterns of human diversity, within and among continents, inferred from biallelic DNA polymorphisms. *Genome Res.* **12**: 602–612.
- Schipper, R.G., Silletti, E., and Vingerhoeds, M.H. 2007. Saliva as research material: Biochemical, physicochemical and practical aspects. *Arch. Oral Biol.* **52**: 1114–1135.
- Slatkin, M. 1995. A measure of population subdivision based on microsatellite allele frequencies. *Genetics* **139**: 457–462.
- Turnbaugh, P.J., Ley, R.E., Hamady, M., Fraser-Liggett, C.M., Knight, R., and Gordon, J.I. 2007. The human microbiome project. *Nature* **449**: 804–810.
- Urwin, R. and Maiden, M.C. 2003. Multi-locus sequence typing: A tool for global epidemiology. *Trends Microbiol.* **11**: 479–487.
- Wang, Q., Garrity, G.M., Tiedje, J.M., and Cole, J.R. 2007. Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl. Environ. Microbiol.* **73**: 5261–5267.
- Wirth, T., Wang, X., Linz, B., Novick, R.P., Lum, J.K., Blaser, M., Morelli, G., Falush, D., and Achtman, M. 2004. Distinguishing human ethnic groups by means of sequences from *Helicobacter pylori*: Lessons from Ladakh. *Proc. Natl. Acad. Sci.* **101**: 4746–4751.

Received August 11, 2008; accepted in revised form December 30, 2008.

Global Diversity in the Human Salivary Microbiome

Ivan Nasidze, Jing Li, Dominique Quinque, Kun Tang, and Mark Stoneking

Supplemental Material

Figures S1-S4

Tables S1-S15




Figure S1. Fraction of different bacterial genera identified from different numbers of cloned partial 16S rRNA sequences. The indicated number of cloned sequences were sampled at random from 150 sequences from each of five individuals. Bacterial genera were identified by comparison to the RDPII database, as described in the Materials and Methods. Each different color line represents the results from one individual; the total number of genera observed from all 150 sequences for each individual is set equal to 100%. Since for all five individuals 120 sequences suffice to capture 94-100% of the bacterial genera observed in 150 sequences, an average of 120 sequences per individual were obtained for the 120 individuals in this study.




Figure S2. Phylogenetic (UPGMA) tree of the "unknown" sequences, based on pairwise estimates of dissimilarity (i.e., the fraction of different sites). These sequences fall into 64 different clusters at a similarity level of 90% (corresponding to a branch length of 0.1), and hence define 64 putative new genera.




Figure S3. Frequency distribution of the number of times a genus was observed. Note that the scale along the X-axis runs from 1-10, then from 10-100, 100-1000, and finally 1000-4000. For example, the value for 200 indicates that 5 genera were each observed between 200-300 times in the collection of 13,862 sequences for which a genus could be assigned.




Figure S4. MDS plot based on linearized F_{st} distances between each pair of individuals, calculated from the genus composition.

Table S1. Frequency (count) of each genus in each of the 10 saliva samples (Arg_a through Arg_j) from Buenos Aires, Argentina.

Genus	Arg_a	Arg_b	Arg_c	Arg_d	Arg_e	Arg_f	Arg_g	Arg_h	Arg_i	Arg_j	TOTAL
Actinobacillus	2	1	0	0	0	2	0	0	0	4	9
Actinomyces	0	2	0	4	4	0	0	0	6	5	21
Aggregatibacter	1	1	0	0	0	1	0	1	1	0	5
Alysiella	1	0	0	0	0	0	0	0	0	0	1
Anaerovorax	0	0	0	0	0	0	0	0	0	1	1
Atopobium	0	0	1	0	2	0	0	0	0	0	3
Bulleidia	0	0	0	1	0	0	0	0	0	0	1
Campylobacter	0	0	2	0	1	1	3	1	0	0	8
Capnocytophaga	0	4	0	0	0	0	0	0	1	0	5
Catonella	0	0	0	0	0	0	0	1	0	0	1
Chryseobacterium	0	0	0	0	0	1	0	0	0	0	1
Coprococcus	0	0	0	2	0	0	0	0	0	0	2
Corynebacterium	1	1	0	0	0	0	0	1	0	0	3
Derxia	0	1	0	0	0	1	4	1	0	0	7
Dialister	0	1	0	0	0	0	1	0	0	0	2
Enterobacter	0	0	0	0	0	0	5	0	0	0	5
Enterococcus	0	1	0	1	0	0	0	0	0	2	4
Eubacterium	2	1	0	1	0	2	1	1	1	1	10
Filifactor	0	0	0	0	0	1	0	0	0	0	1
Fusobacterium	3	2	0	2	6	4	2	3	3	1	26
Gemella	0	0	0	0	1	3	1	0	1	0	6
Granulicatella	1	4	1	2	0	2	2	0	3	0	15
Haemophilus	25	9	3	3	10	15	10	13	3	1	92
Kingella	0	0	0	0	0	0	1	1	0	0	2
Leptotrichia	1	1	3	5	1	3	2	7	2	2	27
Megasphaera	0	0	2	1	5	0	0	0	0	5	13
Micromonas	2	1	0	0	0	0	0	0	0	0	3
Moraxella	0	13	0	0	0	0	0	0	0	0	13
Neisseria	39	30	0	4	1	6	16	24	27	1	148

Genus	Arg_a	Arg_b	Arg_c	Arg_d	Arg_e	Arg_f	Arg_g	Arg_h	Arg_i	Arg_j	TOTAL
Oribacterium	0	0	0	3	0	4	0	4	0	2	13
Parascardovia	0	0	1	0	0	0	0	0	0	0	1
Peptostreptococcus	0	0	0	0	0	0	0	0	1	0	1
Porphyromonas	1	4	0	0	1	11	4	3	1	0	25
Prevotella	3	10	32	43	55	25	12	11	15	40	246
Rothia	12	5	6	8	8	1	7	8	13	3	71
Ruminococcus	0	0	0	0	1	0	0	0	0	2	3
Salmonella	0	0	0	0	0	0	1	0	0	0	1
Selenomonas	0	0	1	3	1	1	2	1	0	3	12
Simonsiella	0	0	0	0	0	0	1	0	0	0	1
Sneathia	0	0	0	0	0	2	0	0	0	0	2
Streptobacillus	0	0	0	0	0	2	0	0	0	1	3
Streptococcus	17	17	15	4	19	24	28	20	29	16	189
Syntrophococcus	0	0	0	0	0	0	0	0	0	1	1
TM7	0	0	0	1	0	1	2	0	1	0	5
Treponema	0	1	0	0	0	2	0	0	0	0	3
Veillonella	4	0	54	29	7	6	6	15	15	25	161
Unclassified	0	1	1	0	0	2	1	0	0	1	6
Unknown	0	2	0	2	1	1	0	1	1	1	9
TOTAL	115	113	122	119	124	124	112	117	124	118	1188

Table S2. Frequency (count) of each genus in each of the 10 saliva samples (Bol_a through Bol_j) from La Paz, Bolivia.

Genus	Bol_a	Bol_b	Bol_c	Bol_d	Bol_e	Bol_f	Bol_g	Bol_h	Bol_i	Bol_j	TOTAL
Actinobacillus	0	0	0	0	0	2	1	2	1	0	6
Actinomyces	0	0	0	2	0	0	0	0	1	0	3
Aggregatibacter	0	0	0	0	2	0	2	0	0	0	4
Alysiella	0	0	0	1	0	0	0	0	0	0	1
Atopobium	2	0	0	0	0	0	0	0	0	0	2
Bulleidia	0	1	1	0	0	0	1	0	1	0	4
Buttiauxella	0	0	0	5	0	0	0	0	0	0	5
Campylobacter	0	0	1	0	0	0	1	0	4	0	6
Capnocytophaga	0	0	0	0	1	0	0	0	0	0	1
Catonella	1	0	2	0	0	0	1	0	0	0	4
Chryseobacterium	0	0	0	0	1	0	0	0	0	0	1
Citrobacter	0	1	0	1	0	0	0	0	0	0	2
Cloacibacterium	1	0	1	0	1	0	0	0	0	0	3
Enterobacter	0	84	0	2	0	0	0	0	0	18	104
Enterococcus	0	0	0	0	2	0	0	0	0	0	2
Eubacterium	0	0	1	0	0	0	0	0	0	1	2
Fusobacterium	1	0	3	0	4	2	2	0	6	1	19
Gemella	0	2	0	0	8	3	0	0	0	3	16
Granulicatella	0	1	0	0	4	3	0	0	1	0	9
Haemophilus	9	1	10	4	3	7	14	3	5	4	60
Klebsiella	0	7	0	3	2	0	0	0	0	2	14
Leptotrichia	2	0	6	1	1	0	2	1	4	1	18
Mannheimia	0	0	0	0	1	0	0	0	0	0	1
Megasphaera	0	1	2	0	0	0	0	0	0	0	3
Mitsuokella	0	0	1	0	0	0	0	0	0	0	1
Neisseria	1	1	10	2	4	17	13	2	9	4	63
Oribacterium	0	0	1	1	1	3	0	1	1	0	8
Pasteurella	0	0	0	0	0	0	1	0	0	0	1
Porphyromonas	0	0	0	0	0	4	1	1	6	0	12

Genus	Bol_a	Bol_b	Bol_c	Bol_d	Bol_e	Bol_f	Bol_g	Bol_h	Bol_i	Bol_j	TOTAL
Prevotella	42	5	25	13	9	19	21	21	21	1	177
Rahnella	0	0	0	0	0	0	0	4	0	15	19
Rothia	5	0	2	0	2	2	5	1	4	2	23
Ruminococcus	1	0	0	0	1	0	0	0	0	0	2
Selenomonas	3	0	3	0	0	0	0	0	0	0	6
Serratia	0	2	0	79	8	0	0	63	0	61	213
Streptobacillus	0	0	0	0	0	0	0	1	0	0	1
Streptococcus	33	7	36	4	61	52	48	5	32	7	285
Syntrophococcus	1	0	1	0	0	0	0	0	0	0	2
TM7	0	0	1	0	0	0	0	0	0	0	1
Thermovirga	0	0	0	0	1	0	0	0	0	0	1
Treponema	0	0	1	0	0	0	1	0	0	0	2
Veillonella	32	2	18	6	3	1	5	10	18	6	101
Yersinia	0	0	0	0	0	0	0	0	0	1	1
Unclassified	0	0	0	0	0	0	0	1	0	0	1
Unknown	3	0	1	0	1	3	0	1	1	1	11
TOTAL	137	115	127	124	121	118	119	117	115	128	1221

Table S3. Frequency (count) of each genus in each of the 10 saliva samples (Cal_a through Cal_j) from Oakland, California.

Genus	Cal_a	Cal_b	Cal_c	Cal_d	Cal_e	Cal_f	Cal_g	Cal_h	Cal_i	Cal_j	TOTAL
Abiotrophia	0	1	0	4	0	0	0	1	0	1	7
Actinobacillus	0	1	0	0	0	1	0	0	0	0	2
Actinomyces	1	2	0	1	0	0	0	3	3	1	11
Aggregatibacter	0	0	0	4	0	0	0	0	1	0	5
Alysiella	0	0	0	0	0	0	1	0	0	0	1
Atopobium	0	1	0	1	0	2	0	0	3	0	7
Bulleidia	0	0	1	1	0	1	0	0	2	1	6
Campylobacter	2	0	1	0	0	1	0	0	1	0	5
Capnocytophaga	0	0	1	0	0	0	0	0	3	0	4
Catonella	0	0	0	0	0	0	0	0	0	1	1
Chryseobacterium	1	0	0	0	0	0	0	0	0	0	1
Cloacibacterium	0	0	0	0	0	0	0	1	0	0	1
Corynebacterium	0	0	0	0	0	0	0	0	1	2	3
Derxia	0	2	0	0	0	0	0	0	0	0	2
Enterococcus	0	1	1	1	0	1	0	0	0	0	4
Eubacterium	0	0	0	0	0	1	0	0	0	0	1
Fusobacterium	2	4	0	0	1	2	0	2	1	0	12
Gemella	0	1	0	3	0	0	0	7	1	4	16
Granulicatella	1	5	2	1	0	2	1	4	2	1	19
Haemophilus	32	14	10	2	0	5	0	5	12	1	81
Kingella	0	0	0	1	0	0	0	0	0	0	1
Leptotrichia	1	3	0	1	0	0	0	0	0	1	6
Megasphaera	0	1	2	0	1	2	0	0	0	0	6
Micromonas	0	0	0	1	0	0	0	0	1	0	2
Neisseria	4	16	14	4	0	3	0	23	8	1	73
Nesterenkonia	0	0	0	1	0	0	0	0	0	0	1
Oribacterium	5	2	0	0	1	1	0	2	0	1	12
Parascardovia	0	0	0	0	1	0	0	0	0	0	1
Peptostreptococcus	1	0	3	0	0	0	0	0	0	0	4

Genus	Cal_a	Cal_b	Cal_c	Cal_d	Cal_e	Cal_f	Cal_g	Cal_h	Cal_i	Cal_j	TOTAL
Porphyromonas	1	2	6	2	0	2	0	2	0	0	15
Prevotella	25	10	24	10	19	53	1	1	9	11	163
Rothia	14	17	13	20	2	1	1	8	2	8	86
Ruminococcus	0	0	0	0	0	0	0	0	0	1	1
Selenomonas	0	0	1	0	2	0	0	0	2	0	5
Serratia	0	0	0	0	0	0	118	0	0	0	118
Streptococcus	13	28	17	36	48	16	2	49	31	91	331
Syntrophococcus	0	0	1	0	1	0	0	0	1	0	3
TM7	0	0	2	3	0	2	0	1	1	1	10
Veillonella	8	9	17	24	45	11	0	4	19	6	143
Unclassified	0	0	0	0	1	0	0	1	2	0	4
Unknown	1	0	0	2	0	2	0	1	2	0	8
TOTAL	112	120	116	123	122	109	124	115	108	133	1182

Table S4. Frequency (count) of each genus in each of the 10 saliva samples (Lou_a through Lou_j) from Baton Rouge, Louisiana.

Genus	Lou_a	Lou_b	Lou_c	Lou_d	Lou_e	Lou_f	Lou_g	Lou_h	Lou_i	Lou_j	TOTAL
Abiotrophia	0	0	0	0	0	3	0	0	0	0	3
Actinobacillus	0	1	2	2	5	1	0	1	1	0	13
Actinomyces	2	2	4	0	0	0	0	1	1	7	17
Aggregatibacter	0	2	0	0	1	0	3	0	1	0	7
Atopobium	1	0	1	0	0	0	0	0	1	2	5
Bergeyella	0	2	0	0	0	0	0	0	0	0	2
Branhamella	0	2	0	0	0	0	0	0	0	0	2
Bulleidia	0	2	0	0	0	0	0	0	0	1	3
Campylobacter	1	0	0	0	0	0	0	0	1	2	4
Capnocytophaga	0	0	0	8	0	0	1	0	1	0	10
Catonella	0	0	0	1	0	0	0	0	0	0	1
Citrobacter	2	0	0	0	0	0	0	0	0	0	2
Derxia	0	0	0	13	0	0	0	0	0	1	14
Enterobacter	0	0	0	0	0	0	0	0	0	1	1
Enterococcus	1	0	0	0	0	1	1	0	0	0	3
Eubacterium	1	0	0	0	0	0	1	0	0	1	3
Fusobacterium	6	1	1	4	0	1	0	0	5	1	19
Gemella	1	7	0	0	0	2	1	2	11	0	24
Granulicatella	1	2	2	0	1	3	3	6	1	0	19
Haemophilus	2	10	23	0	16	14	11	24	4	1	105
Kingella	0	0	0	0	0	0	0	0	0	2	2
Lactobacillus	0	0	0	0	0	0	0	0	0	1	1
Lactococcus	0	0	0	0	0	0	0	0	0	1	1
Leptotrichia	6	0	2	2	0	2	2	0	0	6	20
Megasphaera	1	0	0	0	0	0	0	0	0	0	1
Micromonas	0	0	0	0	0	1	0	0	0	0	1
Mitsuokella	0	0	0	0	0	0	0	0	0	1	1
Mogibacterium	1	0	0	0	0	0	0	0	0	0	1

Genus	Lou_a	Lou_b	Lou_c	Lou_d	Lou_e	Lou_f	Lou_g	Lou_h	Lou_i	Lou_j	TOTAL
Neisseria	0	18	10	50	4	11	24	7	5	2	131
Oribacterium	0	0	1	2	1	0	0	0	0	0	4
Peptococcus	0	0	0	1	0	0	0	0	0	0	1
Porphyromonas	0	0	0	1	2	1	4	1	0	0	9
Prevotella	7	6	13	2	1	5	9	0	4	16	63
Propionibacterium	0	0	0	0	0	0	0	0	0	1	1
Pseudomonas	0	0	0	0	0	0	0	2	0	0	2
Rothia	6	5	11	4	18	0	1	23	2	11	81
Selenomonas	0	0	0	0	0	0	0	0	0	2	2
Streptococcus	50	51	18	25	69	42	28	31	86	37	437
Syntrophococcus	1	0	0	0	0	0	0	0	0	0	1
TM7	1	0	2	1	0	1	0	0	1	0	6
Veillonella	6	2	17	1	1	5	7	0	1	17	57
Unclassified	1	0	0	1	0	0	3	0	0	1	6
Unknown	4	2	0	3	0	4	7	7	2	0	29
TOTAL	102	115	107	121	119	97	106	105	128	115	1115

Table S5. Frequency (count) of each genus in each of the 10 saliva samples (Ger_a through Ger_j) from Dessau/Leipzig, Germany.

Genus	Ger_a	Ger_b	Ger_c	Ger_d	Ger_e	Ger_f	Ger_g	Ger_h	Ger_i	Ger_j	TOTAL
Actinobacillus	0	1	0	1	0	0	2	0	0	2	6
Actinomyces	0	1	0	1	7	3	2	1	2	6	23
Aggregatibacter	0	0	3	1	0	0	1	0	2	3	10
Atopobium	0	0	0	1	3	0	0	1	2	0	7
Bulleidia	1	0	0	0	1	0	0	0	0	0	2
Campylobacter	0	1	6	0	2	3	2	0	0	2	16
Capnocytophaga	3	2	0	0	1	0	0	0	0	1	7
Cardiobacterium	0	0	1	0	0	0	0	0	0	0	1
Catonella	1	1	0	1	0	0	0	0	0	1	4
Chryseobacterium	0	2	0	0	0	0	0	0	0	0	2
Corynebacterium	0	0	0	1	0	0	0	0	0	0	1
Derxia	3	0	0	2	0	0	0	0	0	0	5
Dialister	0	0	0	0	0	0	0	0	0	1	1
Enterococcus	0	0	0	0	2	0	0	0	1	0	3
Eubacterium	1	0	1	0	0	0	0	0	1	0	3
Fusobacterium	8	3	3	3	2	1	4	3	1	1	29
Gemella	1	1	0	0	0	0	0	0	0	1	3
Granulicatella	1	2	1	2	1	0	1	5	0	4	17
Haemophilus	7	15	8	11	10	0	17	0	15	3	86
Hallella	4	0	0	0	0	0	0	0	0	0	4
Lactobacillus	1	0	0	0	0	1	0	0	0	0	2
Leptotrichia	2	2	0	4	3	0	0	0	7	1	19
Megasphaera	0	0	0	0	0	1	0	0	1	0	2
Micrococcus	0	0	0	0	0	0	0	0	1	0	1
Micromonas	0	0	0	1	0	0	0	0	0	0	1
Mitsuokella	0	0	0	0	0	0	0	0	1	0	1
Mogibacterium	1	0	0	0	0	0	0	0	0	0	1
Mycoplasma	3	0	0	0	0	0	0	0	0	0	3
Neisseria	8	27	5	10	8	3	9	0	2	4	76

Genus	Ger_a	Ger_b	Ger_c	Ger_d	Ger_e	Ger_f	Ger_g	Ger_h	Ger_i	Ger_j	TOTAL
Nesterenkonia	0	0	0	0	1	0	0	0	0	0	1
Oribacterium	2	0	3	0	0	0	0	0	0	1	6
Paludibacter	0	0	0	0	0	0	0	0	0	1	1
Peptostreptococcus	1	0	0	1	0	0	0	0	0	0	2
Porphyromonas	16	0	2	0	5	0	12	0	1	3	39
Prevotella	7	4	42	13	32	49	4	11	13	14	189
Rothia	11	15	13	29	5	16	19	27	23	11	169
Selenomonas	1	0	0	0	1	1	2	0	0	2	7
Streptococcus	28	33	17	30	7	26	25	66	26	46	304
Syntrophococcus	0	0	0	1	0	0	0	1	1	0	3
TM7	1	0	0	1	1	1	1	0	1	4	10
Tannerella	0	0	0	0	0	0	2	0	0	0	2
Treponema	0	0	0	0	0	0	2	0	0	0	2
Veillonella	2	3	10	2	19	7	5	3	11	6	68
Unclassified	1	0	0	0	0	0	0	0	1	1	3
Unknown	1	0	1	0	3	2	3	0	2	1	13
TOTAL	116	113	116	116	114	114	113	118	115	120	1155

Table S6. Frequency (count) of each genus in each of the 10 saliva samples (Pol_a through Pol_j) from Warsaw, Poland.

Genus	Pol_a	Pol_b	Pol_c	Pol_d	Pol_e	Pol_f	Pol_g	Pol_h	Pol_i	Pol_j	TOTAL
Actinobacillus	0	0	0	0	0	1	0	0	1	0	2
Actinomyces	6	15	0	3	0	1	1	1	0	2	29
Aggregatibacter	0	0	0	0	0	1	0	5	0	0	6
Alysiella	0	0	0	0	0	1	0	0	0	0	1
Atopobium	4	2	1	1	0	0	0	0	1	2	11
Bifidobacterium	3	0	0	0	0	0	0	0	0	0	3
Bulleidia	1	0	0	0	0	1	1	0	0	1	4
Butyrivibrio	1	0	1	0	0	0	0	0	0	0	2
Campylobacter	2	4	0	0	0	2	1	2	3	0	14
Capnocytophaga	0	0	0	0	0	0	1	1	0	0	2
Cardiobacterium	0	0	0	0	0	0	0	2	0	0	2
Catenibacterium	0	0	0	0	0	0	0	1	0	0	1
Cloacibacterium	0	0	0	0	0	0	0	1	1	0	2
Collimonas	0	0	0	0	0	0	0	0	1	0	1
Coprococcus	0	0	0	1	0	0	0	0	0	0	1
Derxia	0	0	1	0	0	1	1	1	0	0	4
Dialister	0	0	0	0	0	0	1	0	0	0	1
Enterococcus	0	0	1	0	0	0	1	0	0	0	2
Eubacterium	0	1	0	0	0	0	2	0	0	0	3
Faecalibacterium	0	0	0	0	0	1	0	0	0	0	1
Fusobacterium	0	3	2	1	1	0	5	0	6	3	21
Gemella	0	0	0	0	1	0	2	1	1	0	5
Granulicatella	3	0	0	0	1	0	1	1	0	0	6
Haemophilus	2	3	5	1	3	2	12	15	13	2	58
Johnsonella	0	0	0	0	0	0	0	1	0	0	1
Kingella	0	0	1	0	0	0	0	0	0	0	1
Leptotrichia	2	4	2	1	0	3	1	3	3	2	21
Megasphaera	0	5	3	2	0	3	0	0	0	0	13
Mitsuokella	5	0	0	0	0	0	0	0	0	0	5

Genus	Pol_a	Pol_b	Pol_c	Pol_d	Pol_e	Pol_f	Pol_g	Pol_h	Pol_i	Pol_j	TOTAL
Neisseria	0	0	4	1	0	16	15	7	6	2	51
Oribacterium	1	2	1	1	0	0	4	0	1	1	11
Paludibacter	0	0	0	0	0	0	1	1	0	0	2
Porphyromonas	0	0	1	0	0	0	3	2	2	1	9
Prevotella	50	31	46	52	1	43	9	9	29	41	311
Rahnella	0	0	0	0	4	0	0	0	0	0	4
Rothia	4	4	9	2	1	5	4	2	2	1	34
Ruminococcus	0	1	0	3	0	0	0	0	1	0	5
Schwartzia	0	0	0	0	0	0	1	0	0	0	1
Selenomonas	3	0	3	5	0	5	0	0	0	2	18
Serratia	0	0	0	0	94	0	0	0	0	0	94
Shuttleworthia	0	0	0	0	0	0	1	0	0	0	1
Streptococcus	9	13	7	22	9	12	36	54	29	35	226
Syntrophococcus	1	0	1	0	0	0	0	0	0	0	2
TM7	2	0	0	1	0	1	1	0	0	0	5
Treponema	0	1	0	0	0	0	1	0	0	0	2
Veillonella	14	27	28	22	2	16	5	3	10	18	145
Unclassified	2	1	0	0	0	2	1	0	1	0	7
Unknown	0	1	1	0	1	0	4	4	3	2	16
TOTAL	115	118	118	119	118	117	116	117	114	115	1167

Table S7. Frequency (count) of each genus in each of the 10 saliva samples (Geo_a through Geo_j) from Batumi, Georgia.

Genus	Geo_a	Geo_b	Geo_c	Geo_d	Geo_e	Geo_f	Geo_g	Geo_h	Geo_i	Geo_j	TOTAL
Abiotrophia	0	0	1	1	0	0	0	0	0	0	2
Actinobacillus	0	0	1	0	1	1	0	0	2	3	8
Actinomyces	4	3	0	2	0	0	2	0	0	0	11
Aggregatibacter	0	0	1	2	0	1	0	0	1	1	6
Atopobium	1	0	1	0	2	0	2	0	0	0	6
Bulleidia	0	1	1	0	1	0	0	0	0	0	3
Buttiauxella	0	0	0	0	0	0	0	0	14	14	28
Campylobacter	3	1	0	0	0	0	0	0	1	1	6
Capnocytophaga	1	2	0	0	1	0	0	0	0	0	4
Cardiobacterium	0	1	0	0	0	0	0	0	0	0	1
Catonella	0	0	0	0	0	0	0	1	1	1	3
Citrobacter	0	0	0	0	0	0	2	0	0	0	2
Cloacibacterium	0	0	2	0	0	1	0	0	0	0	3
Collimonas	0	0	0	1	0	0	0	0	0	0	1
Corynebacterium	0	0	0	1	0	0	0	0	0	0	1
Derxia	0	0	0	0	1	0	0	0	0	0	1
Enterobacter	22	0	0	0	0	0	16	31	2	1	72
Enterococcus	0	0	0	1	0	0	0	0	0	0	1
Erwinia	0	0	0	0	0	0	0	0	1	0	1
Eubacterium	0	0	0	1	0	1	1	0	0	0	3
Faecalibacterium	0	0	0	0	0	0	0	0	1	1	2
Fusobacterium	2	2	1	0	0	5	1	2	6	6	25
Gemella	0	0	1	2	0	0	1	0	0	0	4
Granulicatella	3	1	2	7	2	1	2	1	3	3	25
Haemophilus	2	16	7	11	15	17	10	2	2	2	84
Johnsonella	0	0	0	0	0	1	0	0	0	0	1
Kingella	0	0	2	0	0	1	0	0	1	1	5
Klebsiella	0	0	0	0	0	0	0	2	0	0	2
Lactobacillus	0	0	1	0	0	0	0	0	0	0	1

Genus	Geo_a	Geo_b	Geo_c	Geo_d	Geo_e	Geo_f	Geo_g	Geo_h	Geo_i	Geo_j	TOTAL
Leptotrichia	6	0	3	0	0	1	3	3	4	4	24
Megasphaera	1	0	0	0	0	2	0	0	1	1	5
Micromonas	0	0	0	0	0	2	0	0	1	1	4
Moraxella	0	0	0	0	0	0	0	0	1	1	2
Neisseria	7	14	3	14	25	10	9	16	6	6	110
Oribacterium	1	1	1	0	1	0	0	2	0	0	6
Paludibacter	0	0	0	0	0	0	0	0	2	2	4
Pasteurella	0	0	1	0	0	0	0	0	0	0	1
Peptostreptococcus	0	0	0	0	0	1	0	0	2	2	5
Plesiomonas	1	0	0	0	0	0	0	0	0	0	1
Porphyromonas	4	3	4	5	9	3	1	0	6	6	41
Prevotella	21	49	31	13	32	44	30	12	27	27	286
Proteus	0	0	0	0	0	0	0	0	1	1	2
Raoultella	0	0	0	0	0	0	3	0	0	0	3
Rothia	2	1	1	6	8	1	5	1	2	2	29
Selenomonas	0	1	1	0	0	0	0	0	2	2	6
Serratia	9	0	0	0	0	0	0	24	0	0	33
Simonsiella	0	0	0	0	0	1	0	1	0	0	2
Sneathia	0	0	0	0	0	0	0	0	1	1	2
Streptobacillus	1	0	0	0	0	0	0	0	1	1	3
Streptococcus	20	8	16	48	7	9	16	16	17	14	171
Syntrophococcus	0	0	0	0	1	1	0	0	0	0	2
TM7	1	1	1	0	2	0	1	0	0	0	6
Tannerella	0	0	0	0	0	1	0	0	1	1	3
Treponema	0	0	0	2	0	0	0	3	2	2	9
Veillonella	11	9	18	8	12	6	10	6	2	2	84
Unclassified	0	1	2	1	2	0	2	0	2	0	10
Unknown	1	3	0	0	3	1	2	1	4	5	20
TOTAL	124	118	103	126	125	112	119	124	120	115	1186

Table S8. Frequency (count) of each genus in each of the 10 saliva samples (Tur_a through Tur_j) from Ankara, Turkey.

Genus	Tur_a	Tur_b	Tur_c	Tur_d	Tur_e	Tur_f	Tur_g	Tur_h	Tur_i	Tur_j	TOTAL
Actinobacillus	1	0	0	0	5	0	0	0	0	0	6
Actinomyces	0	2	3	3	3	0	0	1	3	0	15
Aggregatibacter	0	1	0	1	0	0	0	0	0	0	2
Arthrobacter	0	0	2	0	0	0	0	0	0	0	2
Atopobium	1	0	2	0	1	1	2	1	1	0	9
Bulleidia	0	1	0	0	1	1	0	3	0	5	11
Butyrivibrio	0	1	0	0	0	0	0	0	0	0	1
Campylobacter	3	1	0	1	1	1	1	2	0	1	11
Capnocytophaga	0	0	0	0	1	0	1	1	0	1	4
Catonella	0	1	0	0	0	0	1	0	0	1	3
Chryseobacterium	0	0	0	0	0	0	0	1	0	0	1
Corynebacterium	0	0	1	0	0	0	1	0	0	0	2
Derxia	0	0	0	0	0	0	2	1	0	0	3
Desulfobulbus	0	0	0	0	0	1	0	0	0	0	1
Dialister	0	0	1	0	0	3	0	0	0	0	4
Enterococcus	0	0	0	0	1	0	0	0	0	0	1
Eubacterium	0	0	0	0	0	0	1	0	0	2	3
Fusobacterium	6	1	0	5	1	2	15	7	19	5	61
Gemella	2	0	0	2	0	0	0	0	1	1	6
Granulicatella	1	2	0	2	2	4	0	0	1	2	14
Haemophilus	4	2	1	11	5	9	4	3	11	3	53
Hallella	0	0	0	0	1	0	0	0	0	0	1
Kingella	1	0	1	0	0	1	0	0	0	0	3
Lactobacillus	0	0	2	0	0	0	0	0	0	0	2
Leptotrichia	2	2	11	2	1	1	0	5	0	0	24
Megasphaera	0	3	1	0	1	0	0	0	1	0	6
Micromonas	0	0	0	0	0	0	3	0	0	0	3
Mycoplasma	0	0	2	0	0	0	0	0	0	0	2
Neisseria	6	6	1	2	0	3	8	28	3	0	57

Genus	Tur_a	Tur_b	Tur_c	Tur_d	Tur_e	Tur_f	Tur_g	Tur_h	Tur_i	Tur_j	TOTAL
Oribacterium	2	8	0	0	0	0	1	0	0	0	11
Peptostreptococcus	0	0	0	0	0	0	0	0	1	0	1
Porphyromonas	5	8	1	1	1	7	5	3	5	7	43
Prevotella	30	57	28	15	40	17	30	43	26	40	326
Rikenella	0	0	0	0	0	2	0	0	0	0	2
Rothia	6	3	4	6	9	6	4	3	3	1	45
Ruminococcus	1	0	0	0	1	0	0	0	4	1	7
Selenomonas	1	1	2	1	1	0	1	4	0	1	12
Sneathia	0	0	0	0	0	0	2	0	0	0	2
Streptococcus	26	8	36	53	18	27	6	4	18	19	215
Syntrophococcus	1	1	0	3	0	0	1	0	1	1	8
TM7	0	0	0	0	0	0	5	0	0	5	10
Tannerella	0	0	0	0	0	1	0	0	0	0	1
Thermovirga	0	0	0	0	0	0	0	0	0	2	2
Treponema	0	0	0	0	0	5	0	2	0	0	7
Veillonella	14	7	15	6	18	14	6	1	16	15	112
Unclassified	0	2	0	0	0	0	0	0	0	0	2
Unknown	1	2	0	0	2	6	8	5	3	1	28
TOTAL	114	120	114	114	114	112	108	118	117	114	1145

Table S9. Frequency (count) of each genus in each of the 10 saliva samples (Con_a through Con_j) from Pointe Noire, Congo.

Genus	Con_a	Con_b	Con_c	Con_d	Con_e	Con_f	Con_g	Con_h	Con_i	Con_j	TOTAL
Abiotrophia	0	1	0	0	0	0	0	0	0	0	1
Achromobacter	0	0	0	0	0	0	0	0	0	2	2
Acinetobacter	0	0	0	0	0	0	0	0	3	0	3
Actinomyces	0	0	1	0	1	1	0	3	2	3	11
Aggregatibacter	0	2	0	2	1	0	0	1	0	0	6
Arthrobacter	0	0	0	0	0	1	0	0	0	0	1
Bulleidia	0	1	0	0	0	0	0	0	3	0	4
Burkholderia	0	0	0	0	2	0	0	22	0	0	24
Buttiauxella	0	0	0	0	0	0	0	0	0	76	76
Campylobacter	0	0	0	0	0	0	0	0	1	0	1
Capnocytophaga	0	2	0	1	1	0	0	0	0	0	4
Cedecea	0	0	0	0	0	0	48	1	0	0	49
Chryseobacterium	0	0	1	0	0	0	0	0	0	0	1
Citrobacter	2	0	0	0	0	0	3	1	0	0	6
Derxia	0	0	0	1	0	2	1	0	0	0	4
Enterobacter	8	0	45	0	111	62	27	53	0	27	333
Eubacterium	0	0	0	0	0	0	1	0	0	0	1
Filifactor	0	0	0	1	0	0	1	0	0	0	2
Fusobacterium	0	4	0	4	1	0	2	0	5	0	16
Gemella	0	0	1	2	0	1	0	1	4	1	10
Granulicatella	0	2	7	1	0	1	0	5	4	1	21
Haemophilus	0	5	2	4	1	5	1	1	4	0	23
Hallella	0	0	0	2	0	0	0	0	0	0	2
Johnsonella	0	0	0	0	0	0	1	0	0	0	1
Klebsiella	9	0	9	0	1	24	1	5	0	2	51
Leptotrichia	0	1	1	0	0	0	1	0	0	0	3
Megasphaera	0	1	0	0	0	0	0	0	5	0	6
Micromonas	0	3	0	0	0	0	0	0	0	0	3
Moraxella	0	0	0	0	1	0	3	0	1	0	5

Genus	Con_a	Con_b	Con_c	Con_d	Con_e	Con_f	Con_g	Con_h	Con_i	Con_j	TOTAL
Morganella	0	0	0	0	0	0	0	0	21	0	21
Neisseria	0	5	9	6	1	4	1	0	6	2	34
Oribacterium	1	0	0	5	2	0	4	2	0	2	16
Plesiomonas	0	0	1	0	0	0	0	0	0	0	1
Porphyromonas	0	14	2	8	0	0	2	0	2	0	28
Prevotella	0	12	2	9	3	0	10	1	22	2	61
Rothia	0	2	2	5	0	2	2	0	15	4	32
Salmonella	0	0	0	0	0	1	0	0	0	0	1
Serratia	63	0	0	0	2	1	1	1	0	0	68
Streptococcus	2	65	38	42	2	11	6	14	14	5	199
Syntrophococcus	0	0	0	1	0	0	0	0	0	0	1
TM7	0	0	0	0	0	0	0	0	1	0	1
Veillonella	0	1	2	4	1	3	5	3	7	0	26
Unclassified	0	3	0	0	0	2	0	0	0	0	5
Unknown	0	4	1	3	1	1	0	1	3	0	14
TOTAL	85	128	124	101	132	122	121	115	123	127	1178

Table S10. Frequency (count) of each genus in each of the 10 saliva samples (Saf_a through Saf_j) from Johannesburg, South Africa.

Genus	Saf_a	Saf_b	Saf_c	Saf_d	Saf_e	Saf_f	Saf_g	Saf_h	Saf_i	Saf_j	TOTAL
Actinobacillus	1	4	1	0	2	3	1	2	4	0	18
Actinomyces	4	11	2	6	3	0	0	2	1	0	29
Aggregatibacter	1	1	0	0	1	0	0	0	0	0	3
Atopobium	0	0	0	1	0	0	0	0	0	0	1
Bulleidia	0	0	1	2	1	1	0	0	0	0	5
Buttiauxella	0	0	0	1	0	0	0	0	0	0	1
Campylobacter	0	4	0	1	0	3	0	0	0	0	8
Capnocytophaga	1	1	1	0	1	0	0	2	0	0	6
Catonella	0	0	0	0	0	1	0	0	0	0	1
Citrobacter	0	0	0	4	0	0	0	0	0	0	4
Cloacibacterium	0	0	0	0	0	3	0	0	0	0	3
Corynebacterium	0	2	0	0	0	2	0	0	0	0	4
Derxia	0	3	0	0	0	0	0	1	0	0	4
Enterobacter	0	0	0	12	0	0	0	0	0	0	12
Eubacterium	0	1	0	0	1	1	0	0	0	0	3
Fusobacterium	1	1	4	4	6	0	0	4	1	1	22
Gemella	2	1	0	0	1	1	1	0	3	0	9
Granulicatella	1	1	1	1	0	2	1	0	2	0	9
Haemophilus	2	3	3	1	4	5	13	17	9	0	57
Leptotrichia	1	4	6	0	0	2	0	0	0	0	13
Megasphaera	0	1	3	2	0	2	0	1	0	0	9
Mitsuokella	1	0	0	1	0	0	0	0	0	0	2
Neisseria	2	2	2	5	15	5	2	4	14	3	54
Nesterenkonia	0	0	1	0	0	0	0	0	0	0	1
Oribacterium	2	4	3	0	0	3	2	0	1	0	15
Pantoea	0	0	0	1	0	0	0	0	0	0	1
Porphyromonas	2	0	2	0	21	0	0	5	2	0	32
Prevotella	41	26	46	23	34	28	13	30	5	1	247
Pseudomonas	0	0	0	0	0	0	1	2	0	0	3

Genus	Saf_a	Saf_b	Saf_c	Saf_d	Saf_e	Saf_f	Saf_g	Saf_h	Saf_i	Saf_j	TOTAL
Rahnella	0	0	0	1	0	0	0	0	0	9	10
Rothia	6	4	3	6	1	9	1	0	10	0	40
Ruminococcus	0	0	0	1	0	0	0	0	0	0	1
Selenomonas	1	0	0	1	0	0	0	0	0	0	2
Serratia	0	0	0	6	0	0	0	0	0	116	122
Streptococcus	23	40	16	17	12	56	58	22	54	1	299
Syntrophococcus	0	0	0	1	1	0	0	1	0	0	3
TM7	2	0	3	2	1	0	0	1	0	0	9
Treponema	1	0	1	0	0	0	0	0	0	0	2
Vagococcus	0	0	0	0	1	0	0	0	0	0	1
Veillonella	24	22	9	17	15	23	25	12	10	2	159
Yersinia	0	0	0	1	0	0	0	0	0	0	1
Unclassified	0	1	0	0	0	0	2	0	0	0	3
Unknown	1	1	0	4	4	4	2	1	1	1	19
TOTAL	120	138	108	122	125	154	122	107	117	134	1247

Table S11. Frequency (count) of each genus in each of the 10 saliva samples (Chi_a through Chi_j) from Shanghai, China.

Genus	Chi_a	Chi_b	Chi_c	Chi_d	Chi_e	Chi_f	Chi_g	Chi_h	Chi_i	Chi_j	TOTAL
Abiotrophia	0	0	0	0	1	1	0	0	1	1	4
Actinobacillus	0	1	0	2	14	0	2	4	0	0	23
Actinomyces	0	1	1	2	1	1	1	1	0	0	8
Aggregatibacter	0	2	2	0	1	0	0	4	0	2	11
Alysiella	0	0	0	0	0	0	3	0	0	0	3
Anaerovorax	0	0	0	0	0	0	0	1	0	0	1
Branhamella	0	0	3	0	0	0	0	0	0	0	3
Bulleidia	2	0	2	0	1	0	0	0	1	0	6
Campylobacter	2	0	1	0	0	0	0	0	0	3	6
Capnocytophaga	2	2	0	1	0	1	2	2	1	4	15
Catonella	0	1	0	1	0	0	1	1	0	0	4
Chryseobacterium	1	0	0	0	0	0	0	0	0	0	1
Cloacibacterium	0	0	0	0	0	0	0	0	5	4	9
Corynebacterium	0	0	1	0	0	0	0	1	0	0	2
Derxia	0	8	0	0	0	1	1	2	0	5	17
Enterococcus	0	0	0	0	0	0	0	0	3	0	3
Eubacterium	0	2	1	0	1	2	0	0	0	0	6
Filifactor	2	4	0	0	0	0	0	1	0	0	7
Fusobacterium	6	9	3	3	3	0	1	2	1	4	32
Gemella	1	0	3	0	0	0	0	0	4	1	9
Granulicatella	0	5	5	0	0	2	3	2	2	2	21
Haemophilus	0	9	8	7	6	11	12	7	6	7	73
Johnsonella	0	0	0	1	0	0	0	0	0	0	1
Kingella	0	0	0	0	1	1	0	0	0	0	2
Leptotrichia	5	2	1	3	1	0	0	3	1	4	20
Micromonas	1	0	0	0	0	0	0	0	0	0	1
Mogibacterium	0	1	0	0	1	0	0	0	0	0	2
Moraxella	1	0	6	0	0	0	0	0	0	0	7
Mycoplasma	0	3	0	0	0	0	0	0	0	0	3

Genus	Chi_a	Chi_b	Chi_c	Chi_d	Chi_e	Chi_f	Chi_g	Chi_h	Chi_i	Chi_j	TOTAL
Neisseria	16	13	9	26	25	22	26	25	16	48	226
Oribacterium	0	1	0	3	1	0	1	0	0	0	6
Paludibacter	0	0	0	0	0	1	1	0	0	0	2
Peptococcus	0	0	0	0	1	0	0	0	0	0	1
Peptostreptococcus	3	0	0	0	1	0	0	0	0	0	4
Porphyromonas	6	4	8	0	9	2	5	7	2	6	49
Prevotella	42	7	23	31	16	0	0	22	5	5	151
Rothia	2	5	4	5	4	0	2	0	0	1	23
Ruminococcus	1	0	0	0	0	0	0	0	0	0	1
Scardovia	0	1	1	0	0	0	0	0	0	0	2
Selenomonas	2	1	0	0	0	0	0	1	0	0	4
Serratia	0	0	0	0	0	0	16	0	0	0	16
Sneathia	0	0	0	0	0	0	0	0	2	0	2
Streptococcus	5	27	27	12	21	68	32	20	61	18	291
TM7	1	0	0	0	0	0	0	0	0	0	1
Thermovirga	0	1	0	0	0	0	0	0	1	0	2
Treponema	0	4	1	2	0	0	0	3	0	1	11
Veillonella	3	1	3	14	4	0	1	1	2	0	29
Wolinella	0	1	0	0	0	0	0	0	0	0	1
Yersinia	0	0	0	0	0	0	1	0	0	0	1
Unclassified	1	1	0	1	1	0	0	0	0	0	4
Unknown	5	1	4	0	1	1	3	4	1	2	22
TOTAL	110	118	117	114	115	114	114	114	115	118	1149

Table S12. Frequency (count) of each genus in each of the 10 saliva samples (Phi_a through Phi_j) from Surigao, Philippines.

Genus	Phi_a	Phi_b	Phi_c	Phi_d	Phi_e	Phi_f	Phi_g	Phi_h	Phi_i	Phi_j	TOTAL
Acidaminococcus	0	1	0	0	0	0	0	0	0	0	1
Actinobacillus	1	0	0	0	0	2	0	0	0	0	3
Actinomyces	2	6	7	4	3	1	1	0	0	10	34
Aggregatibacter	1	0	0	5	0	0	0	0	0	0	6
Arthrobacter	1	0	0	0	0	0	0	0	0	0	1
Atopobium	0	1	0	0	0	1	0	0	0	0	2
Bergeriella	0	0	0	0	0	0	0	1	0	0	1
Bulleidia	2	0	0	0	0	0	0	0	1	0	3
Buttiauxella	0	0	0	0	0	0	0	0	2	0	2
Campylobacter	0	0	0	3	0	0	0	1	0	0	4
Capnocytophaga	0	0	0	1	1	0	0	3	0	0	5
Catonella	0	0	0	0	1	0	1	0	0	0	2
Citrobacter	0	0	48	0	0	0	0	0	0	0	48
Cloacibacterium	0	0	0	0	0	0	0	1	0	0	1
Clostridium	0	0	0	0	0	0	0	0	0	1	1
Corynebacterium	1	1	0	0	0	0	0	1	0	1	4
Derxia	0	0	0	0	0	0	1	1	0	0	2
Dialister	0	2	0	0	0	0	0	0	0	0	2
Enterobacter	0	0	51	0	0	0	0	4	56	0	111
Eubacterium	0	0	0	0	0	0	0	1	0	1	2
Filifactor	0	1	0	0	0	2	0	0	0	5	8
Fusobacterium	2	3	0	4	1	1	4	3	2	1	21
Gemella	0	0	0	1	0	1	2	0	0	1	5
Granulicatella	2	4	0	0	2	1	2	5	2	1	19
Haemophilus	9	1	0	6	0	13	2	9	1	9	50
Kineosphaera	0	0	0	0	0	0	1	0	0	0	1
Klebsiella	0	0	3	0	0	0	0	0	5	0	8
Lactobacillus	0	2	0	0	0	0	0	0	0	1	3
Leptotrichia	3	5	1	1	0	2	2	2	0	1	17

Genus	Phi_a	Phi_b	Phi_c	Phi_d	Phi_e	Phi_f	Phi_g	Phi_h	Phi_i	Phi_j	TOTAL
Megasphaera	0	0	0	0	9	1	0	0	0	3	13
Micromonas	0	0	0	0	0	0	0	0	0	1	1
Mitsuokella	0	0	0	0	0	0	0	1	0	0	1
Mogibacterium	0	1	0	0	0	0	0	0	0	0	1
Mycoplasma	0	0	0	0	0	0	1	0	0	0	1
Neisseria	36	3	0	3	1	8	4	9	5	20	89
Oribacterium	0	0	0	1	0	0	0	0	1	1	3
Porphyromonas	7	1	0	9	0	3	3	0	4	2	29
Prevotella	9	37	4	25	23	45	28	9	8	12	200
Rothia	0	2	0	5	3	0	4	16	2	15	47
Ruminococcus	0	2	0	0	0	0	0	0	0	0	2
Scardovia	0	1	0	0	0	0	0	0	0	0	1
Selenomonas	0	2	2	0	0	2	1	2	0	0	9
Streptococcus	34	22	4	41	20	16	43	30	20	21	251
Syntrophococcus	2	2	0	0	0	0	0	0	0	0	4
TM7	0	0	0	1	0	3	2	1	1	0	8
Treponema	0	1	0	5	0	0	3	1	0	0	10
Veillonella	1	17	0	6	45	14	14	16	6	13	132
Unclassified	0	0	0	0	6	0	0	0	0	0	6
Unknown	2	2	0	0	1	2	0	0	0	0	7
TOTAL	115	120	120	121	116	118	119	117	116	120	1182

Table S13. Frequency (%) of each bacterial genus in each location. The number assigned to each genus corresponds to the numbers in Figure 2. Genus names in **boldface** vary significantly in frequency across locations (chi-square tests, $p < 0.05$ with a Bonferroni correction for multiple tests). *, genus not previously reported in the human oral cavity (Human Oral Microbiome Database, <http://homd.org>).

Genus	Number	Arg	Bol	Cal	Lou	Ger	Pol	Geo	Tur	Con	Saf	Chi	Phi	TOTAL
Abiotrophia	1	0.00	0.00	0.59	0.27	0.00	0.00	0.17	0.00	0.08	0.00	0.35	0.00	0.12
Achromobacter	2	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.17	0.00	0.00	0.00	0.01
Acidaminococcus	3	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.08	0.01
Acinetobacter	4	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.25	0.00	0.00	0.00	0.02
Actinobacillus	5	0.76	0.49	0.17	1.17	0.52	0.17	0.67	0.52	0.00	1.44	2.00	0.25	0.68
Actinomyces	6	1.77	0.25	0.93	1.52	1.99	2.49	0.93	1.31	0.93	2.33	0.70	2.88	1.50
Aggregatibacter	7	0.42	0.33	0.42	0.63	0.87	0.51	0.51	0.17	0.51	0.24	0.96	0.51	0.50
Alysiella*	8	0.08	0.08	0.08	0.00	0.00	0.09	0.00	0.00	0.00	0.00	0.26	0.00	0.05
Anaerovorax*	9	0.08	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.09	0.00	0.01
Arthrobacter*	10	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.17	0.08	0.00	0.00	0.08	0.03
Atopobium	11	0.25	0.16	0.59	0.45	0.61	0.94	0.51	0.79	0.00	0.08	0.00	0.17	0.38
Bergeriella*	12	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.08	0.01
Bergeyella	13	0.00	0.00	0.00	0.18	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01
Bifidobacterium	14	0.00	0.00	0.00	0.00	0.00	0.26	0.00	0.00	0.00	0.00	0.00	0.00	0.02
Branhamella*	15	0.00	0.00	0.00	0.18	0.00	0.00	0.00	0.00	0.00	0.00	0.26	0.00	0.04
Bulleidia	16	0.08	0.33	0.51	0.27	0.17	0.34	0.25	0.96	0.34	0.40	0.52	0.25	0.37
Burkholderia	17	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	2.04	0.00	0.00	0.00	0.17
Buttiauxella*	18	0.00	0.41	0.00	0.00	0.00	0.00	2.36	0.00	6.45	0.08	0.00	0.17	0.79
Butyrivibrio	19	0.00	0.00	0.00	0.00	0.00	0.17	0.00	0.09	0.00	0.00	0.00	0.00	0.02
Campylobacter	20	0.67	0.49	0.42	0.36	1.39	1.20	0.51	0.96	0.08	0.64	0.52	0.34	0.63

Genus	Number	Arg	Bol	Cal	Lou	Ger	Pol	Geo	Tur	Con	Saf	Chi	Phi	TOTAL
Capnocytophaga	21	0.42	0.08	0.34	0.90	0.61	0.17	0.34	0.35	0.34	0.48	1.31	0.42	0.47
Cardiobacterium	22	0.00	0.00	0.00	0.00	0.09	0.17	0.08	0.00	0.00	0.00	0.00	0.00	0.03
Catenibacterium*	23	0.00	0.00	0.00	0.00	0.00	0.09	0.00	0.00	0.00	0.00	0.00	0.00	0.01
Catonella	24	0.08	0.33	0.08	0.09	0.35	0.00	0.25	0.26	0.00	0.08	0.35	0.17	0.17
Cedecea*	25	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	4.16	0.00	0.00	0.00	0.35
Chryseobacterium*	26	0.08	0.08	0.08	0.00	0.17	0.00	0.00	0.09	0.08	0.00	0.09	0.00	0.06
Citrobacter*	27	0.00	0.16	0.00	0.18	0.00	0.00	0.17	0.00	0.51	0.32	0.00	4.06	0.45
Cloacibacterium*	28	0.00	0.25	0.08	0.00	0.00	0.17	0.25	0.00	0.00	0.24	0.78	0.08	0.16
Clostridium	29	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.08	0.01
Collimonas*	30	0.00	0.00	0.00	0.00	0.00	0.09	0.08	0.00	0.00	0.00	0.00	0.00	0.01
Coprococcus*	31	0.17	0.00	0.00	0.00	0.00	0.09	0.00	0.00	0.00	0.00	0.00	0.00	0.02
Corynebacterium	32	0.25	0.00	0.25	0.00	0.09	0.00	0.08	0.17	0.00	0.32	0.17	0.34	0.14
Derxia*	33	0.59	0.00	0.17	1.26	0.43	0.34	0.08	0.26	0.34	0.32	1.48	0.17	0.45
Desulfobulbus	34	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.09	0.00	0.00	0.00	0.00	0.01
Dialister	35	0.17	0.00	0.00	0.00	0.09	0.09	0.00	0.35	0.00	0.00	0.00	0.17	0.07
Enterobacter	36	0.42	8.52	0.00	0.09	0.00	0.00	6.07	0.00	28.27	0.96	0.00	9.39	4.52
Enterococcus	37	0.34	0.16	0.34	0.27	0.26	0.17	0.08	0.09	0.00	0.00	0.26	0.00	0.16
Erwinia*	38	0.00	0.00	0.00	0.00	0.00	0.00	0.08	0.00	0.00	0.00	0.00	0.00	0.01
Eubacterium	39	0.84	0.16	0.08	0.27	0.26	0.26	0.25	0.26	0.08	0.24	0.52	0.17	0.28
Faecalibacterium*	40	0.00	0.00	0.00	0.00	0.00	0.09	0.17	0.00	0.00	0.00	0.00	0.00	0.02
Filifactor	41	0.08	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.17	0.00	0.61	0.68	0.13
Fusobacterium	42	2.19	1.56	1.02	1.70	2.51	1.80	2.11	5.33	1.36	1.76	2.79	1.78	2.15
Gemella	43	0.51	1.31	1.35	2.15	0.26	0.43	0.34	0.52	0.85	0.72	0.78	0.42	0.80
Granulicatella	44	1.26	0.74	1.61	1.70	1.47	0.51	2.11	1.22	1.78	0.72	1.83	1.61	1.37
Haemophilus	45	7.74	4.91	6.85	9.42	7.45	4.97	7.08	4.63	1.95	4.57	6.35	4.23	5.82
Hallella*	46	0.00	0.00	0.00	0.00	0.35	0.00	0.00	0.09	0.17	0.00	0.00	0.00	0.05
Johnsonella	47	0.00	0.00	0.00	0.00	0.00	0.09	0.08	0.00	0.08	0.00	0.09	0.00	0.03
Kineospaera*	48	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.08	0.01

Genus	Number	Arg	Bol	Cal	Lou	Ger	Pol	Geo	Tur	Con	Saf	Chi	Phi	TOTAL
Kingella	49	0.17	0.00	0.08	0.18	0.00	0.09	0.42	0.26	0.00	0.00	0.17	0.00	0.11
Klebsiella	50	0.00	1.15	0.00	0.00	0.00	0.00	0.17	0.00	4.33	0.00	0.00	0.68	0.53
Lactobacillus	51	0.00	0.00	0.00	0.09	0.17	0.00	0.08	0.17	0.00	0.00	0.00	0.25	0.06
Lactococcus	52	0.00	0.00	0.00	0.09	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01
Leptotrichia	53	2.27	1.47	0.51	1.79	1.65	1.80	2.02	2.10	0.25	1.04	1.74	1.44	1.50
Mannheimia*	54	0.00	0.08	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01
Megasphaera	55	1.09	0.25	0.51	0.09	0.17	1.11	0.42	0.52	0.51	0.72	0.00	1.10	0.55
Micrococcus*	56	0.00	0.00	0.00	0.00	0.09	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01
Micromonas*	57	0.25	0.00	0.17	0.09	0.09	0.00	0.34	0.26	0.25	0.00	0.09	0.08	0.13
Mitsuokella	58	0.00	0.08	0.00	0.09	0.09	0.43	0.00	0.00	0.00	0.16	0.00	0.08	0.08
Mogibacterium	59	0.00	0.00	0.00	0.09	0.09	0.00	0.00	0.00	0.00	0.00	0.17	0.08	0.04
Moraxella	60	1.09	0.00	0.00	0.00	0.00	0.00	0.17	0.00	0.42	0.00	0.61	0.00	0.19
Morganella*	61	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.78	0.00	0.00	0.00	0.15
Mycoplasma	62	0.00	0.00	0.00	0.00	0.26	0.00	0.00	0.17	0.00	0.00	0.26	0.08	0.06
Neisseria	63	12.46	5.16	6.18	11.75	6.58	4.37	9.27	4.98	2.89	4.33	19.67	7.53	7.88
Nesterenkonia*	64	0.00	0.00	0.08	0.00	0.09	0.00	0.00	0.00	0.00	0.08	0.00	0.00	0.02
Oribacterium	65	1.09	0.66	1.02	0.36	0.52	0.94	0.51	0.96	1.36	1.20	0.52	0.25	0.79
Paludibacter*	66	0.00	0.00	0.00	0.00	0.09	0.17	0.34	0.00	0.00	0.00	0.17	0.00	0.06
Pantoea*	67	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.08	0.00	0.00	0.01
Parascardovia	68	0.08	0.00	0.08	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01
Pasteurella*	69	0.00	0.08	0.00	0.00	0.00	0.00	0.08	0.00	0.00	0.00	0.00	0.00	0.01
Peptococcus	70	0.00	0.00	0.00	0.09	0.00	0.00	0.00	0.00	0.00	0.00	0.09	0.00	0.01
Peptostreptococcus	71	0.08	0.00	0.34	0.00	0.17	0.00	0.42	0.09	0.00	0.00	0.35	0.00	0.12
Plesiomonas*	72	0.00	0.00	0.00	0.00	0.00	0.00	0.08	0.00	0.08	0.00	0.00	0.00	0.01
Porphyromonas	73	2.10	0.98	1.27	0.81	3.38	0.77	3.46	3.76	2.38	2.57	4.26	2.45	2.35
Prevotella	74	20.71	14.50	13.79	5.65	16.36	26.65	24.11	28.47	5.18	19.81	13.14	16.92	17.14
Propionibacterium	75	0.00	0.00	0.00	0.09	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01
Proteus	76	0.00	0.00	0.00	0.00	0.00	0.00	0.17	0.00	0.00	0.00	0.00	0.00	0.01

Genus	Number	Arg	Bol	Cal	Lou	Ger	Pol	Geo	Tur	Con	Saf	Chi	Phi	TOTAL
Pseudomonas	77	0.00	0.00	0.00	0.18	0.00	0.00	0.00	0.00	0.00	0.24	0.00	0.00	0.04
Rahnella*	78	0.00	1.56	0.00	0.00	0.00	0.34	0.00	0.00	0.00	0.80	0.00	0.00	0.23
Raoultella*	79	0.00	0.00	0.00	0.00	0.00	0.00	0.25	0.00	0.00	0.00	0.00	0.00	0.02
Rikenella*	80	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.17	0.00	0.00	0.00	0.00	0.01
Rothia	81	5.98	1.88	7.28	7.26	14.63	2.91	2.45	3.93	2.72	3.21	2.00	3.98	4.82
Ruminococcus*	82	0.25	0.16	0.08	0.00	0.00	0.43	0.00	0.61	0.00	0.08	0.09	0.17	0.16
Salmonella*	83	0.08	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.08	0.00	0.00	0.00	0.01
Scardovia	84	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.17	0.08	0.02
Schwartzia*	85	0.00	0.00	0.00	0.00	0.00	0.09	0.00	0.00	0.00	0.00	0.00	0.00	0.01
Selenomonas	86	1.01	0.49	0.42	0.18	0.61	1.54	0.51	1.05	0.00	0.16	0.35	0.76	0.59
Serratia*	87	0.00	17.44	9.98	0.00	0.00	8.05	2.78	0.00	5.77	9.78	1.39	0.00	4.70
Shuttleworthia	88	0.00	0.00	0.00	0.00	0.00	0.09	0.00	0.00	0.00	0.00	0.00	0.00	0.01
Simonsiella	89	0.08	0.00	0.00	0.00	0.00	0.00	0.17	0.00	0.00	0.00	0.00	0.00	0.02
Sneathia	90	0.17	0.00	0.00	0.00	0.00	0.00	0.17	0.17	0.00	0.00	0.17	0.00	0.06
Streptobacillus*	91	0.25	0.08	0.00	0.00	0.00	0.00	0.25	0.00	0.00	0.00	0.00	0.00	0.05
Streptococcus	92	15.91	23.34	28.00	39.19	26.32	19.37	14.42	18.78	16.89	23.98	25.33	21.24	22.66
Syntrophococcus*	93	0.08	0.16	0.25	0.09	0.26	0.17	0.17	0.70	0.08	0.24	0.00	0.34	0.21
Tannerella	94	0.00	0.00	0.00	0.00	0.17	0.00	0.25	0.09	0.00	0.00	0.00	0.00	0.04
Thermovirga*	95	0.00	0.08	0.00	0.00	0.00	0.00	0.00	0.17	0.00	0.00	0.17	0.00	0.04
TM7	96	0.42	0.08	0.85	0.54	0.87	0.43	0.51	0.87	0.08	0.72	0.09	0.68	0.51
Treponema	97	0.25	0.16	0.00	0.00	0.17	0.17	0.76	0.61	0.00	0.16	0.96	0.85	0.34
Vagococcus*	98	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.08	0.00	0.00	0.01
Veillonella	99	13.55	8.27	12.10	5.11	5.89	12.43	7.08	9.78	2.21	12.75	2.52	11.17	8.62
Wolinella*	100	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.09	0.00	0.01
Yersinia	101	0.00	0.08	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.08	0.09	0.00	0.02
Unclassified	102	0.51	0.08	0.34	0.54	0.26	0.60	0.84	0.17	0.42	0.24	0.35	0.51	0.40
Unknown	103	0.76	0.90	0.68	2.60	1.13	1.37	1.69	2.45	1.19	1.52	1.91	0.59	1.39

Table S14. UniFrac (above diagonal) and linearized F_{st} (below diagonal) distances between each pair of locations. Location abbreviations are as in Table 1.

	Arg	Bol	Cal	Lou	Ger	Pol	Geo	Tur	Con	Saf	Chi	Phi
Arg		0.210	0.161	0.239	0.157	0.144	0.152	0.168	0.332	0.127	0.191	0.136
Bol	0.032		0.146	0.256	0.207	0.199	0.185	0.247	0.204	0.149	0.207	0.143
Cal	0.019	0.011		0.185	0.145	0.167	0.196	0.219	0.288	0.116	0.209	0.120
Lou	0.050	0.049	0.023		0.217	0.284	0.286	0.307	0.315	0.245	0.234	0.234
Ger	0.017	0.033	0.012	0.023		0.184	0.209	0.185	0.346	0.157	0.190	0.159
Pol	0.011	0.020	0.016	0.063	0.024		0.163	0.134	0.337	0.114	0.252	0.150
Geo	0.007	0.024	0.025	0.062	0.023	0.009		0.178	0.283	0.154	0.196	0.153
Tur	0.009	0.036	0.026	0.064	0.020	0.005	0.007		0.372	0.166	0.248	0.191
Con	0.076	0.044	0.069	0.090	0.073	0.081	0.056	0.086		0.292	0.298	0.267
Saf	0.014	0.010	0.004	0.042	0.017	0.004	0.015	0.013	0.068		0.206	0.102
Chi	0.020	0.034	0.023	0.023	0.021	0.036	0.024	0.033	0.075	0.027		0.188
Phi	0.010	0.020	0.016	0.039	0.016	0.016	0.009	0.015	0.042	0.012	0.022	

Table S15. Values of the four variables tested for associations with the UniFrac distances.

Location	Pop. size (millions)	Dist. from eq. (km)	Avg. temp. (C)	Avg. rainfall (cm)
Arg	13	3837	15.8	115
Bol	1	1832	7.5	60
Cal	7	4194	15.3	58
Lou	0.8	3389	19.8	155
Ger	0.5	5698	8.8	61
Pol	3	5791	7.7	52
Geo	0.1	4623	14.5	272
Tur	4	4435	11.7	42
Con	0.7	533	24.7	151
Saf	8	2908	16.4	72
Chi	17	3463	15.8	111
Phi	0.1	1088	27.2	320