



PM₁₀, and children's respiratory symptoms and lung function in the PATY study

Gerard Hoek, Sam Pattenden, Saskia Willers, Temenuga Antova, Eleonora Fabianova, Charlotte Braun-Fahrländer, Francesco Forastiere, Ulrike Gehring, Heike Luttmann-Gibson, Leticia Grize, Joachim Heinrich, Danny Houthuijs, Nicole Janssen, Boris Katsnelson, Anna Kosheleva, Hanns Moshammer, Manfred Neuberger, Larisa Privalova, Peter Rudnai, Frank Speizer, Hana Slachtova, Hana Tomaskova, Renata Zlotkowska and Tony Fletcher

ABSTRACT: Studies of the impact of long-term exposure to outdoor air pollution on the prevalence of respiratory symptoms and lung function in children have yielded mixed results, partly related to differences in study design, exposure assessment, confounder selection and data analysis.

We assembled respiratory health and exposure data for >45,000 children from comparable cross-sectional studies in 12 countries. 11 respiratory symptoms were selected, for which comparable questions were asked. Spirometry was performed in about half of the children. Exposure to air pollution was mainly characterised by annual average concentrations of particulate matter with a 50% cut-off aerodynamic diameter of 10 µm (PM₁₀) measured at fixed sites within the study areas.

Positive associations were found between the average PM₁₀ concentration and the prevalence of phlegm (OR per 10 µg·m⁻³ 1.15, 95% CI 1.02–1.30), hay fever (OR 1.20, 95% CI 0.99–1.46), bronchitis (OR 1.08, 95% CI 0.98–1.19), morning cough (OR 1.15, 95% CI 1.02–1.29) and nocturnal cough (OR 1.13, 95% CI 0.98–1.29). There were no associations with diagnosed asthma or asthma symptoms. PM₁₀ was not associated with lung function across all studies combined.

Our study adds to the evidence that long-term exposure to outdoor air pollution, characterised by the concentration of PM₁₀, is associated with increased respiratory symptoms.

KEYWORDS: Child, lung function, nitrogen dioxide, particulate matter, respiratory symptoms

Short-term increases in outdoor air pollution have been associated with respiratory symptoms and temporary lung function decreases [1, 2]. The impact of long-term exposure to outdoor air pollution on prevalence of respiratory symptoms and lung function in children has been investigated in studies around the world [3–15]. Results concerning symptoms have been mixed, with more evidence for significant effects of outdoor air pollution on bronchitis or symptoms such as cough and phlegm than on asthma or asthmatic symptoms such as wheeze. Results from cross-sectional studies of lung function in children were also mixed [14]. Several prospective studies documented significant effects of outdoor air pollution on lung function development, e.g. the Californian Children's Health Study [14, 15]. Some of the inconsistencies in reported associations between air pollution and lung function may be due to differences in study design [14]. Studies differ in

their study area (air pollution exposure contrasts between communities *versus* within a community), measured pollutants (particulate matter, sulphur dioxide, nitrogen dioxide and ozone), wording of symptom questions, study population (either population based or high risk), inclusion of potential confounders and statistical methods. Particulate matter air pollution has been represented with various indices, including total suspended particulate matter (TSP), particulate matter with a 50% cut-off aerodynamic diameter of 10 µm (PM₁₀) or 2.5 µm (PM_{2.5}). Some studies have reported higher air pollution effect estimates on lung function for females [7, 12], but the evidence is currently not convincing [14]. In the Children's Health study, air pollution effects on symptoms were stronger in males [8], whereas the effects on lung function were stronger in females [7]. Interpretation of subgroup analysis in single studies is difficult as the power to detect significant interactions is limited.

AFFILIATIONS

Author affiliations are listed in the Acknowledgements section.

CORRESPONDENCE

G. Hoek
Institute for Risk Assessment
Sciences
University of Utrecht
PO Box 80178
3508 TD Utrecht
The Netherlands
E-mail: g.hoek@uu.nl

Received:

Jan 07 2011

Accepted after revision:

Dec 09 2011

First published online:

April 20 2012

This article has supplementary material available from www.erj.ersjournals.com

European Respiratory Journal
Print ISSN 0903-1936
Online ISSN 1399-3003

To overcome some of these shortcomings, the Pollution and the Young (PATY) project assembled health and exposure data for 58,561 children from comparable cross-sectional studies in 12 countries on children's respiratory health. Pooling original data allowed harmonisation of data analysis and definition of confounders, pursuit of research questions not addressed originally, and inclusion of unpublished studies. We made further use of the large dataset to assess effect modification with more precision than single studies. We already published on associations between outdoor NO₂ and respiratory symptoms in a subset of the five PATY studies with NO₂ exposure data [16]. In the current article, we examine the association between ambient fine particulate matter (PM₁₀) in all PATY studies and prevalence of respiratory symptoms and lung function.

METHODS

Study design

Cross-sectional studies of children were sought that assessed respiratory symptoms and individual risk factors by questionnaire, included cough and wheeze as outcomes, and allowed calculation of annual mean particulate matter measures by study area. We included all cross-sectional studies that were available at the onset of the study. From the USA and the Netherlands, we included only the most recent and largest study, excluding the Six City and Six School studies [14]. Table 1 describes the studies contributing to this paper. More details on the individual studies can be found elsewhere [4, 10, 11, 13, 16, 18–20]. The studies from Russia, Central and Eastern Europe and Italy have not previously published results of air pollution analysis in the English language. The study areas were of substantially different magnitude. The Austrian and Czech study was performed in one city, and the North American study included a very large area in the USA and Canada. We therefore anticipated some heterogeneity in effects, which was taken into account in the analysis. All studies had obtained permission from the relevant medical ethical committees.

Health data

Original questionnaires were translated into English and critically examined for comparability of wording. 11 comparable symptom outcomes were identified: wheeze in the past 12 months, asthma ever, bronchitis in the past 12 months, phlegm, nocturnal dry cough in the past 12 months, morning cough in the past 12 months, sensitivity to inhaled allergens, hay fever ever, itchy rash ever, woken by wheeze in the past 12 months and allergy to pets. The exact wording of the symptom questions in all studies has been reported previously [21].

Several of the studies also performed spirometry (table 1). All studies used equipment that fulfilled either the 1987 criteria of the American Thoracic Society (ATS) or the 1993 European Respiratory Society (ERS) criteria. Spirometry was performed using the 1987 protocol of the ATS (North American, German and Austrian studies) or the 1993 ERS (CESAR study and the Netherlands). Further details have been published previously [17]. The main lung function measures of interest were forced vital capacity (FVC), forced expiratory volume in 1 s (FEV₁), peak expiratory flow and the forced expiratory flow at 25–75% of the FVC (FEF_{25–75%}).

Exposure assessment methods

The main exposure of interest was the annual mean concentration of PM₁₀ in the corresponding study area. Data on the gaseous pollutants NO₂ and SO₂ were also obtained. In all studies, air pollution concentrations were measured at fixed monitoring sites in the study area. To assess the comparability of monitoring sites and monitoring methods, a standard questionnaire was discussed with the investigators of the studies (online supplementary material pages 2 and 3 and tables S1 and S2). Briefly, PM₁₀ was directly measured specifically for the study with Harvard impactors in Switzerland, the CESAR study, and for Russia and North America. In the Netherlands, Austria and Germany, particulate matter measurements were converted to gravimetric PM₁₀ using co-located measurements with a standard particulate matter sampler in the study area. In the Italian study, multiple particulate matter monitoring methods were used and there was insufficient co-location with standard PM₁₀ equipment to allow calculation of a consistent PM₁₀ concentration. The evaluation further resulted in a few modifications of the original exposure estimates and exclusions of a few areas because of unrepresentative monitoring sites (online supplementary material pages 2 and 3).

Data analysis

A priori, we assessed that air pollution exposure contrasts between countries could not be exploited, as differences in language and (unmeasured) major risk factors were likely to be more dominant determinants for the health outcomes (especially with regards to symptoms). Hence, a two-stage analysis approach was used. In stage one, study-specific PM₁₀ effects were estimated using logistic regression for symptoms and linear regression for lung function. This approach also has the advantage that systematic differences in PM₁₀ sampler performance between studies do not affect the results. An area-level random intercept was included to account for clustering within the study areas. The CESAR study was conducted in five countries and, although common methods were used, the study was analysed per country. In stage two, effect estimates and standard errors were entered into a meta-analysis, obtaining a mean estimate, and a measure and Cochran Chi-squared test of heterogeneity using the STATA (StataCorp, College Station, TX, USA) metan command. Estimation of this mean and its confidence interval takes into account both between-study variation in effects and uncertainty of study-specific estimates [22]. In the first stage, we controlled for age, sex, maternal education, paternal education, household-crowding, current parental smoking, mother smoking during pregnancy, gas cooking, unvented gas/oil/kerosene heater, mould, nationality, birth order and "ever had a pet" [16]. Lung function analyses were additionally adjusted for age, height, weight, technician, instrument, season of testing and reported infection on the day of the test. The natural logarithm of lung function variables was used as the dependent variable to allow for non-normal distribution and nonlinear associations with anthropometric variables [17]. The natural logarithm of age, weight and height (sex-specific using an interaction term) were included as predictors following the North American study [13]. We further calculated predicted values for FVC, FEV₁ and FEF_{25–75%} using recently published prediction equations and used the % predicted lung function as the dependent variable in an additional analysis [23].

TABLE 1 Overview of design, respiratory health and exposure data of studies included in the Pollution and the Young (PATY) project							
Study name, location [ref.]	Study areas	Main age range yrs	Health end-points	Health data collection	Exposure data collection	PM10 [#] µg·m ⁻³	NO ₂ [#] µg·m ⁻³ SO ₂ [#] µg·m ⁻³
Scarpol, Switzerland [11]	10 communities, major cities (Bern, Geneva, Zürich) and small towns	6–12	Symptoms	October 1992–March 1993	1992 [†]	24 (10–33)	32 (16–50) 13 (2–23)
Linz, Austria [17]	Schools assigned to 8 monitors in the city of Linz	6–8	Symptoms, lung function	January 1996–December 1998	1996–1998	32 (24–42)	26 (20–31) 6 (4–14)
CESAR, Central and Eastern Europe [18]	3 areas in 3 towns in Bulgaria	9–12	Symptoms, lung function	Spring 1996	October 1995–October 1996	67 (62–71)	NA NA
	4 areas in the city of Ostrava, Czech Republic	9–12	Symptoms, lung function			76 (65–89)	NA NA
	5 towns spread throughout Hungary	9–12	Symptoms, lung function			61 (56–72)	NA NA
	4 towns spread throughout Poland	9–12	Symptoms, lung function			74 (60–85)	NA NA
Germany [5]	4 areas in three towns in Slovakia	9–12	Symptoms, lung function			49 (41–57)	NA NA
	3 towns former East Germany (Hettstedt, Zerbst, Bitterfeld)	6–12	Symptoms, lung function	September 1992–July 1993	1991–1993	43 (33–53)	NA 60 (46–75)
SIDRIA, Italy [19]	46 areas in 22 towns including major cities (Rome, Turin) and small towns	6–10	Symptoms	October–December 1994	October 1993–October 1994	NA	52 (14–93) 13 (2–32)
24-school, the Netherlands [10]	24 schools located within 400 m of a major road in mid-west Netherlands	7–12	Symptoms, lung function	April 1997–July 1998	April 1997– May 1998	34 (30–39)	35 (27–44) NA
10-city, Russia [20]	13 areas in 10 towns of different size (largest Ekaterinburg) and industrialisation	8–12	Symptoms	April–May 1999	November 1998–November 1999	24 (20–28)	20 (13–35) 26 (7–65)
24-city, USA [4, 13]	24 medium-sized communities in the USA and Canada	8–12	Symptoms, lung function	September–May 1988–1991 [†]	Year prior to health evaluation	24 (15–33)	NA 12 (1–34)
Data are presented as mean (range), unless otherwise stated. PM10 was directly measured in Switzerland, Central European Study on Air pollution and Respiratory Health (CESAR) study, Russia and USA; it was estimated using co-located measurements from total suspended particulate matter in Austria and Germany, and from PM2.5 in the Netherlands. PM10: particulate matter with a 50% cut-off aerodynamic diameter of 10 µm; NA: not available. #: mean and range of study area-specific annual average concentrations; †: 1993 for PM10; ‡: study in three consecutive years including eight, nine and seven cities, respectively. SIDRIA: Studi Italiani sui Disturbi Respiratori nell'Infanzia e l'Ambiente.							

Meta-regressions assessed associations between study-specific estimates and study characteristics. These *a priori*-defined potential sources of heterogeneity between estimates were study period, design type (between-city, within-city or mixed design), location of monitoring station, proportion of younger children (6–8 yrs), questionnaire-date variability across study areas, high response rate (>80%) and response-rate variability across study areas.

Subgroup analyses were conducted in the first stage to assess individual subject characteristics as a source of heterogeneity. We stratified by sex, age and reporting of wheeze, and sensitisation (the latter for lung function only).

All analyses were performed using STATA version 8 (StatCorp).

RESULTS

Associations between air pollution and respiratory symptoms

Data were available from 11 studies on PM10 and symptoms (table 1). The highest PM10 concentrations were measured in the Central and Eastern European areas. The range in concentration within a study was largest in the North American and Swiss studies (ratio of maximum to minimum larger than two) (fig. 1).

Complete information on health and covariates was available for 45,788 children. About one-third of the children were from the North American study (table 2). Several-fold differences in symptom prevalence were found between countries, probably partly related to cultural differences and subtle differences in wording of the question.

After adjustment for confounders, PM10 was significantly associated with phlegm and morning cough (table 3 and fig. 2). Associations between PM10 and a doctor diagnosis of bronchitis, nocturnal cough and hay fever were borderline significant. Significant heterogeneity between study-specific estimates was found for most outcomes. The most consistent pattern was found for phlegm and hay fever, for which most study-specific effect estimates were either positive or slightly negative (fig. 2). For bronchitis and the two cough variables, both positive and negative effect estimates were found. A diagnosis of asthma and the symptoms wheeze and being woken by wheeze were not associated with PM10. Meta-regression could not identify significant factors at the study level explaining the observed




FIGURE 1. Annual average concentrations of particulate matter with a 50% cut-off aerodynamic diameter of 10 µm (PM10) per study area, within each country.

TABLE 2 Prevalence (%) of respiratory symptoms in each country												
	Children [#] n	Wheeze [†]	Asthma	Bronchitis	Phlegm	Nocturnal cough	Morning cough	Sensitivity to inhaled allergens	Hay fever	Itchy rash	Woken by wheeze	Allergy to pets
Austria	3904	13.6	8.6	+		10.4	5.3	12.8	5.2	8.2		5.8
Bulgaria	2765	15.8	15.9	23.7	4.0	5.3	16.3	11.3		9.3	6.6	4.7
Czech Republic	3356	20.6	9.8	41.5	18.5	19.6	26.8	17.3		10.8	15.5	4.6
Germany	1903	10.1	8.5				13.8		4.5	15.0		2.1
Hungary	3460	9.6	22.2	32.9	2.3	9.3	6.4	14.0		6.9	8.2	5.0
The Netherlands	1916	9.5	8.1	7.9	9.5	21.6		15.2	7.2	22.3	4.9	9.3
North America	14496	19.4	9.7	6.2	5.9		5.9	20.7	14.1		5.1	
Poland	2821	12.3	10.5	34.9	11.0	13.7	46.3	13.9		7.4	8.9	6.2
Russia	5453	13.4	1.9	14.7	7.2		11.4	6.4	1.2		4.4	
Slovakia	2975	9.9	6.7	31.3	5.4	13.7	14.6	14.4		9.7	8.9	3.2
Switzerland	2739	10.4	9.0	18.3		21.5	11.9	13.7	10.2	13.3	4.6	4.6
Total	45788	14.9	9.7	18.6	7.3	13.9	12.7	15.3	9.3	10.7	6.8	5.0
Data are presented as %, unless otherwise stated. [#] : number with no missing covariates; [†] : symptoms for the past 12 months, except asthma, hay fever and itchy rash, for which the questions referred to lifetime (ever); ⁺ : some outcomes not included in the questionnaire of all countries.												

Data are presented as %, unless otherwise stated. [†]: symptoms for the past 12 months, except asthma, hay fever and itchy rash, for which the questions referred to lifetime (ever); ⁺: some outcomes not included in the questionnaire of all countries.

TABLE 3 Association between prevalence of respiratory symptoms and long-term average concentration of particulate matter with a 50% cut-off aerodynamic diameter of 10 µm (PM₁₀)

	Mean OR (95% CI)	
	Age-sex adjusted	Fully adjusted [#]
Wheeze	1.03 (0.95–1.11) [†]	1.01 (0.95–1.09)
Asthma	0.99 (0.92–1.08)	1.03 (0.97–1.10)
Bronchitis	1.06 (0.94–1.21) [†]	1.08 (0.98–1.19) [†]
Phlegm	1.16 (1.02–1.31) [†]	1.15 (1.02–1.30) [†]
Nocturnal cough	1.15 (1.00–1.32) [†]	1.13 (0.98–1.29) [†]
Morning cough	1.16 (1.02–1.32) [†]	1.15 (1.02–1.29) [†]
Sensitivity to inhaled allergens	1.01 (0.90–1.13) [†]	1.02 (0.93–1.11)
Hay fever	1.18 (0.96–1.44)	1.20 (0.99–1.46)
Itchy rash	1.04 (0.98–1.11)	1.07 (0.96–1.19) [†]
Woken by wheeze	1.04 (0.94–1.14) [†]	1.01 (0.92–1.12)
Allergy to pets	1.18 (0.95–1.46) [†]	1.18 (0.96–1.45) [†]

Odds ratios (ORs) are combined effect estimates from single pollutant models calculated from country-specific estimates using random effects model. Mean ORs and 95% confidence intervals are given per 10 µg·m⁻³ increase in PM₁₀.

[#]: adjusted for age, sex, maternal education, paternal education, household-crowding, current parental smoking, mother smoking during pregnancy, gas-cooking, unvented gas/oil/kerosene heater, mould, nationality, birth order and "ever had a pet"; [†]: evidence of between-study heterogeneity (p<0.10).

heterogeneity in effect estimates. Table S3 presents the data for the symptom phlegm.

Two-pollutant models were employed for symptoms showing (borderline) significant associations with PM₁₀ in the single-pollutant model (table 4). Adjustment for SO₂ made little difference to the PM₁₀ effect estimates with the exception of bronchitis, which showed a substantial decrease (single *versus* two pollutant PM₁₀ estimates table 4). Adjustment for NO₂ made little difference to the PM₁₀ effect estimates for bronchitis, phlegm, morning cough and hay fever. For the symptoms of nocturnal cough, sensitivity to inhaled allergens, itchy rash and allergy to pets, after inclusion of NO₂, the PM₁₀ effect estimates were reduced. Correlations (r) between the annual average concentration of NO₂ and PM₁₀ in the studies with ≥10 study areas ranged from 0.48 to 0.84. Correlations between SO₂ and PM₁₀ ranged from 0.36 to 0.82.

PM₁₀ effect estimates did not differ significantly between males and females, or between the younger and older children in the study population (table S4). Associations for hay fever and allergy to pets were stronger in the older children, but the difference with the younger children was not significant. Effect of PM₁₀ on nocturnal cough and allergy to pets were stronger in boys than in girls but the differences did not reach statistical significance.

Associations between air pollution and lung function

Valid lung function data were available for 22,809 children (table S5). About 60% of the children with a valid lung function

test were from the North American study. While in the North American, German, Austrian and Dutch studies, the percentage of children with a valid lung function test was well over 80%, it ranged from between 26 and 60% for the four CESAR study countries. This was related to too early termination of the test, resulting in too low FVC values. In all studies except the Austrian study, the mean percentage predicted lung function using the Stanojevic equations was close to 100.

On average, no association was found between the average PM₁₀ concentration of the study area and lung function (fig. 3). In individual studies, both increases (Slovakia) and decreases (North America and Poland) in lung function were found with increasing PM₁₀. Removing the CESAR countries from the analysis did not change the associations substantially (table S6). Analysis of the data using the recent Stanojevic prediction equations also resulted in no significant associations between PM₁₀ and lung function (table S6).

There were no significant differences in PM₁₀ effect estimates between males and females or younger and older children (table 5).

DISCUSSION

Statistically (borderline) significant positive associations were found between PM₁₀ and the prevalence of phlegm, morning cough, hay fever, bronchitis and nocturnal cough. There were no associations with diagnosed asthma and asthma symptoms. PM₁₀ effect estimates did not differ between males and females. PM₁₀ was not associated with lung function.

The main strength of our study is the large number of children (45,000) taken from studies in 12 countries. This reduces the risk of finding spurious associations due to, for example, unmeasured confounding at the study area level as in single studies. The large study size also allowed for analyses of subgroups with more precision. Compared with a standard meta-analysis of studies, our study offers several advantages related to having the original data of all the studies available instead of only the air pollution effect estimates. First, we selected end-points for a common analysis that were considered sufficiently similar in wording. Secondly, the same data analysis procedures were used, including a common set of confounders. Thirdly, the comparability of exposure data collection could be assessed in detail, which resulted in the removal of some study areas from the epidemiological analysis. Finally, several unpublished studies were included in the analysis.

Comparison with previous studies

Our findings of significant associations between PM₁₀ and respiratory symptoms and no associations with lung function are in agreement with the Harvard Six City study and two German studies, which were not included in the current analysis [3, 9, 24]. In contrast, other studies not included in the PATY study did find associations between long-term average air pollution exposure and lung function [14]. For the studies included in the PATY study the large North American study found associations with both lung function and bronchitis [13], whereas the Dutch, German and CESAR studies [5, 10, 18] did not find lung function associations.

The effect of air pollution may have been too modest to be reflected in detectable changes of lung function. The (asthma)




FIGURE 2. Forest plots of study-specific and mean odds ratios for effects of particulate matter with a 50% cut-off aerodynamic diameter of 10 μm (PM₁₀) on a) wheeze, b) asthma, c) bronchitis, d) phlegm, e) nocturnal cough, f) morning cough, g) sensitivity to inhaled allergens, h) hay fever, i) itchy rash, j) being woken by wheeze and k) allergy to pets. Odds ratios per 10 $\mu\text{g}\cdot\text{m}^{-3}$ increase in PM₁₀ are from single pollutant models, adjusted for individual risk factors. The vertical line indicates odds ratio of 1 and horizontal lines represent 95% confidence intervals of estimates. The diamond shape at the bottom of each graph indicates position, and confidence interval, of the mean of the estimates. Symptoms are for the past 12 months, except asthma, hay fever and itchy rash, which refer to lifetime (ever).

symptoms related to the largest deficits in lung function were also not associated with outdoor air pollution. Alternatively, different biases in the analysis of symptoms and lung function may explain the inconsistencies. First, the typically small effects of air pollution on lung function may have been masked by factors such as variability in coaching and judgement by the technician, instrument, subtle shifts in instrument calibration, short-term weather factors and (past) infections [10]. In the CESAR study, good-quality tests were only obtained in <50% of the children, despite a rigorous quality assurance/quality control protocol. Exclusion of the CESAR study countries from the analysis, however, did not change our results. Secondly, reporting bias of symptoms may explain some of the positive associations in studies where parents of the children are aware of high air pollution exposures. This is an unlikely explanation, especially in the North American study where none of the included cities was highly polluted by local sources, and the Dutch study where all schools were located within 400 m from a major road. The pattern of associations, with bronchitis but not asthma, also makes reporting bias unlikely. Thirdly, lung function data were available for a smaller subset of children, but the precision of estimated effects on lung function was

good because of pooling data from a large number of children. Fourthly, pollution levels were moderate. Air pollution may affect lung function at higher pollution levels, such as in many developing countries. Longitudinal studies in highly polluted Mexico City support this [14]. Fifthly, spirometry was performed with now updated guidelines. The updates in performance of testing are probably not sufficiently major to explain the lack of overall associations in our study. We further showed that the use of recent prediction equations for spirometry also resulted in no association between PM₁₀ and lung function. Finally, we only included cross-sectional studies. Several prospective studies have found associations between air pollution and lung function growth, although a series of Austrian studies did not find consistent effects of particulate matter [14].

Heterogeneity of effects

Despite the efforts to select symptoms with similar wording and harmonise confounder data and data analysis, significant heterogeneity was present in both the symptom and lung function country-specific effect estimates. *A priori*-defined study-level factors could not explain this heterogeneity, possibly related to the relatively small number of studies.

TABLE 4 Combined effect estimates of particulate matter with a 50% cut-off aerodynamic diameter of 10 µm (PM10) on respiratory symptoms from single- and two-pollutant models

	Studies n	Mean OR (95% CI) [#]	
		Single-pollutant model	Two-pollutant model
Bronchitis			
In studies with SO ₂ data	3	1.17 (0.93–1.48)	0.81 (0.44–1.51) [‡]
In studies with NO ₂ data	3	1.35 (0.96–1.91)	1.49 (0.85–2.61)
Phlegm			
In studies with SO ₂ data	2	1.23 (0.81–1.89) [‡]	1.24 (0.80–1.94) [‡]
In studies with NO ₂ data	2	1.55 (1.11–2.18)	1.55 (1.01–2.38)
Nocturnal cough			
In studies with SO ₂ data	2	1.41 (0.68–2.93) [‡]	1.43 (0.61–3.36) [‡]
In studies with NO ₂ data	3	1.14 (0.68–1.91) [‡]	0.79 (0.53–1.19)
Morning cough			
In studies with SO ₂ data	5	1.25 (0.93–1.67) [‡]	1.24 (0.90–1.70) [‡]
In studies with NO ₂ data	3	1.42 (1.04–1.92)	1.36 (0.87–2.13)
Hay fever			
In studies with SO ₂ data	5	1.13 (0.98–1.31)	1.19 (1.01–1.40)
In studies with NO ₂ data	4	1.22 (0.84–1.78) [‡]	1.18 (0.70–1.97)
Itchy rash			
In studies with SO ₂ data	3	1.05(0.93–1.19)	1.13 (0.97–1.33)
In studies with NO ₂ data	3	1.18 (0.94–1.50)	0.91 (0.48–1.71) [‡]
Allergy to pets			
In studies with SO ₂ data	3	1.31 (0.90–1.90) [‡]	1.16 (0.88–1.52)
In studies with NO ₂ data	3	1.25 (0.92–1.70)	0.87 (0.42–1.80) [‡]

Combined effect estimates calculated from country-specific estimates using random effects model. [#]: adjusted for age, sex, maternal education, paternal education, household crowding, current parental smoking, mother smoking during pregnancy, gas cooking, unvented gas/oil/kerosene heater, mould, nationality, birth order, and “ever had a pet” and expressed per 10-µg·m⁻³ increase in PM10; [‡]: evidence of between study heterogeneity (p<0.10).

Differences in study design may have contributed to the observed heterogeneity. Six of the included studies assessed exposure contrasts between communities (Switzerland, Germany, North America, Poland, Hungary and Bulgaria), two studies assessed within-community contrasts (Austria and the Czech Republic), and in the remaining three studies exposure contrasts derived from a mixture of between and within community contrasts. The major common exposure variable PM10 is likely to represent different air pollution mixtures in the different studies, with long-range transported particles contributing only in studies with a between-community component and freshly emitted (ultrafine) particles contributing more to the contrast in the within-city studies. However, particles emitted by motorised traffic have contributed to the exposure contrast in all studies. Within the PATY study, there was no consistent pattern of stronger effects for those studies with a strong traffic-exposure component.

We did not find any difference in effect estimates between males and females, either for symptoms or for lung function. In previous single studies, stronger effects were reported for females in some studies [7, 12] and males in other studies [8].

Limitations

The small number of study areas is a limitation in some of our studies, although in Switzerland, Russia, the Netherlands and North America, the number of areas was fairly large (≥10).

The annual average concentration of PM10 was the main exposure variable within the PATY study. We also evaluated NO₂ and SO₂ as pollutants representing the urban air pollution mixture, but the number of studies with data on multiple pollutants was limited. Hence, the ability to assess effects of PM10 independent of the gaseous pollutants was limited. Two-pollutant models are difficult to interpret, especially when the same source affects both pollutants, as is the case for PM10 and NO₂, for which motorised traffic is an important source. We cannot exclude that other pollutants such as the soot or elemental carbon content of particulate matter or the ultrafine particle concentration were associated more strongly with the health outcomes.

Several studies have assessed indicators of motorised-traffic emissions at a fine spatial scale (<100–300 m) [6, 10, 12, 24]. Our central monitoring data do not reflect these fine-scale air pollution variations. Because we did not have access to individual addresses, we were unable to generate individual exposure estimates at the residential address. Hence, our study does not provide information on the role of primary traffic emissions. Although this is a limitation, studies comparing the average concentration across communities remain valuable, as they allow estimation of the potential health effects of more aged pollution mixtures. Furthermore, a large fraction of the population does not live directly on major roads. Personal monitoring studies have shown that the population average




FIGURE 3. Forest plots of study-specific and mean effects of particulate matter with a 50% cut-off aerodynamic diameter of 10 µm (PM₁₀) for a) forced vital capacity (FVC), b) forced expiratory volume in 1 s (FEV₁), c) peak expiratory flow (PEF) and d) forced expiratory flow at 25–75% of the FVC (FEF_{25–75%}). Figures are given as percentage difference in lung function per 10-µg·m⁻³ increase in PM₁₀ from single pollutant models, adjusted for individual risk factors. The vertical line indicates null, i.e. 0% change (no effect), and horizontal lines represent 95% confidence intervals of estimates. “Combined” indicates the mean and its confidence interval of the individual estimates.

TABLE 5 Combined estimates for the fully adjusted effect of a 10-µg·m⁻³ increase in particulate matter with a 50% cut-off aerodynamic diameter of 10 µm on lung function in different subgroups

	FVC	FEV ₁	FEF _{25–75%}	PEF
Sex				
Male	0.1 (-0.8–0.9)	0.3 (-0.6–1.2)	1.0 (-0.8–2.7)	-0.1 (-1.0–0.8)
Female	-0.1 (-0.7–0.6)	0.1 (-0.6–0.8)	0.2 (-1.2–1.7)	-0.3 (-1.8–1.2)
Age yrs				
6–9	0.0 (-1.4–1.4)	0.1 (-1.2–1.4)	0.1 (-2.4–2.6)	-0.3 (-2.3–1.7)
10–12	-0.1 (-0.8–0.6)	0.4 (-0.6–1.4)	1.2 (-0.6–3.0)	0.1 (-1.4–1.6)
Wheeze				
No	0.2 (-0.8–1.2)	0.4 (-0.6–1.4)	1.0 (-0.8–2.8)	-0.1 (-1.7–1.5)
Yes	-0.3 (-1.2–0.5)	-0.4 (-1.3–0.5)	-0.3 (-3.9–3.3)	0.4 (-1.3–2.1)
Sensitivity to inhaled allergens				
No	0.3 (-0.8–1.4)	0.6 (-0.8–2.0)	1.2 (-0.7–1.1)	0.3 (-1.7–2.3)
Yes	-0.4 (-1.1–0.3)	0.0 (-0.9–0.9)	0.6 (-1.8–3.0)	-0.3 (-1.6–1.0)

Data are presented as % difference (95% confidence interval). FVC: forced vital capacity; FEV₁: forced expiratory volume in 1 s; FEF_{25–75%}: forced expiratory flow at 25–75% of the FVC; PEF: peak expiratory flow. Combined effect estimates calculated from country-specific estimates using random effects model.

personal exposure of subjects is strongly related to the outdoor air pollution level measured at a central site in the community [25]. For individual subjects, differences in exposure from the population average may occur. However, much of the measurement error is likely to be Berkson error, which generally does not lead to bias [26]. Finally, the Swiss SAPALDIA (Swiss study on Air Pollution and Lung Disease In Adults) and the Children's Health study study reported an effect of both within and between community contrasts in NO₂ exposure on lung function of adults and children's respiratory symptoms [7, 27, 28]. Chronic bronchitis of adults in the European Community Respiratory Health Survey was associated with individual level variables representing traffic, but not centre level variables such as the central site PM_{2.5} concentration [29]. In that study, the between-community contrast, however, was largely due to differences between countries, a contrast that we specifically decided not to assess because of the potential for too many differences for which we did not have data. Despite the value of assessing community-level pollution, there is a clear need for assessment of the health effects of near-traffic exposures. There is, for example, some evidence that asthma incidence in children and adults may be associated with near-traffic exposures and not with urban background pollution [30].

In conclusion, our study adds to the evidence that long-term exposure to outdoor air pollution, characterised by the concentration of PM₁₀, is associated with increased respiratory symptoms (phlegm and morning cough) in children. We did not find an association between PM₁₀ and lung function, possibly due to modest pollution levels and heterogeneity across studies.

SUPPORT STATEMENT

The PATY study was funded by the EU Fifth Framework Quality of Life Programme (contract number QLRT-2001-02544).

STATEMENT OF INTEREST

None declared.

ACKNOWLEDGEMENTS

The study was coordinated by T. Fletcher (London School of Hygiene and Tropical Medicine, London, UK).

Author affiliations are as follows. G. Hoek: Institute for Risk Assessment Sciences (IRAS), Utrecht University, Utrecht, the Netherlands; S. Pattenden: London School of Hygiene and Tropical Medicine, London, UK; S. Willers: Institute for Risk Assessment Sciences (IRAS), Utrecht University, Utrecht, the Netherlands; T. Antova: Environmental Health Unit, NCPHP, Sofia, Bulgaria; E. Fabianova: Regional Authority of Public Health, Banska Bystrica, Slovakia; C. Braun-Fahrlander: Swiss Tropical and Public Health Institute and University of Basel, Basel, Switzerland; F. Forastiere: Dept of Epidemiology, ASL Rome, Rome, Italy; U. Gehring: Institute for Risk Assessment Sciences (IRAS), Utrecht University, Utrecht, the Netherlands and Helmholtz Zentrum München, Institute of Epidemiology, Neuherberg, Germany; H. Luttmann-Gibson: Dept of Environmental Health, Harvard School of Public Health, Boston, MA, USA; L. Grize: Swiss Tropical and Public Health Institute and University of Basel, Basel, Switzerland; J. Heinrich: Helmholtz Zentrum München, Institute of Epidemiology, Neuherberg, Germany; D. Houthuijs: National Institute Public Health and the Environment (RIVM), Bilthoven, the Netherlands; N. Janssen: Institute for Risk Assessment Sciences (IRAS), Utrecht University, Utrecht and the National Institute Public Health and the Environment (RIVM), Bilthoven, the Netherlands; B. Katsnelson: Ural Regional Centre for Environmental Epidemiology, Ekaterinburg, Russia; A. Kosheleva: Ural Regional Centre for Environmental Epidemiology, Ekaterinburg, Russia; H. Moshhammer: Institute of Environmental

Health, Medical University of Vienna, Vienna, Austria; M. Neuberger: Institute of Environmental Health, Medical University of Vienna, Vienna, Austria; L. Privalova: Ural Regional Centre for Environmental Epidemiology, Ekaterinburg, Russia; P. Rudnai: National Institute of Environmental Health, "Fodor Jozsef" National Center for Public Health, Budapest, Hungary; F. Speizer: Dept of Environmental Health, Harvard School of Public Health, Boston, MA, USA; H. Slachetova: Institute of Public Health, Center of Health Services, Ostrava, Czech Republic; H. Tomaskova: Institute of Public Health, Center of Health Services, Ostrava, Czech Republic; R. Zlotkowska: Epidemiology Dept, Institute of Occupational Medicine and Environmental Health, Sosnowiec, Poland; and T. Fletcher: London School of Hygiene and Tropical Medicine, London, UK.

REFERENCES

- Weinmayr G, Romeo E, De Sario M, *et al.* Short-term effects of PM₁₀ and NO₂ on respiratory health among children with asthma or asthma-like symptoms: a systematic review and meta-analysis. *Environ Health Perspect* 2010; 118: 449–457.
- World Health Organization. Systematic review of air pollution, a global update. Geneva, World Health Organization, 2006.
- Dockery DW, Speizer FE, Stram DO, *et al.* Effects of inhalable particles on respiratory health of children. *Am Rev Respir Dis* 1989; 139: 587–594.
- Dockery DW, Cunningham J, Damokosh AI, *et al.* Health effects of acid aerosols on North American children: respiratory symptoms. *Environ Health Perspect* 1996; 104: 500–505.
- Heinrich J, Hoelscher B, Wjst M, *et al.* Respiratory diseases and allergies in two polluted areas in East Germany. *Environ Health Perspect* 1999; 107: 53–62.
- Kramer U, Behrendt H, Dolgner R, *et al.* Airway diseases and allergies in East and West German children during the first 5 years after reunification: time trends and the impact of sulphur dioxide and total suspended particles. *Int J Epidemiol* 1999; 28: 865–873.
- Peters JM, Avol E, Gauderman WJ, *et al.* A study of twelve Southern California communities with differing levels and types of air pollution. II. Effects on pulmonary function. *Am J Respir Crit Care Med* 1999; 159: 768–775.
- Peters JM, Avol E, Navidi W, *et al.* A study of twelve Southern California communities with differing levels and types of air pollution. I. Prevalence of respiratory morbidity. *Am J Respir Crit Care Med* 1999; 159: 760–767.
- Hirsch T, Weiland SK, von Mutius E, *et al.* Inner city air pollution and respiratory health and atopy in children. *Eur Respir J* 1999; 14: 669–677.
- Janssen NA, Brunekreef B, van Vliet P, *et al.* The relationship between air pollution from heavy traffic and allergic sensitization, bronchial hyperresponsiveness, and respiratory symptoms in Dutch schoolchildren. *Environ Health Perspect* 2003; 111: 1512–1518.
- Braun-Fahrlander C, Vuille JC, Sennhauser FH, *et al.* Respiratory health and long-term exposure to air pollutants in Swiss schoolchildren. SCARPOL Team. Swiss Study on Childhood Allergy and Respiratory Symptoms with Respect to Air Pollution, Climate and Pollen. *Am J Respir Crit Care Med* 1997; 155: 1042–1049.
- Oftedal B, Brunekreef B, Nystad W, *et al.* Residential outdoor air pollution and lung function in schoolchildren. *Epidemiology* 2008; 19: 129–137.
- Raizenne M, Neas LM, Damokosh AI, *et al.* Health effects of acid aerosols on North American children: pulmonary function. *Environ Health Perspect* 1996; 104: 506–514.
- Gotschi T, Heinrich J, Sunyer J, *et al.* Long-term effects of ambient air pollution on lung function: a review. *Epidemiology* 2008; 19: 690–701.
- Gauderman WJ, Avol E, Gilliland F, *et al.* The effect of air pollution on lung development from 10 to 18 years of age. *N Engl J Med* 2004; 351: 1057–1067.

- 16 Pattenden S, Hoek G, Braun-Fahrlander C, *et al.* NO₂ and children's respiratory symptoms in the PATY study. *Occup Environ Med* 2006; 63: 828–835.
- 17 Moshhammer H, Hoek G, Luttmann-Gibson H, *et al.* Parental smoking and lung function in children: an international study. *Am J Respir Crit Care Med* 2006; 173: 1255–1263.
- 18 Pattenden S, Armstrong BG, Houthuijs D, *et al.* Methodological approaches to the analysis of hierarchical studies of air pollution and respiratory health – examples from the CESAR study. Central European Study on Air pollution and Respiratory Health. *J Expo Anal Environ Epidemiol* 2000; 10: 420–426.
- 19 Agabiti N, Mallone S, Forastiere F, *et al.* The impact of parental smoking on asthma and wheezing. SIDRIA Collaborative Group. Studi Italiani sui Disturbi Respiratori nell'Infanzia e l'Ambiente. *Epidemiology* 1999; 10: 692–698.
- 20 Katsnelson B, Kuzmin SV, Privalova L, *et al.* [Chronic respiratory disease risk factors in younger schoolchildren dwelling in towns with different levels of ambient air pollution]. *Vestnik Uralskoy Meditsinskoy Akademicheskoy Nauki* 2007; 16: 27–37.
- 21 Gehring U, Pattenden S, Slachetova H, *et al.* Parental education and children's respiratory and allergic symptoms in the Pollution and the Young (PATY) study. *Eur Respir J* 2006; 27: 95–107.
- 22 DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986; 7: 177–188.
- 23 Stanojevic S, Wade A, Cole TJ, *et al.* Spirometry centile charts for young Caucasian children: the Asthma UK Collaborative Initiative. *Am J Respir Crit Care Med* 2009; 180: 547–552.
- 24 Nicolai T, Carr D, Weiland SK, *et al.* Urban traffic and pollutant exposure related to respiratory outcomes and atopy in a large sample of children. *Eur Respir J* 2003; 21: 956–963.
- 25 Monn C, Brandli O, Schindler C, *et al.* Personal exposure to nitrogen dioxide in Switzerland. SAPALDIA team. Swiss Study on Air Pollution and Lung Diseases in Adults. *Sci Total Environ* 1998; 215: 243–251.
- 26 Armstrong BG. Effect of measurement error on epidemiological studies of environmental and occupational exposures. *Occup Environ Med* 1998; 55: 651–656.
- 27 Schindler C, Ackermann-Lieblich U, Leuenberger P, *et al.* Associations between lung function and estimated average exposure to NO₂ in eight areas of Switzerland. The SAPALDIA Team. Swiss Study of Air Pollution and Lung Diseases in Adults. *Epidemiology* 1998; 9: 405–411.
- 28 Gauderman WJ, Avol E, Lurmann F, *et al.* Childhood asthma and exposure to traffic and nitrogen dioxide. *Epidemiology* 2005; 16: 737–743.
- 29 Sunyer J, Jarvis D, Gotschi T, *et al.* Chronic bronchitis and urban air pollution in an international study. *Occup Environ Med* 2006; 63: 836–843.
- 30 Perez L, Rapp R, Künzli N. The Year of the Lung: outdoor air pollution and lung health. *Swiss Med Wkly* 2010; 140: w13129.

Online supplement

PM₁₀ and children's respiratory symptoms and lung function in the PATY study

Gerard Hoek¹, Sam Pattenden², Saskia Willers¹, Temenuga Antova³, Eleonora Fabianova⁴, Charlotte Braun-Fahrländer^{5,6}, Francesco Forastiere⁷, Ulrike Gehring^{1,8}, Heike Luttmann-Gibson⁹, Leticia Grize^{5,6}, Joachim Heinrich⁸, Danny Houthuijs¹⁰, Nicole Janssen^{1,10}, Boris Katsnelson¹¹, Anna Kosheleva¹¹, Hanns Moshammer¹², Manfred Neuberger¹², Larisa Privalova¹¹, Peter Rudnai¹³, Frank Speizer⁹, Hana Slachtova¹⁴, Hana Tomaskova¹⁴, Renata Zlotkowska¹⁵, Tony Fletcher²

- 1 Institute for Risk Assessment Sciences (IRAS), Utrecht University, Utrecht, the Netherlands
- 2 London School of Hygiene and Tropical Medicine, London, United Kingdom
- 3 Environmental Health unit, NCPHP Sofia, Bulgaria
- 4 Regional Authority of Public Health, Banska Bystrica, Slovak Republic
- 5 Swiss Tropical and Public Health Institute, Basel, Switzerland
- 6 University of Basel, Basel, Switzerland
- 7 Department of Epidemiology, ASL Rome, Italy
- 8 Helmholtz Zentrum München, Institute of Epidemiology, Neuherberg, Germany
- 9 Department of Environmental Health, Harvard School of Public Health, Boston, MA, USA
- 10 National Institute Public Health and the Environment (RIVM), Bilthoven, the Netherlands
- 11 Ural Regional Centre for Environmental Epidemiology, Ekaterinburg, Russia
- 12 Institute of Environmental Health, Medical University of Vienna, Vienna, Austria
- 13 National Institute of Environmental Health, "Fodor Jozsef" National Center for Public Health, Budapest, Hungary
- 14 Institute of Public Health, Center of Health Services, Ostrava, Czech Republic
- 15 Epidemiology Department, Institute of Occupational Medicine and Environmental Health, Sosnowiec, Poland

The online supplement contains in order:

1. More specific explanation of exposure assessment
2. Tables 1 – 2 with more detail on methods of exposure assessment
3. Tables 3 -6 additional PM₁₀ effect estimates not in the main text
4. References for the supplement

METHODS

Exposure assessment

Exposure assessment methods were assessed for comparability, including an evaluation of site selection for the monitors and the monitoring methods, especially for particulate matter. Tables 1 and 2 present the details for the included studies. In the evaluation it was taken into account that in the epidemiological analysis, analyses of the relationships between air pollution and respiratory health were made per country. Hence modest systematic differences in exposure assessment between studies do not directly affect the effect estimates. Although the overall conclusion was that exposure assessment was sufficiently comparable to allow summarizing air pollution effect estimates given a specific study design, some components of the studies were excluded from the analysis:

- Italian particle data, as a different fraction than PM₁₀ was measured using different methods within the study, for which insufficient collocation with PM₁₀ monitors existed
- One of the original four Bulgarian study areas was excluded because measurements used to representing an entire quarter of the city were made at a curbside of a major road
- One of the Russian study areas was excluded because the monitor was located on the premises of a major industrial site and too far away from the study area. A correlation analysis of daily samples supported this exclusion, as the PM concentrations from the excluded site did not correlate with other sites, whereas significant correlations were present among the other sites
- For some of the Austrian and Swiss sites recommendations were made to test the sensitivity of the epidemiological associations for presence of these sites, because there were concerns that these sites were too much affected by traffic on the nearest road

For those studies that did not directly measure PM₁₀ with gravimetric methods, conversion factors were derived. In the Netherlands, PM_{2.5} was measured at all 24 schools. At some of the sites, PM_{2.5} measurements were co-located with PM₁₀ measurements during the study. Therefore, PM_{2.5} could be converted into PM₁₀ using the formula: $PM_{10} = 9.37 + 1.21 * PM_{2.5}$ ($R^2 = 0.93$). (Janssen, 2001)¹. In the German and Austrian study, TSP (total suspended particles) was measured in a

routine monitoring network with beta attenuation continuous monitors. Site-specific comparisons were made with actual PM₁₀ measurements shortly after the study, thus the original particle concentration could be transformed into PM₁₀ concentrations. In Germany, the average PM₁₀/TSP was 0.77. In Austria, the average PM₁₀/TSP ratio was 0.77, 0.93 and 0.94 at three sites used in the current study.

We further recalculated the annual averages for the Russian study from the original daily data because in some areas the study period was shorter than the full year, resulting in potential bias in the exposure estimates because of seasonal variation. We used the average ratio of the annual average for the sites with complete study periods to the more restricted period to make adjustments.

Table S1 Overview of exposure assessment

Center	Data source	Assignment	Monitoring sites	Temporal coverage PM ₁₀ /TSP	Temporal coverage gases	Particle fraction	Particle composition	Other pollutants
Switzerland	Gases: routine network PM ₁₀ : study specific	1 fixed site in each community ^A	Background/traffic	Weekly samples; 50/year	Continuous monitor, entire year	PM ₁₀		SO ₂ , NO ₂ , O ₃
Austria	Routine network	School assigned to nearest monitor	Background ^B	Continuous monitor, entire year	Continuous monitor, entire year	TSP		SO ₂ , NO ₂ , O ₃ , CO
CESAR	Study specific	1 fixed site in each community	Background	Daily samples; 1 per 6 days, one year	NA	PM ₁₀	PM _{2.5} , soot	-
Germany	Routine network	1 fixed site in each community	Background/industrial	Continuous monitor, entire year	Continuous monitor, entire year	TSP	--	SO ₂
Italy	Routine networks	Schools within 1 km of a monitor	Background/traffic	Continuous monitor, entire year	Continuous monitor, entire year	SP/PM ₁₀		SO ₂ , NO ₂
Netherlands	Study specific	1 fixed site per school	Traffic	Weekly samples; 5-10 per year	Weekly samples; 5-10 per year	PM _{2.5}	Soot	NO ₂ , VOC
Russia	Study specific	1 fixed site per study area	Background/industrial	Weekly samples; period 7-12 months (differed per site)	Weekly samples, Nov. 1998 – May 1999	PM ₁₀	PM _{2.5}	SO ₂ , NO ₂ , VOC
USA	Study specific / routine monitoring (ozone some sites)	1 fixed site in each community	Background	Daily samples, every other day	Continuous monitor, entire year	PM ₁₀	PM _{2.5} , H ⁺ , SO ₄ ²⁻	SO ₂ , O ₃

CESAR includes study areas in Bulgaria, Czech Republic, Hungary, Poland and Slovakia

A More sites for NO₂ (five categories, from questionnaire); multiple sites for gaseous pollutants

B Two sites more traffic impact, sensitivity analysis agreed with investigators

Table S2 Monitoring methods

	PM principle	Monitor	Conversion to PM ₁₀	SO ₂	NO ₂
Switzerland	Gravimetric	Harvard impactor	NA	UV Fluorescence*	Chemiluminescence**
Austria	Beta attenuation	EberlineFH63I-N	Co-located measurements several sites	UV Fluorescence	Chemiluminescence
CESAR	Gravimetric	Harvard impactor	NA	NA	NA
Germany	Beta attenuation	FAGFH-62-IN	Co-located measurements with PM ₁₀ in two of the areas one year later and Erfurt	UV Fluorescence	NA
Italy	Diverse	Diverse	Not possible, too diverse		
Netherlands	Gravimetric	Harvard impactor	Co-located measurements PM _{2.5} and PM ₁₀ several schools during study	NA	Palmes tube
Russia	Gravimetric	Harvard impactor	NA	Ogawa badge	Ogawa badge
USA	Gravimetric	Harvard impactor	NA	Denuder	NA

NA not available

* DOAS ([Differential Optical Absorption Spectroscopy](#)) at one site (Biel) and no data at one site (Langnau)

** DOAS ([Differential Optical Absorption Spectroscopy](#)) at one site (Biel) and passive samplers (Langnau)

Table S3 Results from meta-regression analyses stratified by study characteristic for phlegm (8 studies). Odds ratios per 10 $\mu\text{g}/\text{m}^3$ PM_{10} .

Phlegm	No. of studies	Mean odds ratio (95% CI)	p-heterogeneity
Original evidence of heterogeneity			0.03
Study design: Between- or within-towns (or a mixture)			0.30
between towns	4	1.09 (0.91 - 1.30)	
within towns	1	1.06 (0.96 - 1.16)	
mixture	3	1.46 (1.17 - 1.82)	
Season of questionnaire			0.03
<2/3 in spring	3	1.22 (0.83 - 1.80)	
2/3+ in spring	5	1.15 (1.00 - 1.33)	
Variability of date of questionnaire			0.01
high variability	1	1.41 (0.64 - 3.13)	
low variability	7	1.14 (1.01 - 1.30)	
East or West⁵			0.03
East	6	1.21 (1.04 - 1.40)	
West	2	0.98 (0.84 - 1.14)	
Period of study			0.04
pre 95 studies	1	0.97(0.83 - 1.13)	
95 onwards	7	1.21(1.05 - 1.39)	
Response rate			0.01
response rate 80+%	3	1.18(0.84 - 1.66)	
response rate <80%	5	1.16(1.00 - 1.35)	
Response rate variability			0.01
low variability	5	1.18(0.98 - 1.41)	
med. variability	2	1.16(0.86 - 1.57)	
high variability	1	1.41(0.64 - 3.13)	
Proportion of young children (aged 6-8)			0.01
low (<20%)	2	1.20(0.74 - 1.94)	
Medium	6	1.15(1.01 - 1.32)	

Table S4 Combined estimates of PM₁₀ effect by age-group and gender.

Symptom	Age		Gender	
	Older children (9 - 12 yr)	Young children (6- 8 yr)	Boys	Girls
Wheeze	1.00(0.95-1.07)	0.99(0.89-1.11)	1.02 (0.92-1.13) ^H	0.99 (0.92-1.06)
Asthma	1.05(0.98-1.14)	0.98(0.88-1.09)	1.01 (0.90-1.13)	1.03 (0.94-1.13)
Bronchitis	1.06(1.00-1.13)	1.12(0.98-1.27)	1.08 (0.98-1.18)	1.06 (0.96-1.18)
Phlegm	1.16(1.00-1.35) ^H	1.13(0.91-1.41) ^H	1.09 (0.99-1.20)	1.13 (0.98-1.32) ^H
Nocturnal cough	1.14(1.02-1.29) ^H	1.15(0.97-1.36) ^H	1.20 (1.02-1.41) ^H	1.08 (0.93-1.27) ^H
Morning cough	1.16(1.03-1.31) ^H	1.11(0.95-1.29) ^H	1.14 (1.02-1.27) ^H	1.12 (0.98-1.28) ^H
Hay fever	1.03(0.95-1.11)	0.97(0.84-1.11) ^H	1.08 (0.90-1.30)	1.36 (1.02-1.83)
Sensitivity to inhaled allergens	1.30(1.00-1.68)	1.04(0.79-1.36)	1.01 (0.91-1.13) ^H	1.03 (0.95-1.12)
Itchy rash	1.07(0.96-1.20)	1.03(0.87-1.22) ^H	1.06 (0.91-1.24) ^H	1.06 (0.95-1.18)
Woken by wheeze	0.99(0.89-1.10)	1.07(0.96-1.21)	1.07 (0.95-1.21)	0.94 (0.86-1.04)
Allergy to pets	1.29(0.95-1.74) ^H	1.00(0.81-1.23)	1.26 (0.97-1.63) ^H	1.08 (0.91-1.28)

Combined effect estimates calculated from country-specific estimates using random effects model.

‘H’ indicates evidence of between study heterogeneity (p<0.10)

Odds ratios and 95% confidence intervals are per 10 µg/m³ PM₁₀

Table S5 Description of lung function data

	Austria	Czech Republic	Germany	Hungary	Netherlands	North America	Poland	Slovakia
N*	2,898	806	1,788	1,260	1,735	12,737	615	970
Age								
6-9 years (%)	100	17.5	56.8	11.9	50.1	42.3	34.2	24.0
10-12 yr (%)	0	82.5	43.2	88.1	49.9	47.7	65.8	76.0
Height (m)**	1.25 (0.64)	1.44 (0.69)	1.41 (0.15)	1.46 (0.71)	1.43 (0.10)	1.42 (0.76)	1.41 (0.70)	1.45 (0.73)
Weight (kg)**	26 (5)	38 (8)	35 (12)	38 (9)	36 (9)	39 (10)	36 (8)	37 (8)
FVC (l)**	1.44 (0.26)	2.49 (0.37)	2.39 (0.71)	2.50 (0.41)	2.38 (0.51)	2.49 (0.43)	2.31 (0.40)	2.48 (0.40)
FEV ₁ (l)**	1.35 (0.23)	2.18 (0.31)	2.20 (0.59)	2.23 (0.35)	2.11 (0.43)	2.13 (0.36)	2.04 (0.33)	2.21 (0.34)
FEF ₂₅₋₇₅ (l.s ⁻¹)**	1.92 (0.48)	2.54 (0.58)	-	2.73 (0.59)	2.35 (0.65)	2.37 (0.62)	2.46 (0.57)	2.68 (0.62)
PEF (l.s ⁻¹)**	3.03 (0.62)	4.50 (0.78)	4.30 (1.27)	4.78 (0.78)	4.56 (1.07)	4.65 (0.92)	4.43 (0.78)	4.62 (0.89)
FVC%pred.)***	86 (10)	100 (9)	100 (11)	97 (11)	98 (11)	104 (11)	98 (11)	99 (11)
FEV ₁ %pred***	90 (11)	101 (10)	103 (12)	100 (11)	99 (11)	102 (12)	100 (10)	101 (11)
FEF ₂₅₋₇₅ %pred***	102 (24)	98 (21)		103 (20)	93 (21)	95 (23)	99 (21)	103 (22)

* number of children with valid lung function test

** Mean and standard deviation in parentheses

*** Mean and standard deviation in parentheses of percent predicted lung function. Predicted using equations from Stanojevic (2009)²

Table S6 **Combined estimates for the fully adjusted effect of 10 µg/m³ increase in PM₁₀ on lung function**

	FVC (% diff and 95% CI)	FEV₁ (% diff and 95% CI)	FEF₂₅₋₇₅ (% diff and 95% CI)	PEF (% diff and 95% CI)
Original analysis, all centres	0.1 (-0.6, 0.8)	0.3 (-0.5, 1.1)	0.7 (-0.8, 2.3)	0.0 (1.2, 1.3)
Original analysis, without CESAR centres**	0.2 (-1.0, 1.3)	0.0 (-0.9, 0.9)	-0.5 (-1.6, 0.6)	-0.4 (-2.0, 1.2)
Analysis with Stanojevic prediction equations, all centres***	-0.0, (-0.7, 0.7)	0.2 (-0.6, 1.1)	0.8 (-0.7, 2.2)	NA

* p < 0.05

** exclusion because of small number of valid tests

*** percent predicted lung function dependent variable. Prediction using equations from Stanojevic (2009)²

Combined effect estimates calculated from country-specific estimates using random effects model.

NA=not available

References online supplement

1. Janssen NAH, van Vliet PHN, Aarts F, Harssema H, Brunekreef B. Assessment of exposure to traffic related air pollution of children attending schools near motorways. *Atmospheric Environment* 2001;35, 22: 3875-3884.
2. Stanojevic S, Wade A, Cole TJ, Lum S, Custovic A, Silverman M, Hall GL, Welsh L, Kirkby J, Nystad W, Badier M, Davis S, Turner S, Piccioni P, Viložni D, Eigen H, Vlachos-Mayer H, Zheng J, Tomalak W, Jones M, Hankinson JL, Stocks J. Spirometry centile charts for young Caucasian children: The asthma UK collaborative initiative. *American Journal of Respiratory and Critical Care Medicine* 2009; 180: 547-552.