

Letters to the Editor

Disruptive anti-IgE inhibitors prevent mast cell-dependent early airway response in viable atopic lung tissue

To the Editor

IgE plays a central role in the pathophysiology of allergic asthma. The therapeutic anti-IgE antibody omalizumab has been shown to inhibit allergic asthma symptoms by neutralizing free serum IgE and thereby preventing early and late phases of IgE-mediated allergic reactions after 12 to 16 weeks of treatment. Unfortunately, use of omalizumab is restricted to the treatment of moderate-to-severe asthma, ineffective to approximately one third of patients, and associated with high costs. Thus there is an urgent need for the development of new and more efficient therapeutic options to treat severe types of asthma. Here we report that the disruptive anti-IgE designed ankyrin repeat protein (DARPin, a registered trademark of Molecular Partners AG, Switzerland) bi53_79 (hereafter referred to as E002) might significantly improve treatment efficacy in a therapeutic setting in which allergic effector cells in the lung are sensitized with allergen-specific IgE.

In patients with IgE-mediated allergic asthma, mast cells are loaded with allergen-specific IgE, which is bound to the cell surface through the high-affinity IgE receptor (FcεRI).¹ Inhalation of allergens might lead to rapid activation of these airway mast cells through allergen-dependent cross-linking of FcεRI-bound IgE, followed by degranulation and release of mediators, including histamine. Histamine binds to the H1 receptor on airway smooth muscle cells and induces an allergen-specific immediate airway constriction defined as the early airway response (EAR).

The previously described disruptive anti-IgE DARPin E2_79 (hereafter referred to as E001) not only neutralizes free serum IgE with high specificity but further actively dissociates IgE-FcεRI complexes on human primary allergic effector cells.² Compared with classical antibody formats, DARPins represent alternative binding scaffolds. Their small molecular size (approximately 12 kDa) and rigid repetitive ankyrin structure lacking disulfide bridges allows highly efficient bacterial expression. E001 specifically targets human IgE, blocks its binding to FcεRI, and disrupts preformed IgE-FcεRI complexes through a mechanism termed facilitated dissociation in which subsite attachment points within IgE become partially exposed and accessible for E001 binding before full dissociation from FcεRI. In a humanized FcεRI transgenic mouse model, both E001 and its improved biparatopic variant, E002, inhibited allergic skin reactions with high potency after stimulation of passively sensitized cutaneous mast cells *in vivo*.³

In this study we sought to assess the efficacy of E002 to prevent mast cell activation in atopic precision-cut lung slices (PCLS), which represent a well-established technique with a high predictive value for the *in vivo* situation. PCLS contain all cells of the lower respiratory tract, such as epithelial cells, smooth muscle cells, fibroblasts, macrophages, and mast cells. Lung tissue remains viable and maintains functionality for days. Previous studies demonstrated the efficacy of bronchoconstrictors in a multispecies PCLS study and reported that histamine induces

airway constriction in lung tissue of human subjects and nonhuman primates, whereas serotonin leads to airway constriction only in rodents.⁴ Because this pathomechanism is critical for EAR, we functionally compared E002 to the therapeutic anti-IgE antibody omalizumab in human PCLS cultures *ex vivo*.

Allergen-induced airway constriction is mast cell dependent and occurs within the first 10 minutes on allergen exposure. This airway smooth muscle contraction is mainly induced by mediators, such as histamine, leukotrienes, and prostanoids, released from activated mast cells, which reach maximal concentrations a few minutes after allergen challenge.⁵ Initially, we investigated allergen-induced airway constriction in human lung tissue *ex vivo*. Human PCLS that were passively sensitized with plasma of donors with house dust mite (HDM) allergy showed extensive bronchoconstriction after allergen challenge with HDM extract (Fig 1, A and C). Maximal reduction of the initial airway area ($68\% \pm 10\%$) was achieved 20 minutes after allergen challenge (Fig 1, A). Airway constriction was significantly greater in PCLS that were sensitized with plasma of allergic donors when compared with PCLS that were sensitized with plasma from nonallergic subjects.

We further assessed whether the amount of released histamine correlates with the observed bronchoconstriction. Indeed, tissue culture supernatants of PCLS sensitized with plasma of allergic donors contained significantly greater histamine levels (32 ± 10 ng/mL) compared with control sensitized PCLS after allergen challenge (2 ± 0 ng/mL; Fig 1, C). Similar to the findings in bronchoalveolar lavage fluid of human asthmatic patients, we detected greater histamine quantities in supernatants of sensitized and allergen-provoked PCLS compared with those of nonsensitized tissue control subjects.

Omalizumab neutralizes free IgE with high potency, but it is not very efficient in accelerating the dissociation of preformed IgE-FcεRI complexes.³ In an experimental setting in which omalizumab was added to the plasma of an allergic donor before it was incubated with PCLS, omalizumab efficiently inhibited airway constriction and significantly reduced histamine release from 51 ± 6 to 7 ± 5 ng/mL (Fig 2, A and B). Thus preventive treatment with omalizumab during passive PCLS sensitization completely abolished allergen-induced bronchoconstriction and histamine release.

These findings confirm the efficacy of omalizumab in neutralizing free IgE and thus in preventing IgE binding and sensitization of airway mast cells.¹ However, when we treated presensitized human PCLS with $50 \mu\text{mol/L}$ omalizumab for 2 hours before HDM challenge, we observed strong bronchoconstriction with a reduction of initial airway area to $66\% \pm 17\%$ and prominent histamine release (48 ± 21 ng/mL; Fig 2, C and D). Thus when omalizumab was added after passive sensitization of PCLS with plasma of an allergic donor, it showed no inhibition of the EAR, confirming previous reports that omalizumab is a poor disruptor of pre-existing IgE-FcεRI complexes. In contrast, when we used the therapeutic setting, in which human PCLS were first sensitized with human plasma from an allergic donor and subsequently had E002 added, we observed a marked reduction of allergen-induced histamine release and bronchoconstriction (Fig 2, E and F). Allergen-dependent bronchoconstriction in passively sensitized human PCLS was significantly decreased at




FIG 1. A, EAR in a sensitized human PCLS. PCLS were incubated overnight with 1% plasma of a donor with HDM allergy or plasma from a nonallergic donor. Bronchoconstriction provoked with HDM was monitored by using videomicroscopy. Reduction of the airway area was calculated as a percentage of the initial airway area. Representative microscopic images of the EAR in passively sensitized PCLS versus appropriate controls. Images of cross-sectioned airways before and 10 and 20 minutes after allergen (HDM) challenge. B, HDM provoked a reduction in airway area by $68\% \pm 10\%$ in PCLS incubated with allergic plasma. PCLS incubated with plasma of a nonallergic donor showed no bronchoconstriction. C, HDM challenge of passively sensitized PCLS or PCLS treated with plasma of a nonallergic donor led to an average histamine release of 32 ± 10 and 2 ± 0 ng/mL, respectively. Scale bar = 200 μ m. PCLS in duplicates, $n = 3$. *** $P < .001$, t test using the Mann-Whitney method. Data are expressed as means $\pm$ SEMs. A, 1% Allergic plasma; NA, 1% nonallergic plasma.

0.5 μ mol/L E002 and was completely abolished at 50 μ mol/L. The total amount of 32 ± 10 ng/mL histamine in control DARPIn-treated allergen-challenged human PCLS was reduced to 3 ± 1 ng/mL with 0.5 μ mol/L E002. With increasing concentrations, the measured histamine levels further decreased to 2 ± 1 ng/mL at 5 μ mol/L E002 and 0.5 ± 0 ng/mL at 50 μ mol/L E002. Differences in airway response were significant between the treatment groups of E002 and control DARPIn.

These data are in line with the previously reported disruptive activity of E002, and the results confirm a previous study reporting that intradermal injections of E002 six hours before antigen challenge are effective at inhibiting allergic skin reaction in passively sensitized humanized Fc ϵ RI transgenic mice.³ In the same study it was described that omalizumab was only effective to inhibit the allergic skin reaction in mice when used at high concentrations. Indeed, omalizumab also accelerates the dissociation of receptor-bound IgE from isolated primary human blood basophils, although only at high concentrations or extended incubation periods. In contrast, E002 is 10,000-fold more efficient than omalizumab in dissociating IgE complexes.³ Thus our data confirm dissociation of preformed IgE-Fc ϵ RI complexes considering that a 2-hour treatment at a relatively low concentration of 0.5 μ mol/L E002 was sufficient to significantly inhibit the EAR in presensitized PCLS. Although E002 features lower IgE-binding

affinity compared with omalizumab,³ its main advantage is the efficacy with which it actively disrupts preformed IgE-Fc ϵ RI complexes through a facilitated dissociation mechanism.²

Currently, omalizumab is the only anti-IgE therapy approved to treat moderate-to-severe allergic asthma. It is well established that omalizumab efficiently decreases the concentration of free serum IgE and thereby decreases Fc ϵ RI expression on circulating basophils.¹ Moreover, there is evidence that omalizumab reduces IgE in inflamed tissues, such as the airway mucosa.⁶ Previous studies have reported that patients with mild-to-moderate persistent asthma treated with omalizumab for 16 weeks had significant reduction in numbers of IgE-positive cells, as well as cells expressing Fc ϵ RI in the bronchial submucosa.⁶ However, despite a significant decrease in Fc ϵ RI expression on mast cells within the tissue, it is suggested that the time course for this decrease is very slow. For instance, a significant reduction in mast cell Fc ϵ RI expression in skin biopsy specimens from patients with moderately severe allergic rhinitis was only observed after 10 weeks of omalizumab treatment.⁷ These findings might be explained by either poor tissue penetrance of omalizumab or the remarkably slow dissociation rate of IgE from Fc ϵ RI on mast cells on omalizumab treatment.⁸ This latter observation is crucial because the half-life of cell-bound IgE is in the range of weeks and therefore provides persistent sensitization of mast cells.⁸




FIG 2. **A**, EAR is abolished with a preventive approach of omalizumab in sensitized human peripheral airways. One percent allergic plasma was pretreated with or without omalizumab (5 μ mol/L) before adding to the PCLS. Bronchoconstriction provoked with HDM was monitored by using videomicroscopy. Reduction of the airway area was calculated as a percentage of the initial airway area. HDM provoked a reduction of the airway area by $84\% \pm 9\%$ in PCLS incubated with allergic plasma. PCLS incubated with allergic plasma pretreated with omalizumab (5 μ mol/L) showed only a maximal airway constriction of $12\% \pm 9\%$. **B**, HDM treatment of passively sensitized PCLS treated with or without omalizumab led to an average histamine release of 7 ± 5 and 51 ± 6 ng/mL, respectively. PCLS in duplicates, $n = 3$. $^{**}P < .004$ and $^{***}P < .001$, t test using the Mann-Whitney method. Data are expressed as means $\pm$ SEMs. **C**, Omalizumab does not prevent the EAR in a therapeutic setting. Passively sensitized PCLS were treated with or without omalizumab (5 or 50 μ mol/L) 2 hours before allergen challenge. Bronchoconstriction provoked with HDM was monitored by using videomicroscopy. Reduction of the airway area was calculated as a percentage of the initial airway area. Passively sensitized PCLS untreated or treated with 5 or 50 μ mol/L omalizumab showed maximal airway constriction of $84\% \pm 9\%$, $78\% \pm 8\%$, and $66\% \pm 17\%$, respectively. **D**, Passively sensitized PCLS untreated or treated with 5 or 50 μ mol/L omalizumab led to an average histamine release of 51 ± 6 , 40 ± 13 , and 48 ± 21 ng/mL, respectively. PCLS in duplicates, $n = 3$. $^{*}P < .02$, t test using the Mann-Whitney method. Data are expressed as means $\pm$ SEMs. **E**, E002 prevents EAR in a therapeutic setting. Passively sensitized PCLS were treated with or without E002 (0.5 or 50 μ mol/L) 2 hours before the allergen challenge. Bronchoconstriction provoked with HDM was monitored by using videomicroscopy. Reduction of the airway area was calculated as a percentage of the initial airway area. Passively sensitized PCLS untreated or treated with 0.5 or 50 μ mol/L E002 showed maximal airway constriction of $68\% \pm 10\%$, $30\% \pm 3\%$, and $0\% \pm 5\%$, respectively. Passively sensitized PCLS treated with control DARPin showed maximal airway constriction of $62\% \pm 10\%$. **F**, Passively sensitized PCLS untreated or treated with 0.5 or 50 μ mol/L E002 led to an average histamine release of 32 ± 10 , 3 ± 1 , and 0.5 ± 0 ng/mL, respectively. Passively sensitized PCLS treated with control DARPin led to an average histamine release of 32 ± 10 ng/mL. PCLS in duplicates, $n = 3$. $^{**}P < .004$ and $^{***}P < .001$, t test using the Mann-Whitney method. Data are expressed as means $\pm$ SEMs.

Because of its smaller molecular size, E002 might feature better tissue penetrance than antibodies and might therefore show an advantage to dissociate preformed IgE-Fc ϵ RI complexes at local sites of inflammation. From this point of view, combined therapy of omalizumab with E002 might represent an interesting alternative to improve treatment efficacy.

In summary, we provide evidence that disruptive IgE inhibitors, such as E002, might have a therapeutic advantage over conventional anti-IgE compounds that primarily aim at neutralizing free serum IgE. Our data indicate that active desensitization of mast cells in human lung tissue using E002 is remarkably rapid and efficient. Active disruption of IgE-Fc ϵ RI complexes on the cell surfaces of allergic effector cells represents a novel and

interesting way to interfere with critical atopic pathways. Based on these characteristics, E002 might be a suitable candidate for the treatment of allergic asthma.

We thank all colleagues of the Pathology Unit and Primate Husbandry of the German Primate Center and staff members of the Department of Preclinical Pharmacology and *In Vitro* Toxicology of the Fraunhofer ITEM for their cooperative assistance during the experiments. We also thank them, as well as the members of the Eggel Laboratory, for their technical support and exchange of experiences. We further extend our acknowledgements to Molecular Partners AG for giving us access to the DARPin protein technology.

Sharon Jiménez Delgado, MSc^{a*}
Susann Dehmel, PhD^{a*}

Elaine Twisterling, MSc^{a,b}
 Judy Wichmann, PhD^{a,b}
 Danny Jonigk, MD^c
 Gregor Warnecke, MD^c
 Peter Braubach, Dr Med^c
 Hans-Gerd Fieguth, MD^d
 Ludwig Wilkens, MD^d
 Franziska Dahlmann, PhD^a
 Franz-Josef Kaup, PhD^b
 Alexander Eggel, PhD^e
 Sascha Knauf, PhD^{b*}
 Katherina Sewald, PhD^{a*}
 Armin Braun, PhD^{a*}

From ^athe Fraunhofer Institute for Toxicology and Experimental Medicine, Member of the German Center for Lung Research (DZL), Biomedical Research in Endstage and Obstructive Lung Disease (BREATH) research network, Hannover, Germany; ^bGerman Primate Center GmbH, Leibniz-Institute for Primate Research, Goettingen, Germany; ^cthe Institute for Pathology, Hannover Medical School, Hannover, Germany, Member of the German Center for Lung Research (DZL), Hannover, Germany, Biomedical Research in Endstage and Obstructive Lung Disease Hannover (BREATH) Research Network, Hannover, Germany; ^dthe Institute of Pathology, Klinikum Region Hannover (KRH), Hannover, Germany; and ^ethe Department of Rheumatology, Immunology and Allergology, University Hospital Bern, Bern, Switzerland. E-mail: katherina.sewald@item.fraunhofer.de.

*These authors contributed equally to this work.

This study was funded by the German Research Foundation (DFG), Bonn, Germany, and part of the programme SPP 1394 "Mast-cells—promoters of health and modulators of disease," as well as a grant from Fondation Acteria, a Swiss National Science Foundation Ambizione grant (PZ00P3_148185); the Research Fund of the Swiss Lung Association, Berne, Switzerland and the Uniscientia foundationa (to A.E.).

Disclosure of potential conflict of interest: Fraunhofer ITEM is an independent, public,

nonprofit research organization for applied research performing contract research with many companies in biotechnology and the pharmaceutical, chemical, and cosmetic industries. The authors declare that they have no relevant conflicts of interest.

REFERENCES

- Holgate S, Casale T, Wenzel S, Bousquet J, Deniz Y, Reisner C. The anti-inflammatory effects of omalizumab confirm the central role of IgE in allergic inflammation. *J Allergy Clin Immunol* 2005;115:459-65.
- Kim B, Eggel A, Tarchevskaya SS, Vogel M, Prinz H, Jardetzky TS. Accelerated disassembly of IgE-receptor complexes by a disruptive macromolecular inhibitor. *Nature* 2012;491:613-7.
- Eggel A, Baravalle G, Hobi G, Kim B, Buschor P, Forrer P, et al. Accelerated dissociation of IgE-FcεRI complexes by disruptive inhibitors actively desensitizes allergic effector cells. *J Allergy Clin Immunol* 2014;133:1709-19.e8.
- Seehase S, Schlepütz M, Switalla S, Mätz-Rensing K, Kaup FJ, Zöllner M, et al. Bronchoconstriction in nonhuman primates: a species comparison. *J Appl Physiol* 2011;111:791-8.
- Kunitomo M, Otsuka H. Comparison of antigen-induced leukotriene and histamine release from nasal scrapings in allergic rhinitis. *Rhinology* 2005;43:199-204.
- Djukanović R, Wilson SJ, Kraft M, Jarjour NN, Steel M, Chung KF, et al. Effects of treatment with anti-immunoglobulin E antibody omalizumab on airway inflammation in allergic asthma. *Am J Respir Crit Care Med* 2004;170:583-93.
- Beck LA, Marcotte GV, MacGlashan D, Togias A, Saini S. Omalizumab-induced reductions in mast cell FcεRI expression and function. *J Allergy Clin Immunol* 2004;114:527-30.
- Wan T, Beavil RL, Fabiane SM, Beavil AJ, Sohi MK, Keown M, et al. The crystal structure of IgE Fc reveals an asymmetrically bent conformation. *Nat Immunol* 2002;3:681-6.

<https://doi.org/10.1016/j.jaci.2019.11.002>

METHODS

Source of lung material

The study included human lung tissue from 6 donors. Tissue was obtained from patients with cancer who underwent surgery because of lung tumors. Use of human lung tissue for research was approved by the ethics committee of the Hannover Medical School, Hannover, Germany. Experiments complied with the Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments involving human subjects. The privacy rights of human subjects were observed, and written consent was obtained from all patients. Personal data were not recorded to protect anonymity.

Preparation of PCLS

Preparation of PCLS was performed, as recently published in detail.^{E1} Briefly, only lung lobes were used and filled up inside with medium and agarose solution. Soon after the agarose had solidified, lung tissue was sliced into equal parts to generate approximately 250- μ m-thin PCLS using a Krumdieck tissue slicer (Alabama Research and Development, Munford, Ala). PCLS were then incubated under cell-culture conditions (37°C, 5% CO₂, and 100% air humidity) for 2 hours. PCLS were washed and used for further

experiments. Tissue was passively sensitized in human plasma of allergic donors (HDM-IgE > 100 kU/L, specific IgE class 6; PlasmaLab International, Everett, Wash) overnight. For negative reference, PCLS were also incubated in human plasma of nonallergic donors. Treatment with inhibitor was performed after sensitization. Airway constriction was done as previously published.^{E2} Single PCLS were watched by using stereomicroscopy. HDM extract was added, and airway constriction was watched for 20 minutes. Airway area before addition of allergen was set at 100%. Bronchoconstriction was expressed as the percentage decrease in airway area in comparison with initial airway area. Histamine release was analyzed by means of ELISA (De-meditec Diagnostics GmbH, Kiel, Germany). Viability was checked as published previously.^{E1}

REFERENCES

- E1. Neuhaus V, Danov O, Konzok S, Obernolte H, Dehmel S, Braubach P, et al. Assessment of the cytotoxic and immunomodulatory effects of substances in human precision-cut lung slices. *J Vis Exp* 2018;(135):.
- E2. Kunitomo M, Otsuka H. Comparison of antigen-induced leukotriene and histamine release from nasal scrapings in allergic rhinitis. *Rhinology* 2005;43:199-204.