Food Chemistry 419 (2023) 135993

Contents lists available at ScienceDirect

Food Chemistry

journal homepage: www.elsevier.com/locate/foodchem

Review

RuBisCO as a protein source for potential food applications: A review

Madalena Grácio [a], Sabrina Oliveira [a], Ana Lima [b,c], Ricardo Boavida Ferreira [a,b,*]

[a] Instituto Superior de Agronomia, Lisbon University, 1349-017 Lisbon, Portugal
[b] Linking Landscape, Environment, Agriculture and Food – LEAF, Instituto Superior de Agronomia, Universidade de Lisboa, 1349-017 Lisbon, Portugal
[c] Faculty of Veterinary Medicine, Lusófona University, Campo Grande, 376, 1749-024 Lisbon, Portugal

ARTICLE INFO

Keywords:
Plant-based
RuBisCO
Non-allergenic
Complete protein
Bioactive compounds
Sustainability

ABSTRACT

RuBisCO is a complete protein, widely abundant and recognized as ideal for human consumption. Further, its biochemical composition, organoleptic and physical features mean RuBisCO has potential as a nutritionally beneficial food additive. Nonetheless, despite growing plant-based market trends, there is a lack of information about the applications of this protein. Here, we explored the biochemical features of RuBisCO as a potential food additive and compared it with other plant protein sources currently available. We describe potential advantages, including nutritional content, digestibility, non-allergenicity and, potential bioactivities. Despite the lack of industrial procedures for RuBisCO purification, a growing number of novel methods are emerging, justifying discussion of their feasibilities. Overall, this information can help both researchers and industry to review the use RuBisCO as a sustainable source of protein for plant-based food products or formulation of novel functional foods.

1. Introduction

The world is current witnessing an unprecedented demand for plant-based protein sources. The global plant protein market was valued at $29.4 billion in 2020, and is projected to reach $162 billion by 2030, according to Bloomberg Intelligence (New York magazine, August 2020). Moreover, plant-based diets have gained popularity in recent decades due to their potential health advantages and environmental and ethical considerations (Aimutis, 2022). This growing interest has contributed to the surge in demand for plant-based proteins. Consumers worldwide are encouraged to consume more protein, with a preference for plant-based sources and a potential shift away from animal protein. This is motivated by several factors, including the belief that plant-based diets may improve health, concerns about the negative effects of animal protein on human health, the environmental impact and ethical concerns surrounding animal protein production, and the popularity of high protein diets like the ketogenic and Atkins diets (Hertzler et al., 2020). Accordingly, the food industry swiftly adjusted to the growing demand for plant protein by introducing a whole new line of food products. However, these commercially available plant-based proteins have several drawbacks. One of the major challenges is their inconsistent functional characteristics, as the nutritional, molecular, and physicochemical properties of plant proteins are significantly distinct from those of animal proteins. Compared to animal-derived food products, plant-derived products typically have a lower percentage (w/w) of protein. From a nutritional point of view, plant-based products may contain naturally occurring substances that can impact nutrient absorption and digestion (Popova & Mihaylova, 2019). In addition, plant-based proteins are frequently deficient in essential amino acids, such as lysine in cereal seeds, cysteine and methionine in legume seeds, and tryptophan in general (McGuire & Beerman, 2013). Another major problem related to both animal- and plant-based proteins may pose a significant problem in terms of potential allergenicity. It is estimated that around 220 to 250 million individuals suffer from food allergies, with developed countries reporting an incidence of 5 to 8% in children and 1 to 2% in adults (Dorsan & Chaudaty, 2013; Jones, 2014). Food allergies have a significant impact on people's quality of life and can also affect the national and global economy due to the costs associated with diagnosis and treatment, as well as the production of allergen-free foods (Mills et al., 2007; Dorsan & Chaudaty, 2013). One of the most consumed plant-based protein derives from soy seeds, which is among the eight major food allergens according to the Food and Agriculture Organization of the United Nations (FAO). As a result, it is crucial to identify suitable botanical sources and extraction methods to produce consistent and functional plant proteins.

The food industry currently lacks a plant-based protein that can offer

* Corresponding author.
E-mail address: rbferreira@isa.ulisboa.pt (R. Boavida Ferreira).

https://doi.org/10.1016/j.foodchem.2023.135993
Received 2 December 2022; Received in revised form 13 March 2023; Accepted 18 March 2023
Available online 24 March 2023
0308-8146/© 2023 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

all the benefits of animal protein without the previously mentioned drawbacks. The enzyme ribulose bisphosphate carboxylase/oxygenase (EC.4.1.1.39; RuBisCO), a protein that plays a vital role in photosynthesis, could be an excellent option as a plant protein source, as it overcomes most of the serious constraints referred before. RuBisCO's potential for human consumption relates to its high nutritional value, as it is highly digestible and contains a balanced proportion of essential amino acids that align with the recommendations of the Food and Agriculture Organization (FAO) and the World Health Organization (WHO), as noted by Pouvreau et al. (2013). In addition to its natural abundance, RuBisCO offers various nutritional benefits compared to other plant-based proteins. It is regarded as a non-allergenic protein (Jeurink & Savelkoul, 2005; Cirkovic Veličković & Gavrovic-Jankulović, 2014) and possesses numerous organoleptic, functional, and physical properties that are highly promising for the food industry. However, there is currently a lack of efficient methods for large-scale RuBisCO purification and subsequent utilization in the food sector.

In this review, we examine the potential of RuBisCO as a safe and efficient plant protein source for human consumption by highlighting its nutritional and biochemical value, functional properties compared to other commercially available plant-based proteins, and bioactivities for human health. Additionally, we discuss its major industrial applications and evaluate current methods for large-scale purification, weighing the advantages and disadvantages of each. To the best of our knowledge, this is the first review about RuBisCO's features as a food additive. Given the current trend towards innovation in plant-based proteins and the growing number of methods reported for isolating RuBisCO for food purposes, this review can make a significant contribution to both the food industry and researchers.

2. Biochemical features of RuBisCO

RuBisCO is responsible for one crucial step in photosynthesis that makes most life as we know possible. Its primary function is to catalyze various chemical reactions, with the most significant being the first reaction of the Calvin cycle. This sequence of reactions begins with the absorption of carbon dioxide and ends with the synthesis of carbohydrates, the carbon skeletons upon which virtually all life on Earth depends. As a result, RuBisCO is responsible for fixating more than 90% of the inorganic carbon which is converted into biomass. However, RuBisCO can also react with oxygen instead of carbon dioxide, leading to a chain of reactions called photorespiration. This process consumes energy and oxygen while releasing carbon dioxide and ammonia, making RuBisCO an inefficient catalyst: in the presence of oxygen, RuBisCO inevitably spends a minimum of 25% of the time reacting with oxygen. In addition, RuBisCO is a sluggish catalyst, catalyzing its reactions at a rate that is ca. 1 million times lower than other enzymes such as catalase (Erb & Zarzycki, 2018). To overcome these inefficiencies and to grow at appropriate rates, plants must accumulate large amounts of RuBisCO in their leaves, making it the most abundant protein in nature (Spreitzer & Salvucci, 2002). In a plant leaf, there exist more than 10,000 diverse proteins, out of which RuBisCO constitutes a significant portion, ranging from 30 to 65% of the total protein content. It is also one of the most ubiquitous proteins in nature, being found in the photosynthetic tissues of all "green" plants, algae, cyanobacteria, a variety of photosynthetic bacteria and some non-photosynthetic bacteria as well (i.e., chemolitotrophic bacteria; Raven, 2013). It has been estimated that for every person on earth there is 5 kg of RubisCO, most of which synthesized yearly (Erb & Zarzycki, 2018). Altogether, this makes RuBisCO the most abundant protein in the global carbon cycle that literally feeds life on earth.

RuBisCO has a molecular weight of ca. 550 kDa, consisting of eight large subunits (LSU – "Large SubUnit") of approximately 51 to 58 kDa each and eight small subunits (SSU – "Small SubUnit") of molecular weight between 12 and 18 kDa (Jensen & Bahr, 1977), as described in Fig. 1.



Fig. 1. RubisCO from photosynthetic organisms consists of a set of eight large subunits (L or LSU) of 51 to 58 kDa each, and eight small subunits (S or SSU) of 12 to 18 kDa each, forming an L_8S_8 quaternary structure. A: SDS-PAGE separation of RuBisCo subunits from spinach leaves. The estimated molecular masses of standards are indicated on the left and are expressed in kDa. B: Schematic structure of RubisCO showing the spatial arrangement of the 16 subunits; left: side view; right: top view.

Although the amino acid sequence of the large subunit remains relatively consistent among higher plant species, exhibiting over 80% homology, the small subunit's amino acid sequence displays significant differences, with around 70% homology (Udenigwe et al., 2017). Consequently, various crops may exhibit slight differences in the amino acid profiles of RuBisCO and, in turn, experience differences in certain biochemical characteristics, including isoelectric point, molecular weight, bioactivities, and others. The large subunit is the product of a single gene, which is encoded in the chloroplast genome (and therefore inherited from the female progenitor), whereas the small subunit is the product of a gene family and is encoded in the nucleus (meaning that it is inherited from both progenitors). These characteristics significantly impact both the amino acid composition and overall yield of extracted RuBisCO from plant leaves. Therefore, when aiming to extract this protein on a large scale for food purposes, they must be taken into consideration.

RuBisCO is present in the soluble fraction of chloroplasts and constitutes over 50% of the foliar protein in C3 plants (Kawashima & Wildman, 1970). In C4 plants, however, RuBisCO accounts for a lower percentage, ranging from 8% to 23% (Ku et al., 1979). The effective concentration of RuBisCO in foliage depends on various factors, including the availability of nitrogen, the plant species, its age, and the intensity of light (Barbeau & Kinsella, 1988).

3. Advantages of RuBisCO as a protein source for human consumption

3.1. Animal proteins vs RuBisCO

The most prominent animal protein sources that are currently in use by the food industry are creatine, milk (caseins), whey and egg (Table 1). Creatine, a non-protein amino acid, has several advantages in terms of muscle metabolism and sports performance (Balsom, Söderlund, and Ekblom, 1994). According to Greenhaff et al. (1993), supplementing with creatine increases the energy reserve in the form of adenosine triphosphate (ATP). However, it is worth noting that creatine is also associated with weight gain and several side effects, including muscle cramps and gastrointestinal disorders (Stone et al., 1999; Schilling et al., 2000). Caseins are milk proteins that are often used as a positive control model in food allergy studies. Additionally, the two most potent milk allergens, α-lactalbumin and β-lactoglobulin, are present in cheese whey, a by-product of cheese production that is rich in proteins, minerals, and lactose. Cheese whey can be used directly or after fermentation, in the preparation of many products, incorporated as ingredient or additive, and as a sports supplement. The presence of lactose is associated with intolerances, whereas α-lactalbumin and β-lactoglobulin, also present in cheese whey, are associated with allergies. Whey protein concentrates (WPC) find application in a wide range of food products such as confectionery items, nutrition bars, cereals, processed cheeses, baked goods, sports beverages, and muscle gain formulations. Egg protein is considered as a model protein because it is quite complete in essential amino acids and displays several useful functional properties. Despite being a good source of proteins, it has a high allergenic potential, having an incidence in children from 1.6% to 2.6% (Croegaert et al., 2013). The egg contains 24 major proteins, of which ovomucoid, ovalbumin, conalbumin and lysozyme are the ones with the highest allergenic potential (Tosca et al., 2014).

Apart from its plant origin, RuBisCO has two significant advantages, as presented in Table 1: a) it is a complete protein with a balanced proportion of all the essential amino acids, and b) it is a non-allergenic protein that is commonly used as a negative control in allergenic assays (Astwood et al., 1996; Thomas et al., 2004; Jeurink & Savelkoul, 2005; Hoff et al., 2007; Jongh et al., 2013; Cirkovic Veličković & Gavrovic-Jankulović, 2014). These nutritional features make RuBisCO a viable substitute for animal proteins commercially available in the current market.

3.2. Plant-based proteins vs RuBisCO

People's awareness about healthy foods continues to increase, with the recognition that many serious human diseases (e.g., cancer and neurodegenerative disorders) may be delayed, minimized or prevented by the selection of appropriate diets. Furthermore, the current demand to reduce the environmental impact of agrifood systems and the improvement of food production capacity are problems that need to be solved (Godfray et al., 2010). Plant-based proteins could be a promising alternative. However, the plant-based proteins currently available in the market are insufficient to solve the aforementioned problems. Most plant-based proteins are allergenic, have unbalanced compositions of essential amino acids, and/or possess unpleasant tastes. The large-scale extraction and purification of RuBisCO, which is a non-allergenic, complete and readily digestible protein found in very high abundance in all photosynthetic tissues, could potentially address several global issues, such as protein malnutrition, which is a widely recognized public health concern (Müller & Krawinkel, 2005; Stefano et al., 2018).

Given that RuBisCO appears to address all the constraints mentioned earlier and possesses the ideal characteristics required for a protein intended for both animal and human consumption, a critical question

Table 1
Advantages and disadvantages of animal proteins commercially available vs RuBisCO.

Animal Proteins	Commercially available	Advantages	Disadvantages
	Creatine	Improves muscle metabolism and sports performance	A non-protein amino acid that may cause: - Weight gain - Muscle cramps - Gastrointestinal disorders
	Caseins	Balanced amino acid profiles containing all the essential amino acids in their composition	Allergenic proteins. They are present in milk and are associated with lactose intolerances
	Cheese whey proteins	Balanced amino acid profiles containing all the essential amino acids in their composition	Allergenic proteins. They contain lactose in their composition, which is associated with lactose intolerances
	Whey protein concentrates (WPC)	Balanced amino acid profiles containing all the essential amino acids in their composition	Allergenic proteins
	Albumin	Balanced amino acid profile containing all the essential amino acids in its composition	Allergenic protein
RuBisCO		Balanced amino acid profile containing all the essential amino acids in its composition. It is a non-allergenic protein. It is a plant-derived protein, present in all photosynthetic tissues, where it occurs in very high abundance. Good digestibility	None

arises: Why has it not yet become commercially available in the market?

3.2.1. RuBisCO nutritional properties

RuBisCO appears to have the most potential among all proteins for commercial applications and fits the "ideal" protein profile for both human and animal nutrition. The composition of RuBisCO is well-balanced in essential and non-essential amino acids, being the closest (when compared to soy protein and whey protein) to the egg protein, but it is not over-rich in essential amino acids (Table 2) – It is considered as the model protein in nutritional terms. Lysine, the sulphur-containing amino acids, and tryptophan, regarded as critical in low-meat diets, are present at high concentrations in RuBisCO (Pouvreau et al., 2013). In comparison to other plant proteins, spinach RuBisCO exhibits higher quality according to the protein quality model established by FAO/WHO in 2007, with a higher proportion of essential amino acids. The quality of a protein is measured using the chemical index, which is the ratio of the amount of an essential amino acid in a food protein to the amount of the same amino acid in a reference protein. The reference protein used by FAO/WHO (2007) is the pattern of essential amino acids in egg protein. A higher chemical index indicates a better protein quality. The chemical index is defined by:

Chemical index = (mg of limiting amino acid per g of the protein under analysis)/(mg of amino acid per g of the reference protein).

It is common for plant-based proteins available in the market, mostly derived from pulses and cereals, to be deficient in some essential amino acids. When compared to animal-based proteins, plant-based proteins present a lower percentage of essential amino acid content (Gorissen et al., 2018). All cereal seeds, like rice and wheat contain much less protein than animal protein sources or legume seeds and are often incomplete, mainly in lysine (an essential amino acid; McGuire & Beerman, 2013), a problem that demands genetically modified plants and consequently discussions about transgenic cultures (Alink et al., 2008; Nguyen et al., 2012). The supplementation of poultry feedstock with lysine is a well-known example of addressing amino acid deficiencies in animal feed. On the other hand, legume seeds such as soy, pea, and lupin are deficient in the sulfur-containing amino acids, namely cysteine and methionine, while both legume and cereal seeds generally lack sufficient tryptophan (McGuire & Beerman, 2013).

Table 2
Comparison between the content in essential amino acids of RuBisCO and those of other proteins with the chemical index and nutritional recommendation of FAO/WHO ().

Essential amino acids	'Ideal' protein according to FAO/WHO (mg.g^{-1} protein)	Egg protein (mg.g^{-1} protein)	Cheese whey protein (mg.g^{-1} protein)	Soy protein (mg.g^{-1} protein)	Spinach RuBisCO (mg.g^{-1} protein)
Histidine	15	22	13	26	26
Isoleucine	30	54	50	49	43
Leucine	59	86	94	82	82
Lysine	45	70	68	63	71
Methionine + Cysteine	22	57	34	26	40
Phenylalanine + Tyrosine	38	93	128	90	96
Threonine	23	47	68	67	52
Tryptophan	6	17	27	14	17
Valine	39	66	67	50	60
Chemical index [a]		100	87	100	100
Chemical index [b]		100	60	46	70

[a] Calculated in comparison to the 'ideal' protein as recommended by FAO/WHO (2007).
[b] Calculated in comparison to the egg protein.
adapted from Pouvreau et al., 2013

3.3. Allergenicity and digestibility

Another major advantage of RuBisCO as a food protein source of great potential is that it is considered a non-allergenic protein (Jeurink & Savelkoul, 2005; Cirkovic Veličković and Gavrovic-Jankulović, 2014). Surprisingly, RuBisCO has even been utilized as a negative control in several allergy trials and tests (Astwood et al., 1996; Thomas et al., 2004; Hoff et al., 2007; Jongh et al., 2013). As far as we are aware, there is a single case study in which a 23-year-old woman is reported to have developed an allergic reaction after ingestion of spinach leaves. Based on proteomic analysis, the authors stated that RuBisCO was responsible for the sensation of asphyxia and angioedema, both in the lips and in the tongue (Foti et al., 2012).

It is important to note that all proteins commercially available for human and animal consumption are integral members of the lists of major allergens, both in Europe and the US. In soy protein alone, 15 allergenic proteins were found, which implies a very significant nutritional restriction on the use of this well-known protein source by people suffering from allergies (Ogawa et al., 1991). In addition, most of the soy produced worldwide is transgenic, an issue that has been the subject of many scientific and ethical discussions about its environmental, economic and social effects on present and future societies (Damato, 2009). Also, soy protein contains a high concentration of isoflavones that exhibit estrogenic bioactivity. So, soy consumption in large amounts is harmful, especially to men and children (Setchell et al., 2002). Erectile dysfunction and male infertility are some of the side effects that may arise as a result of excessive soy consumption (Mollakhalili-Meybodi et al., 2021). The Kunitz and Bowman-Birk inhibitors present in soybeans can interfere with the digestion of protein and the absorption of nutrients. However, heat treatment, such as cooking or roasting, can denature these inhibitors and improve the digestibility of soy protein.

As for the other pulses, lupin for instance is included in the allergen lists in the EU and Australia (Jappe & Vieths, 2010) and according to Food and Drug Administration (FDA), people who suffer from peanut allergies are more susceptible to suffer from lupin allergies as well, and the reactions could be severe and life-threatening. Rice and peas are generally considered to be hypoallergenic foods, which means that they are less likely to cause allergic reactions than other foods. Ca. 80% of patients suffering from food allergies have immunoglobulins E (IgE) to rice protein (Goliáš et al., 2013). Some rice proteins, including α-globulin (26 kDa), glyoxalase (33 kDa), and a glycoprotein (56 kDa), have been identified as allergenic, with the α-amylase/trypsin inhibitors family (with molecular weights between 14 and 16 kDa) having the highest incidence of allergenicity (Krishnan & Chen, 2013). Even though it is not common, cases of people who have allergy to peanut reacting to pea protein have been described in the literature (Wensing et al., 2003; Lavine & Ben-Shoshan, 2019).

Wheat grains, which contain over 100 main proteins, have been identified as allergenic, largely due to the presence of gluten (Tatham & Shewry, 2008). Gluten intolerance is a well-known issue associated with wheat consumption (Pasha et al., 2016). Celiac disease is another important consideration, which consists of chronic inflammation that affects the gastrointestinal tract, resulting in an atrophy of intestinal villi and consequent malabsorption of nutrients (Gasbarrini & Mangiola, 2014). On this topic, several studies show that the novel gluten-free foods, which have been arising in the market, often contain a low percentage of protein and a higher percentage of carbohydrates and lipids compared to gluten-equivalent products (Berti et al., 2004; Caponio et al., 2008; Segura & Rosell, 2011).

In terms of digestibility, RuBisCO is easily and readily digested by pepsin (digestive protease present in the gastric juice responsible for the hydrolysis of dietary proteins into peptides; Thomas et al., 2004; Jong & Nieuwland, 2011). RuBisCO has been used as a positive control in digestibility assays due to its high susceptibility to proteolytic attack (pronase, trypsin, proteinase K, α-chymotrypsin or subtilisin; see for example Fig. 3 in reference Monteiro et al., 2015). This is not the case for

many plant proteins, especially those that are allergenic and may not be fully hydrolyzed by digestive enzymes before reaching the intestines, reducing their digestibility. Soy protein, for example, is deficient in methionine, which can further lower its digestibility (Friedman & Brandon, 2001). Protein digestibility can also decrease up to 50% due to the high concentration of trypsin inhibitors (which are proteins themselves) present in this legume seed (Friedman & Brandon, 2001; Gilani et al., 2012). Wheat protein extraction involves an alkaline treatment, decreasing its digestibility (up to 40%) and causing a drastic reduction (up to 100%) in protein quality. This extraction has a low yield in protein production, meaning that it takes a large amount of this cereal to obtain a small amount of protein (Gilani et al., 2012). Pea, lupin, and rice proteins are easily digested (House et al. 2010; Zuber et al. 2018). However, a study showed that animal protein digestion is quicker than rice protein isolate (Jäger et al., 2013). The process of cooking was also demonstrated to decrease significantly the gastric and gastrointestinal protein digestibility of rice (Liu et al., 2019).

Among the hundreds of millions of proteins known to exist, only a few % (which still comprise a considerable number) have such a well-balanced amino acid composition as RuBisCO. If we now consider other protein properties, such as complete absence of toxicity, including allergenicity (something that cannot be said for the egg protein, soybean protein and whey protein), high digestibility and good functional properties, very few proteins (if any) besides RuBisCO will remain. In addition, for commercial viability, the protein must occur naturally in high concentrations and be suitable for extraction and purification at an industrial scale. Therefore, it is apparent that RuBisCO stands out among all the proteins in nature in terms of its potential to be used as a protein source for human and animal consumption.

The discussion and information provided in Sections 4 and 5 reinforce the question raised in the last paragraph of Section 3.2: despite its exceptional properties, why hasn't RuBisCO become commercially available yet?

4. Functional and organoleptic properties of RuBisCO

Proteins are incorporated in food products for different reasons, with the most significant being their nutritional value and functional properties. These properties are closely related to solubility, water and fat retention, emulsion, gelatinization, foaming capacity, flavor, and durability. As mentioned earlier, the plant-based food industry faces a significant challenge in creating products with desirable appearance, texture, flavor, mouthfeel, nutrition, and functionality. Therefore, the ability of plant proteins to thicken, gel, emulsify, foam, and retain fluids are some of the most important features for their use in plant-based foods (McClements & Grossmann, 2021). In this respect, RuBisCO presents several physical and chemical properties, as described below and in Table 3.

4.1. Solubility

Protein solubility is one the most important factors because it dictates the other protein functionalities. RuBisCO's high solubility is defined by multiple factors, such as pH, ionic strength, protein concentration, temperature, the presence of divalent cations and the species from which the protein was purified (Sheen, 1991). When treated with magnesium, RuBisCO extracted from tobacco leaves tends to precipitate at pH values lower than 7.5, suggesting the influence of divalent cations and pH value (Prevot-D'Alvise *et al.*, 2004). However, when the same conditions were applied to RuBisCO purified from spinach leaves, it did not exhibit the same behavior, indicating the influence of the species of origin. RuBisCO from alfalfa leaves was soluble when the pH was closer to its isoelectric point (pH 5.0 to 6.5), but almost insoluble under acidic solutions (only 20% w/v soluble). When the solution was alkaline (pH 7 to 12) the solubility increased from 10% to 90% (Prévot-D'Alvise et al., 2004). According to Van del Velde (2011), RuBisCO purified from

Table 3
Advantages and disadvantages of plant-based proteins commercially available vs RuBisCO.

Plant-based proteins	Commercially available	Advantages	Disadvantages
	Rice	Considered an hypoallergenic food Easily digested High water/fat retention Good gelling capacity	Low percentage of protein Incomplete protein, mainly in lysine Low solubility Poor emulsifying capacity Low foaming capacity Off-flavors
	Wheat	Great foaming capacity No flavor	Low percentage of protein Incomplete protein, mainly in lysine and tryptophan Allergenic protein Low protein digestibility Low solubility Low water/fat retention Poor emulsifying capacity
	Soy	High percentage of protein High water/fat retention Good gelling capacity	Incomplete protein, mainly in cysteine, methionine and tryptophan Allergenic protein Low protein digestibility Low solubility Poor emulsifying capacity Low foaming capacity Off-flavors
	Pea	Considered an hypoallergenic food Easily digested	Incomplete protein, mainly in cysteine, methionine and tryptophan Low solubility Low water/fat retention Poor emulsifying capacity Low gelling capacity Low foaming capacity Off-flavors
	Lupin	High percentage of protein Easily digested High water/fat retention Good foaming capacity	Incomplete protein, mainly in cysteine, methionine and tryptophan Allergenic protein Low solubility Poor emulsifying capacity Low gelling capacity Off-flavors
RuBisCO		Well-balanced in essential and non-essential amino acids Non-allergenic protein Easily and readily digested by pepsin High solubility High water/fat retention Great emulsifying capacity Good gelling capacity Great foaming capacity No flavor	None

spinach leaves is highly soluble at pH values commonly found in food, ranging from pH 6 to 8. Concerning other proteins commonly used for human consumption, the protein extracted from different pea seed genotypes revealed to be almost insoluble at pH 5.0. However, its solubility increased below and above this pH value and the proteins exhibited excellent solubility at pH 7 and 8 (Barac et al., 2010). Based on these findings, it can be inferred that pea proteins are suitable for use in food products with a neutral or basic pH. Despite being a commonly used plant-based protein, soy protein is known to have a significant drawback of low solubility (Malhotra & Coupland, 2004). Only 10% of soy proteins are soluble at pH values close to their isoelectric points (4.6; Freitas et al., 2017). Lupin protein, being a legume, shares similarities with soy protein. Meanwhile, rice protein, despite being considered a hypoallergenic alternative, has low solubility at acidic environments, with a solubility of less than 2% (w/v), similar to soy protein, which can affect its other functional properties as well (Wang et al., 2018).

4.2. Water and fat retention capacities

The water binding capacity of a protein can greatly influence its potential for use in food applications, as it ultimately affects the texture of the final product. A protein with low water binding capacity can result in a dry product or may lose water during storage. On the other hand, fat retention is linked to flavor in food products. RuBisCO's water and fat retention capacities have been tested in a few works. Results showed that RuBisCO had a capacity of water retention of 428% by volume, while soy protein, which was already partially denatured, had a capacity of 612% by volume (Douillard & Mathan, 1994). Regarding the fat retention capacity, RuBisCO from alfalfa showed a 208% increase in volume whereas soy seed protein showed an increase of 159% (Douillard & Mathan, 1994). Pea protein water/fat retention is low. Recently, in an attempt to increment pea protein water/fat retention, an assay was performed where pea protein was modified via sequential deaminidation and conjugation to enhance these functionalities (Shen et al., 2022). In the case of rice proteins, the water retention capacity ranged from 387% (w/w) to 560% (w/w), and the fat retention from 374% (w/w) to 918% (w/w), values that are comparable to the well-known casein (Chandi & Sogi, 2007). In what concerns total seed preparations, Raikos and colleagues (2014) demonstrated that lupin flour had a great water holding capacity due to its high protein content, and the results were identical to other studies reported (Khalid & Elharadallou, 2013). On the other hand, wheat flour did not show good water holding retention capacity due to its low protein content (Raikos et al., 2014).

4.3. Emulsifying capacity

Emulsion is the functional property which dictates the protein ability to emulsify oil (Kaushal et al., 2012). RuBisCO from spinach leaves shows a great emulsifying capacity after pH precipitation followed by lyophilization. However, heat coagulation has a negative effect on this property (Rault et al., 1993). RuBisCO showed emulsion stability comparable to the popular egg white protein (Lamsal et al., 2007). Rice proteins have a poor emulsifying capacity due to their low water solubility and high molecular weight (Agboola et al., 2005). As mentioned above (in the solubility section), pea protein is almost insoluble at pH 5.0, which poorly influences its emulsifying capacity at this pH value. However, the emulsion ability increases significantly below and above this pH value (Barac et al., 2010). The emulsifying ability of wheat protein is also low, which negatively affects it's used in the food industry, so to improve this function a conjugation of gluten with fructose via Maillard reaction was performed (Song et al., 2018). The emulsifying capacity of soy protein is limited at pH 5.0 (pH value where protein solubility is lower), something which compromises its utilization in food matrixes (Deng, 2021). Soy protein emulsion capacity can be improved by modifying certain physico-chemical properties (Tang, 2017). Lupin protein has a similar behavior as soy and pea proteins. To improve this function, high pressure treatments have been applied to lupin protein (Chapleau & Lamballerie-Anton, 2003).

4.4. Gelation

Protein denaturation followed by network formation is the process that triggers gelation, which is a functional property that can have a significant impact on the texture of food products (Abaee et al., 2017). The gel formation capacity of RuBisCO has also been examined, with the most favorable outcomes observed for alfalfa-derived RuBisCO after being subjected to heat treatment at 80 °C and pH 12 for 30 min (Barbeau & Kinsella, 1988). The gel produced by RuBisCO derived from spinach and tobacco leaves has also demonstrated favorable outcomes, and similarly to its emulsifying capacity, these gels exhibit greater resistance when compared with those formed by soy protein (Van del Velde et al., 2011). Soy seed protein has been found to own a strong gelling capacity, with as little as 10% (w/w) of soy protein showing excellent rheological properties (Batista et al., 2005). Pea protein gelling properties showed some limitations and were weaker and less elastic when compared with soy protein (Shand et al., 2007). Another constraint of pea protein is its high gelation temperature, which requires temperatures of approximately 95 °C to induce gelation. As typical cooking temperatures reach only around 75 °C, this makes it challenging to utilize pea protein as a gelling agent (Health and Human Services, 2017). Lupin protein exhibits a behavior similar to pea protein: the gels formed with lupin protein were weaker and more deformable when compared to the ones made with soy protein, resulting in a poor gelling ingredient for the food industry (Berghout et al., 2015). Agboola et al. (2005) demonstrated that isolated Australian rice proteins had good gelling properties (See Table 4).

4.5. Foaming capacity

Proteins exhibit great foaming capacities because they contain both hydrophilic and hydrophobic moieties. When comparing the foaming capacity of RuBisCO to the well-known protein found in egg whites, it was observed that both proteins produced similar volumes of foam. However, the foam created by RuBisCO was 33% more stable (Douillard & Mathan, 1994). Recent trials have demonstrated that RuBisCO exhibits a more desirable foaming capacity than soy and whey proteins at pH values of 4.5 and 7.0 (Van del Velde et al., 2011). Foaming capacity of soy protein isolates in their native form was low, which can be explained by their tertiary and quaternary structures (Martínez et al., 2009; Martínez et al., 2011). Lupin protein's noteworthy foaming ability in acidic solutions sets it apart from soy protein. This characteristic makes lupin protein more appropriate for use in food products such as yoghurts or salad dressings (Jayasena et al., 2010). Rice proteins foaming capacity was improved when pressure was applied to these proteins (Zhu et al., 2016). At pH 5.0, pea protein had the lowest foaming capacity, but this property increased significantly when the pH was increased to 7.0 and 8.0 (Barac et al., 2010). Gluten (wheat storage proteins) is composed by glutenin and gliadin, and the last mentioned is responsible for the foaming properties of gluten (Keller et al., 1997; Banc et al., 2007). This protein has a great foaming capacity at neutral and alkaline pH values, but low foaming capacity at pH 2, which can be improved by adding NaCl (Thewissen, Celus, Brijs, and Delcour, 2011).

4.6. Flavor

A significant obstacle in substituting animal proteins for plant-based proteins is the occurrence of undesirable flavors, known as off-flavors. Aldehydes, ketones, and alcohols are the main compounds responsible for the "green" and "grassy" flavor in pea proteins (Heng, 2005). Off-flavors can arise during the harvesting, storage, and subsequent processing of plant-based proteins due to the oxidation and enzymatic degradation of unsaturated fatty acids (Azarnia et al., 2011; Roland

Table 4
Current industrial methods for RuBisCO isolation as a food additive and corresponding advantages and disadvantages.

		Plant source	Method	Advantages	Disadvantages
Company	France Lucerne	Alfalfa	Not disclosed	Capacity to purify large amounts of RuBisCO at a low price	Low degree of purification. Potential allergenic character
	TNO - Innovation For Life	Sugar beet leaves	The process involves pressing, centrifuging and ultrafiltration	Capacity to produce 10 kg protein/h	Low degree of purification. Potential allergenic character
	NIZO	Spinach	The process involves protein extraction, with decantation and pressing, followed by purification by aggregation, precipitation and affinity separation, with a final concentration step by filtration and evaporation	The final product contains 90% RuBisCO. Low % of chlorophylls in the final product	Complex procedure, not suitable to the food industry
	Green Project	*Lemna* spp. (aquatic flowering plants)	Not disclosed	Capacity to purify large amounts of RuBisCO at a low price	The final product has green colour
Project	GreenProtein	Sugar beet leaves	The process involves heat precipitation, decantation, micro-filtration and ultra-filtration, with a final concentration step by spray-drying	The final product is a white powder. Capacity to purify large amounts of RuBisCO at a low price	The use of organic solvents to remove pigments. Low degree of purification. Potential allergenic character

et al., 2017; Lan et al., 2019). Lupin protein, as the majority of legume seed proteins, has a unique flavor profile that has been described as moldy, beetroot-like, grassy, metallic, cooked potato-like and pea-like (Schlegel et al., 2019). The taste perceptions pattern was identified as bitter, salty and astringent (Schlegel et al., 2019). Despite being one of the most commonly utilized plant-based proteins for human consumption, soy protein has significant flavor-related limitations. The presence of isoflavones, saponins, phenolic acids, etc. is responsible for soy protein bitter and astringent tastes (Damodaran & Arora, 2013). Also, residual amounts of phospholipids, which contain polyunsaturated fatty acids (PUFAs), can suffer oxidation and create volatile compounds that are accountable for the beany, grassy, or green odors (Damodaran & Arora, 2013). Rice bran contains a high quantity of lipids, which can trigger the activity of lipases and ultimately lead to the development of a rancid flavor in rice (Champagne, 2004). Degermed wheat and corn flours (endosperm products) have little or no flavor (Rackis et al., 1979). In contrast, RuBisCO has already been highly purified as a tasteless, odorless and colorless powder (Van del Velde et al., 2011).

5. Bioactivities of RuBisCO

Aside from all the advantages mentioned above, RuBisCO has been acknowledged for its functional properties related to human health, especially those that arise from the peptides produced during its proteolytic breakdown, making it a potentially valuable food additive (Udenigwe et al., 2017). Indeed, some authors have postulated that the ability of RuBisCO to produce bioactive peptides upon digestion is one of the features that makes this protein a suitable source for food applications, particularly to produce functional foods (Udenigwe et al., 2017) with important health benefits. Arguably, bioactive peptides have been defined as specific protein fragments that have beneficial influences on the body function and on health in general (Kitts and Weiler, 2003). A comprehensive review of the bioactive peptides generated from RuBisCO has been documented elsewhere (Udenigwe et al., 2017). The list encompasses a wide array of activities, including antihypertensive properties, opioid and anxiolytic activities, antibacterial and antioxidant activities, among others. For instance, a specific bioactive peptide derived from RuBisCO has demonstrated antihypertensive properties by regulating blood pressure (Kapel et al., 2006; Udenigwe & Aluko, 2012). This particular peptide has been shown to inhibit angiotensin I-converting enzyme (ACE) under *in vitro* conditions. ACE plays a vital role in the renin-angiotensin system (RAS), which regulates blood pressure. Additionally, this bioactive peptide has been found to decrease blood pressure in hypertensive animals (Kapel et al., 2006). Systolic blood pressure of hypertensive rats was reduced after oral administration of RuBisCO's hydrolysates (500 mg/kg body weight; Kapel et al., 2006). The dipeptide responsible for this bioactivity was identified as *N*-valyl-tryptophan (VW) and found to be located within RuBisCO's large subunit at different positions (f67-68, f382-383 and f459-460), as well as in the small subunit (at f60-61; Kapel et al., 2006). The bioactive peptides produced from RuBisCO have also been found to possess opioid functionality (Teschemacher, 2003; Perlikowska & Janecka, 2017). Rubiscolin-6, a peptide derived from RuBisCO extracted from spinach leaves, has demonstrated anxiolytic activity (Yang et al., 2001; Hirata et al., 2007). Another interesting effect of rubiscolin-6, detected in mice, was the ability to boost memory consolidation (Yang et al., 2003). Additional studies have indicated that selected RuBisCO-derived peptides possess antimicrobial properties. The hydrolysis of RuBisCO's small subunit has been found to produce bacteriostatic activity against pathogenic Gram-negative bacteria such as *Salmonella arizonae* and *Shigella sonnei* (Trovaslet et al., 2007). Bioactive peptides obtained after hydrolysis of RuBisCO extracted from alfalfa green juice also exhibited antimicrobial activity against Gram negative (*Escherichia coli*) and Gram positive (*Micrococcus luteus*, *Listeria innocus* and *Bacillus subtilis*) bacteria (Kobbi et al., 2015). RuBisCO's derived peptides have also been studied for their potential antioxidative capacity. An *in vitro* assay showed a potent antioxidant activity of RuBisCO's bioactive peptides (Kobbi et al., 2017). The activity was dose-dependent and the mechanism behind it was the inhibition of linoleic acid oxidation, which led to the reduction of ferric ion and scavenging of a radical cation (Kobbi et al., 2017). *Solieria chordalis,* a red seaweed, has been identified as a source of antioxidative peptides in previous research (Bondu et al., 2015). More recently, it has been reported that 60% of the identified antioxidative peptides in *Solieria chordalis* were derived from RuBisCO (Bondu et al., 2015).

6. Technologies for RuBisCO extraction and food applications

The available literature describes many methods for obtaining highly pure RuBisCO, but these methods are either economically unfeasible for industrial-scale production or involve the use of reagents that are harmful to human and animal health.

Despite its natural abundance and ubiquity among photosynthetic cells and organisms, its potential use in food products has been limited

due to the difficulty of efficiently purifying it on a large scale in a cost-effective form that is safe for human consumption, similar to other typical proteins. In the last decade however, several patented methods and projects have yielded several procedures, more or less expensive, to produce industrial amounts of RuBisCO, either as a white, tasteless powder or as a green protein additive (Table 3). The first company that produced RuBisCO on a large scale was France Luzerne, a French company participating in Fralupro, a European project whose objective was to discover a possible use for the leaves from 32 million ha of alfalfa each year. In 1998, this company started to produce 1,200 tons of alfalfa RuBisCO per year, at the same cost of producing soy protein. However, the project failed to achieve a significantly higher degree of RuBisCO purification beyond that already present in the total soluble protein extract. It is worth recalling that RuBisCO is already ca. 50% pure (relative to the total leaf protein) in the leaves of C3 plants. This extract contains thousands of different proteins, many potentially allergenic and/or otherwise undesirable, in addition to tannins and other phenolic compounds, which can negatively contribute to the nutritional quality of the protein concentrate.

TNO – Innovation For Life is a Dutch company that also produces RuBisCO from sugar beet leaves on a large scale. The process involves pressing, centrifugation and ultrafiltration, and has a production capacity of 10 kg protein/h, with a low degree of purification, around 35 to 40%. Therefore, RuBisCO is not obtained in a purified state. Another Dutch company, NIZO, filed a patent application for its RuBisCO purification process in 2010. The end-product resulting from the purification process described in the patent application by NIZO contains at least 90% of the leaf RuBisCO in a soluble form and less than 0.1% (w/v) chlorophylls. However, the process is complex and has not yet been able to be scaled-up to an industrial level. RuBisCO Foods, another Dutch company that produces protein gels and powders for the purpose of their application in food and feed, has developed and patented a unique technology that allows the extraction and purification of soluble proteins from *Lemna* spp. (aquatic flowering plants), including RuBisCO. However, the protein extracts still contain chlorophylls in their composition.

GreenProteins is a European project that presents, as its main objective, the production of high-quality proteins for food applications. Proteins are extracted from sub-products produced by the food industry as a way to promote sustainability. This project, which began in 2016 and ended in 2021, was based on a new methodology used for purifying RuBisCO from sugar beet leaves, as described in patent US20150335043A1. Although the final product achieved is a white powder, which indicates complete removal of chlorophylls, the method described uses organic solvents for the removal of the pigments, which is neither compatible with healthy food nor feasible for further applications in the food industry. It is also important to note that although the disclosure to the public states that RuBisCO is purified, the end-product actually consists of the total extract of soluble proteins found in sugar beet leaves.

The large number of patents and industry-level projects focused on RuBisCO extraction and purification indicates a significant level of interest and demand for this protein. Nonetheless, whilst there are several methods described in the industry, most of these are based on fractionation that yield highly impure RuBisCO, whereas more complex methods typically hinder RuBisCO production at a reasonable price. Therefore, some do not produce the protein in a purified form and those that do are not cost-effective. Some of these methods produce RuBisCO in a form that is insoluble and/or green. These observations may well explain the absence of commercially available RuBisCO in the food markets. Nevertheless, there has been a rising number in novel patents submitted with new methods that claim to purify RuBisCO, which are scalable and suitable for the food industry. Hopefully, with the emerging novel methods, RuBisCO may fulfill its potential as the future of commercialized plant-based proteins.

As a food additive, RuBisCO has already been tested in several of the above-mentioned projects, mostly as a protein additive in foods, using several food matrixes. Noteworthy, the isolated RuBisCO produced by GreenProteins has been successfully used as a protein additive in wheat dough, with prominent published results. Ducrocq, Boire, Anton, Micard, & Morel (2020) supplemented wheat doughs with RuBisCO, gluten and pea protein and the results were quite promising. Whilst gluten and pea protein decreased dough stiffening during heat treatment, RuBisCO showed the ability to bind (weakly and covalently) to the dough and remarkably, the dough treated with RuBisCO had a higher concentration of large polymers after being exposed to heat, hence enhancing its functional properties (Ducrocq et al., 2020).

7. Advantages to the food industry

Overall, RuBisCO seems to present all the advantages to render it the perfect plant-based protein for today's market, including its sustainability, nutritional composition, digestibility, organoleptic and physical properties and even bioactivities for human health, in the absence of allergenicity. Fig. 2 summarizes all the major advantages of RuBisCO as a food additive.

Whilst the production of other proteins available in the market (especially those of animal origin) have typically negative environmental consequences, RuBisCO is a very good candidate from a circular economy and sustainability point of view. Due to its capacity to react with O_2 and catalytic inefficiency, this protein is present in very large amounts in all photosynthetic cells and green tissues, thus allowing its isolation from green food waste produced by the industry. Although its isolation has been mostly reported from alfalfa green juice and sugar beet leaves (Tenorio et al., 2016; Kobbi et al., 2017, respectively), RuBisCO purification can be achieved from a wide variety of crops, a good proportion of which generate common industrial sub-products, like carrot or tomato leaves, hence becoming a very sustainable source of plant-based protein. Other photosynthetic options exist as potential sources of RuBisCO, including macro and microalgae. These algae can also serve as an excellent protein source due to their fast growth and ability to produce high masses in short periods of time.

8. Conclusion

Overall, RuBisCO possesses unique characteristics, a favorable nutritional profile, beneficial bioactivities, and desirable physical attributes, making it a promising candidate for use in food applications. Based on all its well-studied advantages, it seems clear that RuBisCO may be regarded as the ideal protein for both human and animal consumption. The potential applications of this protein for the food industry are vast. To date, what is missing is a good and reliable procedure for RuBisCO purification, i.e., cost-effective, suitable to be scaled-up to an industrial level and relatively easy to perform. Novel methods to purify RuBisCO with all the characteristics described above have been arising and could help to place this protein on a higher position in the plant-based food market. Given the current trends in the plant-based market and RuBisCO's unique features, ubiquity and quantity, it seems plausible that new and improved methods for its large-scale purification will continue to emerge in the coming years. These developments will likely be accompanied by further research uncovering new bioactivities and functions associated with RuBisCO.

Most probably, in the near future, RuBisCO will become a common protein additive used to enhance the nutritional value of various food products such as protein bars, bread, beverages, yogurts, and puddings. Furthermore, since RuBisCO is also known for its bioactive peptides, it could not only enhance the nutritional value of different food products but also add health-promoting value and functional food features to the final product.



Fig. 2. Advantages of RuBisCO as a food additive.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

No data was used for the research described in the article.

References

Abaee, A., Mohammadian, M., & Jafari, S. M. (2017). Whey and soy protein-based hydrogels and nano hydrogels as bioactive delivery systems. *Trends in Food Science & Technology, 70*, 69–81. https://doi.org/10.1016/j.tifs.2017.10.011

Agboola, S., Ng, D., & Mills, D. (2005). Characterisation and functional properties of Australian rice protein isolates. *Journal of Cereal Science, 41*, 283–290. https://doi.org/10.1016/j.jcs.2004.10.007

Aimutis, W. R. (2022). Plant-based proteins: The good, bad, and ugly. *Annual Review of Food Science and Technology, 13*, 1–17. https://doi.org/10.1146/annurev-food-092221-041723

Alink, S., Barlow, S., Cockburn, A., & Flachowsky, G. (2008). Safety and nutritional assessment of GM plants and derived food and feed: The role of animal feeding trials. *Food and Chemical Toxicology, 46*, 2–70. https://doi.org/10.2903/j.efsa.2008.1057

Astwood, J. D., Leach, J. N., & Fuchs, R. L. (1996). Stability of food allergens to digestion *in vitro*. *Nature, 14*, 1269–1273. https://doi.org/10.1038/nbt1096-1269

Azarnia, S., Boye, J. I., Warkentin, T., Malcolmson, L., Sabik, H., & Bellido, A. S. (2011). Volatile flavour profile changes in selected field pea cultivars as affected by crop year and processing. *Food Chemistry, 124*, 326–335. https://doi.org/10.1016/j.foodchem.2010.06.041

Balsom, P. D., Söderlund, K., & Ekblom, B. (1994). Creatine in humans with special reference to creatine supplementation. *Sports, Medicine, 18*, 268–280. https://doi.org/10.2165/00007256-199418040-00005

Banc, A., Desbat, B., Renard, D., Popineau, Y., Mangavel, U., & Navailles, L. (2007). Structure and orientation changes of ω- and γ-gliadins at the air-water interface: A PM-IRRAS spectroscopy and Brewster angle microscopy study. *Langmuir, 23*, 13066–13075. https://doi.org/10.1021/la702037k

Barac, M., Cabrilo, S., Pesic, M., Stanojevic, S., Zilic, S., Macej, O., et al. (2010). Profile and functional properties of seed proteins from six pea (Pisum sativum) genotypes. *International Journal of Molecular Sciences, 11*, 4973–4990. https://doi.org/10.3390/ijms11124973

Barbeau, W. E., & Kinsella, J. E. (1988). Ribulose bisphosphate carboxylase/oxygenase (rubisco) from green leaves - potential as food protein. *Food Reviews International, 4*, 93–127. https://doi.org/10.1080/87559128809540823

Batista, A. P., Portugal, C. A. M., Sousa, I., Crespo, J. G., & Raymundo, A. (2005). Accessing gelling ability of vegetable proteins using rheological and fluorescence techniques. *International Journal of Biological Macromolecules, 36*, 135–143. https://doi.org/10.1016/j.ijbiomac.2005.04.00

Berghout, J. A. M., Boom, R. M., & Goot, A. J. (2015). Understanding the differences in gelling properties between lupin protein isolate and soy protein isolate. *Food Hydrocolloids, 43*, 465–472. https://doi.org/10.1016/j.foodhyd.2014.07.003

Berti, C., Riso, P., Monti, L. D., & Porrini, M. (2004). *In vitro* starch digestibility and *in vitro* glucose response of gluten-free foods and their gluten counterparts. *European Journal of Nutrition, 43*, 198–204. https://doi.org/10.1007/s00394-004-0459-1

Bondu, S., Bonnet, C., Gaubert, J., Deslandes, E., Turgeon, S. L., & Beaulieu, L. (2015). Bioassay-guided fractionation approach for determination of protein precursors of proteolytic bioactive metabolites from macroalgae. *Journal of Applied Phycology, 27*, 2059–2074. https://doi.org/10.1007/s10811-014-0425-0

Chandi, G. K., & Sogi, D. S. (2007). Functional properties of rice bran protein concentrates. *Journal of Food Engineering, 79*, 592–597. https://doi.org/10.1016/j.jfoodeng.2006.02.01

Chapleau, N., & Lamballerie-Anton, M. (2003). Improvement of emulsifying properties of lupin proteins by high pressure induced aggregation. *Food Hydrocolloids, 17*, 273–280. https://doi.org/10.1016/s0268-005x(02)00077-2

Caponio, F., Summo, C., Clodoveo, M. L., & Pasqualone, A. (2008). Evaluation of the nutritional quality of lipid fraction of gluten-free biscuits. *European Food Research and Technology, 223*, 135–139. https://doi.org/10.1007/s00217-007-0702-0

Cirkovic Veličković, T., & Gavrovic-Jankulović, M. (2014). Food allergens digestibility. In T. Cirkovic Veličković, & M. Gavrovic-Jankulović (Eds.), *Food Allergens: Biochemistry and Molecular Nutrition* (pp. 95–140). Springer.

Croegaert, K. A., Ithman, M. M., Spurgin, A. L., Vos, S. S., & Milavetz, G. (2013). Influenza vaccine safety in patients with egg allergy. *Journal of the American Pharmacists Association, 53*, 214–216. https://doi.org/10.1331/JAPhA.2013.13511

Damato, G. (2009). GM-Soy: Destroy the earth and humans for profit. Natural News. Retrieved from: http://www.naturalnews.com/026334_soy_research_Roundup.html. Accessed on February 22, 2022.

Damodaran, S., & Arora, A. (2013). Off-flavor precursors in soy protein isolate and novel strategies for their removal. *Annual Review of Food Science and Technology, 4*, 327–346. https://doi.org/10.1146/annurev-food-030212-182650)

Deng, L. (2021). Current progress in the utilization of soy-based emulsifiers in food applications - A review. *Foods, 10*, 1354. https://doi.org/10.3390/foods10061354

Dorsan, S., Chaudaty, T. (2013). World allergy week emphasizes need for food labeling and standardized action plans children especially affected by food sensitivity. World Allergy Organization. Retrieved from: https://www.worldallergy.org/UserFiles/file/PressReleaseWorldAllergyWeek2013Embargoedto8April.pdf. Accessed on March 28, 2022.

Douillard, R., & Mathan, O. (1994). In *Leaf protein for food use: potential of Rubisco. In New and Developing Sources of Food Proteins* (pp. 307–342). Boston, MA: Chapman & Hall. https://doi.org/10.1007/978-1-4615-2652-0_10.

Ducrocq, M., Boire, A., Anton, M., Micard, V., & Morel, M. H. (2020). Rubisco: A promising plant protein to enrich wheat-based food without impairing dough viscoelasticity and protein polymerisation. *Food Hydrocolloids, 109*, 106101. https://doi.org/10.1016/j.foodhyd.2020.106101

FAO/WHO. Protein and amino acid requirements in human nutrition. World Health Organization, WHO Technical Report Series, 935. 2007. Retrieved from: https://apps.who.int/iris/bitstream/handle/10665/43411/WHO_TRS_935_eng.pdf?sequence=1&isAllowed=y. Accessed on January 28, 2022.

Erb, T. J., & Zarzycki, J. (2018). A short history of RuBisCO: The rise and fall (?) of Nature's predominant CO_2 fixing enzyme. *Current Opinion in Biotechnology, 49*, 100–107. https://doi.org/10.1016/j.copbio.2017.07.017

Foti, C., Damiani, E., Zambonin, C. G., Cassano, N., Nettis, E., Ferrannini, A., et al. (2012). Urticaria and angioedema to rubisco allergen in spinach and tomato. *Annals of Allergy, Asthma & Immunology, 108*, 60–61. https://doi.org/10.1016/j.anai.2011.09.011

Freitas, M. L. F., Albano, K. M., & Telis, V. R. N. (2017). Characterization of biopolymers and soy protein isolate-high-methoxyl pectin complex. *Polímeros, 27*, 62–67. https://doi.org/10.1590/0104-1428.2404

Friedman, M., & Brandon, D. K. (2001). Nutritional and health benefits of soy proteins. *Journal of Agricultural and Food Chemistry, 49*, 1069–1086. https://doi.org/10.1021/jf0009246

Gasbarrini, G. B., & Mangiola, F. (2014). Wheat-related disorders: A broad spectrum of 'evolving' diseases. *United European Gastroenterology Journal, 2*, 254–262. https://doi.org/10.1177/2050640614535929

Gilani, G. S., Xiao, C. W., & Cockell, A. (2012). Impact of antinutritional factors in food proteins on the digestibility of protein and the bioavailability of amino acids and on protein quality. *British Journal of Nutrition, 108*, 315–332. https://doi.org/10.1017/S0007114512002371

Godfray, H. C. J., Beddington, J. R., Crute, I. R., Haddad, L., Lawrence, D., Muir, J. F., et al. (2010). Food security: The challenge of feeding 9 billion people. *Science, 327*, 812–818. https://doi.org/10.1126/science.1185383

Goliáš, J., Humlová, Z., Halada, P., Hábová, V., Janatková, I., & Tučková, L. (2013). Identification of rice protein recognized by the IgE antibodies of patients with food allergies. *Journal of Agricultural and Food Chemistry, 61*, 8851–8860. https://doi.org/10.1021/jf402759f

Gorissen, S. H. M., Crombag, J. J. R., Senden, J. M. G., Waterval, W. A. H., Bierau, J., Verdijk, L. B., et al. (2018). Protein content and amino acid composition of commercially available plant-based protein isolates. *Amino Acids, 50*, 1685–1695. https://doi.org/10.1007/s00726-018-2640-5

Greenhaff, P. L., Bodin, K., Harris, R., Hultman, E., Jones, D. D., & McIntyre, D. (1993). (1993) The influence of oral creatine supplementation on muscle creatine resynthesis following intense contration in man. *Journal of Physiology., 467*, 75P. https://doi.org/10.1113/jphysiol.1993.sp019746

Health and Human Services. (2017). Retrieved from: https://www.betterhealth.vic.gov.au/health/healthyliving/food-safety-when-cooking. Accessed on January 25, 2022.

Heng, L. (2005). Flavour Aspects of Pea and its Protein Preparations in Relation to Novel Protein Foods. Wageningen University, Wageningen. Retrieved from: https://library.wur.nl/WebQuery/wurpubs/343588. Accessed on February 15, 2022.

Hertzler, S. R., Lieblein-Boff, J. C., Weiler, M., & Allgeier, C. (2020). Plant proteins: Assessing their nutritional quality and effects on health and physical function. *Nutrients, 12*, 3704. https://doi.org/10.3390/nu12123704

Hirata, H., Sonoda, S., Agui, S., Yoshida, M., Ohinata, K., & Yoshikawa, M. (2007). Rubiscolin-6, a δ opioid peptide derived from spinach Rubisco, has anxiolytic effect via activating s1 and dopamine D1 receptors. *Peptides, 28*, 1998–2003. https://doi.org/10.1016/j.peptides.2007.07.024

Hoff, M., Son, D. Y., Gubesch, M., Ahn, K., Lee, S. I., Vieths, S., et al. (2007). Serum testing of genetically modifies soybeans with special emphasis on potential allergenicity of the heterologous protein CP4 EPSPS. *Molecular Nutrition & Food Research, 51*, 946–955. https://doi.org/10.1002/mnfr.200600285

House, J. D., Neufeld, J., & Leson, G. (2010). Evaluating the quality of protein from hemp seed (*Cannabis sativa L.*) products through the use of the protein digestibility-corrected amino acid score method. *Journal of Agricultural and Food Chemistry, 58*, 11801–11807. https://doi.org/10.1021/jf102636b

Jäger, R., Dudeck, J. E., Joy, J. M., Lowery, R. P., McCleary, S. A., Wilson, S. M., et al. (2013). Comparison of rice and whey protein osolate digestion rate and amino acid absorption. *Journal of the International Society of Sports Nutrition, 10*, 12. https://doi.org/10.1186/1550-2783-10-s1-p12

Jappe, U., & Vieths, S. D. A. T. A. (2010). Lupine, a source of new as well as hidden food allergens. *Molecular Nutrition & Food Research, 54*, 113–126. https://doi.org/10.1002/mnfr.200900365

Jayasena, V., Chih, H. J., & Nasar-Abbas, S. M. (2010). Functional properties of sweet lupin protein isolated and tested at various pH levels. *Research Journal of Agriculture and Biological Sciences, 6*, 130–137.

Jensen, R. E., & Bahr, J. T. (1977). Ribulose 1,5-bisphosphate carboxylase oxygenase. *Annual Review of Biochemistry, 28*, 378–400. https://doi.org/10.1146/annurev. pp.28.060177.002115

Jeurink, P. V., & Savelkoul, F. J. (2005). Induction and regulation of allergen-specific IgE. In L. J. E. J. Gilissen, H. J. Wichers, H. F. J. Savelkoul, & R. J. Bogers (Eds.), *Allergy Matters: New Approaches to Allergy Prevention and Management* (pp. 13–27). Springer - Life Sciences.

Jones, T. (2014) Alternative proteins. Future agenda. Retrieved from: http://www.futureagenda.org/pg/cx/view#390. Accessed February 4, 2022.

Jong, A., & Nieuwland, M. (2011). *Literature study on the properties of Rubisco*. The Netherlands: TNO report.

Jongh, H. H. J., Robles, C. L., Timmerman, E., Nordlee, J. A., Lee, P. W., Baumert, J. L., et al. (2013). Digestibility and IIE-binding of glycosylated codfish parvalbumin. *BioMed Research International, Article, ID 756789*, 10 pages. https://doi.org/10.1155/2013/756789

Kawashima, N., & Wildman, S. G. (1970). Fraction I protein. *Annual Review of Plant Physiology., 21*, 325–358. https://doi.org/10.1146/annurev. pp. 21.060170.001545

Kapel, R., Chabeau, A., Lesage, J., Riviere, G., Ravallec-Ple, R., Lecouturier, D., et al. (2006). Production, in continuous enzymatic membrane reactor, of an antihypertensive hydrolysate from an industrial alfalfa white protein concentrate exhibiting ACE inhibitory and opioid activities. *Food Chemistry, 98*, 120–126. https://doi.org/10.1016/j.foodchem.2005.05.062

Kaushal, P., Kumar, V., & Sharma, H. K. (2012). Comparative study of physico-chemical, functional, anti-nutritional and pasting properties of taro (*Colocasia esculenta*), rice (*Oryza sativa*), pegion pea (*Cajanus cajan*) flour and their blends. *LWT- Food Science and Technology, 48*, 59–68. https://doi.org/10.1016/j.lwt.2012.02.028

Keller, R. C. A., Orsel, R., & Hamer, R. J. (1997). Competitive adsorption behaviour of wheat flour components and emulsifiers at an air-water interface. *Journal of Cereal Science, 25*, 175–183. https://doi.org/10.1006/jcrs.1996.0089

Khalid, I. I., & Elharadallou, S. B. (2013). Functional Properties of cowpea (*Vigna Ungiculata L.Walp*), and lupin (*Lupinus Termis*) flour and protein isolates. *Journal of Nutrition & Food Sciences, 3*, 1–6. https://doi.org/10.4172/2155-9600.1000234

Kitts, D. D., & Weiler, K. (2003). Bioactive proteins and peptides from food sources. Applications of bioprocesses used in isolation and recovery. *Current Pharmaceutical Design, 9*, 1309–1323. https://doi.org/10.2174/1381612033454883

Kobbi, S., Balti, R., Bougatef, A., Le Flem, G., Firdaous, L., Bigan, M., et al. (2015). Antibacterial activity of novel peptides isolated from protein hydrolysates of RuBisCO purified from green juice alfalfa. *Journal of Functional Foods, 18*, 703–713. https://doi.org/10.1016/j.jff.2015.09.007

Kobbi, S., Bougatef, A., Leflem, G., Balti, R., Mickael, C., Fertin, B., et al. (2017). Purification and recovery of RuBisCO protein from alfalfa green juice: Antioxidative properties of generated protein hydrolysate. *Waste and Biomass Valorization, 8*, 493–504. https://doi.org/10.1007/s12649-016-9589-y

Krishnan, H. B., & Chen, M. H. (2013). Identification of an abundant 56 kDa protein implicated in food allergy as granule-bound starch synthase. *Journal of Agricultural and Food Chemistry, 61*, 5404–5409. https://doi.org/10.1021/jf4014372

Ku, M. S. B., Schmitt, M. R., & Edwards, G. E. (1979). Quantitative determination of RuBP carboxylase oxygenase protein in leaves of several C3 and C4 plants. *Journal of Experimental Botany., 30*, 89–98. https://doi.org/10.1093/jxb/30.1.89

Lamsal, B. P., Koegel, R. G., & Gunasekaran, S. (2007). Some physicochemical and functional properties of alfalfa soluble leaf proteins. *LWT - Food Science and Technology, 40*, 1520–1526. https://doi.org/10.1016/j.lwt.2006.11.010

Lan, Y., Xu, M., Ohm, J. B., Chen, B., & Rao, J. (2019). Solid dispersion-based spray-drying improves solubility and mitigates beany flavour of pea protein isolate. *Food Chemistry, 278*, 665–673. https://doi.org/10.1016/j.foodchem.2018.11.07

Lavine, E., & Ben-Shoshan, M. (2019). Anaphylaxis to hidden pea protein: A Canadian pediatric case series. *The Journal of Allergy and Clinical Immunology. In Practice, 7*, 2070–2071. https://doi.org/10.1016/j.jaip.2019.02.010

Liu, K., Zheng, J., & Chen, F. (2019). Effect of domestic cooking on rice protein digestibility. *Food Science & Nutrition, 7*, 608–616. https://doi.org/10.1002/fsn3.884

Malhotra, A., & Coupland, J. N. (2004). The effect of surfactants on the solubility, zeta potential, and viscosity of soy protein isolates. *Food Hydrocolloids, 18*, 101–108. https://doi.org/10.1016/S0268-005X(03)00047-X

Martínez, K. D., Farías, M. E., & Pilosof, A. M. R. (2011). Effects of soy protein hydrolysis and polysaccharides addition on foaming properties studied by cluster analysis. *Food Hydrocolloid, 25*, 1667–1676. https://doi.org/10.1016/j.foodhyd.2011.03.005

Martínez, K. D., Sánchez, C. C., Patino, J. M. R., & Pilosof, A. M. R. (2009). Interfacial and foaming properties of soy protein and their hydrolysates. *Food Hydrocolloid, 23*, 2149–2157. https://doi.org/10.1016/j.foodhyd.2009.03.015

McClements, D. J., & Grossmann, L. (2021). A brief review of the science behind the design of healthy and sustainable plant-based foods. *npj Science of Food, 5*, 17. https://doi.org/10.1038/s41538-021-00099-y

McGuire, M., & Beerman, K. A. (2013). Protein. In *Nutritional Sciences: From Fundamentals to Food* (pp. 166–174). Cengage Learning.

Mills, E. N. C., Mackie, A. R., Burney, P., Beyer, K., Frewer, L., Madsen, C., et al. (2007). The prevalence, cost and basis of food allergy across Europe. *Blackwell Munksgaard, 62*, 717–722. https://doi.org/10.1111/j.1398-9995.2007.01425.x

Mollakhalili-Meybodi, N., Arab, M., & Zare, L. (2021). Harmful compounds of soy milk: Characterization and reduction strategies. *Journal of Food Science and Technology, 59*, 3723–3732. https://doi.org/10.1007/s13197-021-05249-4

Monteiro, S. A., Carreira, A., Freitas, R., Pinheiro, A. M., & Ferreira, R. B. (2015). A nontoxic polypeptide oligomer with a fungicide potency under agricultural conditions which is equal or greater than that of their chemical counterparts. *PLoS ONE, 10*(4), e0122095.

Müller, O., & Krawinkel, M. (2005). Malnutrition and health in developing countries. *Canadian Medical Association Journal, 173*, 279–286. https://doi.org/10.1503/cmaj.050342

Nguyen, H. C., Hoefgen, R., & Hesse, H. (2012). Improving the nutritive value of rice seeds: Elevation of cysteine and methionine contents in rice plants by ectopic expression of a bacterial serine acetyltransferase. *Journal of Experimental Botany, 63*, 5991–6001. https://doi.org/10.1093/jxb/ers253

Ogawa, T., Bando, N., Tsuji, H., Nishikawa, K., & Sasaoka, K. (1991). Investigation of the IgE-binding proteins in soybeans by immunoblotting with the sera of the soybean-sensitive patients with atopic dermatitis. *Journal of Nutritional Science and Vitaminology, 37*, 555–565. https://doi.org/10.3177/jnsv.37.555

Pasha, I., Saeed, F., Sultan, M. T., Batool, R., Aziz, M., & Ahmed, W. (2016). Wheat allergy & intolerance, recent updates and perspectives. *Critical Reviews in Food Science and Nutrition, 56*, 13–24. https://doi.org/10.1080/10408398.2012.659818

Perlikowska, R., & Janecka, A. (2017). Rubiscolins-highly potent peptides derived from plant proteins. *Mini Reviews in Medicinal Chemistry, 18*, 104–112. https://doi.org/10.2174/1389557517666170426160703

Popova, A., & Mihaylova, D. (2019). Antinutrients in plant-based foods: A review. *The Open Biotechnology Journal, 13*, 68–76. https://doi.org/10.2174/1874070701913010068

Pouvreau, L., Smit, B., & Van de Velde, F. (2013). Securing food proteins: From by-products to functional ingredients. In P. Williams, & G. Phillips (Eds.), *Gums and stabilisers for the food industry 17: The changing face of food manufacture: The role of hydrocolloids* (pp. 46–51). Royal Society of Chemistry.

Prévot-D'Alvise, N., Lesueur-Lambert, C., Fertin-Bazus, A., Fertin, B., Dhulster, P., & Guillochon, D. (2004). Continuous enzymatic solubilization of alfalfa proteins in an

ultrafiltration reactor. *Enzyme and Microbial Technology, 34*, 380–391. https://doi.org/10.1016/j.enzmictec.2003.05.001

Rackis, J. J., Sessa, D. J., & Honig, D. H. (1979). Flavor problems of vegetable food proteins. *Journal of the American Oil Chemists' Society, 56*, 262–271.

Raikos, V., Neacsu, M., Russell, W., & Duthie, G. (2014). Comparative study of the functional properties of lupin, green pea, fava bean, hemp, and buckwheat flours as affected by pH. *Food Science & Nutrition, 2*, 802–810. https://doi.org/10.1002/fsn3.143

Rault, M., Giudiciorticoni, M. T., Gontero, B., & Ricard, J. (1993). Structural and functional-properties of a multienzyme complex from spinach chloroplasts. Stoichiometry of the polypeptide-chains. *European Journal of Biochemistry, 217*, 1065–1073. https://doi.org/10.1111/j.1432-1033.1993.tb18338.x

Raven, J. A. (2013). Rubisco: Still the most abundant protein of earth? *New Phytologist, 198*, 1–3. https://doi.org/10.1111/nph.12197

Roland, W. S. U., Pouvreau, L., Curran, J., van de Velde, F., & Kok, P. M. T. (2017). Flavor aspects of pulse ingredients. *Cereal Chemistry, 94*, 58–65. https://doi.org/10.1094/CCHEM-06-16-0161-Fl

Schilling, B. K., Stone, W. H., Utter, A., Kearney, J. T., Johnson, M., Coglianese, R., et al. (2000). Creatine supplementation and health variables: A retrospective study. *Official Journal of American College of Sport Medicine, 33*, 183–188. https://doi.org/10.1097/00005768-200102000-00002

Schlegel, K., Sontheimer, K., Hickisch, A., Wani, A. A., Eisner, P., & Schweiggert-Weisz, U. (2019). Enzymatic hydrolysis of lupin protein isolates—Changes in the molecular weight distribution, technofunctional characteristics and sensory attributes. *Food Science & Nutrition, 7*, 2747–2759. https://doi.org/10.1002/fsn3.1139

Setchell, K. D., Brown, N. M., & Lydeking, O. E. (2002). The clinical importance of the metabolite equol- A clue to the effectiveness of soy and its isoflavones. *Journal of Nutrition, 132*, 3577–3584. https://doi.org/10.1093/jn/132.12.3577

Shand, P., Ya, H., Pietrasik, Z., & Wanasundara, P. (2007). Physicochemical and textural properties of heat-induced pea protein isolate gels. *Food Chemistry, 102*, 1119–1130. https://doi.org/10.1016/j.foodchem.2006.06.060

Sheen, S. J. (1991). Comparison of chemical and functional-properties of soluble leaf proteins from 4 plant-species. *Journal of Agricultural and Food Chemistry, 39*, 681–685. https://doi.org/10.1021/JF00004A011

Shen, Y., Hong, S., Du, Z., Chao, M., O'Quinn, T., & Li, Y. (2022). Effect of adding modified pea protein as functional extender on the physical and sensory properties of beef patties. *LWT - Food Science and Technology, 154*, Article 112774. https://doi.org/10.1016/j.lwt.2021.112774

Segura, M. E., & Rosell, C. M. (2011). Cheminal composition and starch digestibility of different gluten-free breads. *Plant Foods for Human Nutrition, 3*, 224–230. https://doi.org/10.1007/s11130-011-0244-2

Song, Y., Qin, R., Yang, S., Li, J., & Wang, R. (2018). Improvement of foaming and emulsifying properties of gluten by conjugation with fructose through maillard reaction. *Grain & Oil Science and Technology, 1*, 119–125. https://doi.org/10.3724/sp.j.1447.gost.2018.18017

Spreitzer, R. J., & Salvucci, M. E. (2002). RuBisCo: Structure, regulatory interactions, and possibilities for a better enzyme. *Annual Review of Plant Biology, 53*, 449–475. https://doi.org/10.1146/annurev.arplant.53.100301.135233

Stefano, E., Agyei, D., Njoku, E. N., & Udenigwe, C. C. (2018). Plant RuBisCo: An underutilized protein for food applications. *Journal of the American Oil Chemists' Society, 95*, 1063–1074. https://doi.org/10.1002/aocs.12104

Stone, M. H., Sanborn, K., Smith, L. L., O'Bryant, H. S., Hoke, T., Utter, A. C., et al. (1999). Effects of in-season (5 weeks) creatine and pyruvate supplementation on anaerobic performance and body composition in American football players. *Journal of the International Society of Sports Nutrition, 9*, 146–165. https://doi.org/10.1123/ijsn.9.2.146

Tatham, A. S., & Shewry, P. R. (2008). Allergy to wheat and related cereals. *Clinical & Experimental Allergy, 38*, 1712–1726. https://doi.org/10.1111/j.1365-2222.2008.03101.x

Tang, C. H. (2017). Emulsifying properties of soy proteins: A critical review with emphasis on the role of conformational flexibility. *Critical Reviews in Food Science and Nutrition, 57*, 2636–2679. https://doi.org/10.1080/10408398.2015.1067594

Tenorio, A. T., Gieteling, J., De Jong, G. A., Boom, R. M., & Van Der Goot, A. J. (2016). Recovery of protein from green leaves: Overview of crucial steps for utilisation. *Food Chemistry, 203*, 402–408. https://doi.org/10.1016/j.foodchem.2016.02.092

Teschemacher, H. (2003). Opioid receptor ligands derived from food proteins. *Current Pharmaceutical Design, 9*, 1331–1344. https://doi.org/10.2174/1381612033454856

Thewissen, B. G., Celus, I., Brijs, K., & Delcour, J. A. (2011). Foaming properties of wheat gliadin. *Journal of agricultural and food chemistry, 59*(4), 1370–1375. https://doi.org/10.1021/jf103473d

Thomas, K., Aalbers, M., Bannon, G. A., Bartels, M., Dearman, R. J., Esdaile, D. J., et al. (2004). A multi-laboratory evaluation of a common *in vitro* pepsin digestion assay protocol used in assessing the safety of novel proteins. *Regulatory Toxicology and Pharmacology, 39*, 87–98. https://doi.org/10.1016/j.yrtph.2003.11.003

Tosca, M. A., Pistorio, A., Accogli, A., Silvestri, M., Rossi, G. A., Ciprandi, G. (2014). Egg allergy: the revelance of molecular-based allergy diagnostics. *Clinical & Experimental Allergy*, 44, 1094-1095. https://doi.org/ 10.1111/cea.12360.

Trovaslet, M., Kapel, R., Ravallec-Ple, R., Mouni, F., Clarisse, M., Faille, C., et al. (2007). Secretagogue and bacteriostatic active fractions derived from a peptic hydrolysate of alfalfa RuBisCO small purified subunit. *Journal of the Science of Food and Agriculture, 87*, 534–540. https://doi.org/10.1002/jsfa.2754

Udenigwe, C. C., & Aluko, R. E. (2012). Food protein-derived bioactive peptides: Production, processing, and potential health benefits. *Journal of Food Science, 71*, 11–24. https://doi.org/10.1111/j.1750-3841.2011.02455.x

Udenigwe, C. C., Okolie, C. L., Qian, H., Ohanenye, I. C., Agyei, D., & Aluko, R. E. (2017). Ribulose-1,5-bisphosphate carboxylase as a sustainable and promising plant source of bioactive peptides for food applications. *Trends in Food Science & Technology, 69*, 74–82. https://doi.org/10.1016/j.tifs.2017.09.001

Van del Velde, F. V., Alting, A., Pouvreau, L. (2011). From waste product to food ingredient: The extraction of the abundant plant protein RuBisCO. *New Food*, 14, 10-13. Retrieved from: https://www.newfoodmagazine.com/article/4461/from-waste-product-to-food-ingredient-the-extraction-of-abundant-plant-protein-rubisco/. Accessed January 20, 2022.

Wang, T., Xu, P., Chen, Z., Zhou, X., & Wang, R. (2018). Alteration of the structure of rice proteins by their interaction with soy protein isolates to design novel protein composites. *Food & Function, 9*, 4282–4291. https://doi.org/10.1039/c8fo00661j

Wensing, M., Knulst, A. C., Piersma, S., O'Kane, F., Knol, E. F., & Koppelman, S. J. (2003). Patients with anaphylaxis to pea can have peanut allergy caused by cross-reactive IgE to vicilin (Ara h 1). *The Journal of Allergy and Clinical Immunology, 111*, 420–424. https://doi.org/10.1067/mai.2003.61

Yang, S., Kawamura, Y., & Yoshikawa, M. (2003). Effect of rubiscolin, a δ opioid peptide derived from Rubisco, on memory consolidation. *Peptides, 24*, 325–328. https://doi.org/10.1016/s0196-9781(03)00044-5

Yang, S., Yunden, J., Sonoda, S., Doyama, N., Lipkowski, A. W., Kawamura, Y., et al. (2001). Rubiscolin, a δ selective opioid peptide derived from plant Rubisco. *FEBS Letters, 509*, 213–217. https://doi.org/10.1016/s0014-5793(01)03042-3

Zhu, S. M., Lin, S. L., Ramaswamy, H. S., Yu, Y., & Zhang, Q. T. (2016). Enhancement of functional properties of rice bran proteins by high pressure treatment and their correlation with surface hydrophobicity. *Food and Bioprocess Technology, 10*, 317–327. https://doi.org/10.1007/s11947-016-1818-7

Zuber, T., Siegert, W., Salehi, H., Hummel, F., & Rodehutscord, M. (2018). Variability of amino acid digestibility of lupin and pea grains in caecectomised laying hens. *British Poultry Science, 60*, 229–240. https://doi.org/10.1080/00071668.2018.1556389