

PHENOTYPES

Extracellular Cyclophilin Levels Associate with Parameters of Asthma in Phenotypic Clusters

ERIK J. STEMMY, PH.D.,¹ ANGELA S. BENTON, B.A.S.,² JENNIFER LERNER, B.S.,² SARAH ALCALA, B.S.,^{1,2} STEPHANIE L. CONSTANT, PH.D.,¹ AND ROBERT J. FREISHTAT, M.D., M.P.H.^{2,3,*}

¹*Department of Microbiology, Immunology, and Tropical Medicine, The George Washington University Medical Center, Washington, DC, USA.*

²*Center for Genetic Medicine Research, Children's National Medical Center, Washington, DC, USA.*

³*Departments of Pediatrics, Emergency Medicine, and Integrative Systems Biology, The George Washington University Medical Center, Washington, DC, USA.*

Objective. Leukocyte persistence during chronic (quiescent) phases of asthma is a major hallmark of the disease. The mechanisms regulating these persistent leukocyte populations are not clearly understood. An alternative family of chemoattracting proteins, cyclophilins (CyPs), has recently been shown to contribute to leukocyte recruitment in animal models of allergic asthma. The goals of this study were to determine whether CyPs are present in asthma patients during the chronic phase of the disease and to investigate whether levels of CyPs associate with clinical parameters of disease severity. **Methods.** Nasal wash samples from an urban cohort of 137 patients of age 6–20 years with physician-diagnosed asthma were examined for the presence of cyclophilin A (CypA), cyclophilin B (CypB), as well as several other classical chemokines. Linear, logistic, or ordinal regressions were performed to identify associations between CyPs, chemokines, and clinical parameters of asthma. The asthma cohort was further divided into previously established phenotypic clusters (cluster 1: $n = 55$; cluster 2: $n = 31$; and cluster 3: $n = 51$) and examined for associations. **Results.** Levels of CypB in the asthma group were highly elevated compared to nonasthmatic controls, while a slight increase in Monocyte Chemoattractant Protein-1 (MCP-1) was also observed. CypA and MCP-1 were associated with levels of eosinophil cationic protein (ECP; a marker of eosinophil activation). Cluster-specific associations were found for CypA and CypB and clinical asthma parameters [e.g. forced expiratory volume in 1 second (FEV₁) and ECP]. **Conclusions.** CyPs are present in nasal wash samples of asthma patients and may be a novel biomarker for clinical parameters of asthma severity.

Keywords asthma, chemokine, cluster analysis, cyclophilin, phenotype

INTRODUCTION

Cyclophilins (CyPs) are ubiquitous intracellular proteins commonly known as the target of the immunosuppressive drug cyclosporine A (1). In addition to their intracellular role, increasing evidence has suggested a role for CyPs in intercellular communication (2). We (3–6) and others (7–9) have shown that cyclophilin A (CypA) and/or cyclophilin B (CypB) are potent chemoattractants for a variety of human and mouse leukocyte subsets. In light of these findings, we have previously proposed that extracellular CyPs may be important alternative regulators of leukocyte migration (2, 10). Indeed, elevated levels of CyPs have been reported in several inflammatory diseases, including rheumatoid arthritis (11, 12), vascular smooth muscle disease (13, 14), and severe sepsis (15). Evidence that extracellular CyPs not only are present, but also contribute to leukocyte recruitment during inflammatory responses was demonstrated in several

animal models of inflammation, including acute lung injury (3) and acute allergic asthma (4, 6).

A major hallmark of chronic allergic asthma is the persistence of proinflammatory leukocytes within the airways, submucosa, and mucosal epithelium during these quiescent (or chronic) periods of the disease. The factors that regulate leukocyte persistence in human asthma are not clearly understood. Levels of classical chemokines responsible for leukocyte influx into the lung, such as eotaxins 1–3, Macrophage Inflammatory Protein 1 α (MIP-1 α), and MCP-1, have been shown to increase rapidly in airways after acute allergen exposure, but return to baseline levels by 24 hour after exposure (16, 17). In other studies, levels of eotaxin were found to be equivalent between asthma patients in the chronic phase of the disease and healthy controls, despite the former group having elevated airway eosinophil numbers (18). Such findings suggest that factors other than classical chemokines may contribute to the persistent recruitment of proinflammatory leukocytes during chronic asthma.

The presence of cytokines, chemokines, and other inflammatory proteins has been demonstrated in nasal wash samples from infants and children with several types of respiratory inflammation (19–22). It has also been shown that changes in these protein levels in nasal wash samples correlate well with changes seen in the lower

The authors Stephanie L. Constant and Robert J. Freishtat contributed equally to the oversight of these studies.

*Corresponding author: Robert J. Freishtat, M.D., M.P.H., Center for Genetic Medicine Research, Children's National Medical Center, 111 Michigan Avenue, NW, Washington, DC 20010, USA; Tel: (202) 476-2971; Fax: (202) 476-6014; E-mail: rfreishtat@childrensnational.org

airways [for review see Frischer and Baraldi (23)]. Studies in our laboratory examining the role of Cyps in acute allergic asthma demonstrate that blocking the activity of extracellular Cyps reduced leukocyte influx into airways and tissues by up to 80% (4). Other recent studies in our laboratory using a new mouse model of chronic allergic asthma show elevated levels of extracellular Cyps present in airways, not only following acute allergen exposure, but also during the chronic phase of the disease (24). Based on these observations, extracellular Cyps are potential candidates for regulating the persistent leukocyte infiltration observed during chronic phases of human asthma. Further, the association shown between Cyps and inflammation severity in other inflammatory diseases may suggest similar associations for Cyps in asthma. In the current study we investigated whether extracellular Cyps are present in nasal wash samples collected from asthma patients in the chronic phase of their disease and whether those levels demonstrate any specific associations with parameters of disease phenotype and/or severity.

METHODS

Study Cohort

Study subjects were enrolled in the Asthma Severity Modifying Polymorphisms (AsthMaP) Project, a study of asthma in urban children and adolescents based at the Children's National Medical Center (Washington, DC, USA). The AsthMaP Project is described in detail elsewhere (25–28). Briefly, 154 participants aged 6–20 years, with physician-diagnosed asthma for at least 1 year (with varying chronic asthma therapy), were recruited from the Children's National Emergency Department. Participants then returned to the Clinical Research Center for a study visit at least 4 weeks after completion of their most recent oral steroid dose. If subjects arrived at the Clinical Research Center with current symptoms, they were deferred to return when asymptomatic. Informed consent and assent were obtained from participants and/or guardians. Healthy nonsmoking, nonasthmatic subjects without seasonal nasal allergies ($n = 22$) were recruited from adult volunteers to provide control nasal wash samples. These studies were approved by our Institutional Review Board.

Clinical Data Collection

Clinical characteristics were measured in each study participant. Parental interviews using the Integrated Therapeutics Group (ITG) Child Asthma Short Form (29) and the National Institutes of Health National Asthma Education and Prevention Program (NAEPP) 2007 criteria (30, 31) were conducted at least 4 weeks after the most recent oral corticosteroid dose. Spirometry measurements were performed with a MedGraphics CPSF/DTM USB PC-based system (MedGraphics, St. Paul, MN, USA) before and after short-acting

β -agonist treatment. Levels of serum immunoglobulin-E (IgE) were measured by Quest Diagnostics (Herndon, VA, USA). Details of β -agonist, inhaled corticosteroid, and leukotriene agonist use were also recorded. Only limited clinical data were collected from the anonymous healthy control volunteers ($n = 22$) in whom nasal washes were performed.

Nasal washes were performed by instilling 3 mL of sterile isotonic saline into each nare, holding for 10 seconds, and then collecting in a specimen collection container. For comparison, nasal washes were also performed on 22 control individuals without asthma. The collected material was then centrifuged to separate cells and soluble factors. Analyses were conducted on individual samples. Nasal washes were performed as a minimally invasive alternative to bronchoalveolar lavage. Other studies have shown that nasal samples correlate well with samples collected from the lower airways (23).

Eosinophil and neutrophil numbers were counted from nasal wash samples on slides stained with Wright's stain. P-selectin, MIP-1 α Chemokine (C-C motif) Ligand (CCL3), MCP-1 (CCL2), and Regulated upon Activation, Normal T-cell Expressed, and Secreted (RANTES) (CCL5) were measured in the soluble fraction (previously frozen) using a FlowCytomix Simplex Kit and analyzed using FlowCytomix Pro software v2.3 (eBioscience, San Diego, CA, USA). Eotaxin-1 (CCL11), eotaxin-2 (CCL24), and eosinophil cationic protein (ECP) were measured by ELISA (eotaxins: R&D Systems, Minneapolis, MN, USA; ECP: Medical & Biological Laboratories, Nagoya, Japan). Levels of extracellular CypA and CypB were measured by Western blot analysis: several concentrations of recombinant CypA and B were used to establish a standard curve of Cyp concentrations, which was then used to quantify the amount of Cyp present in each sample.

Asthma Phenotypic Clustering

The 154 AsthMaP participants were grouped into four phenotypic clusters based on groupings of clinical variables as we previously described (26). One cluster consisted of six subjects displaying a mild asthma phenotype and was excluded from further analyses. An additional 11 participants lacked nasal wash samples and were also excluded from analyses. The remaining 137 participants were grouped into the clusters summarized in Table 1. The first cluster ($n = 55$) was predominantly male (80%), with a preponderance of neutrophils in their nasal wash. Cluster 2 ($n = 31$) was predominantly female (67.7%) with a high mean body mass index (BMI) percentile (86), as well as later onset asthma. The third cluster ($n = 51$) is a typical allergic asthma phenotype characterized by elevated eosinophils in nasal washes (Table 1). The phenotype of cluster 2 is important because the combination of high BMI and late age of onset (8.0-year-olds, compared to 2.0 and 3.3 for clusters 1 and 3, respectively) of asthma is frequently seen in the clinic. Further, studies have suggested a

TABLE 1.—Associations between cyclophilins, classical chemokines, and asthma.

Variable	Healthy controls (n = 22)	All asthma (n = 137)	(neutrophilic) Cluster 1 (n = 55)	(female late onset) Cluster 2 (n = 31)	(allergic asthma) Cluster 3 (n = 51)	p-value [†]
Sex, % Male	41	58	80	32	51	<.001
Age, years (SE)	31.6 (2.4)	11.5 (0.3)	11.2 (0.6)	13.5 (0.6)	10.0 (1.0)	.002
Age of onset, years (SE)	—	3.9 (0.3)	3.3 (0.4)	8.0 (0.7)	2.0 (0.4)	<.001
Body Mass Percentile, % (SE)	—	72 (2)	50 (3)	86 (3)	87 (2)	<.001
Prebronchodilator FEV ₁ , %predicted	—	87 (2)	86 (3)	99 (3)	82 (2)	<.001
FEV ₁ change with bronchodilator (SE)	—	8 (1)	7 (2)	6 (1)	10 (2)	.38
NAEPP severity classification (IQR)	—	3 (2,4)	3 (2,3)	3 (2,4)	3 (2,4)	.28
ITG-composite score (IQR)	—	81 (67,92)	83 (67,94)	81.0 (67,92)	79 (65,88)	.26
Nasal eosinophils, % (IQR)	—	52 (3,92)	10 (0,63)	56 (14,91)	82 (45,96)	<.001
Nasal neutrophils, % (IQR)	—	33 (5,71)	50 (8,95)	38 (3,49)	13 (3,50)	.014
Nasal wash ECP, µg/mL (IQR)	—	9.0 (1.3,47.2)	27.0 (1.3,51.8)	3.6 (0.6,38.1)	6.9 (1.3,37.5)	.06
Nasal wash P-selection, µg/mL (SE)	—	1.4 (0.2)	1.3 (0.2)	14 (0.13)	1.0 (0.2)	.31
Nasal wash CypA, µg/mL (IQR)	0 (0,0)	0 (0,0)	0 (0,0)	0 (0,0)	0 (0,0)	.14
Nasal wash CypB, µg/mL (IQR)	1.2 (0.9,3.9)	7.9 (5.3,14.4)*	8.6 (5.1,14.5)	8.9 (5.3,15.5)	7.6 (5.6,12.0)	.75
Nasal wash MCP-1, pg/mL (IQR)	16.3 (0,104.5)	58.9 (0,149.8)*	80.4 (20.6,172.2)	51.9 (0,105.0)	48.7 (0,113.7)	.16
Total serum IgE, IU/mL (IQR)	—	248 (70,512)	210 (59,202)	265 (28,531)	248 (95,577)	.55
Current β-agonist use (%)	—	69	62	74	75	.29
Current inhaled steroid use (%)	—	34	24	36	45	.07
Current leukotriene antagonist use (%)	—	4	4	7	4	.81

Notes: SE, standard error; FEV₁, forced expiratory volume in 1 second; NAEPP, National Asthma Education and Prevention Program, 2007 criteria; IQR, interquartile range; ITG, Integrated Therapeutics Group's Child Asthma Short Form; ECP, eosinophil cationic protein.

*Indicates significant difference ($p < .05$) compared to control group.

**Significant values among the three clusters were derived using one-way ANOVA for normally distributed continuous variables, or Kruskal–Wallis for nonparametric continuous variables, and chi-square for nominal variables.

connection between obesity and asthma, particularly in females during adolescence (32, 33). β-Agonist, inhaled corticosteroid, and leukotriene agonist use was similar among the three clusters (Table 1). There were no significant differences in the variables summarized in Table 1 between the clustered participants and those lacking nasal wash samples.

Statistical Analyses

Associations between the variables summarized in Table 1 were investigated between the asthma and healthy nonasthmatic control groups as a whole, as well as between the clusters within the asthma group. Data were log₁₀ transformed when not normally distributed. *t*-Tests were used to compare Cyp and chemokine levels between the asthma and control groups. To identify associations between the variables and cluster groups we used logistic, linear, or ordinal regression. All beta coefficients and *p*-values were corrected for age, race, gender, and BMI percentile during analysis. Statistical tests were performed using SPSS Statistics 18 (SPSS, Chicago, IL, USA).

RESULTS

Associations between Cyclophilins, Classical Chemokines, and Asthma

This study examined nasal wash samples from 137 participants in the AsthMaP project for whom nasal wash samples were available (Table 1). In the asthma cohort as a whole, 58.4% were male and the average age was 11.5

[standard error (SE), 0.3] years. The mean BMI percentile was 72 (2.0). Of the participants, 90% self-identified as African American, and 89% had persistent asthma as defined by NAEPP 2007 criteria.

We first measured levels of extracellular CypA and CypB, as well as the classical chemokines MCP-1, MIP-1α, RANTES, eotaxin-1, and eotaxin-2, in nasal wash samples collected from the AsthMaP cohort and 22 healthy nonasthmatic control participants. As shown in Figure 1, levels of both CypA and CypB were elevated in asthma patients, relative to control donors. In the case of CypB, this difference was significant ($p < .001$); however, in the case of CypA, levels of the protein were undetectable in many samples. Levels of MCP-1 were also significantly increased in the asthma group ($p < .05$). RANTES, MIP-1α, and eotaxin-2 were not different between the two groups (Figure 1) and levels of eotaxin-1 were below the sensitivity threshold of the assay in all samples (data not shown).

We next compared levels of CypA, CypB, and MCP-1 between the asthma and control groups in the context of the phenotypic clusters of asthma. We observed no significant associations between any of these three chemotactic agents and the AsthMaP clusters, nor were there any significant increases in these variables in the cluster groups (Figure 2).

Associations between Cyclophilins, MCP-1, and Parameters of Asthma Severity

We next examined whether the observed levels of CypA, CypB, or MCP-1 associate with parameters associated with disease severity (Table 2). After correcting for




FIGURE 1.—CypB and MCP-1 increased in asthma patients during the chronic phase of the disease. Levels of extracellular CypA, CypB, and other classical chemokines were assessed in nasal wash samples from asthma patients and healthy control volunteers. CypA and CypB were measured by Western blot analysis, and classical chemokines were measured by cytometric bead array or ELISA (eotaxin-2). Asthma group: $n = 137$; and control group: $n = 22$. For the analysis, data were $\log_{10}$ transformed, and a t -test was used to identify significant differences from the control group. (A) CypA, (B) CypB, (C) MCP-1, (D) MIP-1 α , (E) RANTES, and (F) eotaxin-2.

Notes: CypA, cyclophilin A; CypB, cyclophilin B; MCP.

* $p < .05$; ** $p < .0001$.

age, gender, BMI, and race, we found significant associations with several clinical parameters. Despite a failure to observe a statistically significant difference in levels between asthma and control groups, we observed associations between levels of CypA and several different parameters of the disease. For every $\log_{10}$ unit increase in CypA there was a 0.204 unit increase in $\log_{10}$ ECP [$\beta = 0.204$ [95% confidence interval (CI): 0.089, 0.319], adjusted $p = .001$], a marker of eosinophil activation. CypA levels were also negatively associated with a spirometry measurement: forced expiratory volume in 1 second (FEV_1), change with bronchodilator. For every $\log_{10}$ unit increase in CypA there was a 2.973 unit decrease in $\log_{10}$ FEV_1 change with bronchodilator [-2.973 (-4.887 , -1.059); adjusted $p = .003$], indicating a worse response to the bronchodilator drug. Based on recently published observations that platelet activation correlates with airway eosinophilia in asthma (25), we also examined whether levels of CypA or CypB might be associated with platelet activation in asthmatic airways.

Both Cyps were associated with levels of P-selectin, which is a measurement of platelet activation [CypA: 0.418 (0.153, 0.683), adjusted $p = .002$; CypB: 0.842 (0.018, 1.665); adjusted $p = .045$]. For every $\log_{10}$ unit increase in Cyp, there was a corresponding increase in P-selectin (0.418 for CypA; 0.842 for CypB), indicating that greater levels of Cyps are associated with more platelet activation. In the case of CypB, this was the only significant association in the asthma cohort as a whole.

Levels of MCP-1 were associated with ECP [0.209 (0.10, 0.318); adjusted $p < .001$] and P-selectin [0.349 (0.113, 0.585); adjusted $p = .004$]. Additionally, MCP-1 was associated with increases in NAEPP step level. For every $\log_{10}$ unit increase in MCP-1 there was a 0.232 unit increase in NAEPP step level [(0, 0.64) adjusted $p = .05$].

We next looked within each phenotypic cluster for associations between levels of CypA, CypB, MCP-1, and disease parameters associated with asthma severity (Table 2). In cluster 1 we observed associations between CypA and P-selectin [0.553 (0.142, 0.965), adjusted




FIGURE 2.—Compared levels of CypA, CypB, and MCP-1 by asthma cluster. Levels of extracellular CypA, CypB, and MCP-1 were measured in nasal wash samples and then grouped according to asthma phenotype (Table 1). Asthma cluster 1: $n = 55$; cluster 2: $n = 31$; cluster 3: $n = 51$; control group: $n = 22$. Data were $\log_{10}$ transformed, and t -tests were used to identify significant differences from the control group. No significant differences were observed. (A) CypA, (B) CypB, and (C) MCP-1 clusters.
Note: CypA, cyclophilin A; CypB, cyclophilin B; MCP.

$p = .01$] and between CypB and P-selectin [2.685 (0.497, 4.874), adjusted $p = .018$], indicating that for each $\log_{10}$ unit increase in Cyp there was a corresponding increase in P-selectin (0.553 for CypA, 2.685 for CypB). Also in cluster 1, for every $\log_{10}$ unit increase in MCP-1 there was a 0.215 $\log_{10}$ unit increase in ECP [(0.051, 0.380), adjusted $p = .011$]. In cluster 2 we also observed a similar association between MCP-1 and nasal wash ECP [0.322 (0.033, 0.612), adjusted $p = .032$]. The only significant association in cluster 3 was a 0.274 $\log_{10}$ increase in ECP for every $\log_{10}$ unit increase in CypA [(0.007, 0.478), adjusted $p = .01$].

DISCUSSION

Asthma is characterized by recurrent periods of acute airway constriction, mucus hypersecretion, and lung inflammation in response to inhaled allergen or other environmental stimuli (34). This inflammation is driven by specific subsets of leukocytes, such as antigen-specific T-cells and IgE-switched B-cells, mast cells, and eosinophils (35). CD4⁺ T-cells expressing the activation markers CD25 and major histocompatibility complex (MHC) class II have been reported to be increased in the airways of asthmatic patients and have also been shown to correlate with disease severity (36, 37). These activated CD4⁺ T-cells promote and perpetuate the asthmatic response by secreting TH2-associated cytokines such as interleukin (IL)-4, IL-5, and IL-13, which drive isotype switching in B-cells and help recruit activated eosinophils into airways (35, 38, 39).

However, asthma is also a chronic disease in that the inflammation observed during acute responses never completely resolves, even in the absence of allergen stimulation. A major hallmark of chronic allergic asthma is the persistence of proinflammatory leukocytes within the submucosa and mucosal epithelium during these chronic periods of the disease. Indeed, bronchial biopsies in asthma patients sampled during the chronic phase show elevated numbers of eosinophils, activated lymphocytes, and mast cells, compared to biopsies of healthy control subjects (18). These persistent inflammatory cells are thought to contribute to the tissue injury and remodeling that mediate the pathology of chronic asthma (40). Importantly, activated effector leukocytes recruited to the lung have a limited lifespan (41–43) and do not proliferate at sites of inflammation (44). Therefore the persistent inflammation seen during chronic phases of asthma likely requires recruitment stimuli to replenish these populations of leukocytes.

The specific factors regulating leukocyte persistence in the absence of allergen stimulation are not known. Obvious candidates are chemokines typically associated with allergic responses, including eotaxins 1–3, RANTES, MIP-1 α , and MCP-1. However, while levels of these chemokines are greatly increased after acute allergen challenge, they have been shown to return to baseline levels by 24 hour after exposure (16, 17). In other studies, levels of eotaxin were found to be equivalent between asthma patients in clinical remission and healthy controls, despite the former group having elevated airway eosinophil numbers (18). Such findings suggest that

TABLE 2.—Associations between cyclophilins, MCP-1, and parameters of asthma severity.

Continuous variables	<i>n</i> = 137		Cluster 1 (<i>n</i> = 55)		Cluster 2 (<i>n</i> = 31)		Cluster 3 (<i>n</i> = 51)	
	(95% CI)	<i>p</i>	(95% CI)	<i>p</i>	(95% CI)	<i>p</i>	(95% CI)	<i>p</i>
Associations between CypA levels and asthma parameters								
Log ECP	0.204 (0.089,0.319)	.001	0.123 (−0.027,0.277)	.106	0.333 (−0.075,0.741)	.101	0.274 (0.07,0.478)	.01
P-selectin	0.418 (0.153,0.683)	.002	0.553 (0.142,0.965)	.01	0.346 (−0.363,1.054)	.298	0.224 (−0.140,0.588)	.214
Log FEV ₁ , prebronchodilator	2.157 (−0.212,4.526)	.074	3.749 (0.218,7.28)	.038	1.882 (−4.432,8.196)	.544	0.13 (−3.631,3.892)	.945
Log FEV ₁ change with bronchodilator	−2.973 (−4.887, −1.059)	.003	−4.009 (−7.007, −1.012)	.01	−1.937 (−5.410,1.535)	.261	−1.823 (−5.336,1.69)	.31
NAEPP Severity Classification	0.148 (−0.093,0.389)	.229	0.270 (−0.079,0.619)	.129	−0.72 (−0.812,0.668)	.849	0.018 (−0.410,0.446)	.935
Associations between CypB levels and asthma parameters								
Log ECP	0.31 (−0.152,0.772)	.186	−0.21 (−0.833,0.791)	.958	0.638 (−0.709,1.984)	.325	0.562 (−0.107,1.232)	.096
P-selectin	0.842 (0.018,1.665)	.045	2.685 (0.497,4.874)	.018	−0.239 (−2.558,2.081)	.821	0.590 (−0.527,1.707)	.284
Log FEV ₁ , prebronchodilator	3.668 (−5.45,12.786)	.428	−0.689 (−19.982,18.604)	.943	12.889 (−5.959,31.736)	.171	0.576 (−10.856,12.008)	.92
Log FEV ₁ change with bronchodilator	−0.268 (−0.7828,0.7292)	.944	2.124 (−14.672,18.919)	.8	−0.364 (−11.356,10.628)	.946	0.02 (−10.792,6.6910.83)	.997
NAEPP Severity Classification	0.462 (−1.367,0.444)	.318	0.370 (−1.321,2.061)	.668	−1.548 (−3.973,8.77)	.211	−0.279 (−1.562,1.003)	.669
Associations between MCP-1 levels and asthma parameters								
Log ECP	0.209 (0.10,0.318)	< .001	0.215 (0.51,0.380)	.011	0.322 (0.033,0.612)	.032	0.066 (−0.169,0.302)	.566
P-selectin	0.349 (0.113,0.585)	.004	0.463 (−0.034,0.961)	.067	0.340 (−0.178,0.858)	.172	0.170 (−0.208,0.547)	.36
Log FEV ₁ , prebronchodilator	−0.1906 (−4.171,0.360)	.098	−2.404 (−6.566,1.757)	.251	−0.484 (−5.356,4.387)	.839	−2.458 (−6.215,1.299)	.194
Log FEV ₁ change with bronchodilator	0.54 (−1.352,2.431)	.573	0.672 (−2.997,4.342)	.714	−0.335 (−3.064,2.394)	.802	0.188 (−3.434,3.810)	.917
NAEPP Severity Classification	0.232 (0,0.464)	.05	−0.042 (−0.409,0.325)	.823	0.239 (−0.442,0.919)	.492	0.498 (0.045,0.095)	.031

Notes: ECP, eosinophil cationic protein; FEV₁, forced expiratory volume in 1 second; NAEPP, National Asthma Education and Prevention program, 2007 criteria.
p < .05 are listed in bold.

factors other than classical chemokines may instead contribute to the persistent recruitment of proinflammatory leukocytes during chronic asthma.

Our investigations of extracellular Cyps in nasal wash samples from human asthma patients revealed significant increases in CypB compared to classical chemokines in the asthma group, relative to healthy controls (Figure 1). In fact, the levels of most classical chemokines measured were either comparable between control and asthmatic groups or below the sensitivity threshold of the assay used for detection. One interesting finding was that although levels of CypA were not significantly elevated in the asthma group compared to the control group, CypA showed significant associations with several parameters of the disease (Table 2). Of particular interest is that levels of CypA were negatively associated with a parameter of lung function: as levels of CypA increased, patients' response to bronchodilator drugs decreased. It should be noted that while interesting, the biological and clinical relevance of associations with CypA remains unclear since there was no difference in CypA between the control group and the asthma groups as a whole. Nonetheless, in light of our recent findings in animal models of asthma (24, 45), we believe these interesting associations warrant further investigation.

These associations and findings of elevated levels of Cyps in clinical patients provide the first evidence that Cyps may play a role in human asthma, particularly in the chronic phase of the disease. This may extend beyond leukocyte recruitment to other mediators of asthma such as airway remodeling. For example, we observed that increases in CypA and CypB associated with increased levels of eosinophil (measured by ECP levels) and platelet (measured by P-selectin levels) activation. Studies have demonstrated not only that CypA can be released from activated platelets (46), but also that CypB increases platelet adhesion to collagen (47). Further, platelet activation has also been shown to be important for leukocyte activation and recruitment into tissues (25, 48). Thus, Cyps have the potential to be involved in many aspects of asthma pathology, from leukocyte activation and recruitment, to airway remodeling. This could also extend to lung function, since CypA was also associated with a worse response to bronchodilator drugs. An important caveat to these studies is the relatively small sample size available for our analyses, particularly after subdividing the asthma group into phenotypic clusters. Studies of larger cohorts will allow a more powerful analysis of clinically relevant asthma parameters and could further investigate the role of Cyps in human disease.

CONCLUSIONS/KEY FINDINGS

Extracellular Cyps have been shown to associate with inflammation in a variety of human diseases. Further, recent studies have demonstrated a potential role for Cyps in leukocyte recruitment in animal models of acute (45)

and chronic (24) allergic asthma. This study provides the first evidence that Cyps are elevated in human asthma patients and that increases in levels of Cyps associate with several clinical parameters of disease severity.

ACKNOWLEDGMENTS

This project was conducted through the General Clinical Research Center under the CReFF program at the Children's National Medical Center and supported by the National Institutes of Health National Center for Research Resources, grant M01RR020359, with additional funding support from Sheldon C. Siegel Investigator Award Grant from the Asthma and Allergy Foundation of America (RJF), National Institutes of Health grants K23RR020069 (RJF), P20MD000198 (RJF), and American Heart Association Pre-doctoral Award 0815226E (EJS).

DECLARATION OF INTEREST

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

REFERENCES

1. Handschumacher RE, Harding MW, Rice J, Drugge RJ, Speicher DW. Cyclophilin: a specific cytosolic binding protein for cyclosporin A. *Science* 1984; 226(4674):544–547.
2. Bukrinsky MI. Cyclophilins: unexpected messengers in intercellular communications. *Trends Immunol* 2002; 23(7):323–325.
3. Arora K, Gwinn WM, Bower MA, Watson A, Okwumabua I, MacDonald HR, Bukrinsky MI, Constant SL. Extracellular cyclophilins contribute to the regulation of inflammatory responses. *J Immunol* 2005; 175(1):517–522.
4. Balsley MA, Malesevic M, Stemmy EJ, Gigley J, Jurjus RA, Herzog D, Bukrinsky MI, Fischer G, Constant SL. A cell-impermeable cyclosporine derivative reduces pathology in a mouse model of allergic lung inflammation. *J Immunol* 2010; 185(12):7663–7670.
5. Damsker JM, Bukrinsky MI, Constant SL. Preferential chemotaxis of activated human CD4+ T cells by extracellular cyclophilin A. *J Leukoc Biol* 2007; 82(3):613–618.
6. Gwinn WM, Damsker JM, Falahati R, Okwumabua I, Kelly-Welch A, Keegan AD, Vanpouille C, Lee JJ, Dent LA, Leitenberg D, Bukrinsky MI, Constant SL. Novel approach to inhibit asthma-mediated lung inflammation using anti-CD147 intervention. *J Immunol* 2006; 177(7):4870–4879.
7. Sherry B, Yarlett N, Strupp A, Cerami A. Identification of cyclophilin as a proinflammatory secretory product of lipopolysaccharide-activated macrophages. *Proc Natl Acad Sci U S A* 1992; 89(8):3511–3515.
8. Xu Q, Leiva MC, Fischkoff SA, Handschumacher RE, Lyttle CR. Leukocyte chemotactic activity of cyclophilin. *J Biol Chem* 1992; 267(17):11968–11971.
9. Yurchenko V, Zybarth G, O'Connor M, Dai WW, Franchin G, Hao T, Guo H, Hung HC, Toole B, Gallay P, Sherry B, Bukrinsky M. Active site residues of cyclophilin A are crucial for its signaling activity via CD147. *J Biol Chem* 2002; 277(25):22959–22965.
10. Yurchenko V, Constant S, Bukrinsky M. Dealing with the family: CD147 interactions with cyclophilins. *Immunology* 2006; 117(3):301–309.
11. Billich A, Winkler G, Aschauer H, Rot A, Peichl P. Presence of cyclophilin A in synovial fluids of patients with rheumatoid arthritis. *J Exp Med* 1997; 185(5):975–980.

12. Kim H, Kim WJ, Jeon ST, Koh EM, Cha HS, Ahn KS, Lee WH. Cyclophilin A may contribute to the inflammatory processes in rheumatoid arthritis through induction of matrix degrading enzymes and inflammatory cytokines from macrophages. *Clin Immunol* 2005; 116(3):217–224.
13. Jin ZG, Melaragno MG, Liao DF, Yan C, Haendeler J, Suh YA, Lambeth JD, Berk BC. Cyclophilin A is a secreted growth factor induced by oxidative stress. *Circ Res* 2000; 87(9):789–796.
14. Suzuki J, Jin ZG, Meoli DF, Matoba T, Berk BC. Cyclophilin A is secreted by a vesicular pathway in vascular smooth muscle cells. *Circ Res* 2006; 98(6):811–817.
15. Tegeder I, Schumacher A, John S, Geiger H, Geisslinger G, Bang H, Brune K. Elevated serum cyclophilin levels in patients with severe sepsis. *J Clin Immunol* 1997; 17(5):380–386.
16. Brown JR, Kleimberg J, Marini M, Sun G, Bellini A, Mattoli S. Kinetics of eotaxin expression and its relationship to eosinophil accumulation and activation in bronchial biopsies and bronchoalveolar lavage (BAL) of asthmatic patients after allergen inhalation. *Clin Exp Immunol* 1998; 114(2):137–146.
17. Holgate ST, Bodey KS, Janezic A, Frew AJ, Kaplan AP, Teran LM. Release of RANTES, MIP-1 alpha, and MCP-1 into asthmatic airways following endobronchial allergen challenge. *Am J Respir Crit Care Med* 1997; 156(5):1377–1383.
18. van den Toorn LM, Overbeek SE, de Jongste JC, Leman K, Hoogsteden HC, Prins JB. Airway inflammation is present during clinical remission of atopic asthma. *Am J Respir Crit Care Med* 2001; 164(11):2107–2113.
19. Gern JE, Martin MS, Ankam KA, Shen K, Roberg KA, Carlson-Dakes KT, Adler K, Gilbertson-White S, Hamilton R, Shult PA, Kirk CJ, Da Silva DF, Sund SA, Kosorok MR, Lemanske Jr, RF. Relationships among specific viral pathogens, virus-induced interleukin-8, and respiratory symptoms in infancy. *Pediatr Allergy Immunol* 2002; 13(6):386–393.
20. Hurst JR, Perera WR, Wilkinson TM, Donaldson GC, Wedzicha JA. Systemic and upper and lower airway inflammation at exacerbation of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2006; 173(1):71–78.
21. Legg JP, Hussain IR, Warner JA, Johnston SL, Warner JO. Type 1 and type 2 cytokine imbalance in acute respiratory syncytial virus bronchiolitis. *Am J Respir Crit Care Med* 2003; 168(6):633–639.
22. Midulla F, Tromba V, Lo Russo L, Mileto F, Sabatino G, Sgarrella M, Panuska JR, Manganozzi L, Korn D, Moretti C. Cytokines in the nasal washes of children with respiratory syncytial virus bronchiolitis. *Int J Immunopathol Pharmacol* 2006; 19(1):231–235.
23. Frischer T, Baraldi E. Upper airway sampling. *Am J Respir Crit Care Med* 2000; 162(2 Pt 2):S28–30.
24. Stemmy EJ, Balsley MA, Jurjus RA, Damsker JM, Bukrinsky MI, Constant SL. Blocking cyclophilins in chronic phase of asthma reduces leukocyte persistence and disease reactivation. *Am. J. Respir. Cell Mol. Biol.* 2011; published ahead of print on April 14, 2011 as doi:10.1165/rcmb.2011-0007OC.
25. Benton AS, Kumar N, Lerner J, Wiles AA, Foerster M, Teach SJ, Freishtat RJ. Airway platelet activation is associated with airway eosinophilic inflammation in asthma. *J Invest Med* 2010; 58(8):987–990.
26. Benton AS, Wang Z, Lerner J, Foerster M, Teach SJ, Freishtat RJ. Overcoming heterogeneity in pediatric asthma: tobacco smoke and asthma characteristics within phenotypic clusters in an African American cohort. *J Asthma* 2010; 47(7):728–734.
27. Freishtat RJ, Iqbal SF, Pillai DK, Klein CJ, Ryan LM, Benton AS, Teach SJ. High prevalence of vitamin D deficiency among inner-city African American youth with asthma in Washington, DC. *J Pediatr* 2010; 156(6):948–952.
28. Watson AM, Ngor WM, Gordish-Dressman H, Freishtat RJ, Rose MC. MUC7 polymorphisms are associated with a decreased risk of a diagnosis of asthma in an African American population. *J Invest Med* 2009; 57(8):882–886.
29. Bukstein DA, McGrath MM, Buchner DA, Landgraf J, Goss TF. Evaluation of a short form for measuring health-related quality of life among pediatric asthma patients. *J Allergy Clin Immunol* 2000; 105(2 Pt 1):245–251.
30. National Asthma Education and Prevention Program. Expert Panel Report 3 (EPR-3): guidelines for the diagnosis and management of asthma-summary report 2007. *J Allergy Clin Immunol* 2007; 120 (5 Suppl.):S94–S138.
31. Gorelick MH, Brousseau DC, Stevens MW. Validity and responsiveness of a brief, asthma-specific quality-of-life instrument in children with acute asthma. *Ann Allergy Asthma Immunol* 2004; 92(1): 47–51.
32. Castro-Rodriguez JA, Holberg CJ, Morgan WJ, Wright AL, Martinez FD. Increased incidence of asthma like symptoms in girls who become overweight or obese during the school years. *Am J Respir Crit Care Med* 2001; 163(6):1344–1349.
33. Guerra S, Wright AL, Morgan WJ, Sherrill DL, Holberg CJ, Martinez FD. Persistence of asthma symptoms during adolescence: role of obesity and age at the onset of puberty. *Am J Respir Crit Care Med* 2004; 170(1):78–85.
34. Cookson W. The alliance of genes and environment in asthma and allergy. *Nature* 1999; 402(6760 Suppl.):B5–B11.
35. Cohn L, Elias JA, Chupp GL. Asthma: mechanisms of disease persistence and progression. *Annu Rev Immunol* 2004; 22: 789–815.
36. Corrigan CJ, Hartnell A, Kay AB. T lymphocyte activation in acute severe asthma. *Lancet* 1988; 1(8595):1129–1132.
37. Walker C, Kaegi MK, Braun P, Blaser K. Activated T cells and eosinophilia in bronchoalveolar lavages from subjects with asthma correlated with disease severity. *J Allergy Clin Immunol* 1991; 88(6):935–942.
38. Larche M, Robinson DS, Kay AB. The role of T lymphocytes in the pathogenesis of asthma. *J Allergy Clin Immunol* 2003; 111(3):450–463, quiz 64.
39. Wills-Karp M. Immunologic basis of antigen-induced airway hyperresponsiveness. *Annu Rev Immunol* 1999; 17:255–281.
40. Homer RJ, Elias JA. Consequences of long-term inflammation. Airway remodeling. *Clin Chest Med* 2000; 21(2):331–343, ix.
41. Medoff BD, Thomas SY, Luster AD. T cell trafficking in allergic asthma: the ins and outs. *Annu Rev Immunol* 2008; 26: 205–232.
42. Robertson JM, MacLeod M, Marsden VS, Kappler JW, Marrack P. Not all CD4+ memory T cells are long lived. *Immunol Rev* 2006; 211: 49–57.
43. Rothenberg ME, Hogan SP. The eosinophil. *Annu Rev Immunol* 2006; 24:147–174.
44. Harris NL, Watt V, Ronchese F, Le Gros G. Differential T cell function and fate in lymph node and nonlymphoid tissues. *J Exp Med* 2002; 195(3):317–326.
45. Balsley MA, Malesevic M, Stemmy EJ, Gligley J, Jurjus RA, Herzog D, Bukrinsky MI, Fischer G, Constant SL. A cell-impermeable cyclosporine a derivative reduces pathology in a mouse model of allergic lung inflammation. *J Immunol* 2010; 185(12):7663–7670 [Research Support, N.I.H., Extramural Research Support, Non-US Government].
46. Coppinger JA, Cagney G, Toomey S, Kislinger T, Belton O, McRedmond JP, Cahill DJ, Emili A, Fitzgerald DJ, Maguire PB. Characterization of the proteins released from activated platelets leads to localization of novel platelet proteins in human atherosclerotic lesions. *Blood* 2004; 103(6):2096–2104.
47. Allain F, Durieux S, Denys A, Carpentier M, Spik G. Cyclophilin B binding to platelets supports calcium-dependent adhesion to collagen. *Blood* 1999; 94(3):976–983.
48. Pitchford SC, Momi S, Giannini S, Casali L, Spina D, Page CP, Gresele P. Platelet P-selectin is required for pulmonary eosinophil and lymphocyte recruitment in a murine model of allergic inflammation. *Blood* 2005; 105(5):2074–2081.

Copyright of Journal of Asthma is the property of Taylor & Francis Ltd and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.