

Luminex
complexity simplified.



**Flexible, Intuitive, and
Affordable Cytometry.**

Guava® easyCyte™ Flow Cytometers.

LEARN MORE >



The Transcription Factor PLZF Is Necessary for the Development and Function of Mouse Basophils

This information is current as of August 12, 2019.

Sai Zhang, Joshua A. Vieth, Agata Krzyzanowska, Everett K. Henry, Lisa K. Denzin, Mark C. Siracusa and Derek B. Sant'Angelo

J Immunol published online 31 July 2019
<http://www.jimmunol.org/content/early/2019/07/30/jimmunol.1900068>

Supplementary Material <http://www.jimmunol.org/content/suppl/2019/07/30/jimmunol.1900068.DCSupplemental>

Why *The JI*? [Submit online.](#)

- **Rapid Reviews! 30 days*** from submission to initial decision
- **No Triage!** Every submission reviewed by practicing scientists
- **Fast Publication!** 4 weeks from acceptance to publication

**average*

Subscription Information about subscribing to *The Journal of Immunology* is online at:
<http://jimmunol.org/subscription>

Permissions Submit copyright permission requests at:
<http://www.aai.org/About/Publications/JI/copyright.html>

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://jimmunol.org/alerts>



The Transcription Factor PLZF Is Necessary for the Development and Function of Mouse Basophils

Sai Zhang,^{*,†} Joshua A. Vieth,^{*} Agata Krzyzanowska,^{*,†} Everett K. Henry,[‡]
Lisa K. Denzin,^{*,†,§} Mark C. Siracusa,[‡] and Derek B. Sant'Angelo^{*,†,§}

Basophils are innate immune cells associated with type 2 immunity, allergic reactions, and host defense against parasite infections. In this study, we show that the transcription factor PLZF, which is known for its essential role in the function and development of several innate lymphocyte subsets, is also important for the myeloid-derived basophil lineage. PLZF-deficient mice had decreased numbers of basophil progenitors in the bone marrow and mature basophils in multiple peripheral tissues. Functionally, PLZF-deficient basophils were less responsive to IgE activation and produced reduced amounts of IL-4. The altered function of basophils resulted in a blunted Th2 T cell response to a protein allergen. Additionally, PLZF-deficient basophils had reduced expression of the IL-18 receptor, which impacted migration to lungs. PLZF, therefore, is a major player in controlling type 2 immune responses mediated not only by innate lymphocytes but also by myeloid-derived cells. *The Journal of Immunology*, 2019, 203: 000–000.

Despite representing <1% of circulating leukocytes in the peripheral blood of mammals, basophils have been shown to play multiple roles in the immune response, including regulating innate and the adaptive immune responses, protecting against parasitic infection, and contributing to immunological disorders such as allergy and autoimmunity (1–8). Activated basophils can produce a range of Th2 cytokines, including IL-4, IL-6, and IL-13, that can help promote Th2 cell differentiation. Therefore, basophils impact immune responses such as the clearance of parasites, ectoparasites, and the initiation of allergic responses (9–11).

Like NKT cells, basophils have the unusual feature of continuous transcription of the IL-4 gene (12–14). The continuous presence of “preformed” mRNA for IL-4 presumably plays a role in enabling basophils to rapidly release IL-4 after activation (15–17). Depletion of basophils or disrupting IL-4 production by basophils leads to impaired differentiation of T cells into the Th2 lineage (15, 17) and altered type 2 innate lymphocyte (ILC2) responses (18). Basophils are reported to express MHC class II (MHCII) as well as

the costimulation molecules CD40, CD80, and CD86 and have been shown to interact with lymphocytes (19). Therefore, basophils might also influence lymphocyte activation via a contact-dependent mechanism (20–22). Beyond activating ILC2s and T cells, basophils also express IL13, BAFF, and a proliferation-inducing ligand (APRIL), which can lead to B cell activation and class-switch recombination to produce IgD, IgG, IgA, and IgE (23–27).

Basophils primarily mature in the bone marrow before they enter the periphery, where they have a life span of only ~60 h (28). Basophils are currently reported to arise from the common myeloid progenitor, which gives rise to granulocyte/macrophage progenitors (GMPs). GMPs are reported to have the potential to differentiate into neutrophils, eosinophils, basophils, and mast cells. Along the basophil/mast cell lineage, GMPs develop into the prebasophil and mast cell progenitors (MCPs) in the bone marrow or, alternatively, into basophil-MCPs in the spleen (29, 30). Both prebasophil and MCPs and basophil-MCPs can differentiate into basophil progenitors (BaPs) and MCPs, which ultimately differentiate into mature basophils and mast cells, respectively (31).

Studies have demonstrated that basophil development is dependent on the expression of C/EBP α -enhancer-binding proteins C/EBP α and GATA2. GATA2 expression is induced by IRF8. The ordered expression of these two proteins determines whether GMPs develop into basophils, mast cells, or eosinophils (32, 33). GMPs that first downregulate and then upregulate C/EBP α expression whereas remaining high for GATA2 expression will differentiate into BaPs. In contrast, GMPs will differentiate into eosinophils if C/EBP α expression remains high throughout development, but into MCPs if C/EBP α remains low (8). The commitment to the mast cell lineage is also thought to rely on the transcription factor MITF (34, 35). C/EBP α and MITF suppress each other's expression, which secures the basophil and mast cell lineages (29).

The broad-complex, tramtrack and bric-à-brac, zinc finger (BTB-ZF) genes are a family of transcription factors that are characterized by an N-terminal BTB domain that is involved in protein–protein interactions and a domain with C-terminal Krüppel-type zinc fingers enabling binding of DNA (36–38). It is now appreciated that BTB-ZF transcription factors play key roles for the development and/or function of multiple immune

^{*}Child Health Institute of New Jersey, Rutgers Robert Wood Johnson Medical School, New Brunswick, NJ 08901; [†]Rutgers Graduate School of Biomedical Sciences, Piscataway, NJ 08854; [‡]Department of Medicine, Center for Immunity and Inflammation, Rutgers New Jersey Medical School, Newark, NJ 07103; and [§]Department of Pediatrics, Rutgers Robert Wood Johnson Medical School, New Brunswick, NJ 08901

ORCID: 0000-0001-8756-2759 (J.A.V.); 0000-0002-3884-2612 (E.K.H.); 0000-0002-5510-9367 (D.B.S.).

Received for publication January 22, 2019. Accepted for publication July 4, 2019.

This work was supported by National Institutes of Health/National Institute of Allergy and Infectious Diseases Grants R01 AI083988 and AI059739 (to D.B.S.), Robert Wood Johnson Foundation Grant 67038 (to the Child Health Institute of New Jersey), the Dr. James J. O'Connell Fund (to S.Z.), and a New Jersey Health Foundation Excellence in Research Award (to D.B.S.).

Address correspondence and reprint requests to Dr. Derek B. Sant'Angelo, Rutgers Robert Wood Johnson Medical School, 89 French Street, New Brunswick, NJ 08901. E-mail address: santandb@rwjms.rutgers.edu

The online version of this article contains supplemental material.

Abbreviations used in this article: BaP, basophil progenitor; BTB-ZF, broad-complex, tramtrack and bric-à-brac, zinc finger; GMP, granulocyte/macrophage progenitor; HI, heat-inactivated; ILC2, type 2 innate lymphocyte; MAIT, mucosal-associated invariant T; MCP, mast cell progenitor; MFI, mean fluorescence intensity; MHCII, MHC class II; PLZF, promyelocytic leukemia zinc finger.

Copyright © 2019 by The American Association of Immunologists, Inc. 0022-1767/19/\$37.50

cell lineages. For example, the BTB-ZF transcription factor, promyelocytic leukemia zinc finger (PLZF; *zbtb16*) is essential for the normal function of several types of innate lymphoid-derived cells, including $\alpha\beta$ NKT and $\gamma\delta$ NKT cells (39–42), mucosal-associated invariant T (MAIT) cells (43), and ILC2 (44–46). In the absence of PLZF, NKT cells do not acquire their characteristic activated phenotype and fail to produce cytokines upon primary activation (41, 42). Instead, these cells require secondary stimulation to be fully activated and carry out effector functions, making them similar to conventional T cells (41). In humans, in addition to NKT and MAIT cells, PLZF is expressed in a population of CD161^{hi} CD4 and CD8 T cells, all $\gamma\delta$ T cells, and all NK cells (47). Blood samples from a PLZF-deficient person showed that there was an impact on the frequency and/or phenotype of all of these lymphocyte populations (47).

Ectopic expression of PLZF spontaneously induces conventional T cells to be more sensitive to antigenic stimulation and able to produce proinflammatory cytokines, such as IL-17, INF- γ , and TNF, upon primary activation (48–50). Tight regulation of endogenous PLZF expression would, therefore, appear important to prevent aberrant T cell responses that might lead to allergy or autoimmune diseases. Consistent with this, our studies showed that PLZF expression among lymphocytes is restricted to innate-like cells and that it cannot be induced in noninnate T cells (51). Strong TCR signaling in developing thymocytes or in mature T cells, for example, does not induce PLZF (51), and expression in noninnate lymphocytes was also not found in response to inflammatory conditions resulting from bacterial infection. Analysis of the chromatin structure of *Zbtb16* (the gene encoding PLZF) using chromatin immunoprecipitation for H3K4me3 and H3K27me3 also suggests that PLZF expression is suppressed in Th1, Th2, Th17, regulatory T cells, and naive CD4 T cells (51, 52). Repression of expression of PLZF was recently reported to be dependent on Bcl11b and occurs early during thymocyte development (53).

The specificity and tight control of PLZF expression in lymphocytes suggests that other immune cells expressing PLZF might also be dependent on the transcription factor for essential effector functions. To explore this possibility, we examined PLZF expression within the myeloid compartment. Basophils, we find, express PLZF both early in development and as mature cells. Genetic deletion of PLZF showed that the transcription factor plays a role during development and, most interestingly, for the full elaboration of basophil effector functions. PLZF-deficient basophils fail to fully respond to a protein allergen, papain, resulting in blunted Th2 CD4 T cell responses. Therefore, PLZF is important for type 2 responses by directly controlling the function of innate lymphocytes and also, indirectly, through the control of basophil functions.

Materials and Methods

General experiment design and statistical analysis

This study was designed to study the role of the transcription factor PLZF in basophil development and function in Th2 immunity. Age-matched (8–12-wk-old) wild type (WT) and PLZF knockout (KO) mice were used for comparisons in this study. Basophil frequency in different mouse tissues was analyzed. We used both an *in vivo* and an *in vitro* system to study basophil functions. Each experiment contained at least three biological replicates. Data are shown as the mean $\pm$ SEM. Statistical analyses were performed using GraphPad Prism 10 software. All samples were analyzed using an unpaired, two-tailed Student *t* test. A *p* value <0.05 was considered significant.

Mice

Rag1 KO, CD1d KO, IL-18R KO, PLZF-IRES-GFP-Cre, and *Rosa26*-tdTomato mice were purchased from The Jackson Laboratory. The PLZF-eGFP (PEG) reporter and PLZF-Cre mouse lines were generated in our laboratory as previously described (51, 54). PLZF-deficient mice were

previously described (55). All mouse strains were bred and maintained in the animal facility of the Child Health Institute of New Jersey. All experimental protocols and procedures were approved by the Institutional Animal Care and Use Committee of the Child Health Institute of New Jersey. Animal care and experimental procedures were carried out in accordance with the guidelines of the Institutional Animal Care and Use Committee of the Child Health Institute of New Jersey and the National Institutes of Health *Guide for the Care and Use of Laboratory Animals*.

Flow cytometry

Cells were harvested from the tissues and preincubated with the anti-Fc γ R mAb and normal mouse serum to block nonspecific Ab binding for 15 min at 4°C. Surface staining was performed in FACS buffer (PBS with 1% heat-inactivated [HI] FBS) for 25 min at 4°C using the indicated surface Abs. Dead cells were excluded by DAPI staining, and doublet events were eliminated by comparing the forward scatter width and forward scatter height to the side scatter width and side scatter height. IL-4 intracellular staining was performed using the BD Fixation/Permeabilization Solution Kit (BD Biosciences). Intracellular staining of PLZF was performed by fixing cells in 4% paraformaldehyde in PBS for 30 min at 4°C. Cells were then permeabilized for 15 min at 4°C in commercially-available permeabilization buffer (Invitrogen), and incubated with anti-PLZF Ab in permeabilization buffer for 1 h at 4°C. Samples were washed three times with permeabilization buffer and one time with FACS buffer prior to analysis. Events were acquired on an LSRII cytometer (BD Biosciences), and the data were analyzed with the FlowJo software.

Abs

The following Ab clones were used: anti-CD3 (500A2), anti-CD4 (RM45), anti-CD19 (1D3), anti-CD49b (DX5), anti-CD34 (RAM34), anti-CD117 (ACK2), anti-CD200R (OX110), anti-CD40L (MR1), anti-CD62L (MEL-14), anti-CCR2 (SA203G11), anti-CCR7 (4B12), anti-CCR9 (CW-1.2), anti-MHCII (212.A1), anti-ST2 (RMST2-2), IL-18R α (P3TUNYA), anti-Fc ϵ R1 α (MAR-1), anti-PLZF (Mags.21F7), anti-GATA3 (TWAJ), and anti-IL-4 (11b11). Abs were used with different fluorochrome conjugations: FITC, PE, PerCP-Cy5.5, PE-Cy7, allophycocyanin, allophycocyanin-Cy7, Alexa Fluor 700, and Pacific Blue.

Quantitative RT-PCR

Total cellular RNA from cells was isolated by a mirVana miRNA Isolation Kit (Thermo Fisher Scientific). cDNA was generated with reverse transcription using oligo-dT, random primers, and reverse transcriptase (Promega). Quantitative PCR of the cDNA was performed on a QuantStudio 6 Flex Real-Time PCR System (Thermo Fisher Scientific) and the following TaqMan gene expression primer sets (Thermo Fisher Scientific): *Hprt* (Mm03024075_m1) and *zbtb16* (Mm01176868_m1). Gene expression was analyzed using *Hprt* as an endogenous control in each sample.

In vivo basophil activation

Mice were immunized s.c. in the rear footpad with 50 μ g of papain (Calbiochem) dissolved in sterile PBS. As control, the same amount of HI papain was injected into the other side of the footpad. To heat inactivate, papain was dissolved in PBS and placed in boiled water for 10 min. Mice were sacrificed either 3 d (for basophil staining) or 4 d (for GATA3 staining in T cells) after injection and analyzed by flow cytometry.

In vitro basophil activation

One hundred microliters of peripheral blood was drawn from the mice and mixed with 100 μ l of RPMI 1640. The cells were activated with different concentrations of IgE or PMA (50 ng/ml)/ionomycin (500 ng/ml) for 3 h total. Brefeldin A was added at 1 h after the start of the activation. After activation, the cells were washed two times with PBS and spun at 500 g for 5 min, then the cells were treated with RBC lysis buffer to remove RBCs. The cells were then blocked and stained with surface markers, followed by intracellular cytokine staining.

Results

Mapping PLZF expression during the development of myeloid lineage cells

Using our previously described PEG reporter mice (51, 54, 56), we evaluated PLZF expression during early myeloid lineage development. In contrast to PLZF expression in lymphocytes, which has been shown to be largely restricted to specific cell types (51),

we found expression of PLZF in many myeloid precursor populations (Fig. 1). As indicated by fold change as compared with matched WT cells (Fig. 1) and as shown in FACS histograms (Supplemental Fig. 1), PLZF expression levels ranged substantially depending on the subset examined, which were identified as shown in Supplemental Figs. 2–4 (30, 57–60). Expression patterns of PLZF in the different populations were consistent from mouse to mouse (Supplemental Fig. 1). Low levels of expression of the transcription factor, as defined by eGFP expression, were found in common myeloid progenitors, MCPs, and GMPs (Fig. 1). Eosinophil progenitors also expressed low levels of PLZF; however, macrophage/DC precursors and common monocyte precursors were negative. The most pronounced expression of PLZF was in mast cells and basophils as well as the immediate precursors of basophils, BaP. BaP expressed higher levels of PLZF as compared with mature basophils. Downregulation of PLZF as basophils matured is similar to what has previously been observed during NKT cell development (41).

We chose to focus on a potential role for PLZF in basophils because these cells are known to rapidly express IL-4 similarly to NKT cells. The burst of IL-4 produced by NKT cells shortly after activation is completely dependent upon PLZF (41, 42). We found that PLZF was expressed in all basophils in the spleen, peripheral blood, bone marrow, and lung (Fig. 2A). Expression levels were, however, lower than that found in NKT cells (Fig. 2B). PLZF expression was distinctly higher in BaPs in the bone as compared with mature bone marrow basophils (Fig. 2C). Importantly, intracellular

staining with anti-PLZF Ab verified protein expression in basophils (Fig. 2D). The increased concentrations of paraformaldehyde that we found were necessary for reliable retention of proteins in the permeabilized basophils, however, resulted in reduced signal intensity both in the basophils and NKT cells that were stained via the same protocol. Therefore, the conclusions concerning the absolute amounts of PLZF protein in the cells should not be drawn.

To further confirm PLZF expression in basophils, we carried out “fate-mapping” experiments. We used both our PLZF-Cre BAC transgenic mice (PCre), as well as PLZF-IRES-GFPcre knock-in mice (44). These two strains were both crossed to mice that have an allele of Rosa 26 modified with a gene that encodes a fluorescent protein, either ZsGreen or tdTomato, that is preceded by a loxP-flanked STOP cassette. In these mice, Cre expression from the PLZF BAC or from the knock-in allele, even if it is transient, will permanently label cells with the fluorescent protein. Both mouse lines showed uniform expression of the fluorescent protein in basophils from the bone marrow and spleen (Fig. 2E). It should be noted that direct expression of GFP in basophils from the PLZF-IRES-GFP/Cre knock-in construct was not detectable. This, presumably, was due either to lower expression levels of GFP from the knock-in construct as compared with the BAC transgene or, possibly, alterations in expression resulting from the insertion of the IRES-GFP/Cre cassette in the 3' UTR of the *zbtb16* gene. Indeed, microRNA target sites within the 3' UTR have been shown to impact PLZF expression (61). Importantly, we found that use of the floxed-stop ROSA26 ZsGreen allele proved to give results that




FIGURE 1. PLZF expression during myeloid lineage cell development. Myeloid lineage cells were stained and identified in bone marrow from PLZF-eGFP reporter mice (PEG) as shown in Supplemental Figs. 2 and 3. PLZF expression, as defined by eGFP expression, was determined by comparison of the identical population obtained from WT mice (Supplemental Fig. 1). The progression of development was based on publications, as indicated in the text. Color coding in the figure represents MFI values for each subset of cells, based on eGFP expression (MFI of eGFP signal divided by the background fluorescence of the identical cell population from a matched control analyzed in the same experiment). Fold change of MFI key is shown in figure.




FIGURE 2. PLZF is expressed in basophils. **(A)** eGFP expression in basophils from spleen, bone marrow, blood, and lung of PEG mice. Basophils were identified as DAPI⁻CD3⁻CD19⁻FcεRI⁺CD49b⁺ cells. **(B)** Comparison of eGFP expression in basophils, and NKT cells. WT basophils are shown as a fluorescence control. **(C)** Intracellular staining of PLZF in splenic basophils, with CD8 T cells and NKT cells as negative and positive controls, respectively. Quantification of PLZF MFI (fold change over CD8 T cells) is shown ($n = 3$ biological replicates from two individual experiments). **(D)** eGFP expression in BaPs in the bone marrow in PEG mice. BaPs were identified as DAPI⁻CD3⁻CD19⁻CD34⁺FcεRI⁺CD49b⁺cKit⁻ cells. **(E)** ZsGreen and tdTomato expression in basophils from the bone marrow and spleen in Pcre × ZsGreen and PLZF-IRES-GFPCre × tdTomato mice, respectively. Representative FACS plots from one of three independent experiments are shown. * $p < 0.05$, ** $p < 0.01$, determined by Student t test.

are more consistent with known PLZF expression patterns as compared with the previously reported floxed-stop ROSA26 tdTomato allele (44, 62). The mice carrying the ZsGreen version of the allele showed that, as expected, nearly 100% of NKT cells in the thymus and spleen were labeled but that the previously

reported and unexpected labeling of thymocytes, B cells, and T cells was eliminated or greatly reduced (Supplemental Fig. 4). For example, B cells and double positive thymocytes were essentially negative for ZsGreen expression, indicating that “leaky” expression of the PLZF-Cre in hematopoietic stem cells, as previously




FIGURE 3. Reduced numbers of basophils in PLZF-deficient mice. **(A)** Basophil numbers in the bone marrow, spleen, lung, and blood of PLZF-deficient and WT mice. Cells shown are DAPI⁻CD3⁻CD19⁻CD34⁻ population, and basophils were identified as CD49b⁺FCεR1⁺. Numbers indicate the percentage of events within the gated area. Percentage and absolute number of basophils are shown in scatter plot graphs. Each symbol indicates a single mouse ($n = 14$ with ~5 independent experiments for each tissue). **(B)** Basophil progenitor numbers in the bone marrow of PLZF-deficient and WT mice. Cells shown are DAPI⁻CD3⁻CD19⁻CD34⁺ population. Numbers in dot plots show the percentage of events in the gated region. Percentage and absolute number of basophil precursors are shown in scatter plot graphs. Each symbol indicates a single mouse ($n = 3$ with three independent experiments). The horizontal lines indicate the mean ($\pm$ SEM). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ determined by Student t test.

suggested, was unlikely. Overall, these data demonstrate that PLZF is expressed both in BaPs in the bone marrow and in mature basophils in multiple mouse tissues.

PLZF-deficient mice have reduced numbers of basophils

The expression of PLZF in BaPs and mature basophils raised the possibility that, like NKT cells, basophils require the transcription factor for development. To study this, we compared the frequency and absolute numbers of basophils in C57BL/6 and age-matched PLZF-deficient mice. FACS analysis was done by first excluding lymphocytes by use of anti-CD3 and anti-CD19 and then identifying basophils by expression of CD49b and FcεRI (Fig. 3A). PLZF deficiency resulted in an ~50–60% decrease of basophils in multiple tissues, including the bone marrow, spleen, lung, and peripheral

blood (Fig. 3A). The absolute numbers of basophils were reduced by a similar degree. Consistent with the reduced number of basophils in all tissues, the BaP population in the bone marrow showed an ~70% decrease as compared with age-matched WT mice (Fig. 3B). Therefore, similar to NKT cells, albeit to a lesser degree, PLZF-deficient mice have reduced frequencies of basophils. Unlike NKT cells, PLZF deficiency results in the reduction of basophils that is found in all tissues. Combined, these data suggest that PLZF has a direct impact on basophil development.

Papain-induced basophil migration is impaired in PLZF-deficient mice

Proteolytic activities that are associated with some allergens and parasite secretions are thought to be directly involved in the




FIGURE 4. Papain-induced basophil-mediated Th2 responses were decreased in PLZF KO mice. **(A)** Fifty micrograms of papain was injected into the footpads of PLZF KO and WT mice. The same amount of HI papain was injected into the other footpad as control. Three days after injection, the draining lymph nodes were removed and stained for basophils. Cells shown were gated are DAPI⁺CD3⁺CD19⁺. Frequency of basophils identified as CD49b⁺FcεRI⁺ is shown in the dot plot and for multiple experiments in the scatter plot graphs. Each symbol indicates a single mouse ($n = 8$, with three to five independent experiments). **(B)** and **(C)** Fifty micrograms of papain or HI papain was injected into **(B)** RAG1 KO ($n = 3$, three independent experiments) or **(C)** CD1d KO ($n = 5$, three independent experiments) mice and WT controls. Basophil frequency was determined as in **(A)**. The horizontal lines in the scatter plots indicate the mean ($\pm$ SEM). * $p < 0.05$, ** $p < 0.01$ determined by Student t test. n.s., not significant.

activation of basophils (63–66). An excellent model of this type of immune response is the s.c. injection of the cysteine protease, papain. We injected 50 μ g of papain into one rear footpad of mice and the same amount of HI papain into the other footpad as negative control. Three days after injection, the draining lymph nodes were removed and analyzed by FACS. Papain injection resulted in a basophil response that comprised $\sim 0.6\%$ of the $CD3^+$, $CD19^-$ cells in the draining lymph nodes (Fig. 4A, top). In contrast, very few basophils were recruited to the lymph nodes in response to the allergen in PLZF-deficient mice (Fig. 4A, bottom). Although the number of basophils present in PLZF-deficient mice is reduced, the near complete absence of the cells following treatment implies that a migration deficiency is in effect. HI papain did not result in basophil recruitment in either mouse strain.

Basophil responses (both their population expansion and recruitment) to most stimuli, including helminth, parasite, and papain, are dependent on IL-3–IL-3R signaling (67–69). Further, T cells are reported to be the main source of IL-3 required for basophil responses to be initiated (70). Consistent with these data, basophil activation and recruitment are disrupted in lymphocyte-deficient mice as a result of IL-3 deficiency (70). In addition, injection of RAG1-deficient mice with papain resulted in limited basophil migration to the draining lymph nodes (Fig. 4B). This result raised the possibility that the failure of basophil migration was because of loss of functional NKT cells in the PLZF-deficient mice. We tested this by injecting CD1d-deficient mice, which are devoid of invariant NKT cells, with papain. The results showed that there was no difference in the frequency of recruited basophils between CD1d KO and WT mice (Fig. 4C). Combined, these data show that PLZF expression in basophils is important for their response to an allergic stimulus.

After papain injection, naive $CD4^+$ T cells in the lymph nodes have been shown to respond to the IL-4 produced by basophils and undergo Th2 differentiation, which requires the transcription factor GATA3. Therefore, the frequency of $GATA3^+$ T cells indirectly reflects the extent of the Th2 response mediated by basophils (17). Results showed that in WT mice $\sim 6\%$ of T cells in the lymph nodes expressed GATA3 after papain immunization (Fig. 5A). The frequency of $GATA3^+$ $CD4^+$ T cells in the draining lymph nodes following papain injection of PLZF-deficient mice was, however, reduced by nearly half (Fig. 5A). PLZF-expressing

NKT cells do not play a role in this Th2 differentiation response because there was no difference in GATA3-positive $CD4^+$ T cells in the lymph nodes of CD1d KO mice following papain injection (Fig. 5B).

Functional analysis of basophils in PLZF-deficient mice *in vitro*


The reduced basophil-mediated Th2 response in PLZF-deficient mice was a consequence of either the decreased number of basophils or, alternatively, reduced basophil effector functions. We considered the possibility that, like NKT cells, PLZF deficiency in basophils might result in disrupted IL-4 production. To test this, peripheral blood from mice was activated *in vitro* with either IgE or PMA/ionomycin. Intracellular staining for cytokines in basophils was done 3 h after activation. Incubation with 0.05 ng/ml of IgE resulted in activation of basophils as measured by upregulation of CD200R expression (Fig. 6A). Approximately 15% of WT basophils produced IL-4 under these conditions. In contrast, only $\sim 7\%$ of PLZF-deficient basophils produced IL-4 (Fig. 6A). Next, we used titrated concentrations of IgE to determine whether the sensitivity of PLZF-deficient basophils to activating signals was reduced. The percentage of responding basophils that was deficient for PLZF was significantly reduced at nearly every concentration of IgE. Furthermore, we found that the dose of IgE needed to activate PLZF-deficient basophils was ~ 100 times higher than that needed for WT cells (Fig. 6B).

PMA/ionomycin was used to determine the maximal IL-4 production capacity of WT and PLZF-deficient basophils. Fewer PLZF KO basophils responded, as measured by intracellular staining of IL-4 (Fig. 6C). The mean fluorescence intensity (MFI) of IL-4 was also determined as a measure of the amount of cytokine produced by the responding cells and was found to be ~ 2 -fold lower (Fig. 6D). Overall, these data show that in the absence of PLZF, fewer basophils produced IL-4, and those that did produce the cytokine made less.

PLZF-deficient basophils have reduced IL-18R expression

To investigate the potential mechanism for reduced migration of PLZF-deficient basophils after papain immunization, we stained for various cytokine and chemokine receptors on basophils. Loss of PLZF did not impact the expression levels of CD62L, CCR7, or

FIGURE 5. Differentiation of Th2 T cells is reduced in the absence of PLZF. **(A)** Fifty micrograms of papain was injected into the footpads of WT or PLZF KO mice. Four days after papain injection, the draining lymph nodes were removed. Th2 differentiation, as measured by intracellular staining for GATA3 in $CD3^+$ $CD4^+$ T cells, with the percentage positive cells shown in the dot plots. Scatter plots show the frequency of $GATA3^+$ $CD4^+$ T cells ($n = 5$ and 4 independent experiments). **(B)** Fifty micrograms of papain was injected into the footpads of WT or CD1d KO mice. The frequency of $GATA3^+$ $CD4^+$ T cells is shown in the dot plots. FACS is representative of three independent experiments. The horizontal lines in the scatter plots indicate the mean ($\pm$ SEM). $**p < 0.01$ determined by Student *t* test. n.s., not significant.






FIGURE 6. Function of PLZF-deficient basophils. **(A)** Peripheral blood from PLZF KO and WT mice was activated with 0.05 ng/ml of IgE for 3 h. Brefeldin A was added for the last 2 h of activation. The cells were made permeable and stained for intracellular IL-4. CD200R was analyzed to measure activation. Cells shown are CD3⁺CD19⁺FcεRI⁺CD49b⁺. Representative FACS plots from one of four independent experiments are shown. **(B)** Peripheral blood from WT or PLZF KO mice was activated with increasing concentrations of IgE for 3 h. The percentage of basophils expressing IL-4 is shown. **(C)** Peripheral blood samples from PLZF KO and WT mice were activated with PMA/ionomycin for 3 h. Brefeldin A was added for the last 2 h of activation. The cells were then stained for intracellular IL-4. Basophils were identified as the CD3⁺CD19⁺FcεRI⁺CD49b⁺ population. The scatter plot shows the frequency of basophils that produced IL-4 under these conditions. **(D)** Fold change of IL-4 MFI of the cytokine-producing basophils from (C), normalized to WT ($n = 10$ with five independent experiments). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.0001$ determined by Student t test.

CCR9, all of which have been implicated in basophil migration (Fig. 7A) (17, 22, 71, 72). There also were no differences in the expression of other cell surface proteins, such as ST2, MHCII, or

CD200R, in the absence of PLZF (data not shown). Interestingly, however, PLZF-deficient basophils were found to express lower levels of the IL-18 receptor (Fig. 7B). These differences were




FIGURE 7. PLZF impacts IL-18R expression on basophils and their migration to the lung. **(A)** Expression of the indicated proteins was not altered in PLZF-deficient spleen basophils. **(B)** IL-18R expression on basophils from spleen, lung, and bone marrow was reduced on PLZF-deficient basophils. Representative FACS plots from one of four independent experiments is shown. The fold change on basophils from each of these tissues is shown in the scatter plot ($n = 8$ with four independent experiments). **(C)** IL-18R expression is shown for spleen T cells ($CD3^+CD8^+$ or $CD3^+CD4^+$) taken from WT mice (gray filled) or from mice ectopically expressing PLZF in all T cells. **(D)** The frequency of basophils in the lung, spleen, and bone marrow from IL-18R KO and WT mice ($n = 7$ – 12 mice with three to five independent experiments). $**p < 0.01$, $***p < 0.0001$ determined by Student t test.

relatively small but were highly consistent in multiple tissues, including the spleen, lung, and bone marrow (Fig. 7B). Altered expression of IL-18R is consistent with data showing that PLZF binds to and, possibly, directly regulates expression of *Il18r1* (73, 74). Furthermore, ectopic expression of PLZF in conventional T cells resulted in upregulation of IL-18R expression (Fig. 7C).

IL-18R-deficient mice were analyzed to determine whether altered expression of this receptor impacted basophil migration. The frequency (Fig. 7D) and phenotype of the cells from the spleen and bone marrow was not notably altered. In the lung, however, there was an ~30% decrease in the frequency of basophils. Therefore, PLZF control of IL-18R may influence the ability of basophils to migrate to some tissues.

Together, our data shows that expression of PLZF during myeloid lineage development is necessary for full population of the basophil compartment. Furthermore, PLZF-deficient basophils display decreased migration, reduced IL-4 production upon activation, and lower surface levels of IL-18R, indicating that PLZF plays an important role in basophil trafficking and function.

Discussion

The importance of the expression of PLZF in innate-like lymphocytes, such as NKT cells, V γ 1.1; V δ 6.3 $\gamma\delta$ NKT cells, ILCs, and MAIT cells, has been well established (75). Less is known, however, about the transcription factor's function in myeloid lineage subsets. PLZF has been shown to impact the early development of the myeloid lineage cells in humans (76), but its expression and potential function in mature myeloid lineage cell types is largely unexplored. In this study, we found that PLZF is expressed in basophils, where it plays an important role both their development and function.

PLZF was found to be expressed in basophils from all tissues, including the bone marrow, spleen, lung, and peripheral blood. As during NKT cell development, PLZF was downregulated as BaPs matured in the bone marrow. In the absence of PLZF, we found an ~50–60% decrease of the cells. More interestingly, PLZF-deficient basophils proved to have a reduced capacity to migrate and produce IL-4. Nearly 100 times the dose of IgE was required to induce IL-4 release in the absence of PLZF. Furthermore, maximal IL-4 production, as determined by activation with PMA/ionomycin, was reduced. The reduced IL-4 release, therefore, correlated with a diminished Th2 immune response to a protein allergen. It has been reported that IL-13 production by PLZF-deficient ILC2 harvested from the lungs of mice treated with papain for 3 d produce somewhat less IL-13 (45). Therefore, it is possible that reduced function of IL2s in the absence of PLZF also impacts Th2 T cell differentiation. Overall, the data suggest that PLZF impacts the lymphocyte contribution in type 2 immune responses both by cell-intrinsic and cell-extrinsic means.

In humans with fatal asthma, there often is an increased infiltration of basophils into the lung (77). In mouse asthma models, basophils have been shown to mediate early phase airway obstruction (78). Therefore, the control of basophil migration and activation by PLZF might play a role in the pathogenesis of asthma. In support of this, our results showed that PLZF-deficient basophils have reduced levels of IL-18R and that loss of this receptor impacted basophil migration to the lung. IL-18 is a pleiotropic cytokine produced by macrophages, Kupffer cells, as well as epithelial cells such as keratinocytes (79, 80). Previous studies have shown that IL-18 combined with IL-3 stimulates large amounts of IL-4 and IL-13 production by basophils in vitro (79). Injection of IL-18 in mice has been shown to increase IL-4 production by basophils, recruit Ag-induced eosinophils, and increase Th2 cytokines in the lung (12, 81, 82). Recently, it was shown that IL-18 signaling is necessary for the response of

PLZF-expressing NKT cells to influenza infection in the lung (83). Perhaps, therefore, activation by IL-18 signaling is a common feature of lung NKT cells and basophils. This interpretation may, however, not apply to humans, in which it has been shown that, although basophils express high levels of IL18R in vitro, they do not obviously respond to IL-18 (84).

In the absence of PLZF, basophil numbers were significantly decreased. PLZF has been shown to regulate development of human cord blood-derived myeloid progenitor cells by transcriptional modulation of key transcription factors, including GFI-1, C/EBP α , LEF-1, DUSP6, and ID2 (38, 85). The maturation of basophils also requires C/EBP α , suggesting a possible mechanism for the control of basophil development by PLZF.

Collectively, our data showed that the transcription factor PLZF regulates the development and function of basophils. Although, there is not a complete loss of basophil effector functions, the reduction in IL-4 production in response to a protein allergen is sufficient to blunt Th2 immunity. With a cell-intrinsic role in controlling the NKT cells, ILC2, and now a cell-extrinsic role that impacts Th2 cell development, PLZF is emerging as a major regulator of the overall type 2 immune response. Interestingly, data show that variations in the expression levels of PLZF have major impacts on cell function (62, 86). Therefore, subtle variations in PLZF expression in the human population could be associated with the efficiency of clearing various pathogens and parasites. In addition, the control of IL-18R expression, which we show influences migration of basophils to the lung, implies that variations in PLZF expression might also impact susceptibility to asthma. Overall, our results further elucidate the transcriptional regulation of basophil functions, which helps for better understanding of the role of basophils in the prevention and control of disease.

Acknowledgments

We thank A. Rabson for insightful discussions, L. Osorio for technical assistance, and A. Roberts for cell sorting.

Disclosures



The authors have no financial conflicts of interest.

References


- Schroeder, J. T. 2009. Basophils beyond effector cells of allergic inflammation. *Adv. Immunol.* 101: 123–161.
- Mack, M., M. A. Schneider, C. Moll, J. Cihak, H. Brühl, J. W. Ellwart, M. P. Hogarth, M. Stangassinger, and D. Schlöndorff. 2005. Identification of antigen-capturing cells as basophils. *J. Immunol.* 174: 735–741.
- Kubo, M. 2018. Mast cells and basophils in allergic inflammation. *Curr. Opin. Immunol.* 54: 74–79.
- Rigoni, A., M. P. Colombo, and C. Pucillo. 2018. Mast cells, basophils and eosinophils: from allergy to cancer. *Semin. Immunol.* 35: 29–34.
- Siracusa, M. C., B. S. Kim, J. M. Spergel, and D. Artis. 2013. Basophils and allergic inflammation. *J. Allergy Clin. Immunol.* 132: 789–801; quiz 788.
- Karasuyama, H., K. Mukai, K. Obata, Y. Tsujimura, and T. Wada. 2011. Non-redundant roles of basophils in immunity. *Annu. Rev. Immunol.* 29: 45–69.
- Min, B. 2008. Basophils: what they 'can do' versus what they 'actually do'. *Nat. Immunol.* 9: 1333–1339.
- Siracusa, M. C., J. G. Perrigoue, M. R. Comeau, and D. Artis. 2010. New paradigms in basophil development, regulation and function. *Immunol. Cell Biol.* 88: 275–284.
- Falcone, F. H., H. Haas, and B. F. Gibbs. 2000. The human basophil: a new appreciation of its role in immune responses. *Blood* 96: 4028–4038.
- Sokol, C. L., and R. Medzhitov. 2010. Role of basophils in the initiation of Th2 responses. *Curr. Opin. Immunol.* 22: 73–77.
- Min, B., M. A. Brown, and G. Legros. 2012. Understanding the roles of basophils: breaking dawn. *Immunology* 135: 192–197.
- Mohrs, K., A. E. Wakil, N. Killeen, R. M. Locksley, and M. Mohrs. 2005. A two-step process for cytokine production revealed by IL-4 dual-reporter mice. *Immunity* 23: 419–429.
- Min, B., M. Prout, J. Hu-Li, J. Zhu, D. Jankovic, E. S. Morgan, J. F. Urban, Jr., A. M. Dvorak, F. D. Finkelman, G. LeGros, and W. E. Paul. 2004. Basophils produce IL-4 and accumulate in tissues after infection with a Th2-inducing parasite. *J. Exp. Med.* 200: 507–517.

14. Gessner, A., K. Mohrs, and M. Mohrs. 2005. Mast cells, basophils, and eosinophils acquire constitutive IL-4 and IL-13 transcripts during lineage differentiation that are sufficient for rapid cytokine production. *J. Immunol.* 174: 1063–1072.
15. Oh, K., T. Shen, G. Le Gros, and B. Min. 2007. Induction of Th2 type immunity in a mouse system reveals a novel immunoregulatory role of basophils. *Blood* 109: 2921–2927.
16. Hida, S., M. Tadachi, T. Saito, and S. Taki. 2005. Negative control of basophil expansion by IRF-2 critical for the regulation of Th1/Th2 balance. *Blood* 106: 2011–2017.
17. Sokol, C. L., G. M. Barton, A. G. Farr, and R. Medzhitov. 2008. A mechanism for the initiation of allergen-induced T helper type 2 responses. *Nat. Immunol.* 9: 310–318.
18. Kim, B. S., K. Wang, M. C. Siracusa, S. A. Saenz, J. R. Brestoff, L. A. Monticelli, M. Noti, E. D. Tait Wojno, T. C. Fung, M. Kubo, and D. Artis. 2014. Basophils promote innate lymphoid cell responses in inflamed skin. *J. Immunol.* 193: 3717–3725.
19. Sullivan, B. M., H. E. Liang, J. K. Bando, D. Wu, L. E. Cheng, J. K. McKerrow, C. D. Allen, and R. M. Locksley. 2011. Genetic analysis of basophil function in vivo. *Nat. Immunol.* 12: 527–535.
20. Sokol, C. L., N. Q. Chu, S. Yu, S. A. Nish, T. M. Laufer, and R. Medzhitov. 2009. Basophils function as antigen-presenting cells for an allergen-induced T helper type 2 response. *Nat. Immunol.* 10: 713–720.
21. Perrigoue, J. G., S. A. Saenz, M. C. Siracusa, E. J. Allenspach, B. C. Taylor, P. R. Giacomini, M. G. Nair, Y. Du, C. Zaph, N. van Rooijen, et al. 2009. MHC class II-dependent basophil-CD4⁺ T cell interactions promote T(H)2 cytokine-dependent immunity. *Nat. Immunol.* 10: 697–705.
22. Yoshimoto, T., K. Yasuda, H. Tanaka, M. Nakahira, Y. Imai, Y. Fujimori, and K. Nakanishi. 2009. Basophils contribute to T(H)2-IgE responses in vivo via IL-4 production and presentation of peptide-MHC class II complexes to CD4⁺ T cells. *Nat. Immunol.* 10: 706–712.
23. Gauchat, J. F. S., S. Henchoz, G. Mazzei, J. P. Aubry, T. Brunner, H. Blasey, P. Life, D. Talabot, L. Flores-Romo, J. Thompson, et al. 1993. Induction of human IgE synthesis in B cells by mast cells and basophils. *Nature* 365: 340–343.
24. Yanagihara, Y., K. Kajiwara, Y. Basaki, K. Ikizawa, K. Akiyama, and H. Saito. 1997. Induction of human IgE synthesis in B cells by a basophilic cell line, KU812. *Clin. Exp. Immunol.* 108: 295–301.
25. Yanagihara, Y., K. Kajiwara, Y. Basaki, K. Ikizawa, M. Ebisawa, C. Ra, H. Tachimoto, and H. Saito. 1998. Cultured basophils but not cultured mast cells induce human IgE synthesis in B cells after immunologic stimulation. *Clin. Exp. Immunol.* 111: 136–143.
26. Chen, K., W. Xu, M. Wilson, B. He, N. W. Miller, E. Bengtén, E. S. Edholm, P. A. Santini, P. Rath, A. Chiu, et al. 2009. Immunoglobulin D enhances immune surveillance by activating antimicrobial, proinflammatory and B cell-stimulating programs in basophils. *Nat. Immunol.* 10: 889–898.
27. Merluzzi, S., E. Betto, A. A. Ceccaroni, R. Magris, M. Giunta, and F. Mion. 2015. Mast cells, basophils and B cell connection network. *Mol. Immunol.* 63: 94–103.
28. Schwartz, C., J. U. Eberle, and D. Voehringer. 2016. Basophils in inflammation. *Eur. J. Pharmacol.* 778: 90–95.
29. Qi, X., J. Hong, L. Chaves, Y. Zhuang, Y. Chen, D. Wang, J. Chabon, B. Graham, K. Ohmori, Y. Li, and H. Huang. 2013. Antagonistic regulation by the transcription factors C/EBP α and MITF specifies basophil and mast cell fates. *Immunity* 39: 97–110.
30. Arinobu, Y., H. Iwasaki, M. F. Gurish, S. Mizuno, H. Shigematsu, H. Ozawa, D. G. Tenen, K. F. Austen, and K. Akashi. 2005. Developmental checkpoints of the basophil/mast cell lineages in adult murine hematopoiesis. *Proc. Natl. Acad. Sci. USA* 102: 18105–18110.
31. Sasaki, H., D. Kurotaki, and T. Tamura. 2016. Regulation of basophil and mast cell development by transcription factors. *Allergol. Int.* 65: 127–134.
32. Iwasaki, H., and K. Akashi. 2007. Myeloid lineage commitment from the hematopoietic stem cell. *Immunity* 26: 726–740.
33. Iwasaki, H., S. Mizuno, Y. Arinobu, H. Ozawa, Y. Mori, H. Shigematsu, K. Takatsu, D. G. Tenen, and K. Akashi. 2006. The order of expression of transcription factors directs hierarchical specification of hematopoietic lineages. *Genes Dev.* 20: 3010–3021.
34. Kitamura, Y., E. Morii, T. Jippo, and A. Ito. 2002. Regulation of mast cell phenotype by MITF. *Int. Arch. Allergy Immunol.* 127: 106–109.
35. Takemoto, C. M., Y. N. Lee, A. G. Jegga, D. Zablocki, S. Brandal, A. Shahlaee, S. Huang, Y. Ye, S. Gowrisankar, J. Huynh, and M. A. McDevitt. 2008. Mast cell transcriptional networks. *Blood Cells Mol. Dis.* 41: 82–90.
36. Beaulieu, A. M., and D. B. Sant'Angelo. 2011. The BTB-ZF family of transcription factors: key regulators of lineage commitment and effector function development in the immune system. *J. Immunol.* 187: 2841–2847.
37. Siggs, O. M., and B. Beutler. 2012. The BTB-ZF transcription factors. *Cell Cycle* 11: 3358–3369.
38. Suliman, B. A., D. Xu, and B. R. Williams. 2012. The promyelocytic leukemia zinc finger protein: two decades of molecular oncology. *Front. Oncol.* 2: 74.
39. Alonzo, E. S., R. A. Gottschalk, J. Das, T. Egawa, R. M. Hobbs, P. P. Pandolfi, P. Pereira, K. E. Nichols, G. A. Koretzky, M. S. Jordan, and D. B. Sant'Angelo. 2010. Development of promyelocytic zinc finger and ThPOK-expressing innate gamma delta T cells is controlled by strength of TCR signaling and Id3. *J. Immunol.* 184: 1268–1279.
40. Kreslavsky, T., A. K. Savage, R. Hobbs, F. Gounari, R. Bronson, P. Pereira, P. P. Pandolfi, A. Bendelac, and H. von Boehmer. 2009. TCR-inducible PLZF transcription factor required for innate phenotype of a subset of gammadelta T cells with restricted TCR diversity. *Proc. Natl. Acad. Sci. USA* 106: 12453–12458.
41. Kovalovsky, D., O. U. Uche, S. Eladad, R. M. Hobbs, W. Yi, E. Alonzo, K. Chua, M. Eidson, H. J. Kim, J. S. Im, et al. 2008. The BTB-zinc finger transcriptional regulator PLZF controls the development of invariant natural killer T cell effector functions. *Nat. Immunol.* 9: 1055–1064.
42. Savage, A. K., M. G. Constantinides, J. Han, D. Picard, E. Martin, B. Li, O. Lantz, and A. Bendelac. 2008. The transcription factor PLZF directs the effector program of the NKT cell lineage. *Immunity* 29: 391–403.
43. Koay, H. F., N. A. Gherardin, A. Enders, L. Loh, L. K. Mackay, C. F. Almeida, B. E. Russ, C. A. Nold-Petry, M. F. Nold, S. Bedoui, et al. 2016. A three-stage intrathymic development pathway for the mucosal-associated invariant T cell lineage. *Nat. Immunol.* 17: 1300–1311.
44. Constantinides, M. G., B. D. McDonald, P. A. Verhoef, and A. Bendelac. 2014. A committed precursor to innate lymphoid cells. *Nature* 508: 397–401.
45. Verhoef, P. A., M. G. Constantinides, B. D. McDonald, J. F. Urban, Jr., A. I. Sperling, and A. Bendelac. 2016. Intrinsic functional defects of type 2 innate lymphoid cells impair innate allergic inflammation in promyelocytic leukemia zinc finger (PLZF)-deficient mice. *J. Allergy Clin. Immunol.* 137: 591–600.e1.
46. Constantinides, M. G., H. Gudjonson, B. D. McDonald, I. E. Ishizuka, P. A. Verhoef, A. R. Dinner, and A. Bendelac. 2015. PLZF expression maps the early stages of ILC1 lineage development. *Proc. Natl. Acad. Sci. USA* 112: 5123–5128.
47. Eidson, M., J. Wahlstrom, A. M. Beaulieu, B. Zaidi, S. E. Carsons, P. K. Crow, J. Yuan, J. D. Wolchok, B. Horsthemke, D. Wiczorek, and D. B. Sant'Angelo. 2011. Altered development of NKT cells, $\gamma\delta$ T cells, CD8 T cells and NK cells in a PLZF deficient patient. *PLoS One* 6: e24441.
48. Kovalovsky, D., E. S. Alonzo, O. U. Uche, M. Eidson, K. E. Nichols, and D. B. Sant'Angelo. 2010. PLZF induces the spontaneous acquisition of memory/effector functions in T cells independently of NKT cell-related signals. *J. Immunol.* 184: 6746–6755.
49. Savage, A. K., M. G. Constantinides, and A. Bendelac. 2011. Promyelocytic leukemia zinc finger turns on the effector T cell program without requirement for agonist TCR signaling. *J. Immunol.* 186: 5801–5806.
50. Raberger, J., A. Schebesta, S. Sakaguchi, N. Boucheron, K. E. Blomberg, A. Berglöff, T. Kolbe, C. I. Smith, T. Rülke, and W. Ellmeier. 2008. The transcriptional regulator PLZF induces the development of CD44 high memory phenotype T cells. *Proc. Natl. Acad. Sci. USA* 105: 17919–17924.
51. Zhang, S., A. Laouar, L. K. Denzin, and D. B. Sant'Angelo. 2015. Zbtb16 (PLZF) is stably suppressed and not inducible in non-innate T cells via T cell receptor-mediated signaling. *Sci. Rep.* 5: 12113.
52. Wei, G., L. Wei, J. Zhu, C. Zang, J. Hu-Li, Z. Yao, K. Cui, Y. Kanno, T. Y. Roh, W. T. Watford, et al. 2009. Global mapping of H3K4me3 and H3K27me3 reveals specificity and plasticity in lineage fate determination of differentiating CD4⁺ T cells. *Immunity* 30: 155–167.
53. Hosokawa, H., M. Romero-Wolf, M. A. Yui, J. Ungerback, M. L. G. Quilaoan, M. Matsumoto, K. I. Nakayama, T. Tanaka, and E. V. Rothenberg. 2018. Bcl11b sets pro-T cell fate by site-specific cofactor recruitment and by repressing Id2 and Zbtb16. *Nat. Immunol.* 19: 1427–1440.
54. Zhu, L., Y. Qiao, E. S. Choi, J. Das, D. B. Sant'angelo, and C. H. Chang. 2013. A transgenic TCR directs the development of IL-4⁺ and PLZF⁺ innate CD4 T cells. *J. Immunol.* 191: 737–744.
55. Barna, M., N. Hawe, L. Niswander, and P. P. Pandolfi. 2000. Plzf regulates limb and axial skeletal patterning. *Nat. Genet.* 25: 166–172.
56. Thapa, P., J. Das, D. McWilliams, M. Shapiro, R. Sundsbak, M. Nelson-Holte, S. Tangen, J. Anderson, S. Desiderio, S. Hiebert, et al. 2013. The transcriptional repressor NKAP is required for the development of iNKT cells. *Nat. Commun.* 4: 1582.
57. Yang, L., D. Bryder, J. Adolfsson, J. Nygren, R. Månsson, M. Sigvardsson, and S. E. Jacobsen. 2005. Identification of Lin(-)Sca1(+)/kit(+)/CD34(+)/Flt3(-) short-term hematopoietic stem cells capable of rapidly reconstituting and rescuing myeloablative transplant recipients. *Blood* 105: 2717–2723.
58. Akashi, K., D. Traver, T. Miyamoto, and I. L. Weissman. 2000. A clonogenic common myeloid progenitor that gives rise to all myeloid lineages. *Nature* 404: 193–197.
59. Iwasaki, H., S. Mizuno, R. Mayfield, H. Shigematsu, Y. Arinobu, B. Seed, M. F. Gurish, K. Takatsu, and K. Akashi. 2005. Identification of eosinophil lineage-committed progenitors in the murine bone marrow. *J. Exp. Med.* 201: 1891–1897.
60. Hettinger, J., D. M. Richards, J. Hansson, M. M. Barra, A. C. Joschko, J. Krigsveld, and M. Feuerer. 2013. Origin of monocytes and macrophages in a committed progenitor. *Nat. Immunol.* 14: 821–830.
61. Pobeziński, L. A., R. Eitzensperger, S. Jeurling, A. Alag, T. Kadakia, T. M. McCaughy, M. Y. Kimura, S. O. Sharrow, T. I. Guinter, L. Feigenbaum, and A. Singer. 2015. Let-7 microRNAs target the lineage-specific transcription factor PLZF to regulate terminal NKT cell differentiation and effector function. *Nat. Immunol.* 16: 517–524.
62. Lynch, L., X. Michelet, S. Zhang, P. J. Brennan, A. Moseman, C. Lester, G. Besra, E. E. Vornhof-Dekrey, M. Tighe, H. F. Koay, et al. 2015. Regulatory iNKT cells lack expression of the transcription factor PLZF and control the homeostasis of T(reg) cells and macrophages in adipose tissue. *Nat. Immunol.* 16: 85–95.
63. Rosenstein, R. K., J. S. Bezbradica, S. Yu, and R. Medzhitov. 2014. Signaling pathways activated by a protease allergen in basophils. *Proc. Natl. Acad. Sci. USA* 111: E4963–E4971.
64. Miyake, K., and H. Karasuyama. 2017. Emerging roles of basophils in allergic inflammation. *Allergol. Int.* 66: 382–391.

65. Schroeder, J. T. 2011. Basophils: emerging roles in the pathogenesis of allergic disease. *Immunol. Rev.* 242: 144–160.
66. Karasuyama, H., T. Wada, S. Yoshikawa, and K. Obata. 2011. Emerging roles of basophils in protective immunity against parasites. *Trends Immunol.* 32: 125–130.
67. Lantz, C. S., J. Boesiger, C. H. Song, N. Mach, T. Kobayashi, R. C. Mulligan, Y. Nawa, G. Dranoff, and S. J. Galli. 1998. Role for interleukin-3 in mast-cell and basophil development and in immunity to parasites. *Nature* 392: 90–93.
68. Voehringer, D. 2013. Protective and pathological roles of mast cells and basophils. *Nat. Rev. Immunol.* 13: 362–375.
69. Kim, S., H. Karasuyama, A. F. Lopez, W. Ouyang, X. Li, G. Le Gros, and B. Min. 2013. IL-4 derived from non-T cells induces basophil- and IL-3-independent Th2 immune responses. *Immune Netw.* 13: 249–256.
70. Shen, T., S. Kim, J. S. Do, L. Wang, C. Lantz, J. F. Urban, G. Le Gros, and B. Min. 2008. T cell-derived IL-3 plays key role in parasite infection-induced basophil production but is dispensable for in vivo basophil survival. *Int. Immunol.* 20: 1201–1209.
71. Iikura, M., M. Ebisawa, M. Yamaguchi, H. Tachimoto, K. Ohta, K. Yamamoto, and K. Hirai. 2004. Transendothelial migration of human basophils. *J. Immunol.* 173: 5189–5195.
72. Tang, H., W. Cao, S. P. Kasturi, R. Ravindran, H. I. Nakaya, K. Kundu, N. Murthy, T. B. Kepler, B. Malissen, and B. Pulendran. 2010. The T helper type 2 response to cysteine proteases requires dendritic cell-basophil cooperation via ROS-mediated signaling. *Nat. Immunol.* 11: 608–617.
73. Mao, A. P., M. G. Constantinides, R. Mathew, Z. Zuo, X. Chen, M. T. Weirauch, and A. Bendelac. 2016. Multiple layers of transcriptional regulation by PLZF in NKT-cell development. *Proc. Natl. Acad. Sci. USA* 113: 7602–7607.
74. Gleimer, M., H. von Boehmer, and T. Kreslavsky. 2012. PLZF controls the expression of a limited number of genes essential for NKT cell function. *Front. Immunol.* 3: 374.
75. Alonzo, E. S., and D. B. Sant'Angelo. 2011. Development of PLZF-expressing innate T cells. *Curr. Opin. Immunol.* 23: 220–227.
76. Dick, J. E., and S. Doulatov. 2009. The role of PLZF in human myeloid development. *Ann. N. Y. Acad. Sci.* 1176: 150–153.
77. Kopley, C. L., P. J. McFeeley, J. M. Oliver, and M. F. Lipscomb. 2001. Immunohistochemical detection of human basophils in postmortem cases of fatal asthma. *Am. J. Respir. Crit. Care Med.* 164: 1053–1058.
78. Nabe, T., K. Matsuya, K. Akamizu, M. Fujita, T. Nakagawa, M. Shioe, H. Kida, A. Takiguchi, H. Wakamori, M. Fujii, et al. 2013. Roles of basophils and mast cells infiltrating the lung by multiple antigen challenges in asthmatic responses of mice. *Br. J. Pharmacol.* 169: 462–476.
79. Yoshimoto, T., H. Tsutsui, K. Tominaga, K. Hoshino, H. Okamura, S. Akira, W. E. Paul, and K. Nakanishi. 1999. IL-18, although antiallergic when administered with IL-12, stimulates IL-4 and histamine release by basophils. *Proc. Natl. Acad. Sci. USA* 96: 13962–13966.
80. Sims, J. E., and D. E. Smith. 2010. The IL-1 family: regulators of immunity. *Nat. Rev. Immunol.* 10: 89–102.
81. Kumano, K., A. Nakao, H. Nakajima, F. Hayashi, M. Kurimoto, H. Okamura, Y. Saito, and I. Iwamoto. 1999. Interleukin-18 enhances antigen-induced eosinophil recruitment into the mouse airways. *Am. J. Respir. Crit. Care Med.* 160: 873–878.
82. Wild, J. S., A. Sigounas, N. Sur, M. S. Siddiqui, R. Alam, M. Kurimoto, and S. Sur. 2000. IFN-gamma-inducing factor (IL-18) increases allergic sensitization, serum IgE, Th2 cytokines, and airway eosinophilia in a mouse model of allergic asthma. *J. Immunol.* 164: 2701–2710.
83. Gaya, M., P. Barral, M. Burbage, S. Aggarwal, B. Montaner, A. Warren Navia, M. Aid, C. Tsui, P. Maldonado, U. Nair, et al. 2018. Initiation of antiviral B cell immunity relies on innate signals from spatially positioned NKT cells. *Cell* 172: 517–533.e20.
84. Pecaric-Petkovic, T., S. A. Didichenko, S. Kaempfer, N. Spiegel, and C. A. Dahinden. 2009. Human basophils and eosinophils are the direct target leukocytes of the novel IL-1 family member IL-33. *Blood* 113: 1526–1534.
85. Doulatov, S., F. Notta, K. L. Rice, L. Howell, A. Zelent, J. D. Licht, and J. E. Dick. 2009. PLZF is a regulator of homeostatic and cytokine-induced myeloid development. *Genes Dev.* 23: 2076–2087.
86. Constantinides, M. G., D. Picard, A. K. Savage, and A. Bendelac. 2011. A naive-like population of human CD1d-restricted T cells expressing intermediate levels of promyelocytic leukemia zinc finger. *J. Immunol.* 187: 309–315.




Supplemental Figure 1. Comparison of PLZF expression levels in myeloid progenitor lineages. All populations were identified according to the gating outlined in Supplemental figures 2 and 3. Histogram overlays of cells of each lineage from WT and PLZF-eGFP bone marrow are shown. Scatter plots show results from individual mice. Basophil and mast cell histograms show data from experiments with >5 fold change. Three independent experiments; 5 biological repeats.




Supplemental Figure 2. Gating of Myeloid Progenitor Lineages. a) Identification of the individual myeloid progenitor populations included selection of live single cells that were CD45.2⁺ from bone marrow. b) Common myeloid progenitor (CMP) cells were identified as lineage⁻, Sca1⁻, CD34⁺, cKit⁺, and CD16/32⁻. Granulocyte-monocyte progenitors (GMP) were identified as lineage⁻, Sca1⁻, CD34⁺, cKit⁺, and CD16/32⁺. c) Eosinophil progenitors (EoP) were identified as lineage⁻, Sca1⁻, CD34⁺, cKit⁺, and CD125⁺. c) Common monocyte progenitors (CMoP) were identified as lineage⁻, Sca1⁻, CD115⁺, cKit⁺, Flt3⁻, and Ly-6C⁺. Monocyte-macrophage-dendritic cell progenitors (MDP) were identified as lineage⁻, Sca1⁻, CD115⁺, cKit⁺, Flt3⁺, and Ly-6C⁻. Dot plots shown are representative of three independent experiments and 5 biological replicates.


a) Basophil Gating:




b) Mast Cell Gating:



c) Basophil Progenitor Gating:




d) Mast Cell Progenitor Gating:




Supplemental Figure 3. Gating of basophil and mast cell lineages. All populations were first gated according to the strategy outlined in Supplemental Figure 2a. **a)** Basophils (Bas) were identified as lineage⁻, CD49b⁺, cKit⁺, and FcεR1⁺. **b)** Mast cells (Mast) were identified as lineage⁻, CD11b⁺, cKit⁺, and FcεR1⁺. **c)** Basophil progenitors (BaP) were identified as CD34⁺, lineage⁻, cKit⁺, IL-33R^{low}, Intβ7^{low}, and FcεR1⁺. **d)** Mast cell progenitors (McP) were identified as lineage⁻, cKit⁺, Intβ7⁺, FcεR1⁺, IL-33R⁺, and Intβ7⁺. Dot plots shown are representative of three independent experiments and 5 biological replicates.

Thymus



Spleen



Supplemental Figure 4. Fate mapping PLZF expression with PLZF-Cre BAC and the floxed-stop ROSA26 ZsGreen allele. As expected, ZsGreen expression was observed in nearly all PLZF expressing cells, which include thymus and spleen NKT cells. Distinct from data we previously reported using a floxed-stop ROSA26 tdTomato allele, expression of ZsGreen was rare in double positive (CD4⁺CD8⁺) thymocytes, B cells and MHCII⁺ cells. Some CD4SP and CD8SP thymocytes and some CD4⁺ and CD8⁺ T cells were labeled. Data are representative of 5 repeats. Similar experiments with mice carrying a floxed-stop ROSA26 YFP allele showed incomplete expression of YFP in PLZF expressing cells, such as NKT cells (data not shown).