

Epigenomic analysis of primary human T cells reveals enhancers associated with T_H2 memory cell differentiation and asthma susceptibility

Grégory Seumois^{1,3,7}, Lukas Chavez^{1,7}, Anna Gerasimova^{1,7}, Matthias Lienhard¹, Nada Omran², Lukas Kalinke², Maria Vedanayagam², Asha Purnima V Ganesan², Ashu Chawla¹, Ratko Djukanović², K Mark Ansel³, Bjoern Peters¹, Anjana Rao^{1,4-6} & Pandurangan Vijayanand¹⁻³

A characteristic feature of asthma is the aberrant accumulation, differentiation or function of memory CD4⁺ T cells that produce type 2 cytokines (T_H2 cells). By mapping genome-wide histone modification profiles for subsets of T cells isolated from peripheral blood of healthy and asthmatic individuals, we identified enhancers with known and potential roles in the normal differentiation of human T_H1 cells and T_H2 cells. We discovered disease-specific enhancers in T cells that differ between healthy and asthmatic individuals. Enhancers that gained the histone H3 Lys4 dimethyl (H3K4me2) mark during T_H2 cell development showed the highest enrichment for asthma-associated single nucleotide polymorphisms (SNPs), which supported a pathogenic role for T_H2 cells in asthma. *In silico* analysis of cell-specific enhancers revealed transcription factors, microRNAs and genes potentially linked to human T_H2 cell differentiation. Our results establish the feasibility and utility of enhancer profiling in well-defined populations of specialized cell types involved in disease pathogenesis.

The acquisition of immunological memory is the hallmark of a protective immune response¹. During this evolutionarily conserved process, naive T cells and B cells that have not previously encountered antigen differentiate during primary infection into memory cells that have specialized functions in immune system defense, thus permitting the organism to effectively respond to a later infection with the same pathogen. As expected for a process of cell-lineage specification, differentiation of memory T cells and B cells involves extensive epigenetic changes that are required to initiate and maintain a heritable program of gene expression². Adaptive immunity is not without risks: some genetically susceptible individuals develop abnormal memory responses to potentially harmless antigens, which results in a multitude of immunological diseases ranging from autoimmunity to allergies and asthma³⁻⁵. A clear understanding of the molecular and epigenetic mechanisms underlying normal as well as aberrant differentiation of human memory cell types will pave the way to develop new approaches to tackle immune system-mediated diseases.

Asthma is a disease characterized by airway inflammation that is mediated by excessive memory responses to inhaled allergens, such as grass pollen³. The alarming rise in asthma incidence is a major global health concern, not only in the western world but also in large developing countries such as India, China and Brazil⁶. Over 200 million people suffer from asthma world-wide, which causes an economic

burden that exceeds that of tuberculosis and HIV-AIDS combined⁶. At present, there is no cure for asthma, and most patients require long-term, daily nonspecific medication such as corticosteroids to control the underlying inflammation and prevent symptoms and life-threatening asthma attacks⁷. Therapies targeting specific type 2 cytokines are only efficacious in certain types of asthma⁸, which raises the possibility that there are unclassified molecular subtypes of asthma for which different therapies may prove beneficial.

A molecular feature of asthma and other allergic diseases is the excessive differentiation of a subset of CD4⁺ T helper cells known as T_H2 cells, which produce a characteristic spectrum of type 2 cytokines, including the interleukins IL-4, IL-5 and IL-13 (ref. 3). Genes encoding these three cytokines are localized on human chromosome 5, in a conserved grouping known as the T_H2 cytokine locus in which the *IL5* gene is separated from the *IL4* and *IL13* genes by the *RAD50* gene, which encodes a conserved DNA repair protein⁹. The last few introns of the *RAD50* gene contain four conserved enhancers that together constitute a locus control region (LCR) for the cytokine genes; in addition, the T_H2 cell cytokine locus contains many evolutionarily conserved enhancers, silencers and other *cis*-regulatory regions whose functions have been comprehensively analyzed in mice, by monitoring changes in histone modifications and DNase I hypersensitivity during T_H2 cell differentiation as well as by gene disruption^{9,10}. Many

¹La Jolla Institute for Allergy & Immunology, La Jolla, California, USA. ²Clinical and Experimental Sciences, Southampton National Institute for Health Research Respiratory Biomedical Research Unit, University of Southampton, Faculty of Medicine, Southampton, UK. ³University of California San Francisco, San Francisco, California, USA. ⁴Sanford Consortium for Regenerative Medicine, La Jolla, California, USA. ⁵Department of Pharmacology, University of California San Diego, San Diego, California, USA. ⁶University of California San Diego Moores Cancer Center, San Diego, California, USA. ⁷These authors contributed equally to this work. Correspondence should be addressed to P.V. (vijay@liai.org), B.P. (bpeters@liai.org) or A.R. (arao@liai.org).

Received 10 January; accepted 3 June; published online 6 July 2014; doi:10.1038/ni.2937

of these enhancers show sequence conservation in humans, and there is evidence for functional conservation as well¹¹.

Profiling DNA and histone modifications (epigenomic marks) layered over the genome sequence provides detailed information about the location and activity of *cis*-regulatory elements and the likely transcriptional potential of their target genes¹². For instance, chromatin immunoprecipitation (ChIP) of DNA regions associated with the H3K4me1 modification followed by high-throughput sequencing (ChIP-seq) identifies *cis*-regulatory regions of genes that are poised for expression but may not actually be expressed until an appropriate environmental signal is received (for example, stimulation of the T cell antigen receptor by peptide–major histocompatibility complex binding); H3K27 acetylation (H3K27Ac) is associated with enhancers of genes that are actively being expressed¹³. Analysis of the H3K4 dimethyl (H3K4me2) modification has a dual advantage: it reveals both active and poised enhancers and provides a more precise localization of enhancers than analyses of H3K4me1 and H3K27Ac^{14,15}. Epigenetic modifications tend to be more stable than RNA transcripts from the associated gene and provide a better measure of transcriptional status than RNA sequencing (RNA-seq) alone^{14,15}. However, the broader applicability of epigenomic approaches for studying primary human cells is hampered by technical and practical constraints, including the small amount of blood or tissue that can be obtained from human subjects for research purposes, the variability of genome-wide ChIP assays when performed with small numbers of cells, the fact that the cell populations available for genome-wide analysis are typically mixtures of many different cell types, the genetic heterogeneity of humans and the complex nature of most human diseases.

Our goal in this study was to develop methods to profile active and poised enhancers in naive and memory CD4⁺ T cells from healthy individuals and asthmatic patients. Because we wished to extend these methods eventually to very small cell populations isolated from diseased tissues, we chose to profile *cis*-regulatory regions using a single histone modification, H3K4me2. By generating maps of H3K4me2-marked *cis*-regulatory regions, we identified a large number of cell type–specific and disease-specific enhancers. Some of these are linked to genes already implicated in T_H2 cell differentiation, but for many more enhancers our analysis provides the first such indication. Several of the identified enhancers also harbor asthma-associated SNPs, validating our premise that the study of T_H2 subtype–enriched cells from peripheral blood of asthmatic patients would yield information relevant to disease associated with T_H2 cells. We conclude that enhancer profiling for H3K4me2 is a promising approach for studying the functions of well-defined cell populations in health and disease, with particular value when only limiting cell numbers are available.

RESULTS

Mapping *cis*-regulatory DNA in primary human CD4⁺ T cells

To capture the *cis*-regulatory landscape of human T cells, we isolated circulating naive and memory CD4⁺ T cells from peripheral blood of 24 subjects (12 healthy controls and 12 asthma patients; Online Methods). We subdivided CD4⁺ memory T cells on the basis of their surface expression of the chemokine receptor CCR4 (Supplementary Fig. 1). The CCR4⁺ T cells were enriched for T_H2 cells, which are known to have a pathogenic role in asthma, whereas the CCR4[−] T cells were depleted for T_H2 cells but enriched for T_H1 cells, which produce interferon- γ (IFN- γ)^{16–18}; for simplicity, hereafter we refer to CCR4⁺ T cells as ‘T_H2 cells’ and CCR4[−] T cells as ‘T_H1 cells’. Using ChIP-seq, we identified DNA regions associated with H3K4me2. To ensure that comparisons between sample groups were not affected by assay variability, we optimized the H3K4me2 ChIP-seq assay to provide highly reproducible

results in multiple replicate assays (Fig. 1a and Supplementary Fig. 2) and also microscaled the assay (Supplementary Figs. 3 and 4) to enable reproducible detection of *cis*-regulatory DNA regions in as few as 10⁴ cells (Fig. 1b). We performed 120 high-throughput whole-genome ChIP-seq assays, which generated $\sim 2.8 \times 10^9$ reads covering $\sim 1 \times 10^{11}$ bases across three cell types (Supplementary Table 1), and computationally analyzed this data set to identify enhancers associated with CD4⁺ memory T cell differentiation as well as asthma.

H3K4me2 enrichment at most genes was highly comparable in the three cell types across all study subjects, which emphasizes the value of our multisample ChIP-seq assay in reducing technical variation (STIM1, NUP98, SELP and SELL genes; Fig. 1c and Supplementary Fig. 5). As expected, the H3K4me2 mark was depleted in the *cis*-regulatory regions of genes not expressed in T cells (e.g., SELE; Fig. 1c). We found clear evidence of cell type–specific *cis*-regulatory DNA elements in noncoding sequences: for instance, compared to naive T cells, the T_H2 cell subset displayed extensive enrichment of H3K4me2 at promoters and *cis*-regulatory regions in the extended CCR4 and CCR6 loci (Fig. 1c and Supplementary Fig. 5). Merging multiple samples from the same cell types and disease categories reduced the background variability (that is, nonspecifically enriched regions) observed in individual assays, thus improving the sensitivity of our assay for detecting genomic regions with true H3K4me2 enrichment (Fig. 1d,e and Supplementary Fig. 6). Comparing enrichment values between these large sets of samples from the three different cell types, we detected highly significant differences at loci known to be regulated in a cell type–specific fashion (for example, CCL5 and CCR4 loci; Fig. 1e).

Enhancers linked to CD4⁺ T cell differentiation *in vivo*

To identify *cis*-regulatory DNA regions differentially enriched for H3K4me2 in pairwise comparisons between cell types (differentially enriched regions; DERs), we calculated H3K4me2 enrichment based on reads in non-overlapping, consecutive 500-base-pair (bp) windows across the whole genome^{19,20}. To identify cell-specific enhancers present in the general human population independent of disease status, we grouped each cell type from all 24 subjects and calculated differential enrichment for each group by modeling a negative binomial distribution for the data dispersion (Online Methods, Supplementary Note, and Supplementary Figs. 7 and 8). Analogous to gene-expression values in RNA-seq data, enrichment values, expressed as reads per kilobase per million mapped (RPKM), were calculated for each of the $\sim 6 \times 10^6$ 500-bp windows spanning the whole genome. 71,640 unique genomic windows (500-bp size), representing $\sim 1\%$ of the human genome, showed differential enrichment of H3K4me2 in the three cell types compared (Bonferroni-adjusted $P < 0.05$; Fig. 2a and Supplementary Table 2). We detected most of these differences ($\sim 90\%$ of DERs) in the naive to memory T cell comparison (Fig. 2a and Supplementary Table 3) rather than between the T_H2 and T_H1 memory cell types, which confirmed that the activation and consequent differentiation of naive T cells into memory T cells results in major changes in the *cis*-regulatory landscape of chromatin²¹.

Over 90% of the $\sim 71,000$ cell-specific DERs were localized to intergenic and intronic regions; only 4% were present in promoter regions (defined as transcription start site (TSS) ± 1 kilobase (kb)) of annotated RefSeq genes (Supplementary Fig. 9). We termed all DERs outside promoter regions ‘putative differentiation-related enhancers’^{14,15} and classified them into three subgroups (Fig. 2b and Supplementary Table 3): $\sim 30,263$ ‘T_H2 enhancers’ that exhibited the strongest H3K4me2 gain or loss in memory T cells enriched for the T_H2 cell subset; $\sim 17,483$ ‘T_H1 enhancers’ that exhibited the strongest H3K4me2 gain or loss in memory T cells enriched for the T_H1 cell




Figure 1 Reproducibility, microscaling and sensitivity of the H3K4me2 ChIP-seq assay. **(a)** Standard ChIP-seq assays (2×10^6 cells; six replicates) showing H3K4me2 enrichment patterns of the gene loci (top) in D10 cells. **(b)** Standard ChIP-seq assay (2×10^6 cells) and micro-scaled ChIP-seq assay (10^5 , 10^4 and 10^3 cell samples; 3–4 replicates) showing H3K4me2 enrichment patterns in D10 cells. **(c)** ChIP-seq analysis showing H3K4me2 enrichment patterns, for control regions *STIM1*, *NUP98*, *SELP* and *SELL* loci, nonexpressed *SELE* locus, and T_H2 cell-type specific *CCR4* and *CCR6* loci, in the naive cells and T_H2 cells of six healthy subjects. Significant H3K4me2 enrichment (exact test for negative binomial distribution, using edgeR integrated in Bioconductor package MEDIPS) across distal *cis*-regulatory elements and promoters in these loci are highlighted by purple and blue dashed-line boxes, respectively. **(d)** ChIP-seq analysis showing cell-specific H3K4me2 enrichment patterns, for *CCL5* (T_H1 cell-specific), *CCR4* (T_H2 cell-specific) and control region *YWHAZ* (T cell housekeeping gene), in naive, T_H1 cells and T_H2 cells. For each cell type, data were merged from all donors, including assay duplicates. **(e)** H3K4me2 enrichment values for a specific 500-bp window (highlighted in purple dashed line boxes in **d**). Each dot represents data from a single assay; error bars indicate mean $\pm$ s.e.m. $*P < 1 \times 10^{-6}$, exact test for negative binomial distribution (using edgeR integrated in Bioconductor package MEDIPS).

subset; and ~21,659 'shared memory enhancers' that displayed equivalent H3K4me2 gain or loss in both T_H2 and T_H1 memory cells when compared to naive T cells. Several of these putative human enhancer DERs corresponded to well-characterized DNase I hypersensitive (HS) enhancers in the mouse T_H2 locus (LCR-O, LCR-A, HS II, HS V; **Fig. 2c**). Moreover, a human T_H1 DER (green bar) enriched in both T_H2 cells and T_H1 cells relative to naive T cells corresponded to a silencer region (HS IV in the mouse) that is DNase I hypersensitive in both T_H1 cells and T_H2 cells and restrains type 2 cytokine production in T_H1 cells²². Likewise, the region corresponding to mouse hypersensitivity site S3 (HSS3), which contains a binding site for the chromatin insulator binding protein CTCF and exhibits constitutive DNase I hypersensitivity in naive, T_H1 and T_H2 memory cells⁹, appeared as a shared memory enhancer region in the human T_H2 (*IL5–RAD50–IL13–IL4*) locus. Further, a number of evolutionarily conserved enhancers in other relevant gene loci (e.g., *IFNG*, *GATA3* and *TBX21*, encoding the signature T_H1 cell cytokine IFN- γ and the lineage-defining T_H1 and T_H2 cell transcription factors GATA-3 and T-bet, respectively) also exhibited cell type-specific enhancer activity as judged by H3K4me2 enrichment (**Fig. 2d**)²³. However, five evolutionarily conserved regions identified as enhancers in the mouse (LCR-B, LCR-C, CGRE, CNS-1 and HS III) did not display cell-specific enhancer profiles in human T cells. In summary, we identified a number of putative lineage-specific enhancers in human T cells and showed that many but not all of these are conserved between human and mouse (**Fig. 2c,d**). Our findings indicate that at least some of the diverse differentiation-related enhancers identified in our screen may be important in shaping human CD4⁺ memory T cell fate.

Genes and molecular pathways linked to T_H cell differentiation

To connect our list of DERs to target genes, we focused on H3K4me2 at gene promoters (TSS $\pm$ 1 kb). Enrichment of H3K4me2 at promoters reliably marks genes that are either active or poised for expression in response to extrinsic stimuli^{14,15}. We identified ~1,400 protein-coding genes (~7% of all annotated genes), 70 microRNAs (miRNAs) and 78 long noncoding RNAs that showed differential enrichment of H3K4me2 at their promoters (Bonferroni-adjusted $P < 0.05$; **Fig. 3a** and **Supplementary Table 4**). We observed concordant changes in gene expression and H3K4me2 enrichment patterns in many potential enhancers located in introns and within ± 20 kb of TSSs or transcription end sites of these genes (**Supplementary Fig. 10**).

We classified target genes of promoter-localized DERs into six subgroups based on preferential gain or loss of H3K4me2 enrichment during memory differentiation (**Fig. 3a**). Genes encoding many cytokines (*IL4*, *IFNG*, *IL2* and *IL21*), chemokines (*CCL3*, *CCL4* and *CCL5*), chemokine receptors (*CCR3*, *CCR4*, *CCR8* and *CCR10*) and miRNAs (miR-146a, miR-21, miR-155, miR-29b, miR-15b and miR-193b)²⁴ known to be differentially expressed in memory cells partitioned as expected into these subgroups (**Fig. 3a–c** and **Supplementary Table 4**). In addition, we discovered unique miRNAs and genes not previously linked to helper T cell differentiation (**Fig. 3b** and **Supplementary Table 4**).

Analysis of biological process enrichment for genes that gain H3K4me2 during memory differentiation showed overrepresentation of pathways linked to positive regulation of T cell proliferation and activation, positive regulation of tyrosine phosphorylation of STAT proteins, regulation of adaptive immune response, activity of chemokine receptors, homeostasis of cellular calcium ions and regulation of immunoglobulin production (**Supplementary Table 5**). We observed significant enrichment for genes involved in regulation of apoptosis and mitosis (cell cycle) (**Table 1**) for the 'T_H2 gain'


subgroup of genes that gained promoter H3K4me2 enrichment in T_H2 cells. Bioinformatic analysis of regulatory interactions among genes in the 'T_H2 gain' subgroup pointed to three genes, *MYC*, *E2F2* and *E2F4*, as potential major regulators of T_H2 cell growth and survival (**Fig. 4**). *MYC*, *E2F2* and *E2F4* are known to regulate cell cycle, proliferation, apoptosis and metabolic reprogramming of T cells after activation^{25–27}, and their increased expression likely confers a survival (growth) advantage to T_H2 cells over T_H1 cells, as previously described for mouse T_H2 cells²⁸. Thus, beside the well-known differences in cytokine production and tissue homing between human T_H1 cells and T_H2 cells, our data will provide information on genes that regulate other important functional differences between these cell types.

To identify target genes for the vast majority of DERs (~69,000) located outside promoter regions (putative enhancers), we determined the location of the nearest CTCF binding sites, potential insulator elements occupied by CTCF across many different cell types²⁹. On the assumption that genes located in CTCF-bounded regions are controlled by enhancers located in the same region, we connected a total of ~7,600 target genes to the ~69,000 differentiation-related enhancers, which included nearly 50% of genes identified in a recent study of enhancers in *in vitro*-derived T_H1 cells and T_H2 cells³⁰ (**Supplementary Table 6**). We assigned several candidate genes (median = 1.7 genes) to each enhancer and linked many enhancers (median = 4.9 enhancers) to each gene. To determine the functional significance of this large list of candidate genes, we performed a gene set enrichment analysis after classifying the genes into subgroups (as for the promoter analysis). Analysis of the enhancer-linked genes revealed many pathways important for T_H2 and T_H1 memory cell differentiation, and included many genes linked to promoter-localized DERs, **Supplementary Table 7**). An important caveat of pathway analysis performed with enhancer-linked genes is that the results can be confounded by correlation (or clustering) of genes with similar function throughout the genome. Additional experimental and computational analysis will be required to provide definitive evidence for the proposed assignments of genes to enhancers and vice versa.

Comparison of gene expression profiles with H3K4me2 enrichment profiles between cell types showed that differentially expressed genes display concordant changes in H3K4me2 enrichment patterns in the extended gene loci (**Supplementary Fig. 11a** and **Supplementary Table 8**). As expected, some of the genes that showed differential H3K4me2 enrichment did not show a significant change in gene expression (negative binomial test, false discovery rate $\leq 1\%$; **Supplementary Fig. 11a**). At least a fraction of these genes are likely to be poised for expression after an appropriate environmental signal, as these include the cytokine genes *IL4* and *IL21*, which are expressed only by stimulated T cells (**Supplementary Fig. 11b**). Overall, comparison of H3K4me2 profiles as opposed to gene expression profiles between T_H cell subtypes allowed us to define genes poised for expression in a cell-specific manner as well as identify enhancers involved in cell-specific gene regulation.

Transcription factor binding sites and motifs at T_H cell enhancers

We took two approaches to identify *trans*-acting factors that might drive human T_H cell differentiation *in vivo*. First, we identified known transcription factor binding motifs within the differentiation-related enhancers shown in **Figure 2b** (**Fig. 5a**). As expected, binding motifs for GATA-3 and T-bet were enriched in enhancers that gained H3K4me2 in T_H2 cells and T_H1 cells, respectively (**Fig. 5a** and **Supplementary Table 9**). In addition, motifs for a number






Figure 3 Genes and pathways linked to differentiation of CD4⁺ memory cells. **(a)** Z-scores of normalized read counts for each unique promoter-localized DER (rows) obtained from any of the three pairwise comparisons of naive, TH₂ and TH₁ cells; to illustrate samples that are above or below the average across all samples per window (row), a two-color scale (see key) was used for Z-scores; data are from each independent ChIP-seq assay ($n = 120$) (columns). Promoter-localized DERs were classified into six groups based on gain or loss of H3K4me2 enrichment during differentiation of naive T cells into the TH₂ cells or TH₁ cells (**Supplementary Table 3**); in each group, the promoter-localized DERs were arranged by their chromosomal location (from the start of chromosome 1 to the end of chromosome Y); 'shared' denotes 'shared memory enhancers' that displayed equivalent H3K4me2 gain or loss in both TH₂ cells and TH₁ cells when compared to naive cells. Target genes of some promoter-localized DERs are shown to the right. **(b)** miRNAs that showed the strongest gain or loss of H3K4me2 enrichment in their promoter regions for the various DER subgroups. **(c)** miRNAs for which the H3K4me2 enrichment tracks for each cell type are shown, for indicated miRNAs (* in **b**). Purple dashed-line boxes indicate the promoter regions (TSS $\pm$ 1 kb) for each miRNA.

of transcription factors not previously implicated in T cell differentiation were markedly enriched in 'TH₂ enhancers' (**Fig. 5a**). Second, using publicly available ChIP-seq data from cell lines and primary T cells, we confirmed that some of these distinct transcription factors actually bound certain TH₂ cell enhancers (**Fig. 5b,c** and **Supplementary Table 10**). NRF2 (also known as nuclear factor (erythroid-derived 2)-like 2; *NFE2L2*), and the structurally related

factor NFE2 that binds antioxidant response elements, were strong hits in our *in silico* analysis (**Fig. 5b**). Activation of NRF2 has been shown to skew differentiation of mouse naive cells to TH₂ cells and increase TH₂ cell cytokine production³¹. Our results suggest that NRF2 may be a determining factor in human TH₂ cell differentiation; experimental validation of its role in TH₂ cell-driven diseases will be important because NRF2 signaling is activated by synthetic

Table 1 Enrichment for genes involved in regulation of apoptosis and mitosis

Category	GO term	P value	Enrichment score	Genes	Count (% of total)
T _H 2 cell gain	Mitosis	3.45×10^{-7}	6.4	<i>CDK1, KIFC1, KIF22, KIF11, ANAPC4, KIF18A, TPX2, CENPF, NEDD9, NDC80, CENPE, PTTG1, CEP55, UBE2C, LATS1, WEE1, SMC4, CCNB2, CDCA2, SKA2, SKA1, NEK6, TUBB3 and CDCA3</i>	24 (4)
T _H 2 cell gain	Regulation of apoptosis	6.14×10^{-4}	3.1	<i>DLC1, CSF2, NUA2, CLU, TNFSF15, TP63, FASLG, TNFRSF8, AKAP13, PMAIP1, TNFSF12, CALR, MCF2L, NLRC4, G2E3, CHD8, NOD2, CDKN2A, CDKN2C, CD2, HSPA5, BMF, MYC, CASP2, TXNIP, IL4, CFLAR, CDK1, IL2RA, IL7, ACTN3, SAP30BP, RPS6, NFKBIL1, ATM, NCSTN, CD38, NTRK1, CASP12, TNFAIP8, FAIM3, ACVR1 and IL2</i>	43 (7)

Functional biological process enrichment analysis of genes linked to promoter-localized DERs in the T_H2 cell gain category. The Gene Ontology (GO) term of the biological process (BP), enrichment index and P value (calculated using database for annotation, visualization and integrated discovery (DAVID) software; Online Methods), names and number of genes present in the T_H2 cell gain category (including the percentage of all genes in that biological process) are shown (full list in **Supplementary Table 5**).

antioxidants that are commonly used as food preservatives in the United States³¹.

Finally, we observed a highly significant overlap between our list of cell-specific and differentiation-related enhancers with binding sites for the 161 transcription factors profiled by the encyclopedia of DNA elements (ENCODE) project³² as well as 18 other CD4-related transcription factors (odds ratio > 2.5 and $P < 0.01$ empirical test, Online Methods; **Supplementary Table 10**). Over one-third (38%) of all our identified enhancers contained at least one known transcription factor binding site and ~17% of the enhancers had > 3 such sites (**Supplementary Table 10**), which supports the notion that a substantial fraction of the enhancers identified in our study are likely to be functionally important.

Asthma-associated SNPs are enriched in T_H2 cell enhancers

Disease-associated SNPs are enriched in regulatory regions that are active in cell types that contribute directly to disease³³; in this context, we have shown previously that asthma-associated SNPs are enriched in regulatory regions that are active in CD4⁺ T cells³⁴. Given that aberrant accumulation of T_H2 memory cells is a key event in asthma pathogenesis, we asked whether asthma-associated SNPs are more enriched in enhancers linked to differentiation of T_H2 as opposed to T_H1 memory cells. We observed the highest enrichment (~4.99 fold; $P < 0.0001$) of asthma SNPs in enhancers (**Fig. 2b**) that gained H3K4me2 during development of T_H2 cells rather than T_H1 cells (**Fig. 6a** and **Supplementary Table 11**). This enrichment held true when considering haplotype blocks instead of individual SNPs and




Figure 4 Upstream regulators of T_H2 cell genes. Induced gene-regulatory network analysis (performed using version 27 software from the ConsensusPathDB interaction database; Online Methods) of genes in the T_H2 cell gain category (listed in **Supplementary Table 4**) shows that MYC, E2F2, E2F4 are key upstream regulators of this subgroup of genes.

when limiting the source of SNPs that are significant in subjects from multiple study populations³⁵ (**Supplementary Table 11**). These data support a pathogenic role of T_H2 cells in asthma, and imply that SNPs

associated with asthma risk may perturb T_H2 memory cell differentiation by modulating the activity of enhancers that change dynamically during this developmental process. SNPs associated with some other




Figure 5 Enrichment of transcription factor binding motifs and sites in enhancers linked to CD4 memory differentiation. (a) Heat map shows known transcription factor (TF) binding motifs that were significantly enriched in each DER subgroup (analysis performed using HOMER accessing its motif database; Online Methods and **Supplementary Note**). Motifs with $P \leq 1.00 \times 10^{-3}$ and ratio of target sequences with motif versus background sequences with motif > 1.1 were defined as significantly enriched (**Supplementary Table 9**). (b) Selected TFs (independent analysis performed for 161 TFs profiled by the ENCODE project and 18 other CD4⁺ T cell-related TFs, Online Methods) that showed significant enrichment of their binding sites at genomic locations of cell-specific enhancers in the T_H2 cell gain subgroup (**Supplementary Table 10**). Shown are percentage of binding sites that overlap different cell-specific enhancer DER subgroups, the absolute number of binding sites that overlap all DERs and their percentage of the total genome-wide binding sites (in parentheses next to the name). Also shown is an example of a TF (SUZ12) whose binding sites are depleted in T_H2 cell gain DERs (last row). (c) ChIP-seq peak tracks (black bars) of transcription factors (T_H2 cell gain category) for human T_H2 cell cytokine locus (*IL4*, *IL13* and *RAD50*), *IL12A* and *NFE2L2* (encoding NRF2) along with UCSC gene tracks (top), multispecies gene conservation tracks (dark blue tracks), cell type-specific enhancer DERs and H3K27Ac track from ENCODE.




Figure 6 Asthma GWAS SNPs are enriched in T_H2 cell enhancers. **(a)** Enrichment values of asthma GWAS SNPs in T_H cell enhancer subgroups (**Fig. 2b**) and other cell tissue-specific enhancers³² (top) and for SNPs associated with other diseases (bottom; **Supplementary Table 11**). Enrichment values that did not reach significance (Chi-squared test, Online Methods) are shown in gray. ADMSC, adipose-derived mesenchymal stem cells. **(b)** Overlap of cell-specific DERs (shown in **Fig. 2b**) with asthma GWAS SNPs (top) and percentages of overlapping DERs or asthma SNPs in different DER subgroups (bottom). **(c)** UCSC tracks of *IL33–IL18R*, *IL5–RAD50–IL13–IL4* and *RORA* loci containing large haplotype blocks of asthma-associated SNPs (black lines indicate their genomic location, red lines are SNPs that overlap DERs), along with cell-specific DERs tracks and H3K4me2 tracks for each cell type (merged from all assays shown in **Fig. 2**). Graphs show H3K4me2 enrichment values for each asthma-SNP-associated DERs (500-bp regions harboring the asthma SNP; highlighted in purple dashed-line boxes in **c**) in T_H2 cells from the same H3K4me2 ChIP-seq assays shown in **Figure 2**. Each dot represents data from an independent assay; $n = 18$ assays from 10 healthy (HC) subjects, $n = 24$ assays from 12 asthmatic (AS) subjects; error bars indicate mean $\pm$ s.e.m.; *raw $P < 0.05$; **raw $P < 0.01$; NS, nonsignificant, calculated using MEDIPS.


Figure 7 Identification of asthma-associated enhancers. (a) MA plots (vertically displayed) illustrate genomic regions with differences in H3K4me2 enrichment (DERs) between healthy and asthmatic subjects in the three different cell types (Supplementary Table 12). Red dots and orange dots indicate windows with adjusted $P < 0.05$, or with raw $P < 0.005$, respectively (exact test for negative binomial distribution, using edgeR integrated in Bioconductor package MEDIPS). Z-scores (right) of normalized read counts for each asthma-associated DER (rows) identified in the T_H2 cells. (b) Manhattan plot illustrates the genome-wide distribution of asthma-associated DERs in relation to their statistical significance values (P values, MEDIPS; y-axis parameter). Red dashed line sets the threshold for an adjusted $P < 0.05$. (c) Comparison of H3K4me2 enrichment between healthy and asthmatic subjects in indicated cells. H3K4me2 tracks for each cell type were merged from all assays performed in healthy (HC) and asthmatic (AS) donors (same cohort as shown for the analysis above and in Fig. 2). (d) H3K4me2 enrichment values for each asthma-associated DER (highlighted in purple dashed line boxes in c from the same H3K4me2 ChIP-seq assays shown in Fig. 2). Each dot represents data from an independent assay; $n = 18$ assays from 10 healthy subjects (HC), $n = 24$ assays from 12 asthmatic patients (AS); error bars indicate mean $\pm$ s.e.m.; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ (MEDIPS).

immune system-mediated diseases showed preferential enrichment in different enhancer subgroups: SNPs associated with lupus and HIV risk were enriched in enhancers that lose and gain H3K4me2 enrichment in T_H1 cells, respectively, and SNPs associated with ankylosing spondylitis risk were enriched in enhancers that lose H3K4me2 enrichment in T_H2 cells (Fig. 6a and Supplementary Table 11).

Of the ~1,500 SNPs that have been associated with asthma, 52 were within the ~71,000 enhancer DERs associated with the development of $CD4^+$ T cell memory; conversely, 38 DERs harbored one or more SNPs associated with asthma risk (Fig. 6b–d). Most of the 38 enhancers that harbored SNPs associated with asthma risk, including the conserved T_H2 cell locus enhancers LCR-A and LCR-O, gained the H3K4me2 modification during T_H2 cell development; a few showed a trend toward differential enrichment in asthma subjects versus healthy controls (Fig. 6b–d). These data suggest that the underlying SNPs may influence enhancer strength and have a potential role in shaping the pathological gene expression patterns observed in disease.

Enhancers associated with asthma

Finally, we sought to identify $CD4^+$ T cell enhancers that differ in strength between healthy and asthmatic subjects. A total of 200 enhancer regions showed differential enrichment of H3K4me2 in the three cell types when we compared asthmatic versus healthy controls (Bonferroni-adjusted $P < 0.05$; Fig. 7a–c and Supplementary Table 12). As expected, most of these regions (here termed ‘asthma-associated enhancers’) were specific to the potentially pathogenic T_H2 memory



cell population (163 of 200 regions, of which 33 and 130 gained and lost H3K4me2 enrichment, respectively), with smaller numbers (29 and 11 regions) specific to naive T cells and T_H1 cells, respectively (Fig. 7a–d). Nearly 10% of these asthma-associated enhancers overlapped with the ~71,000 enhancer DERs associated with $CD4^+$ T cell memory, a significant enrichment (odds ratio = 8.74; $P = 1.7 \times 10^{-12}$, Online Methods) that suggests an important functional role for at least a subset of the enhancers. 84 of the 200 asthma-associated enhancers (42%) contained at least one transcription factor binding site (Supplementary Table 13), with significant enrichment for binding sites of transcription factors involved in T cell differentiation (for example, *GATA3*, *TBX21* and *RUNX3*; Supplementary Table 13). Enrichment analysis of candidate genes assigned to asthma-associated enhancers demonstrated significant overrepresentation of genes involved in chemokine and Toll-like receptor signaling pathways (KEGG pathway; $P = 0.0009$ calculated using statistical software from the ConsensusPathDB interaction database, Online Methods; Supplementary Table 14). Candidate genes

enriched in this pathway include *CCL3L1*, *CCL3L3* and *CCL4L2*, a cluster of genes on chromosome 17 that encodes chemokines known to bind CCR5 and prevent entry of HIV into T cells³⁶, and *AKT3* and *GNB4*, involved in phosphoinositide 3-kinase and G protein signaling (Fig. 7d). Further studies will clarify the role of these genes and enhancers in asthma pathogenesis.

DISCUSSION

We mapped the genome-wide distribution of H3K4me2 in naive, CD4⁺, T_H1 and T_H2 cells from 24 subjects. Through computational analysis of the data, we identified putative enhancer regions that are associated with specific T_H cell subsets. By comparing T_H cell subsets from healthy and asthmatic individuals, we identified enhancers that reproducibly differ in H3K4me2 enrichment and thus are presumed to be differentially active in T cells from asthma patients compared to healthy controls.

Our strategy can be applied to any accessible cell type with a function relevant to human health or disease. Purified cell populations are not necessarily needed: even though we performed ChIP-seq experiments on enriched cell populations rather than on purified T cell subsets, the majority of H3K4me2-marked enhancers that we identified were clearly associated with the enriched cell type. This was particularly true for T_H2 cells, which were present only in the CCR4⁺ population and depleted from the CCR4[−] population; in contrast, IFN- γ -producing T_H1 cells were present in both the CCR4⁺ and CCR4[−] populations, which explains why overall we identified fewer enhancers as specifically T_H1 cell-associated. To facilitate studies of rare cell populations involved in disease pathogenesis, we optimized our H3K4me2 ChIP-seq assay so that it can be applied to very small numbers of cells. This feature will be valuable not only for studies of the immune system but in cases where only limited amounts of biological materials are available for research. Our method offers a substantial advantage over transcriptional profiling by microarray or conventional RNA-seq, which identifies only those genes that are represented by steady-state transcripts at the time that the cells are lysed. In contrast, H3K4me2 ChIP-seq has the potential to identify promoters and enhancers that control the expression of genes that are not actively being transcribed but are poised for expression in response to extrinsic stimuli.

We showed that *in silico* analyses of enhancer profiles obtained by H3K4me2 ChIP-seq can be used to predict binding sites for transcriptional regulators that set up gene expression programs in specific cell types and disease states. Many such transcription factors are only transiently expressed at specific developmental stages that may be difficult to isolate *in vivo*, which implies that they cannot be identified merely through expression profiling of fully differentiated cells. However, lineage-defining or disease-initiating transcription factors often leave a footprint of their activity on the *cis*-regulatory landscape of a cell, by altering H3K4me2 levels at promoters and enhancers during cellular differentiation or early disease progression. Thus, H3K4me2 profiling of peripheral blood cells is particularly relevant for studies of the immune system in patients: differentiation of memory T cells occurs in relatively inaccessible compartments such as lymph nodes, spleen or thymus, and the majority of patients present in the clinic well after the first disease-initiating events have occurred.

Consistent with the idea that T_H2 cell function is aberrant in asthma, we found that T_H2 cell-specific enhancers were highly enriched for asthma-associated SNPs identified in genome-wide association studies (GWAS). Our approach has narrowed down the large list of SNPs in asthma-associated haplotype blocks to those that are likely to be

functional in cells relevant to asthma pathogenesis by determining whether they are located in a putative enhancer region, and whether that enhancer shows evidence of altered function during memory T cell differentiation. The candidate enhancer-localized SNPs identified in our study can now be formally tested for functional importance in disease pathogenesis.

Separate from GWAS analysis, we identified enhancers associated with asthma susceptibility simply by comparing T_H cell enhancer profiles of healthy and asthmatic subjects as defined by H3K4me2 enrichment patterns. By analogy with GWAS, such approaches have been termed epigenome-wide association studies³⁷. An important consideration is the need to confirm the link between disease-associated enhancers and their target genes, for instance, by using new techniques to detect physical interactions between enhancers and gene promoters^{38,39}; moreover, detailed testing of the candidate genes for function in human helper T cells will be necessary to attribute an important role in asthma pathogenesis. Notably, the 12 asthma patients evaluated in our study are a very small patient cohort; replication of our hits in multiple independent and larger cohorts is needed to strengthen their biological importance and to address the important question of whether epigenome-wide association studies can be used to distinguish molecular subtypes of asthma.

METHODS

Methods and any associated references are available in the [online version of the paper](#).

Accession codes. Gene Expression Omnibus: [GSE53646](#) (sequencing data).

Note: Any Supplementary Information and Source Data files are available in the online version of the paper.

ACKNOWLEDGMENTS

We thank the staff at the Wellcome Trust Clinical Research Facility (University of Southampton) where samples were acquired from volunteers; M. North for assisting in patient recruitment, assessment and sample collection; R. Jewel and C. McGuire for providing assistance in the flow cytometry facility (University of Southampton); J. Day for assistance with high-throughput sequencing at the La Jolla Institute for Allergy and Immunology sequencing facility, and A. Moghaddas Gholami at the La Jolla Institute for Allergy and Immunology bioinformatics core for help with the SNP enrichment analysis. L.C. is funded by a Feodor Lynen Research Fellowship from the Alexander von Humboldt Foundation. This work was supported by the Dana Foundation (K.M.A.), GlaxoSmithKline National Clinician Scientist Fellowship Award and Peel Travel Fellowship Award (P.V.), R01 HL114093 (to B.P., A.R. and P.V.) and U19 AI100275 (to B.P., A.R. and P.V.).

AUTHOR CONTRIBUTIONS

G.S., K.M.A., B.P., A.R. and P.V. conceived the work, designed, performed and analyzed experiments, and wrote the paper; N.O., L.K., M.V. and A.P.V.G. assisted in the performing some of the experiments under the supervision of G.S. and P.V.; R.D. provided support and direction for obtaining and processing clinical specimens; L.C. identified DERs; and A.G., M.L. and A.C. performed the bioinformatic analysis under the supervision of L.C. and B.P.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

Reprints and permissions information is available online at <http://www.nature.com/reprints/index.html>.

1. Ahmed, R. & Gray, D. Immunological memory and protective immunity: understanding their relation. *Science* **272**, 54–60 (1996).
2. Ansel, K.M., Lee, D.U. & Rao, A. An epigenetic view of helper T cell differentiation. *Nat. Immunol.* **4**, 616–623 (2003).
3. Kay, A.B. Allergy and allergic diseases. Second of two parts. *N. Engl. J. Med.* **344**, 109–113 (2001).

4. Ober, C. & Yao, T.C. The genetics of asthma and allergic disease: a 21st century perspective. *Immunol. Rev.* **242**, 10–30 (2011).
5. Gregersen, P.K. & Olsson, L.M. Recent advances in the genetics of autoimmune disease. *Annu. Rev. Immunol.* **27**, 363–391 (2009).
6. WHO fact sheets N206 and N307 <http://www.who.int/mediacentre/factsheets/fs206/en/>; <http://www.who.int/mediacentre/factsheets/fs307/en/index.html> (accessed November 2013).
7. Holgate, S.T. & Polosa, R. Treatment strategies for allergy and asthma. *Nat. Rev. Immunol.* **8**, 218–230 (2008).
8. Wenzel, S.E., Wang, L. & Pirozzi, G. Dupilumab in persistent asthma. *N. Engl. J. Med.* **369**, 1276 (2013).
9. Ansel, K.M., Djuretic, I., Tanasa, B. & Rao, A. Regulation of T_H2 differentiation and I κ B locus accessibility. *Annu. Rev. Immunol.* **24**, 607–656 (2006).
10. Vijayanand, P. *et al.* Interleukin-4 production by follicular helper T cells requires the conserved I κ B enhancer hypersensitivity site V. *Immunity* **36**, 175–187 (2012).
11. Loots, G.G. *et al.* Identification of a coordinate regulator of interleukins 4, 13, and 5 by cross-species sequence comparisons. *Science* **288**, 136–140 (2000).
12. Ernst, J. *et al.* Mapping and analysis of chromatin state dynamics in nine human cell types. *Nature* **473**, 43–49 (2011).
13. Creighton, M.P. *et al.* Histone H3K27ac separates active from poised enhancers and predicts developmental state. *Proc. Natl. Acad. Sci. USA* **107**, 21931–21936 (2010).
14. Zhang, J.A., Mortazavi, A., Williams, B.A., Wold, B.J. & Rothenberg, E.V. Dynamic transformations of genome-wide epigenetic marking and transcriptional control establish T cell identity. *Cell* **149**, 467–482 (2012).
15. Koche, R.P. *et al.* Reprogramming factor expression initiates widespread targeted chromatin remodeling. *Cell Stem Cell* **8**, 96–105 (2011).
16. Vijayanand, P. *et al.* Chemokine receptor 4 plays a key role in T cell recruitment into the airways of asthmatic patients. *J. Immunol.* **184**, 4568–4574 (2010).
17. Mikhak, Z., Strassner, J.P. & Luster, A.D. Lung dendritic cells imprint T cell lung homing and promote lung immunity through the chemokine receptor CCR4. *J. Exp. Med.* **210**, 1855–1869 (2013).
18. Zielinski, C.E. *et al.* Pathogen-induced human TH17 cells produce IFN- γ and are regulated by IL-1 β . *Nature* **484**, 514–518 (2012).
19. Chavez, L. *et al.* Computational analysis of genome-wide DNA methylation during the differentiation of human embryonic stem cells along the endodermal lineage. *Genome Res.* **20**, 1441–1450 (2010).
20. Lienhard, M., Grimm, C., Morkel, M., Herwig, R. & Chavez, L. MEDIPS: genome wide differential coverage analysis of sequencing data derived from DNA enrichment experiments. *Bioinformatics* **30**, 284–286 (2014).
21. Wei, G. *et al.* Global mapping of H3K4me3 and H3K27me3 reveals specificity and plasticity in lineage fate determination of differentiating CD4⁺ T cells. *Immunity* **30**, 155–167 (2009).
22. Ansel, K.M. *et al.* Deletion of a conserved I κ B silencer impairs T helper type 1-mediated immunity. *Nat. Immunol.* **5**, 1251–1259 (2004).
23. Wilson, C.B., Rowell, E. & Sekimata, M. Epigenetic control of T-helper-cell differentiation. *Nat. Rev. Immunol.* **9**, 91–105 (2009).
24. Seumois, G. *et al.* An integrated nano-scale approach to profile miRNAs in limited clinical samples. *Am. J. Clin. Exp. Immunol.* **1**, 70–89 (2012).
25. Douglas, N.C., Jacobs, H., Bothwell, A.L. & Hayday, A.C. Defining the specific physiological requirements for c-Myc in T cell development. *Nat. Immunol.* **2**, 307–315 (2001).
26. Wang, R. *et al.* The transcription factor Myc controls metabolic reprogramming upon T lymphocyte activation. *Immunity* **35**, 871–882 (2011).
27. Zhu, J.W. *et al.* E2F1 and E2F2 determine thresholds for antigen-induced T-cell proliferation and suppress tumorigenesis. *Mol. Cell. Biol.* **21**, 8547–8564 (2001).
28. Pandiyan, P. *et al.* CD152 (CTLA-4) determines the unequal resistance of Th1 and Th2 cells against activation-induced cell death by a mechanism requiring PI3 kinase function. *J. Exp. Med.* **199**, 831–842 (2004).
29. Kim, T.H. *et al.* Analysis of the vertebrate insulator protein CTCF-binding sites in the human genome. *Cell* **128**, 1231–1245 (2007).
30. Hawkins, R.D. *et al.* Global chromatin state analysis reveals lineage-specific enhancers during the initiation of human T helper 1 and T helper 2 cell polarization. *Immunity* **38**, 1271–1284 (2013).
31. Rockwell, C.E., Zhang, M., Fields, P.E. & Klaassen, C.D. Th2 skewing by activation of Nrf2 in CD4⁺ T cells. *J. Immunol.* **188**, 1630–1637 (2012).
32. Gerstein, M.B. *et al.* Architecture of the human regulatory network derived from ENCODE data. *Nature* **489**, 91–100 (2012).
33. Maurano, M.T. *et al.* Systematic localization of common disease-associated variation in regulatory DNA. *Science* **337**, 1190–1195 (2012).
34. Gerasimova, A. *et al.* Predicting cell types and genetic variations contributing to disease by combining GWAS and epigenetic data. *PLoS ONE* **8**, e54359 (2013).
35. Moffatt, M.F. *et al.* A large-scale, consortium-based genomewide association study of asthma. *N. Engl. J. Med.* **363**, 1211–1221 (2010).
36. Dolan, M.J. *et al.* CCL3L1 and CCR5 influence cell-mediated immunity and affect HIV-AIDS pathogenesis via viral entry-independent mechanisms. *Nat. Immunol.* **8**, 1324–1336 (2007).
37. Rakan, V.K., Down, T.A., Balding, D.J. & Beck, S. Epigenome-wide association studies for common human diseases. *Nat. Rev. Genet.* **12**, 529–541 (2011).
38. Jin, F. *et al.* A high-resolution map of the three-dimensional chromatin interactome in human cells. *Nature* **503**, 290–294 (2013).
39. Zhang, Y. *et al.* Chromatin connectivity maps reveal dynamic promoter-enhancer long-range associations. *Nature* **504**, 306–310 (2013).

ONLINE METHODS

Study subjects and sample processing. The ethics committees of the Southampton University Hospitals Trust and La Jolla Institute approved the study, and written informed consent was obtained from all subjects. Twelve nonsmoking subjects with asthma (6 with mild asthma never treated with corticosteroids and 6 subjects with moderate asthma treated with inhaled corticosteroids¹⁶), meeting established diagnostic criteria⁴⁰, and 12 healthy subjects were studied (**Supplementary Table 15**). Subjects with mild asthma had symptoms <3 times a week, with forced expiratory volume in 1 s (FEV₁) >80% of predicted, and used short-acting inhaled β_2 -agonists as needed for symptom relief. Subjects with moderate asthma were additionally treated with regular inhaled corticosteroids and inhaled short-acting β_2 -agonists. All asthmatic subjects were atopic and allergic to house dust mite (*Dermatophagoides pteronyssinus*). All study subjects were Caucasian adults (10 females and 14 males, age range 18–65, inclusion criterion). Healthy controls had no history of smoking or respiratory symptoms suggestive of asthma.

For isolating T cell subtypes from peripheral blood samples, peripheral blood mononuclear cells (PBMCs) were first separated into a CD4⁺ memory cell fraction and remaining cells by use of the memory CD4⁺ T cell isolation kit (Miltenyi Biotec). The CD4⁺ memory cells were then stained with a cocktail of fluorescently conjugated antibodies (anti-CD45RA FITC-conjugated (clone HI100), anti-CD4 APC-Cy7-conjugated (clone SK3), anti-CCR4 PE-conjugated (clone 1G1), and anti-CD25 APC-conjugated (clone M-A251) and sorted on the FACS Aria to obtain two cell populations: CD4⁺CD45RA⁺CCR4⁺ and CD4⁺CD45RA⁺CCR4⁺CD25⁺ (**Supplementary Fig. 1**). Naive T cells were sorted from the remaining cells after CD4⁺ memory cell isolation by staining with anti-CD45RA FITC-conjugated, anti-CD4 PerCP-Cy5.5-conjugated (clone SK3), anti-CD62L APC-Cy7-conjugated (clone DREG-56) antibodies and anti-CD45RO PE-conjugated (clone UCHL1). Sorted cells were washed and fixed for 10 min at 20 °C with 1:10 dilution of 11% formaldehyde (in 50 mM HEPES, pH 7.5, 100 mM NaCl, 1 mM EDTA and 0.5 mM EGTA), neutralized with 1:20 dilution of 2.5 M glycine, washed twice in ice-cold phosphate buffered saline, and cell pellets were snap frozen in liquid nitrogen before storage at –80 °C.

Microscaled multisample chromatin immunoprecipitation–sequencing. Formaldehyde-fixed cell pellets were snap-frozen in liquid nitrogen and stored at –80 °C before the ChIP assay. This storage step gave us the flexibility to batch multiple samples for ChIP-seq. Frozen cells were lysed in 120 μ l of lysis buffer (50 mM Tris-HCl, pH 8.0, 10 mM EDTA, 1% SDS, 1 mM phenyl methane sulfonyl fluoride, 20 mg/ml sodium butyrate and proteinase inhibitor cocktail; Sigma), and chromatin was sheared by sonication, using Bioruptor (Diagenode), to generate 100–500 bp DNA fragments. Shearing multiple samples ($n = 12$) in parallel using the Bioruptor ensured equivalent shearing of chromatin among the samples that were to be compared. For the initial validation of microscaled ChIP-seq, chromatin from 10⁵, 10⁴ or 10³ whole-cell equivalents (D10.G4.1 cells, ATCC, mycoplasma-negative) was immunoprecipitated with anti-H3K4me2 (clone Y47, lot#YH041501C; Abcam). For samples from study subjects, chromatin from 10⁵ cells was used for immunoprecipitation. Standard H3K4me2 ChIP using 2–10 $\times$ 10⁶ cells was performed as described previously¹⁰. For microscaled H3K4me2 ChIP-seq, chromatin was diluted in 1 ml of radioimmunoprecipitation assay (RIPA) buffer and immunoprecipitation was done overnight at 4 °C by incubating 1 μ l of antibody precoated on 10 μ l protein A coated magnetic beads (Invitrogen). To reduce nonspecific DNA binding (background signal in ChIP-seq assays), we used pipette tips and tubes with low DNA and protein binding (Axygen Maximum recovery). Immunocomplexes were captured and washed for 5 min each with RIPA buffer, high-salt buffer (50 mM Tris-HCl, pH 8, 500 mM NaCl, 0.1% SDS, 0.5% Na-deoxycholate, 1% Nonidet-P40 and 1 mM EDTA), LiCl buffer (50 mM Tris-HCl, pH 8, 250 mM LiCl, 1 mM EDTA, 1% Nonidet-P40 and 0.5% Na-deoxycholate), and low-salt buffer (10 mM Tris-HCl, pH 8, 1 mM EDTA, 50 mM NaCl). Beads were resuspended in TE (10 mM Tris-HCl, pH 8, 1 mM EDTA), transferred to fresh tubes, captured and then resuspended in 200 μ l of elution buffer (50 mM Tris-HCl, pH 8, 10 mM EDTA and 1% SDS). Chromatin was detached from the beads by incubating for 15 min at 65 °C, transferred to fresh tubes, then treated with 2 μ l of RNase A (20 mg/ml; Invitrogen) for 30 min at 37 °C and with proteinase K (0.2 mg/ml, Invitrogen)

for 6 h at 55 °C, and incubated overnight at 65 °C. DNA was purified using affinity columns (Zymo Research) and eluted in 12 μ l of TE (10 mM Tris-HCl, pH 8 and 0.1 mM EDTA). Selected DNA sequences from control regions (*IL4* promoter, *IFNG* promoter, *GAPDH* promoter and *HOX7A* promoter) were quantified by real-time quantitative PCR to assess the quality of the ChIP. The amount of DNA in the ChIP samples was quantified using picogreen assay (Invitrogen). One nanogram of ChIP DNA was PCR-amplified for 18–22 cycles using whole genome amplification (WGA) primers (WGA-SEQX, Sigma) following the manufacturer's instructions. 2 μ g of this amplified DNA was treated with restriction enzyme (WGA-SEQX, Sigma), to remove the WGA primer sequences and then purified using AmpureXP beads (Beckman Coulter). Efficiency of removal of WGA primer sequences was assessed by PCR. From this step, 500 ng of purified DNA was diluted with TE to obtain a total volume of 65 μ l. Diluted DNA was sonicated with E220 Covaris multiplex sonicator (Covaris) to generate 100–250 bp DNA fragments. ~250 ng of DNA was used for preparing standard SOLiD sequencing library (5500 SOLiD Fragment 48 Library Core Kit & Fragment Library Barcode Adaptors 1-96). Following emulsion PCR, samples were sequenced on the SOLiD 5500 sequencer to obtain 35-bp single-end reads (SOLiD EZ Bead E120 System kits). Both whole-genome amplification and sequencing-library preparation were performed in a 96-well format, which reduced hands-on time, beside reducing technical and assay-to-assay variability. Multiple quality control steps (**Supplementary Fig. 3**) were included to determine optimal shearing of DNA, ChIP efficiency and number of PCR amplification cycles, to assess removal of whole-genome PCR amplification adaptors, efficiency of ligation of sequencing adaptors and to determine library complexity. Samples that failed quality control were eliminated from downstream steps. Reproducibility and correlation between standard and microscaled ChIP-seq, performed using various cell numbers, is shown in **Figures 1a–c** and **Supplementary Figures 4** and **5**.

RNA sequencing. Total RNA was purified using miRNAeasy kit (Qiagen) and quantified as described previously²⁴. 10–15 ng of purified total RNA was used for poly(A) section (Poly(A)Purist Mag kit, Life Technologies). Poly(A)-selected RNA was amplified with the Whole Transcriptome Amplification Sequencing Technology kit (SEQR, Sigma) as per manufacturer's recommendation. 1 μ g of this amplified cDNA was treated with restriction enzyme (SEQR, Sigma), to remove the primer sequences and then purified using AmpureXP beads (Beckman Coulter). Efficiency of removal of SEQR primer sequences was assessed by PCR. From this step, 250 ng of purified DNA was sonicated with E220 Covaris multiplex sonicator (Covaris) to generate 100–250 bp DNA fragments. Sonicated cDNA was used for preparing standard SOLiD sequencing library as described in the previous section for ChIP-seq.

Mapping and primary data processing of sequencing reads. Human ChIP-seq data (csfasta and qual files) were mapped against the human reference genome hg19 (downloaded from the Genome Bioinformatics database of the University of California Santa Cruz) using bowtie v.0.12.7 (-n 2 -m 1 -C)⁴¹. 120 different ChIP-seq assays were performed and the obtained sam files were converted and merged into separated bam files using samtools v.0.1.18 (ref. 42). For visualization of the ChIP-seq data in public genome browsers, sequencing coverage was calculated at genome-wide 50-bp windows after extending each read to a length of 120 bp along the sequencing direction using MEDIPS v.1.10.0 (extend = 120, uniq = F, window_size = 50, BSgenome = "BSgenome.Hsapiens.UCSC.hg19")^{19,20}, and the resulting coverage profiles (RPKM) were exported as wiggle files. Mouse ChIP-seq data were mapped against the mouse reference genome mm10 as described for the human ChIP-seq data (bowtie v.0.12.7 -n 2 -m 1 -C). Human RNA-seq data was mapped against hg19 using tophat⁴³ (v1.4.1, -library-type fr-secondstrand -C) and the RefSeq gene annotation downloaded from the UCSC genome Bioinformatics site. Sequencing read coverage per gene was counted using HTSeq-count (-m union -s yes -t exon -i gene_id, <http://www-huber.embl.de/users/anders/HTSeq/>).

Correlation between ChIP-sequencing assays and samples. Pairwise Spearman correlation between samples were calculated by comparing the sequencing coverage at genome-wide 500-bp windows using MEDIPS v.1.10.0 (extend = 120, uniq = F, window_size = 500, BSgenome = "BSgenome.Mmusculus.UCSC.mm10") (**Supplementary Figs. 2** and **4**).

Identification of differentially enriched *cis*-regulatory regions comparing cell types. To identify cell type-specific DERs, H3K4me2 ChIP-seq data were grouped according to their cell types: naive CD4⁺ T cells, ($n = 42$), T_H2 cells ($n = 42$) and T_H1 cells ($n = 36$). All technical and biological replicates were independently processed ChIP assays with separate ChIP, whole-genome amplification and sequencing, and each ChIP replicate was sequenced to a targeted depth of $\sim 20 \times 10^6$ mapped reads. Additional details are available in **Supplementary Note**.

Sample sizes for ChIP-seq were chosen to achieve power to detect consistent differences between asthmatic and healthy donors in the presence of variability inherent to the assay. Assay variability was assessed based on biological replicates of mouse cell line data, and differences were introduced computationally at different sites. Recovery of these sites using MEDIPS was used to estimate error rates depending on sample size. To estimate a false discovery rate for our chosen significance threshold to identify DERs (Bonferroni-adjusted $P < 0.05$), we performed a random approach by permuting the 42 naive CD4⁺ T cells and the 42 T_H2 cell ChIP-seq assays resulting in two groups of intermixed samples. Subsequently, we applied the same statistical framework used for our original approach. In ten independent iterations of such random sample permutations, we observed a very small number of DERs ranging from 4 to 11 (data not shown) compared to 51,261 DERs detected for the correct assignment of samples at the same significance level of a Bonferroni adjusted $P < 0.05$. This translates to a very low false discovery rate (FDR) of 0.02%, suggesting that the P -value threshold used for our analysis is stringent and does not result in overcalling of DERs. Further, in our approach, the detectable effect size is dependent on enhancer strength (see MA plots where the 'enhancer strength' is represented on the x axis). With our sample size (naive = 42, T_H2 = 42) and the inherent variation between samples, we can detect differences between cell types with a ratio ('effect size') of at least 1.6 in the lowest 25% quantile and of at least 1.2 in the upper 25% quantile of signal-enhancer strengths.

Assignment of target genes to DERs. To assign DERs to transcripts and genes, we downloaded the hg19 table refGene (track RefSeq Genes via the Table browser at the UCSC Genome Bioinformatics site (17 October 2013), and assigned DERs to promoters if their midpoint fell within ± 1 kb of a transcription start site. This identified 2,235 promoter DERs associated with 1,710 transcripts of 1,529 genes. We classified the remaining 69,405 non-promoter-localized DERs as enhancers. To associate enhancer DERs to genes, we considered CTCF binding sites previously identified in 19 diverse human cell types⁴⁴ as boundaries for defining extended transcript loci. For each transcript, their extended locus was defined as the upstream and downstream region around their start sites limited by the occurrence of a CTCF binding site or by a maximal distance of 200 kb. We observed 25,888 DERs that fell into the extended loci of 8,983 transcripts of 7,647 unique genes.

Correlation of H3K4me2 enrichment patterns at promoters and enhancers. To examine putative concordant changes of H3K4me2 enrichment at promoters of a transcript and enhancers located within or close to that transcript, we considered the following two groups of transcripts: (i) transcripts containing a DER in their promoter region with gain of H3K4me2 in T_H2 cells compared to naive CD4⁺ T cells (T_H2 gain, $n = 534$), (ii) transcripts containing a DER in their promoter region with loss of H3K4me2 in T_H2 cells compared to naive CD4⁺ T cells ($n = 405$, T_H2 cell loss). In addition, we defined the following four groups of genomic regions: upstream: -20 kb to a transcript start site, promoter: ± 1 kb around a transcript start site, transcript body: between a transcript start and end site, and downstream: $+20$ kb from a transcript end site. We divided the upstream and downstream regions into ten equally spaced windows of 2 kb but considered the promoter as one window and the transcript body as another window. Subsequently, we calculated sequencing read coverage at the predefined windows of the selected transcripts for each of the 42 naive CD4⁺ T cell and for each of the 42 T_H2 cell samples using MEDIPS v1.12.0 (function MEDIPS.createROIset, extend = 120, uniq = F, bn = 10 (for upstream and downstream) or bn = 1 (for promoter and transcript bodies), BSgenome = "BSgenome.Hsapiens.UCSC.hg19"). For each transcript and for each tested window, mean counts over all samples per group were calculated. In order to avoid division by 0, we added 1 to each tested window in both groups, before calculation of fold change between groups at individual

transcripts and windows. For each transcript, we calculated the mean fold change over all 2-kb windows in the upstream or downstream regions, respectively. The heat map in **Supplementary Figure 10** shows the $\log_2$ of fold changes at the four tested groups of genomic regions from left to right and at individual transcripts of groups i and ii from top to bottom. Transcripts in groups i and ii are ordered by the sum of $\log_2$ fold changes across the four tested genomic regions, where group i is sorted from highest (top) to lowest (bottom), and group ii is sorted from lowest (top) to highest (bottom).

Principal component analysis of the RNA-sequencing data and differential gene expression analysis. The principal component analysis (**Supplementary Fig. 11**) has been performed by calculating singular value decomposition of the centered, variance-stabilized and library-normalized counts of the 5,000 most variable genes, using the R functions svd (base package, <http://www.r-project.org/>) and varianceStabilizingTransformation (Bioconductor package DESeq⁴⁵). To identify differential gene expression between cell types, we performed negative binomial tests for pairwise comparisons of the naive CD4⁺, T_H2 and T_H1 cells employing the Bioconductor package DESeq⁴⁵ and allowing for a false discovery rate of 1%.

Correlation of gene expression with H3K4me2 enrichment patterns at promoters and enhancers. This is described in **Supplementary Note**.

Functional gene-set enrichment analysis. Gene sets obtained for each promoter-localized DER category (classified based on preferential gain or loss of H3K4me2 enrichment during memory differentiation; **Supplementary Table 3**) were analyzed using DAVID (version 6.750; refs. 46,47). Biological process and pathways enriched for genes present in our data set are shown in **Supplementary Table 5**. As some of the gene sets obtained for enhancer-DER groups exceeded the maximum limit of 3,000 genes for the DAVID software, we additionally used the software from the ConsensusPathDB interaction database (CPDB; version 27; refs. 48,49) for the overrepresentation analysis. Biological pathways and processes enriched for genes present in our data set are shown in **Supplementary Tables 5, 7 and 14**). To determine upstream regulators of genes in the T_H2 cell (CCR4⁺) gain category, we performed induced gene-regulatory network analysis using the software from CPDB^{48,49}. Analysis was performed using default settings on the software and allowing for intermediate nodes with a Z-score threshold of 20 (**Fig. 4**).


Detecting enrichment of transcription factor motifs and binding at cell-specific enhancers. This is described in **Supplementary Note**.

Identification of asthma-associated DERs. To identify differentially enriched regions (DERs) between healthy and asthmatic donors, we grouped the ChIP-seq data of samples into the following six groups: (i) naive CD4⁺ T cells, healthy ($n = 18$), (ii) naive CD4⁺ T cells, asthma ($n = 24$), (iii) T_H2 cells, healthy ($n = 18$), (iv) T_H2 cells, asthma ($n = 24$), (v) T_H1 cells, healthy ($n = 13$), and (vi) T_H1 cells, asthma ($n = 23$). Differential sequence coverage was calculated for each cell type between healthy and asthma samples as described above to identify cell type-specific DERs. This resulted in 29 DERs for naive CD4⁺ T cells (4 asthma > healthy, and 25 asthma < healthy), 163 DERs for T_H2 cells (33 asthma > healthy, and 130 asthma < healthy), and 11 DERs for T_H1 cells (0 asthma > healthy, and 11 asthma < healthy) (Bonferroni-adjusted $P < 0.05$). In total, there were 203 significant windows in the three comparisons falling into 200 distinct asthma DERs. As above, we divided DERs into those that fall into promoters and enhancers. There were 13 promoter-localized DERs associated with 15 transcripts of 14 genes. A total of 86 nonpromoter DERs were located in the extended loci of 129 transcripts of 111 unique genes. Transcription factor binding site enrichment analysis for the asthma-associated DERs has been calculated as described above for the cell-specific enhancers. Owing to the relatively small number of DERs, we considered the entire 500-bp DER and according random regions for calculating overlaps to transcription factor binding sites.

SNPs data sets and enrichment calculation. 62 significant lead SNPs ($P < 1.0 \times 10^{-5}$) from GWAS studies of asthma were obtained by querying the HaploReg database⁵⁰. Cross-referencing the P values of these lead SNPs using

the GWAS Integrator database⁵¹ and the original articles confirmed that all SNPs had $P < 1.0 \times 10^{-5}$. An additional 20 lead SNPs associated with asthma-related traits (plasma eosinophil counts, serum IgE and YKL-40 concentrations) were obtained from the GWAS Integrator database, after manual review of the originating studies. Details of all 82 lead SNPs, P values and source study are provided in **Supplementary Table 11a**. For each lead SNP, SNPs in tight genetic linkage ($r^2 > 0.8$) were retrieved based on data from the phase I of the 1,000 genome project using European (EUR) as the reference population⁵² (calculations performed using HaploReg v2.0, **Supplementary Table 11b**). The total number of asthma-associated SNPs in the combined set (lead SNPs + linked SNPs) was 1,528. For all other diseases, significant lead SNPs ($P < 1.0 \times 10^{-5}$) were obtained from the HaploReg⁵⁰ database, and no additional traits were considered. Linked SNPs were calculated as described for the asthma SNPs above. More detailed description is available in **Supplementary Note**.


40. Bousquet, J. Global initiative for asthma (GINA) and its objectives. *Clin. Exp. Allergy* **30** (suppl. 1), 2–5 (2000).
41. Langmead, B., Trapnell, C., Pop, M. & Salzberg, S.L. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol.* **10**, R25 (2009).
42. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).
43. Trapnell, C., Pachter, L. & Salzberg, S.L. TopHat: discovering splice junctions with RNA-Seq. *Bioinformatics* **25**, 1105–1111 (2009).
44. Wang, H. *et al.* Widespread plasticity in CTCF occupancy linked to DNA methylation. *Genome Res.* **22**, 1680–1688 (2012).
45. Anders, S. & Huber, W. Differential expression analysis for sequence count data. *Genome Biol.* **11**, R106 (2010).
46. Huang da, W., Sherman, B.T. & Lempicki, R.A. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat. Protoc.* **4**, 44–57 (2009).
47. Huang da, W., Sherman, B.T. & Lempicki, R.A. Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res.* **37**, 1–13 (2009).
48. Kamburov, A., Stelzl, U., Lehrach, H. & Herwig, R. The ConsensusPathDB interaction database: 2013 update. *Nucleic Acids Res.* **41**, D793–D800 (2013).
49. Kamburov, A., Wierling, C., Lehrach, H. & Herwig, R. ConsensusPathDB—a database for integrating human functional interaction networks. *Nucleic Acids Res.* **37**, D623–D628 (2009).
50. Ward, L.D. & Kellis, M. HaploReg: a resource for exploring chromatin states, conservation, and regulatory motif alterations within sets of genetically linked variants. *Nucleic Acids Res.* **40**, D930–D934 (2012).
51. Yu, W. *et al.* GWAS Integrator: a bioinformatics tool to explore human genetic associations reported in published genome-wide association studies. *Eur. J. Hum. Genet.* **19**, 1095–1099 (2011).
52. Abecasis, G.R. *et al.* An integrated map of genetic variation from 1,092 human genomes. *Nature* **491**, 56–65 (2012).



Supplementary Figure 1

Schematic model summarizing the study design and cell sorting strategy.


Schematic diagram depicts the cell types isolated from peripheral blood of healthy and asthmatic subjects. Sorting strategy for isolating naïve and CCR4⁻ (T_H1), CCR4⁺ (T_H2) memory T cells from peripheral blood mononuclear cells (PBMC), and FACS plots pre and post-sorting are shown. The number of samples processed (passing quality control checks) for H3K4me2 ChIP-Seq assay is shown below.



Supplementary Figure 2


Reproducibility of standard H3K4me2 ChIP-seq assay.

Density plots show pair-wise comparison of sequencing coverage (number of reads) at genome-wide 500 bp windows (*MEDIPS* v.1.10.0 software, **Methods** and **Supplementary Notes**) obtained from 6 independent standard H3K4me2 ChIP-Seq assays performed with 2×10^6 D10 cells. The amount of DNA (post H3K4me2 ChIP; 15ng or 30ng) used for whole genome amplification (see **Methods** and **Supplementary Notes**) is shown. Pairwise Spearman correlation values (numbers inside boxes) for genome-wide comparison of H3K4me2 enrichment patterns between replicate samples (labeled as Rep 1-6) are illustrated.



Supplementary Figure 3


Workflow of microscaled ChIP-seq assay from sample isolation to sequencing.



Supplementary Figure 4

Reproducibility of the micro-scaled H3K4me2 ChIP-seq assay.


(a) Density plots show pair-wise comparison of sequencing coverage at genome-wide 500bp windows (*MEDIPS* v.1.10.0 software, **Methods** and **Supplementary Notes**) obtained from standard ChIP-Seq (2 x 10⁶ cells) and micro-scaled ChIP-Seq (10⁵, 10⁴ and 10³ cell samples) performed with D10 cells. Pairwise Spearman correlation values (numbers inside boxes) for genome-wide comparison of H3K4me2 enrichment patterns between assays performed with different cell numbers are illustrated. (b) Shows density plots and pairwise Spearman correlation between multiple (n = 11) replicate micro-scaled ChIP-Seq assays (10⁵ cell samples) performed in two separate sequencing runs (Run1 and Run2). (c) ROC analysis (detailed methodology is described in **Methods** and **Supplementary Notes**) shows the percentage of true and false positives identified by micro-scaled ChIP-Seq assay (10⁵ cell sample) when tested for the top 1% of enriched windows (true positives) identified in the standard ChIP-Seq assay.



Supplementary Figure 5

Reproducibility of the microscaled H3K4me2 ChIP-seq assay.


ChIP-Seq analysis showing H3K4me2 enrichment patterns from each assay (rows), for the following gene loci: control regions: *STIM 1*, *NUP98*, *SELP* and *SELL* gene loci; non-expressed gene: *SELE* locus; TH2 cell-type specific regions: *CCR4* and *CCR6* locus; TH1 cell-type specific regions: *TBX21* (encoding T-BET), performed in peripheral blood naïve, TH1 and TH2 memory T cells from all study subjects. The significant cell-specific H3K4me2 enrichment across (enhancers and promoters) in these loci is highlighted in the red dashed line boxes.



Supplementary Figure 6

Sensitivity of the H3K4me2 ChIP-seq assay.

ChIP-Seq analysis showing cell-specific H3K4me2 enrichment patterns, for the following gene loci: *HNRPLL*, *ADAM19*, *miR155* and *miR221-222*, in naive, T_H2 and T_H1 cells. For each specific cell-type, data was merged from all donors including assay duplicates. H3K4me2 enrichment values for specific 500 bp windows (highlighted in red boxes) are shown in the graphs below. Each dot represents data from a single assay-donor; error bars indicate mean $\pm$ (s.e.m.).




Supplementary Figure 7

Bioinformatic analysis of H3K4me2 ChIP-seq data set.

Diagnostic plots examining different characteristics of the ChIP-Seq data based on raw counts (*left*) and


quantile normalized counts (*right*) resulting from the three pairwise cell type comparisons. The MA plots (*top*) contrast $\log_2$ fold changes (*y*-axis) against mean sequencing coverage (*x*-axis) for all genome wide 500 bp windows. Genomic windows with a Bonferroni adjusted *P*-value <0.05 are indicated in red. The density plots (*middle*) show the relative distribution of read counts (*x*-axis) at genome wide 500 bp windows for windows with read counts ≤ 10 . The boxplots (*bottom*) show read counts at genome wide 500 bp windows for individual assays.



Supplementary Figure 8

Bioinformatic analysis of H3K4me2 ChIP-seq data set.


(a) As examples, cell-type specific enhancer DER tracks (pre- and post-normalization (norm)) along with UCSC tracks are shown for the following gene loci: *TBX21*, *IFNG*, *CXCR3*, *CXCR6*, *STAT1* and *STAT4*, where additional DERs were detected following quantile normalization of the ChIP-Seq data (**Supplementary Fig. 7**). (b) The number and size distribution of DERs after merging consecutive DERs (see **Methods** and **Supplementary Notes**).



Supplementary Figure 9

Genomic distribution of differentially enriched *cis*-regulatory regions (DERs).

Pie chart (*left*) shows the distribution of DERs in different genomic regions, and compared to the reference annotation (*right*)




Supplementary Figure 10

Concordant changes in gene expression and H3K4me2 enrichment patterns at promoter and distal *cis*-regulatory elements (enhancers).

(a) Heat map shows the comparison of gene expression (RNA-Seq) and H3K4me2 enrichment patterns (across an extended genomic region) for transcripts having a DER in their promoter region when comparing naïve to T_H2 memory $CD4^+$ T cells (each row represents one transcript). Upstream = -20 kb from transcription start site (TSS); Promoter = +/- 1 kb around TSS; Transcript body = region between TSS and transcript end site (TES); Downstream = +20 kb from TES; RNA = RNA-Seq data. The heat map indicates concordant gain (red) or loss (blue) of H3K4me2 enrichment in T_H2 memory compared to naïve $CD4^+$ T cells, at the promoter and at enhancers located in or close to that transcript (see **Methods** and **Supplementary Notes**). Similarly, the last column indicates concordant up- (red) or down- (blue) regulation of gene expression in T_H2 memory cells compared to naïve $CD4^+$ T cells for the corresponding transcripts. **(b)** As examples, H3K4me2 enrichment and RNA-Seq tracks for naïve and T_H2 memory T cells (data merged from all donors including duplicate assays) are shown for the following gene loci: *CCL20* and *CCR8* (T_H2 gain); *S100B* and *LY86* (T_H1 gain), along with

UCSC gene tracks (*top row*) and cell-type specific enhancer DERs.



Supplementary Figure 11

Concordant changes in gene expression and H3K4me2 enrichment patterns at promoter and distal *cis*-regulatory elements.

(a) Heat map shows the comparison of gene expression (RNA-Seq) and H3K4me2 enrichment patterns (across an extended genomic region) for genes that are differentially expressed in any of the three pairwise comparisons of naïve, T_H2 and T_H1 cells (each row represents one gene). First column of the heat map

(labeled RNA-Seq) displays $\log_2$ fold change in gene expression values. Next to the RNA-Seq data, fold change in H3K4me2 enrichment pattern across each extended gene locus (transcription start site (TSS) $\pm$ 50 kb) is displayed (see **Methods** and **Supplementary Notes**). The heat map indicates concordant changes in (up-*red* or down-*blue*) gene expression and H3K4me2 enrichment at the promoter and distal enhancer regions. *Bottom panel* shows genes with no significant changes in expression levels, but display a promoter-localized differentially enriched H3K4me2 region (DER) in any of the three pairwise comparisons of naïve, T_H2 and T_H1 cells. **(b)** As examples, RNA-Seq and H3K4me2 enrichment tracks for naïve and memory T cells (data merged from all donors including duplicate assays) are shown for the following gene loci: *CCR6*, *CCR8* and *FAM129a* (T_H2 active genes); *CCL5* and *CX3CR1* (T_H1 active genes); *IL4* and *IL21* (T_H2 poised genes), along with UCSC gene tracks (*top row*), cell-type specific DERs and differentially expressed genes (DEX). Red dashed line boxes highlight cell-specific DERs. **(c)** Shows MA plots (vertically displayed) for the pairwise comparisons of naïve, T_H2 and T_H1 cells, and red dots indicate differentially expressed genes (false discovery rate of 1%). **(d-e)** Shows principal component analysis (PCA) for RNA-Seq data from each sample, and the genes contributing to the PCA.

Supplementary Notes

Additional Methods

Micro-scaled multi-sample ChIP-Sequencing. Additional considerations that improved the reproducibility and consistency of our micro-scaled ChIP-Seq assay are: (i) Handling all tissue and cell isolation procedures at 4 °C to minimize changes from epigenetic patterns that are present *in vivo*. (ii) Formaldehyde-fixed cell pellets were snap frozen in liquid nitrogen and stored at –80 °C prior to the ChIP assay. This storage step gave us the flexibility to perform ChIP assays in large batches of samples ($n \sim 24$) from subjects in disease and control groups, thus minimizing assay-to-assay variability. (iii) Validating sonication conditions for reproducible shearing of chromatin from small numbers of cells. We found that using the BioRuptor™ sonicator, which has a 12-tube holder, aided equivalent shearing of chromatin among the multiple samples that were to be compared. The extent of chromatin shearing was verified for all samples by using highly sensitive SYBR gold™ dye to stain DNA run on agarose gels. (iv) We validated a picogreen™-based assay (Life Technologies) for measuring picogram quantities of DNA, to ensure the use of equal starting amounts of template chromatin from all the samples when undertaking multi-sample ChIP assays or whole genome amplification steps post-ChIP. (v) Switching to tubes with low DNA and protein binding, and changing tubes during post IP washes significantly reduced assay background while preserving specific enrichment in ChIP assays. (vi) qPCR-based measurement of enrichment at control genes was used to guarantee comparable ChIP performance across multiple samples. (vii) Using equivalent starting amounts of post-ChIP DNA (1 ng) for whole genome amplification (WGA, Sigma), ensured that the number of PCR cycles for amplification remained similar across the samples to be compared.

Additional considerations that improve the validity of our micro-scaled ChIP-Seq approach (i) The whole-genome amplification step that is essential for preparing libraries from micro-scaled ChIP assays does not introduce random noise. Instead, the observed differences in amplification yield for different regions of the genome are highly reproducible as can be seen in the high correlation between different amplification runs (**Supplementary Fig. 4b**). Therefore, differences between samples (using 10^5 primary

cells) that have undergone the same amplification procedure, which is what we are looking at with our DER analysis, will reflect differences in the un-amplified samples.

(ii) The majority of regions with a high number of reads in the unamplified sample also have a high number of reads in the amplified sample. To quantify this, we have performed an receiver operating characteristic (ROC) analysis in which the top 1% of windows in the standard sample was identified using the read counts in windows from the amplified sample (excluding windows with no reads in either sample to avoid overestimation of performance due to genomic regions in which no reads can be mapped). This gave a very high area under the curve (AUC) of 0.976, with 70% of the high scoring windows in the standard sample recovered in the top 2% of windows in the amplified sample (**Supplementary Fig. 4c**). At the same time, only 5% of the high-scoring windows of the standard sample were not included in the top 10% of the amplified sample, indicating that only a small fraction of genomic regions were amplified poorly.

Identification of differentially enriched *cis*-regulatory regions (DERs) comparing cell types. We have calculated coefficients of variability (CV; standard deviation and average) between replicates from the same donor and between averages between different donors. For all cell types, the CV was higher in the inter-donor comparison but not greatly so, e.g. for naïve T cells, CV = 32.9% in the intra-donor comparison vs. 35.3% in the inter-donor comparison. So there is more variability between donors, yet there is also considerable variability between replicates. To account for both, the biological and ChIP assay-to-assay variability in determining statistically significant DERs, replicate ChIP assays from the same individuals were not pooled into an aggregate profile. For each sample, sequencing coverage was calculated at genome wide 500 bp windows, and pairwise differential coverage between groups of cell-types was calculated using *MEDIPS* v.1.12.0 (*extend* = 120, *uniq* = *F*, *window_size*=500, *BSgenome* = "*BSgenome.Hsapiens.UCSC.hg19*", *minRowSum*=1, *p.adj* = "*bonferroni*", *diff.method*="edgeR")^{1,2}.

Inspection of the resulting MA plots indicated a varying efficiency of immunoprecipitation in the group of T_H1 cell samples compared to the groups of naïve CD4⁺ T cells and T_H2 cell samples (**Supplementary Fig. 7**). To correct for this effect, we have performed quantile normalization on the genome-wide count tables across the samples prior to calculation of pairwise differential coverage between groups. MA plots (**Fig. 2a, left**) as

well as density plots of group-wise mean raw and quantile normalized sequencing coverage (**Supplementary Fig. 7**), indicate improved comparability of genome-wide enrichment distributions after quantile normalization. In addition, several well-known regulatory elements that contrast T_H1 cells from T_H2 cells became significant only after this normalization procedure (**Supplementary Fig. 8a**). Consequently, we performed quantile normalization as a standard prior to identification of DERs, resulting in 51,261 DERs for naïve CD4⁺ T cells vs. T_H2 cells (23,583 naïve > T_H2 and 27,678 naïve < T_H2), 38,665 DERs for naïve CD4⁺ T cells vs. T_H1 cells (22,494 naïve > T_H1, and 16,171 naïve < T_H1), and 9,737 DERs for T_H2 cells vs. T_H1 cells (5,847 T_H2 > T_H1, and 3,890 T_H2 < T_H1) (Bonferroni adjusted *P*-value < 0.05).

To test the influence of potential interdependence between replicate ChIP assays from the same individual, we repeated the comparison between naïve and T_H2 cells by pooling ChIP replicates from the same donor into an aggregate profile. As expected, due to the reduced statistical power caused by a smaller sample size (due to pooling of replicate assays), we detected a slightly lower number of DERs (*n* = 46,912) compared to our original approach (*n* = 51,261). Most importantly, at the same significance level (adjusted *P*-value < 0.05), 44,425 (87%) of DERs identified by our original approach were also detected when pooling ChIP replicates. We have further characterized the location of DERs that were not re-discovered by pooling replicates (*n* = 6,836) and found that 3,922 (57%) are individual 500 bp windows that are part of a group of adjacent significant DERs (*i.e.* consecutive DERs) identified by our approach (data not shown). Consequently, we observed an overall agreement of 94% between the distinct genomic loci identified by our approach compared to the pooled approach. Thus, we conclude that the presented results obtained by separate processing of ChIP replicates are not confounded by a potential interdependence between replicate ChIP assays from the same individuals.

A recently published method (“DiffReps”) for analyzing quantitative differences of broad histone marks across groups of ChIP-Seq samples was compared to a number of other available tools and shown to perform comparably on a small set of ChIP-Seq samples³. One method included in this comparison is very similar to the *MEDIPS* methodology employed in our study and the authors reported high agreement between their “DiffReps” method and this *MEDIPS*-like approach. As a further comparison, we applied the “DiffReps” tool to analyze our larger data set and then compared the results with those obtained by our

method (*MEDIPS*) using the same statistical cut off (Bonferroni adjusted P -value < 0.05). We observed that around 56% of the DERs detected by our method (*MEDIPS*) were also detected by the “DiffReps” method, suggesting a significant overlap between the two methods (P -value $< 1.67\text{e-}179$, *hypergeometric* test). Further, a number of likely false positives and false negatives were also observed with the “DiffRep” method compared to the *MEDIPS* method (data not shown).

Classification of DERs into sub-groups. As described above there was a total of 99,663 DERs identified in the 3 cell type comparisons. However many DERs that were identified in different cell-type comparisons had the same genomic location. The total number of DERs at distinct locations was 71,640. Merging consecutive windows (DERs), we identified a total of 35,428 distinct genome regions ranging in size from 500 bp to 29 Kb (**Supplementary Fig. 8b**). Next, we wanted to assign an overall state to each distinct DER based on the combined results in the 3 cell type comparisons. In each comparison, there are 3 possible outcomes (up, down, or non-significant). Our 71,640 DERs showed sixteen different patterns of outcomes, which are shown in **Supplementary Table 3**. We grouped DERs with patterns of outcomes that have similar biological meaning into six merged groups. Specifically, we defined a group of DERs with T_H2 and T_H1 cells gain comparing to naïve cells but no significant difference between T_H2 and T_H1 cells as *shared gain* (9,317 DERs); a group of DERs with T_H2 and T_H1 cells loss comparing to naïve T cells but no significant difference between T_H2 and T_H1 cells as *shared loss* (12,870 DERs); groups with T_H2 cells gain (20,679) as *T_H2 gain*; groups with T_H2 cells loss (10670) as *T_H2 loss*; groups with T_H1 cells gain (8534) as *T_H1 gain*; and groups with T_H1 cells loss (9570) as *T_H1 loss*.

Assignment of target genes to DERs. There is a considerable overlap between this list of genes and those previously identified by Hawkins *et al.*⁴ which describes H3K4me1-enriched enhancers in T_H1 and T_H2 cells generated *in vitro*: if we use the Hawkins *et al.* criteria to assign genes to enhancers, our study identifies 80% of the genes also identified by Hawkins *et al.* as linked to T_H1 and T_H2 -specific enhancers (significance of the overlap: P -value $< 2.2\text{e-}16$, Fisher’s exact test). However, our large sample size (>100) and our use of *in vivo*-generated T_H1 and T_H2 cells, we have been able to identify a number of additional

novel candidate genes, enhancers and miRNAs that were not described by Hawkins *et al.* (see **Supplementary Table 6**, and data not shown). We attribute these differences to several factors: Hawkins *et al.* used naïve T cells polarized with T_H1 or T_H2 cytokines *in vitro* for a relatively short time (72 h), whereas T_H2 memory development *in vivo* is likely to be a more sustained and lengthy process. Moreover, the Hawkins *et al.* study used only a single sample of T_H1 and T_H2 cells for their analyses, potentially reducing the statistical power to detect differences in enhancer profiles at the genome-wide level.

Correlation of gene expression with H3K4me2 enrichment patterns at promoters and enhancers.

In **Supplementary Figure 10 (right)**, we show gene expression changes between naïve and T_H2 cells for the transcripts included in the heat map of H3K4me2 profiles (*left*). For all genes that are significantly differentially expressed (adjusted *P*-value ≤ 0.01), we have calculated the log₂ fold change of T_H2 / naïve CD4⁺ T cells based on the library size normalized count values per group as reported by *DESeq*. In order to avoid division by zero, we have added a pseudo count of five to each group. Transcripts are given in the same order as in the heat map of H3K4me2 profiles. In cases where a transcript (linked to promoter-localized DERs) is associated with a gene that is not differentially expressed, the expression change for that gene is represented as grey.

To further examine the dependency of H3K4me2 enrichment profiles at potential distal *cis*-regulatory regions (beyond promoter-localized DERs) on gene expression, we have examined H3K4me2 profiles in the extended gene loci (TSS +/- 50 kb) for all differentially expressed genes (**Supplementary Fig. 11**). Here, we have divided the +/- 50 kb region around the transcription start sites of differentially expressed genes into consecutive 2 kb windows. For each 2 kb region, the previously calculated H3K4me2 log₂ fold change of the 500 bp window with the most significant *P*-value was selected, where only windows with a non-adjusted *P*-value ≤ 0.05 were considered. In case a 2 kb window does not contain a significant 500 bp window (non-adjusted *P*-value ≤ 0.05), the window is represented as grey.

Detecting enrichment of transcription factor binding motifs at cell-specific enhancers. We examined if there was a significant enrichment in known TF binding motifs in the subsets of DERs defined above. In each DER category, we retrieved the genomic sequence corresponding to the DERs, and used the `findMotifsGenome.pl` of the HOMER v 4.3 software package⁵ with default parameter settings (**Fig. 5a** and **Supplementary Table 9**). The motifs with P -values less or equal to $1.00\text{E-}0.3$ and ratio of target sequences with motif vs background sequences with motif >1.1 were defined as significantly enriched (**Supplementary Table 9**).

Detecting enrichment of transcription factor binding sites at cell-specific enhancers. To investigate, if DERs occur at sites that are known to be bound by transcription factors, we have downloaded ChIP-Seq derived binding sites of 161 transcription factors from 91 different human cell types generated by the ENCODE project and provided via the Table Browser at the UCSC Genome Bioinformatics Site (track *Txn Fac ChIP V3*, table *wgEncodeRegTfbsClusteredV3*). Moreover, we downloaded ChIP-Seq derived binding sites of 18 additional transcription factors provided via GEO-NCBI [GSM776559, GSM776558, GSM776557, GSE37589, GSM1208648, GSM1208734, GSM1208702, GSM1208676, GSM1208658, GSM1208799, GSM1208792, GSM1208791, GSM1208776, GSM1208775, GSM1208774, GSM1031240, GSM831021, GSE38567]. Binding sites provided in *hg18* were converted to *hg19* using the *liftOver* tool at the UCSC Genome Bioinformatics Site^{6,7}.

For each of the 179 transcription factors, we tested if there is an enrichment of their binding sites within any of the predefined groups of DERs compared to random background. To assemble background datasets for each DER group, we sampled 100 sets of random genome wide 500 bp windows, each with the same amount of regions as the number of DERs. To avoid artificially enhanced enrichments due to inclusion of genomic regions in the background that are inaccessible or for which reads cannot be mapped, we restrict the genomic space for background sampling to windows that show sequencing coverage in any of our H3K4me2 samples, and further exclude genomic regions known to harbor signal artifacts in sequencing experiments as provided by the Table Browser of the UCSC Genome Bioinformatics Site (*hg19*, track *Mapability*, table *DAC Blacklist wgEncodeDacMapabilityConsensusExcludable*). Random regions were

generated using the function *shuffleBed* of the *bedtool* suite v.2.16.2 providing the genomic regions to be excluded via the parameter *excl*. Overlaps between the midpoint of DERs or random regions, respectively, and transcription factor binding sites were calculated using the function *IntersectBed* of the *bedtools* suite v.2.16.2 (ref. 8) For each transcription factor, and for each group of DERs, we calculate the mean and standard deviation of overlaps observed for the 100 background sets of genomic regions. We use the mean and standard deviation obtained from random sampling to calculate an odds ratio and to model a normal distribution and calculate *P*-values for the observed overlap between DERs and binding sites compared to random sampling (**Supplementary Table 10**). We report all significance values with an odds ratio >1 and with a *P*-value ≤ 0.01.

SNPs datasets and enrichment calculation. For a given set of disease associated SNPs, we generated a specific control set of SNPs with similar properties. For each SNP in the disease set, we assigned a 'functional code' similar to the approach in Yu *et al.*⁹ The code was composed of the functional assessment in the snp138Common table (set to one of the following: 0='nonsense', 100='missense', 200='stop-loss', 300='frameshift', 400='cds-indel', 500='coding-synon', 600='intron', 700='ncRNA', 800='splice-3', 900='splice-5', 1000='untranslated-3', 1100='untranslated-5', 1200='near-gene-3', 1300='near-gene-5', 1400='unknown') plus the minor allele frequency divided into intervals of 5 percentage points (e.g. '0' = 0-5%, '5' = 5-10%, '10' = 10-15% etc.). We then randomly chose 40 control SNPs with the same functional code from the SNPs tested for by the Illumina 1 MDuo genotyping array as specified from the SNP Genotyping Array track for *hg18* of the UCSC genome browser^{6,7}. The Illumina platform was chosen as it was most frequently used in the asthma GWAS studies. If less than 40 exactly matching control SNPs were available, we also included control SNPs with up to 10% difference in their allele frequency. For some of the disease datasets other than asthma, there were individual disease-associated SNPs in a functional code category for which there still was not a sufficient number of control SNPs available, and such singletons were excluded from the control set generation.

To calculate SNP enrichment in a given DER category, we determined the fraction of disease-associated SNPs located in the corresponding genomic region, and compared it to the fraction of non-disease

associated SNPs from a background set to calculate enrichments (**Fig. 6a**). To calculate overlapping and intersecting genome coordinates, the BEDTools utilities package⁸ was utilized. Significance was assessed using the chi-square test. In addition to the DER analysis in T cell subsets, we also calculated enrichment of disease-associated SNP sets in H3K4me1 peaks of eight different tissues: adipose nuclei, liver, adipose-derived mesenchymal stem cells (ADMSC), kidney, breast myo-epithelial cells, brain, CD4⁺ T cells and skeletal muscle cells (**Fig. 6a** and **Supplementary Table 11c**). These H3K4me1-peaks were calculated in our previous analysis¹⁰ based on ChIP-Seq data together with input samples obtained from the Human Epigenome Atlas^{11,12}.

Haploblock-based SNP enrichment analysis. We performed a second analysis to examine the overlap of asthma associated SNPs and enhancers of different cell types based on previously published methods¹³ with a few relevant modifications. As a source of SNPs, we used a recent analysis of asthma-associated SNPs across multiple study populations in the European population¹⁴. We collected SNPs that were shown in **Supplementary Table 1** of Moffatt study¹⁴ as showing genome wide significant association with either childhood onset asthma or later onset asthma (66 SNPs in total). We used HapMap release 27 data (compiled from merged genotype data from phases I+II+III (HapMap rel #27, NCBI B36) downloaded from (ftp://ftp.ncbi.nlm.nih.gov/hapmap/ld_data/2009-04_rel27/) as a source for Linkage Disequilibrium (LD) information between SNPs, as this was the only source for which genome wide linkage information was available in a pre-calculated form (which was needed for the automated calculation of a large set of randomized controls as described below). CEU (Utah residents with ancestry from northern and western Europe) was chosen as the reference population. For 3 lead SNPs, no linkage information was included in Hapmap, and they were excluded from the dataset.

To construct haploblocks, we grouped SNPs together that were in linkage with $r^2 > 0.1$, resulting in 11 haploblocks. For each haploblock, we added all SNPs that were in linkage with $r^2 > 0.8$ for any of the lead SNPs. For each asthma-associated haploblock, we calculated a set of 20 randomized controls choosing a set of control-lead SNPs that matched the frequency and properties of the lead-SNPs from meta-analysis. We chose control lead SNPs from the Illumina 1 MDuo genotyping array as specified from the SNP Genotyping Array track for *hg18* of the UCSC genome browser^{6,7}. The Illumina platform was chosen as it

was most frequently used in the asthma GWAS studies. As described in the SNP enrichment above, we assigned a 'functional code, composed of the functional assessment in the snp138Common table and the allele frequency. Control haploblocks were randomly drawn by selecting first a matching control SNP for the asthma-SNP lead-SNP with the highest P -value, and then examining if the control had matching SNPs in linkage with $r^2 > 0.1$. For asthma-haploblocks with up to 4 SNPs, controls matching all these criteria for all SNPs were found. For larger haploblocks, at least 4 SNPs and more than half of the contained lead-SNPs had to perfectly match.

For a given set of haploblocks and a set of enhancers, we determined the number of haploblocks that overlapped with at least one enhancer. We determined the average number of overlapping haploblocks in the 20 control datasets, and used the binomial distribution to calculate P -values for the number of overlapping haploblocks in the meta-analysis to have occurred by chance (**Supplementary Table 11d**).

REFERENCES

1. Chavez, L. et al. Computational analysis of genome-wide DNA methylation during the differentiation of human embryonic stem cells along the endodermal lineage. *Genome Res* **20**, 1441-1450 (2010).
2. Lienhard, M., Grimm, C., Morkel, M., Herwig, R. & Chavez, L. *MEDIPS*: genome wide differential coverage analysis of sequencing data derived from DNA enrichment experiments. *Bioinformatics* **30**, 284-286 (2014).
3. Shen L, Shao NY, Liu X, Maze I, Feng J, Nestler EJ. diffReps: detecting differential chromatin modification sites from ChIP-seq data with biological replicates. *PLoS One*, **8** e65598 (2013).
4. Hawkins, R.D. et al. Global chromatin state analysis reveals lineage-specific enhancers during the initiation of human T helper 1 and T helper 2 cell polarization. *Immunity* **38**, 1271-1284 (2013).
5. Heinz, S. et al. Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**, 576-589 (2010).
6. Karolchik, D. et al. The UCSC Table Browser data retrieval tool. *Nucleic Acids Res* **32**, D493-496 (2004).

7. Karolchik, D., Hinrichs, A.S. & Kent, W.J. The UCSC Genome Browser. *Curr Protoc Bioinformatics* **Chapter 1**, Unit1 4 (2009).
8. Quinlan, A.R. & Hall, I.M. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841-842 (2010).
9. Yu, W. et al. GWAS Integrator: a bioinformatics tool to explore human genetic associations reported in published genome-wide association studies. *Eur J Hum Genet* **19**, 1095-1099 (2011).
10. Gerasimova, A. et al. Predicting cell types and genetic variations contributing to disease by combining GWAS and epigenetic data. *PLoS One* **8**, e54359 (2013).
11. Bernstein, B.E. et al. The NIH Roadmap Epigenomics Mapping Consortium. *Nat Biotechnol* **28**, 1045-1048 (2010).
12. Milosavljevic, A. Emerging patterns of epigenomic variation. *Trends Genet.* **27**, 242-250 (2011).
13. Schaub, M.A., Boyle, A.P., Kundaje, A., Batzoglou, S., Snyder, M. Linking disease associations with regulatory information in the human genome. *Genome Res* **22**, 1748-1759 (2012).
14. Moffatt, M.F. et al. A large-scale, consortium-based genomewide association study of asthma. *N Engl J Med* **363**, 1211-1221 (2010).