

Letter to the Editor

Pilot study of budesonide inhalant suspension irrigations for chronic eosinophilic sinusitis*To the Editor:*

Chronic sinusitis (CS) is often associated with nasal polyposis (NP). CS represents several disease processes including chronic hyperplastic eosinophilic sinusitis (CHES).¹ Many immune and pathological features of CHES are shared with asthma, suggesting that these are similar disease processes involving the upper and lower airways. Historically, patients with CS were approached as having a chronic infectious disorder, and treatment consisted of antibiotics and surgery to promote drainage. Often these approaches proved unsatisfactory for patients with CHES.² Given the marked eosinophilia and pathological similarity of CHES to asthma,³ corticosteroids could be used as a treatment for this disorder. Systemic corticosteroids shrink hyperplastic tissue and associated NP, reduce symptoms, and restore the senses of smell and taste.⁴ However, their side-effect profile has precluded long-term use.

It has been suggested that topical intranasal corticosteroids would prove effective for eosinophilic forms of CS, given their utility for asthma and allergic rhinitis, while maintaining a minimal risk profile. Although intranasal corticosteroids reduce NP,⁵ they have never been shown to improve the sinusitis component of this disorder, likely reflecting their inability to access the sinus cavity. Saline irrigations have proven to be an effective method of addressing sinus disease,⁶ likely through their ability to lavage the nasal and sinus cavities. It has been speculated that the addition of a corticosteroid such as budesonide inhalation suspension (Pulmicort respules; Astra Zeneca, Wilmington, De), a medication already approved for asthma, to the lavage fluid would more effectively treat sinus inflammation. This study was performed to obtain pilot data documenting objective and subjective evidence for improvement in patients who used this approach for a period of ≥ 3 months. For information regarding the study design, see Methods in the Online Repository at www.jacionline.org.

Patients with CHES or aspirin-exacerbated respiratory disease having failed to respond to previous medical therapy and having been treated with budesonide nasal irrigation for ≥ 3 months were recruited. A total of 8 subjects were enrolled (see this article's Table E1 in the Online Repository at www.jacionline.org). All subjects had allergy as defined by a positive skin prick test (≥ 5 -mm wheal) to at least 1 aeroallergen, 4 subjects were classified as having aspirin-exacerbated respiratory disease, and all but 1 had physician-diagnosed asthma. None of the subjects took oral steroids during treatment with budesonide, and 5 were on montelukast, with 2 also taking zileuton.

Our primary outcome measure for sinus improvement was the change in CT score. The median CT score before treatment was 15 (maximum, 30), which improved to 5 ($P < .05$) after treatment (Fig 1). By using our visual analog scale (see this article's Fig E1 in the Online Repository at www.jacionline.org), we calculated scores for each of 16 sinus symptoms on a scale of 0 to 6, with 6 being severe and 0 none (maximum, 96). After budesonide treatment, subjects' sinus scores decreased (mean $\pm$ SD) from 43.1 ± 5.4 to 20.1 ± 3.0 ($P < .02$; Fig 2). Sense of smell was separately examined (0, none, to 6, complete) because of its particular




FIG 1. Change in CT score after budesonide nasal irrigation. Each dot represents an individual with a line connecting the prebudesonide and postbudesonide CT scores. The bold lines to the side of the data points indicate the median, with the interquartile range and inner fence indicated with a box and whisker plot, respectively ($P < .05$ for prebudesonide compared with postbudesonide sinus score).




FIG 2. Self-assessed change in sinus symptom score after budesonide nasal irrigation. Each dot represents an individual, with a line connecting the prebudesonide and postbudesonide sinus scores. Lines to the side of the data points indicate the mean $\pm$ SEM ($P < .01$ for prebudesonide compared with postbudesonide sinus score).

impact on quality of life. Subjects reported a significant improvement in their sense of smell (mean $\pm$ SD) (see this article's Fig E2 in the Online Repository at www.jacionline.org), from 1.1 ± 0.7 to 3.6 ± 0.8 ($P < .05$). Other sinus measures displayed similar improvement. Of the 6 subjects who had prerhinocopy and postrhinocopy, 5 showed improvement after treatment, and 3 of 4 subjects had complete resolution of NP (see this article's Fig E3 and Table E2 in the Online Repository at www.jacionline.org).

It has been suggested that improvements in upper respiratory disease can lead to improvements in lower respiratory symptoms. The asthma visual analog scale (see this article's Fig E4 in the Online Repository at www.jacionline.org) was converted to a scale

ranging from 0 to 6 and scored (maximum, 36). Of the 7 patients in the study with asthma, there was a trend toward a decrease in the asthma score (mean $\pm$ SD) from 15.3 ± 2.9 pretreatment to 11.4 ± 2.2 posttreatment (Table E2). Using the National Institutes of Health definition of asthma control,⁷ 3 of 4 subjects who had poorly controlled disease at the beginning of the study improved to well controlled (Table E2). The remaining 3 patients with asthma had well controlled disease throughout the study.

The current study suggests that the addition of budesonide suspension to nasal saline irrigations produces a significant improvement in sinus symptoms including sense of smell. It is possible, given the construct of the study's methodology, that the findings may overestimate the true benefit of this therapy, especially with regard to recall bias. However, previous works have indicated that patients are able to recall symptoms of smell accurately.⁸ The lack of a control group makes it difficult to determine what role a placebo effect and the natural course of the disease may have played. That being noted, the cohort was one of patients who remained symptomatic on previous therapy and for whom the addition of budesonide irrigations to the sinus regimen represented the only change.

Subjective improvements were corroborated by objective findings on CT scans and nasal endoscopy. CT scanning is validated for accurately quantifying the extent of mucosal inflammation/hypertrophy within the sinuses. Previous studies have shown that, at least in the short term, radiographic findings of CS are stable. The significant improvement in CT scores is therefore compelling. The complete regression of polyps observed in 3 of 4 patients with NPs before initiation of therapy is consistent with reports of the ability of intranasal corticosteroids to reduce polyposis⁵ and is indicative of the role that corticosteroid irrigations may play in the management of this disease.

Modest trends toward improvement in asthma symptom scores were also observed. A lack of significance in the pilot data was likely driven by the inclusion of patients with well controlled asthma; however, 3 of 4 subjects who did not have controlled disease at baseline did show improvement. Given the low probability that topical corticosteroids in the upper airway would access the bronchial airway, improvement in asthma more likely reflects 1 or more of the indirect pathways by which reduced sinusitis has been proposed to affect asthma positively.⁹ Although it remains possible that improvement in asthma symptoms could reflect systemic absorption of the steroid, especially considering the high dose of budesonide used in this study (500 μ g twice a day), the nature of sinus irrigations is that most of the sinus irrigant does not remain. As such, likely <50 μ g of budesonide remains in the sinuses, a dose equivalent to rhinitis therapy. Absorption from the sinuses is not likely to be dissimilar to that from the airways. These studies provide support for the unified airway concept.

This study supports the concept that addition of budesonide inhalation suspension to standard nasal saline irrigation produces subjective and objective benefit in eosinophilic sinus disease. It is hoped that these data will support enthusiasm for performing a double-blind, placebo-controlled study of budesonide inhalation suspension in nasal saline washes in CS.

John W. Steinke, PhD^a

Spencer C. Payne, MD^b

M. Elizabeth Tessier, MD^a

Lori O. Borish, RN^a

Joseph K. Han, MD^c

Larry C. Borish, MD^a

From the Departments of ^aMedicine and ^bOtolaryngology—Head and Neck Surgery, University of Virginia Health Systems, Charlottesville; and ^cthe Department of Otolaryngology—Head and Neck Surgery, Eastern Virginia Medical School, Norfolk. E-mail: js3ch@virginia.edu

Supported by National Institutes of Health grant nos. RO1 AI01793 and PO1 AI50989. Disclosure of potential conflict of interest: J. W. Steinke has received research support from the National Institutes of Health. S. C. Payne has received honoraria from Acclarant, Inc, has received research support from the American Academy of Otolaryngologic Allergy, and has provided legal consultation or expert witness testimony on the topic of tonsillectomies. L. C. Borish is on the speakers' bureau for Merck, is on an advisory board for Genentech and Novartis, and has received research support from Genentech and Merck. The rest of the authors have declared that they have no conflict of interest.

REFERENCES

1. Meltzer EO, Hamilos DL, Hadley JA, Lanza DC, Marple BF, Nicklas RA, et al. Rhinosinusitis: establishing definitions for clinical research and patient care. *J Allergy Clin Immunol* 2004;114:S155-212.
2. Poole MD. Pediatric sinusitis is not a surgical disease. *Ear Nose Throat J* 1992;71:622-3.
3. Steinke JW, Bradley D, Arango P, Crouse CD, Frierson H, Kountakis SE, et al. Cysteinyl leukotriene expression in chronic hyperplastic sinusitis-nasal polyposis: importance to eosinophilia and asthma. *J Allergy Clin Immunol* 2003;111:342-9.
4. Jankowski R, Bodino C. Evolution of symptoms associated to nasal polyposis following oral steroid treatment and nasalization of the ethmoid-radical ethmoidectomy is functional surgery for NPS. *Rhinology* 2003;41:211-9.
5. Stjarne P, Blomgren K, Cayé-Thomasen P, Salo S, Soderstrom T. The efficacy and safety of once-daily mometasone furoate nasal spray in nasal polyposis: a randomized, double-blind, placebo-controlled study. *Acta Otolaryngol* 2006;126:606-12.
6. Harvey R, Hannan SA, Badia L, Scadding G. Nasal saline irrigations for the symptoms of chronic rhinosinusitis. *Cochrane Database Syst Rev* 2007;18:CD006394.
7. National Heart L, and Blood Institute. National Asthma Education and Prevention Program Expert Panel report 3: guidelines for the diagnosis and management of asthma. Bethesda (MD): National Institutes of Health; 2007.
8. Bhattacharyya N. Clinical and symptom criteria for the accurate diagnosis of chronic rhinosinusitis. *Laryngoscope* 2006;116:1-22.
9. Braunstahl GJ, Kleinjan A, Overbeek SE, Prins JB, Hoogsteden HC, Fokkens WJ. Segmental bronchial provocation induces nasal inflammation in allergic rhinitis patients. *Am J Respir Crit Care Med* 2000;161:2051-7.

doi:10.1016/j.jaci.2009.09.018

METHODS

Study design and participants

Patients age 18 to 60 years with CHES who had failed to respond to medical and/or surgical therapy were recruited from the outpatient allergy/immunology and otolaryngology clinics at the University of Virginia. Medical therapy included antibiotics, large-volume saline irrigations, and topical intranasal steroid administration. Patients were subsequently treated with budesonide inhalation suspension 500 µg twice daily administered by diluting the budesonide suspension in a large volume of saline (>100 mL), which was then administered intranasally as part of standard sinus irrigation care. Patients who had been treated with budesonide nasal washes for at least 3 months were invited to participate. Participants were asked to provide a medical history with detailed information regarding symptoms of sinusitis, asthma, allergic rhinitis, aspirin allergy, and medication use. All subjects had a baseline sinus CT scan and rhinoscopy before initiation of treatment. Exclusion criteria included the diagnosis of cystic fibrosis, sinonasal tumor, or immunodeficiency. Informed consent was obtained from all subjects before enrollment under a protocol approved by the University of Virginia Institutional Review Board.

Visual analogue scale

A standardized visual analog scoring system of sinusitis^{E1} (Fig E1) and asthma (Fig E3) symptoms was used to compare related symptoms before and after treatment with budesonide inhalation suspension. In addition, asthma control was determined using National Institutes of Health guidelines.^{E2} Each visual analogue scale (VAS) question was converted to a numerical value (0-6) to allow statistical analysis of symptoms before and after treatment (maximum sinus score for 16 symptoms, 96; maximum asthma score for 6 symptoms, 36). Subjects were asked whether they had hypersensitivity to aspirin or other non-steroidal anti-inflammatory drugs, defined as a severe asthma exacerbation occurring within 2 to 3 hours of ingestion, that would be suggestive of aspirin-exacerbated respiratory disease.

Sinus CT score

Sinus disease was evaluated by using coronal reconstructions of a helical CT scan obtained in the supine position with axial images taken at 1-mm-thickness intervals. All CT scans were assigned a quantitative score (0-30) by a

blind investigator (S.C.P.) according to our previously published and validated methods.^{E3-E6}

Rhinoscopy

Nasal rhinoscopy was performed using a 30° rigid endoscope before starting budesonide suspension and was compared with a rhinoscopy performed after ≥3 months of treatment. Bilateral examination of each nasal cavity was performed with reporting of crusting, erythema/swelling, purulent drainage, and thick mucus (1 point assigned for each with each naris scored separately; maximum score, 8). In addition, the presence of nasal polyps was noted with a score of 1 assigned for each naris if present (Fig E4).

Statistical analysis

Data are expressed as means ± SEMs or medians with interquartile ranges. Statistical significance was determined by using the Wilcoxon signed-rank test for the endoscopy scores and VAS questionnaire and a 1-tailed Wilcoxon signed-rank test for CT scores. A *P* value ≤.05 was considered statistically significant.

REFERENCES

- E1. Meltzer EO, Hamilos DL, Hadley JA, Lanza DC, Marple BF, Nicklas RA, et al. Rhinosinusitis: developing guidelines for clinical trials. *J Allergy Clin Immunol* 2006;118:S31-80.
- E2. 3 EPR. National Asthma Evaluation and Prevention Program: guidelines for the diagnosis and management of asthma. Bethesda (MD): US Department of Health and Human Services; 2007.
- E3. Newman LJ, Platts-Mills TAE, Phillips CD, Hazen KC, Gross CW. Chronic sinusitis: relationship of computed tomographic findings to allergy, asthma, and eosinophilia. *JAMA* 1994;271:363-7.
- E4. Phillips CD, Platts-Mills TAE. Chronic sinusitis: relationship between CT findings and clinical history of asthma, allergy, eosinophilia, and infection. *AJR Am J Roentgenol* 1995;164:185-7.
- E5. Crater SE, Peters EJ, Phillips CD, Platts-Mills TAE. Prospective analysis of CT of the sinuses in acute asthma. *Am J Radiol* 1999;173:127-31.
- E6. Peters E, Hatley TK, Crater SE, Phillips CD, Platts-Mills TAE, Borish L. Sinus computed tomography scan and markers of inflammation in vocal cord dysfunction and asthma. *Ann Allergy Asthma Immunol* 2003;90:316-22.
- E7. Lund VJ, Kennedy DW. Staging for rhinosinusitis. *Otolaryngol Head Neck Surg* 1997;117:S35-40.

None – None to occasional symptoms

Mild – Steady symptoms but easily tolerated

Moderate – Symptoms hard to tolerate, may interfere with activities of daily living/ sleep

Severe – Symptoms are so bad, person can not function virtually all the time

Example:	X			
	None	Mild	Moderate	Severe
Nasal Obstruction	_____			
Discolored nasal drainage (Front)	_____			
Discolored nasal drainage (Back)	_____			
Facial pressure/ fullness	_____			
Fatigue	_____			
Ear pain/ pressure/ fullness	_____			
Cough	_____			
Halitosis (Bad breath)	_____			
Dental/ Tooth pain	_____			
Fever	_____			
Headache	_____			
Itching of nose/ roof of mouth	_____			
Sneezing	_____			
Overall assessment of severity of symptoms	_____			
Sense of Smell	_____			
	(no decrease)	(moderate decrease)	(complete loss)	
Sense of taste	_____			
	(no decrease)	(moderate decrease)	(complete loss)	

FIG E1. VAS for sinus symptoms. Each subject was asked to rate each of 16 symptoms. The range of symptoms was none to severe.




FIG E2. Patient perception of sense of smell after budesonide nasal irrigation. Using a VAS, subjects were asked to assess their sense of smell before and after nasal irrigation. The VAS was converted to a digital scale rated 0 (no smell) to 6 (no impairment in ability to smell). Each *dot* represents an individual with a *line* connecting the prebudesonide and postbudesonide scores, and the *lines* to the side of the data points indicate the means $\pm$ SEMs ($P < .05$ for prebudesonide compared with postbudesonide sense of smell).

Patient Name:
Date of Visit:
Diagnosis:

ENDOSCOPIC FINDINGS

(IF NONE PRESENT, CIRCLE=0)
(IF PRESENT, CIRCLE=1)

1: CRUSTING	0	1
2: ERYTHEMA	0	1
3: SWELLING	0	1
4: SCAR BAND	0	1
5: PURULENT DRAINAGE	0	1
6: THICK MUCOUS	0	1

(IF NO POLYP PRESENT, CIRCLE=0)
(IF POLYP PRESENT IN MIDDLE MEATUS, CIRCLE=1)
(IF POLYP PRESENT OUTSIDE MIDDLE MEATUS, CIRCLE=2)

7: POLYPS	0	1	2
-----------	---	---	---

FIG E3. Endoscopic scoring system. Modified from Lund VJ, Kennedy DW. Staging for rhinosinusitis. Otolaryngol Head Neck Surg 1997;117:S35-S40.^{E7}

None – None to occasional symptoms

Mild – Steady symptoms but easily tolerated

Moderate – Symptoms hard to tolerate, may interfere with activities of daily living/ sleep

Severe – Symptoms are so bad, person can not function virtually all the time

Example	X			
	None	Mild	Moderate	Severe
Shortness of breath				
Wheezing				
Cough				
Sputum/ phlegm production				
Chest tightness				
Exercise-induced shortness of breath				

FIG E4. VAS for asthma symptoms. Each subject was asked to rate each of 6 symptoms.

TABLE E1. Baseline demographics and subject characteristics

Patients	Sex	Age	Race	SPT+ Yes/No	Asthma Yes/No	AERD Yes/No
1	F	52	W	Yes	No	No
2	F	50	W	Yes	Yes	No
3	F	77	W	Yes	Yes	No
4	F	48	B	Yes	Yes	Yes
5	M	59	W	Yes	Yes	No
6	M	53	W	Yes	Yes	Yes
7	F	59	W	Yes	Yes	Yes
8	M	47	W	Yes	Yes	Yes

AERD, Aspirin-exacerbated respiratory disease; *B*, black; *F*, female; *M*, male; *SPT*, skin prick test; *W*, white.

TABLE E2. Prebudesonide and postbudesonide suspension clinical characteristics

Patient	Asthma control		Asthma score		Rhinoscopy score		Polyp score	
	Pre	Post	Pre	Post	Pre	Post	Pre	Post
1	NA	NA	NA	NA	ND	ND	ND	ND
2	Uncontrolled	Well	17	12	6	4	1	0
3	Very poorly controlled	Well	24	10	0	2	0	0
4	Uncontrolled	Uncontrolled	12	12	5	4	1	0
5	Well	Well	12	12	4	0	0	0
6	Well	Well	2	2	2	ND	0	0
7	Well	Well	16	22	3	1	1	2
8	Very poorly controlled	Well	24	10	3	2	1	0

NA, Not asthmatic; ND, not determined.