

ORIGINAL ARTICLE

Skin and Eye Diseases

Birch pollen-related foods can cause late eczematous reactions in patients with atopic dermatitis

A. Wassmann-Otto  | A. Heratizadeh | K. Wichmann | T. Werfel

Division of Immunodermatology and Allergy Research, Department of Dermatology and Allergy, Hannover Medical School, Hannover, Germany

Correspondence

Anja Wassmann-Otto, Division of Immunodermatology and Allergy Research, Department of Dermatology and Allergy, Hannover Medical School, Hannover, Germany.
Email: anja_wassmann@web.de

Abstract

Background: Previous studies have indicated that in patients with atopic dermatitis (AD) and birch pollen allergy pollen-related foods are able to cause late eczematous response. However, the relevance of AD worsening by ingestion of birch pollen-related foods is still a matter of debate.

Objective: The purpose of this retrospective study was to determine how frequently birch pollen-related foods induce a deterioration of eczema. Additionally, the diagnostic value of specific IgE (sIgE) determination was evaluated.

Methods: A total of 182 children and adults with AD and suspected birch pollen-related food allergy underwent 261 double-blind placebo-controlled food challenges (DBPCFC). Total and sIgE levels were determined prior to DBPCFC.


Results: Sixty-five patients developed allergic reactions (responders) upon DBPCFC with birch pollen-related foods ($n = 103$ DBPCFC). Of these, 32 patients exhibited significant deterioration of AD defined as a median increase of 15.4 severity scoring of atopic dermatitis index points (95% CI 12.4–16.3) from baseline making up 37% of all positive reactions. Responders showed significantly higher sIgE levels to birch pollen and apple as well as a higher prevalence of allergic rhinoconjunctivitis compared to nonresponders ($P < .05$). However, patients with late eczematous response could not be differentiated from those with isolated immediate-type reactions by sIgE levels.

Conclusion: In a subpopulation of patients with AD and birch pollen sensitization, related foods should be considered as a trigger for an aggravation of eczema. As sufficient markers for prediction of late eczematous reactions are still lacking, DBPCFC cannot be replaced in diagnosis of birch pollen-related foods in patients with AD.

Clinical implications: In patients with AD and birch pollen allergy, birch pollen-related foods should be considered as a provocation factor for an aggravation of disease signs and symptoms.

KEYWORDS

atopic dermatitis, birch pollen-related food allergy, diagnostic value, food challenge, late eczematous reactions



GRAPHICAL ABSTRACT

Thirty-seven percent of all positive reactions to birch pollen-related foods were associated with a significant deterioration of atopic dermatitis (AD). Responders showed significantly higher sIgE levels to birch pollen and apple as well as a higher prevalence of allergic rhinoconjunctivitis compared to nonresponders ($P < .05$). In patients with AD and birch pollen allergy, birch pollen-related foods should be considered as a provocation factor for an aggravation of dermatitis even in the absence of immediate-type reaction to birch pollen-related foods.

1 | INTRODUCTION

Foods can provoke a flare-up of eczema in a subgroup of sensitized patients with atopic dermatitis (AD).¹⁻³ In children, staple foods such as cow's milk and hen's egg belong to the most important causes of food allergies.^{2,4} By contrast, in adolescents and adults, food allergies commonly occur as a consequence of primary sensitizations to pollen

allergens.² In central Europe, birch pollen-related food antigens are the most frequent elicitor of a pollen-related food allergy. The major birch pollen allergen is Bet v 1, which belongs to the pathogenesis-related protein family 10.⁵ Over the last decades, an abundance of birch pollen-related foods from different plant families such as *Rosaceae*, *Apiaceae*, and *Fabaceae* has been described, and Bet v 1 homologue allergens in fruits, vegetables, legumes, and nuts have been




FIGURE 1 Clinical outcome of 261 double-blind placebo-controlled food challenges (DBPCFC) performed with birch pollen-related foods (apple, carrot, celery, and hazelnut) in a total of 182 patients with atopic dermatitis (AD). In total, 65 patients showed an allergic reaction upon birch pollen-related food ingestion in 103 DBPCFC. Of these, 42 patients showed an allergic immediate-type reaction, and 32 patients displayed an isolated and/or a combined late-type reaction with a deterioration of AD within 6–48 h after DBPCFC. *Some patients underwent several DBPCFC and showed different clinical reaction patterns. Hence, some responders are mentioned multiple times in Figure 2 [Colour figure can be viewed at wileyonlinelibrary.com]

characterized.⁵⁻¹⁴ Among these, apple, carrot, celery, and hazelnut most often induce allergic symptoms.¹⁵⁻¹⁷ Typical symptoms of birch pollen-related food allergy are immediate-type reactions such as oral allergy syndrome (OAS), urticaria, angioedema, rhinoconjunctivitis, and asthma, but also anaphylactic reactions were reported.^{12,13,16-19} In a previous study, birch pollen-related foods were demonstrated to cause significant worsening of dermatitis in birch pollen-sensitized adult patients with AD even in the absence of immediate-type reactions.²⁰ In lesional skin, Bet v 1-specific T-cell responses could be detected, which illustrates that in late eczematous reactions to birch pollen-related foods, Bet v 1-specific T lymphocytes are important key players.²⁰ Likewise, in a subsequent study including children with AD and birch pollen sensitization, Breuer et al²¹ observed deterioration of dermatitis after ingestion of birch pollen-related foods in 4 of 12 children. Further data in this field indicate that even heat-treated birch pollen-related foods can cause a worsening of AD, whereas thermal processing abrogated their ability to elicit local immediate-type reactions in forms of an OAS.²²

Despite increasing evidence, birch pollen-related foods are still a matter of debate as relevant provocation factor of AD.

In this retrospective study, we focused on the relevance of birch pollen-related food allergy in a large cohort of children and adults with AD. We aimed to better define the frequency of birch pollen-related late-type reactions and possible predictors as well as the diagnostic validity of determination of specific IgE (sIgE).

2 | METHODS

2.1 | Patients

A total of 182 patients met the following inclusion criteria and were enrolled in a retrospective data analysis: (i) Physician-diagnosed mild to severe AD measured by severity scoring of atopic dermatitis (SCORAD) index. (ii) Suspected diagnosis of birch pollen-related food allergy based on a history of seasonal respiratory or cutaneous symptoms during the birch pollen season. (iii) All patients already had consecutively undergone oral food challenges with birch pollen-related foods as a routine diagnostic procedure in the Department of Dermatology and Allergy, Hannover Medical School, Germany, according to the guidelines on food allergy.^{23,24} Prior to oral food challenges, written informed consent had been obtained from all patients. As mentioned above, corresponding data were analyzed retrospectively.

2.2 | Medical history and scoring of AD

A detailed medical history of AD, respiratory, and birch pollen-related food allergy was obtained. AD was diagnosed following the criteria of Hanifin and Rajka which include pruritus, a typical morphology and distribution.²⁵

According to the position paper of the European Academy of Allergy and Clinical Immunology (EAACI) and the Global Allergy and Asthma European Network (GA2LEN) regarding eczematous

reactions to food in patients with AD,²⁶ the severity of AD was rated daily during double-blind placebo-controlled food challenges (DBPCFC) (prechallenge). In addition, AD was also scored 1 day after the DBPCFC had been finalized in order not to miss possible late-type reactions. This was done by a trained dermatologist using the SCORAD index²⁷ which assesses signs and subjective symptoms of AD.

2.3 | Total and specific IgE

Total IgE and sIgE against birch pollen and birch pollen-related foods (apple, carrot, celery, and hazelnut) were determined prior to the DBPCFC using the ImmunoCAP® fluorescence enzyme immunoassay (FEIA) according to the manufacturer's instructions (Thermo Fisher Scientific Inc., Waltham, MA, USA; Manufactured: Phadia AB, Uppsala, Sweden). Patients with a sIgE titer of at least 0.35 kU/L were regarded as sensitized.

2.4 | Double-blind placebo-controlled food challenges

Birch pollen-sensitized patients with suspected immediate-type and/or late-type allergy to at least 1 birch pollen-related food (apple, carrot, celery, or hazelnut) underwent DBPCFC as a standard diagnostic procedure according to European guidelines^{23,24} outside the birch pollen season after a 4-week elimination diet including all kinds of birch pollen-related food upon instruction of a dietitian.

Double-blind placebo-controlled food challenges were performed during stable course of AD. If required, an ongoing topical therapy with mild corticosteroids (European class I or II) in addition to emollients was allowed in selected cases. The treatment regimen had to remain unchanged during the complete period of DBPCFC. Antihistamines were withdrawn 72 hours prior to the DBPCFC.

Apple, carrot, celery, and hazelnut were used for DBPCFC as these represent birch pollen-related foods which are commonly consumed in central Europe. In certain cases, only 1-3 birch pollen-related foods were administered. The reasons for this approach were as follows: (i) exclusion of those foods known to induce clear immediate-type reactions in a patient, (ii) the patient refused to be tested with a certain kind of birch pollen-related food, or (iii) consecutive DBPCFC with single food after positive DBPCFC with a meal consisting of a mixture of several birch pollen-related foods.

Verum and placebo were prepared identically regarding color, consistency, and taste as have been published previously.^{20,21} Basically, DBPCFC with raw birch pollen-related food was performed with fresh ground apple, carrot, celery, and hazelnut. For oral food challenge with heat-treated foods, apple, carrot, and celery were pre-cooked for 5-15 minutes, while hazelnut was heated in a microwave oven (700 W for 2 minutes). The amount of birch pollen-related foods was adapted to a representative daily intake as previously published.²⁰ Placebo meal consisted of carob, rice flakes, orange flavor, and beta-carotene.

According to national and international guidelines,^{23,24,26,28,29} placebo and verum were administered each on 2 consecutive days in a random order. The first challenge day was performed titrated, and on the second day, the full dose was given. The severity of AD was rated daily before starting DBPCFC using the SCORAD index.

Double-blind placebo-controlled food challenges were performed in a 2-step procedure built up of a "spit and a swallow phase" as first described by Ballmer-Weber et al.¹⁶ When starting the challenge with a new meal, the patients first had to keep 15 g of the meal in their mouth and spit it out after 10 and 30 seconds, respectively. Subsequently, the swallow phase was performed in a titrated manner with increasing doses of verum/placebo every 30 minutes (15, 45, 90, and 180 g). The challenge was stopped after the full dose had been given or immediately in case of symptoms of an allergic immediate-type reaction. On the following day, the complete dose was administered all at once. On both days, patients were further observed up to 3 hours after the last dose had been given. Allergic reactions that occurred within 6 hours after ingesting the last dose were defined as immediate-type reactions. An increase in ≥ 10 SCORAD index points within 6–48 hours after administration of the last dose was considered as a significant deterioration of AD (late eczematous reaction) as defined in the position paper of EAACI and GA2LEN.²⁶ Combined reactions were defined as an immediate-type reaction prior to a significant worsening of AD.

2.5 | Statistical analysis

As the data showed a non-normal distribution, statistical analysis was performed by nonparametric tests. The medians of metrically scaled variables (age, total and sIgE levels, SCORAD) were compared with the Mann-Whitney *U* test. The history of respiratory atopic diseases of responders and nonresponders was compared with the chi-square test. The prevalence of birch pollen-related food allergies in the different age groups was compared with the Fisher's exact test.

To evaluate the diagnostic validity of birch pollen-specific IgE determination for the diagnosis of late eczematous reactions, sensitivity, specificity, positive predictive value (PPV), and negative predictive value (NPV) were calculated according to the method of Thomas³⁰ using the following equations:

$$\text{Sensitivity (\%)} = \frac{\text{TP}}{\text{TP} + \text{FN}} \times 100$$

$$\text{Specificity (\%)} = \frac{\text{TN}}{\text{TN} + \text{FP}} \times 100$$

$$\text{PPV (\%)} = \frac{P \times \text{Sensitivity} \times 100}{(P \times \text{Sensitivity}) + (100 - P) \times (100 - \text{Specificity})}$$

$$\text{NPV (\%)} = \frac{(100 - P) \times \text{Specificity} \times 100}{(100 - P) \times \text{Specificity} + P \times (100 - \text{Sensitivity})}$$

The prevalence (*P*) was defined as the number of responders with a deterioration of dermatitis divided by total patients who




FIGURE 2 Pattern of clinical reactions (*n* = 103 double-blind placebo-controlled food challenges [DBPCFC]): Immediate reactions including delayed symptoms are defined as reactions occurring within 0–6 h after DBPCFC. Combined reactions are defined as immediate reactions developed within 0–6 h followed by an eczematous reaction within 6–48 h after DBPCFC, and isolated late cutaneous reactions are defined as a deterioration of atopic dermatitis (AD) within 6–48 h after DBPCFC without prior immediate-type reaction

underwent a DBPCFC. True-positive (TP) results were defined as patients who exhibited an eczematous reaction upon DBPCFC and who had birch-specific IgE levels at a cutoff point of 17.5 kU/L. Patients who did not show a worsening of AD after DBPCFC but had birch pollen-specific IgE determination of ≥ 17.5 kU/L were regarded as false-positive (FP) results. True-negative (TN) results were defined as patients without an eczematous reaction after DBPCFC and sIgE determination < 17.5 kU/L. Patients who showed an aggravation of AD after the DBPCFC but had a birch pollen-specific IgE levels < 17.5 kU/L were regarded as false-negative (FN) results.

3 | RESULTS

3.1 | Demographics and baseline characteristics

A total of 182 patients were included in this retrospective analysis (58% female, 42% male). The median age of the study population was 26 years (range 3–70 years; 82% ≥ 12 years and 18% < 12 years). The median disease severity before oral food challenges assessed by SCORAD index was 28.0 points (range 8.3–65.0 points).

Seventy-four percent of the patients suffered from seasonal or perennial respiratory allergic symptoms (allergic rhinoconjunctivitis [RCA] or asthma bronchiale).

3.2 | Clinical outcome of oral food challenges

In these 182 patients, consecutively, 261 DBPCFC with at least 1 birch pollen-related food (apple, carrot, celery, and hazelnut) were performed. One hundred and seven challenges were performed

TABLE 1 Clinical reaction pattern of oral food challenges with the different kinds of raw and heat-treated birch pollen-related foods

	n	Positive reactions (%)	Immediate reactions (%)	Combined reactions (%)	Late reactions (%)
DBPCFC with 4 kinds of birch pollen-related foods (raw and heat-treated)	107	40 (37)	20 (50)	11 (28)	9 (22)
DBPCFC with 2-3 kinds of birch pollen-related foods (raw and heat-treated)	52	17 (33)	9 (53)	5 (29)	3 (18)
	n	Positive reactions (%)	Immediate reactions (%)	Combined reactions (%)	Late reactions (%)
Apple (raw and heat-treated)	14	7 (50)	7 (100)	0 (0)	0 (0)
Carrot (raw and heat-treated)	29	11 (38)	10 (91)	0 (0)	1 (9)
Celery (raw and heat-treated)	30	18 (60)	11 (61)	5 (28)	2 (11)
Hazelnut (raw and heat-treated)	29	10 (35)	8 (80)	0 (0)	2 (20)

DBPCFC, double-blind placebo-controlled food challenge; n, number of oral food challenges.

using all 4 kinds of birch pollen-related foods. Fifty-two challenge tests were performed with 2 or 3 components, and 102 challenge tests were conducted with a single birch pollen-related food.

Sixty-five patients showed a clinical reaction (responders) with immediate and/or late-type allergic symptoms upon ingestion of birch pollen-related foods (103 DBPCFC) (Figure 1).

Together, 32 patients developed a late-type reaction with a significant increase in SCORAD index from prechallenge to postchallenge (median increase 15.4 points, 95% CI 12.4-16.3), whereas no significant increase in SCORAD index has been observed upon placebo challenges.


Of 103 positive oral food challenges, 63% were associated with an immediate-type reaction, and 37% of positive food challenges were accompanied by a late eczematous reaction (Figure 2).

Symptoms of immediate-type reactions in this study were OAS (58%), pruritus (12%), flush (9%), urticaria and/or angioedema (5%), gastrointestinal symptoms (5%), and asthma/dyspnea (3%). In 1 patient, an anaphylactic reaction with flush and hypotension occurred after ingestion of a verum meal consisting of all 4 birch pollen-related foods apple, carrot, celery, and hazelnut. In this case, no late eczematous reaction was observed.

3.3 | Clinical reaction patterns related to different birch pollen-related foods

Forty-nine percent of all positive DBPCFC with a verum meal containing 2-4 kinds of birch pollen-related foods were associated with a deterioration of AD, whereas DBPCFC with only 1 kind of food led to a broad variety of immediate and/or late-type symptoms (Table 1).

With regard to the DBPCFC with single birch pollen-related foods, raw and cooked celery were most frequently a trigger for an exacerbation of dermatitis (39% of positive food challenges with celery), while raw apple as well as raw and cooked carrot predominantly caused immediate-type reactions (100% and 91% of positive DBPCFC with these foods, respectively). Double-blind placebo-

**FIGURE 3** Age comparison between responders and nonresponders. * $P = .018$, Mann-Whitney U test

controlled food challenges with raw and heat-treated hazelnut caused a deterioration of AD in 20% of positive oral food challenges with this birch pollen-related food (Table 1).

3.4 | Oral food challenges with heat-treated birch pollen-related foods

Twenty-four of 261 oral food challenges were performed with heat-treated birch pollen-related foods. In total, 8 (33%) of these were positive with immediate-type symptoms in 6 cases. A combined reaction and an isolated late eczematous reaction occurred in another 2 DBPCFC.

In more detail, celery was particularly associated with a high rate of allergic symptoms despite heat treatment. Six of 7 DBPCFC with heat-treated celery were positive: In 4 cases, an immediate-type reaction was induced, while in 1 patient significant deterioration of




FIGURE 4 Patient's history of allergic rhinoconjunctivitis (RCA) in responders and nonresponders. * $P = .030$; chi-square test

AD and in another patient a combined reaction was observed after ingestion of cooked celery. Heat-treated apple caused an immediate-type allergic reaction in 1 of 3 tested patients while cooked carrot elicited an immediate-type reaction in 1 of 8 performed DBPCFC. Contrarily, no allergic reaction appeared upon DBPCFC with heated hazelnut ($n = 5$ DBPCFC).

3.5 | Characterization of responders and nonresponders

Patients who developed an allergic reaction upon DBPCFC were compared to nonresponders with regard to age, history of respiratory allergic diseases, and total and sIgE levels.

3.5.1 | Age and severity of AD

Responders were significantly older than nonresponders ($P = .018$; median age of responders and nonresponders: 29 [range 3-66 years] and 23 [range 3-70 years] years, respectively; Figure 3).

A significantly higher proportion of responders could be observed in adolescents ≥ 12 years and adults in comparison with children < 12 years of age (DBPCFC response rate 39% vs 14%, Fisher's exact test, $P = .01$). However, no significant difference in age could be

observed between patients with an isolated immediate-type reaction and those with a deterioration of AD. In more detail, there was a trend toward a higher proportion of late eczematous reactions to birch pollen-related foods in adolescent and adults compared to children < 12 years of age without reaching the level of significance (Fisher's exact test, $P = .613$). Likewise, the patients who experienced an aggravation of AD upon DBPCFC did not differ significantly from those with an isolated immediate-type reaction in terms of severity of AD before oral food challenge (Mann-Whitney U test, $P = .309$).

3.5.2 | Patient's history of respiratory atopic diseases

Regarding the patient's history, responders suffered significantly more often from RCA than nonresponders (Figure 4). No significant difference in the prevalence of RCA could be observed between responders with a deterioration of AD and those with an isolated immediate-type reaction (68% vs 84%, Fisher's exact test, $P = .213$).

With regard to both age groups, subjects ≥ 12 years of age showed a significantly higher prevalence of RCA and asthma bronchiale in comparison with subjects < 12 years of age (RCA 76% vs 14%, chi-square test $P < .001$; asthma 54% vs 14%, chi-square test $P = .001$). Responders and nonresponders did not significantly differ in terms of history of asthma bronchiale (data not shown).

3.5.3 | Specific IgE levels to birch pollen and birch pollen-related foods

Specific IgE levels against birch pollen and apple turned out to be significantly higher in responders compared to nonresponders, whereas no significant differences between both groups were found in terms of total IgE levels as well as sIgE to carrot, celery, and hazelnut (Table 2).

Responders with an eczematous late-type reaction ($n = 32$) and those with an isolated immediate-type reaction ($n = 33$) did not significantly differ in serum total IgE and sIgE levels (63.20 kU/L [95% CI 33.30-100.00] vs 53.35 kU/L [95% CI 38.35-100.00], Mann-Whitney U test $P = .928$). Likewise, no differences could be observed in serum birch pollen-specific antibodies between children and adolescent/adult patients (data not shown).

TABLE 2 Comparison of the in vitro diagnostic of responders with nonresponders (medians)

IgE (kU/L)	Responders (95% CI)	Nonresponders (95% CI)	P value*
Total IgE ($n_R = 54$; $n_N = 96$)	858.00 (507.50-1499.50)	1008.50 (431.51-1623.00)	.565
sIgE birch ($n_R = 57$; $n_N = 97$)	58.0 (38.70-88.80)	34.20 (26.00-47.70)	.047
Ratio sIgE birch/total IgE ($n_R = 54$; $n_N = 96$)	0.046 (0.023-0.082)	0.024 (0.017-0.034)	.100
sIgE apple ($n_R = 54$; $n_N = 72$)	5.00 (3.65-7.75)	2.67 (1.24-3.48)	.014
sIgE carrot ($n_R = 54$; $n_N = 70$)	1.87 (1.15-3.79)	1.19 (0.66-2.00)	.099
sIgE celery ($n_R = 53$; $n_N = 72$)	1.61 (0.93-2.54)	1.00 (0.45-1.61)	.070
sIgE hazelnut ($n_R = 55$; $n_N = 75$)	2.70 (1.37-5.68)	1.68 (1.06-2.87)	.087

n_N , number of nonresponders; n_R , number of responders; sIgE, specific IgE.

*Mann-Whitney U test.

3.6 | Diagnostic value of birch pollen-specific IgE

In this retrospective study, IgE sensitization to birch pollen at a cut-off point of 17.5 kU/L showed a high sensitivity (81%) and NPV (88%) for the diagnosis of deterioration of dermatitis, while the specificity (31%) and PPV (21%) here were low.

4 | DISCUSSION

This work confirms previous observations^{20–22} that birch pollen-related foods can cause immediate-type reactions in patients with AD and, moreover, may serve as a relevant provocation factor for an aggravation of AD. In the present study, 37% of 103 positive DBPCFC with birch pollen-related foods resulted in a worsening of AD. With respect to all tested patients, 18% reacted with a late eczematous reaction combined with or without a prior immediate-type reaction upon DBPCFC. This rate is lower than the frequency of eczematous reactions observed in former studies in adult patients (46%)²⁰ and in children with AD (33%).²¹ Such variable incidence of late-type reactions might be due to different study inclusion criteria: Reekers et al²⁰ only included adult patients who were highly sensitized against birch pollen (slgE ≥ 17.5 kU/L), and who did not have experienced an immediate-type reaction after ingestion of birch pollen-related foods before. Breuer et al²¹ only included children who were not supposed to suffer from an immediate-type reaction after ingestion of the tested foods. In the present study, the defined threshold for slgE to birch pollen was lower than in the trial of Reekers et al.²⁰ Moreover, 63% of responders retrospectively analyzed had exhibited an immediate-type reaction after the DBPCFC and, for safety reasons, had finally not undergone a repetitive provocation over 2 consecutive days. Subsequently, it could be assumed that in these patients the amount of allergen uptake was probably too low to elicit an eczematous reaction.

In accordance with former studies,^{31–35} OAS symptoms in the present study appeared as main immediate-type symptom of birch pollen-related food allergy. Only 1 DBPCFC performed with 4 kinds of birch pollen-related foods (apple, carrot, celery, and hazelnut) led to an anaphylactic reaction. It may be speculated that in this patient a late-type reaction was suppressed through antiallergic systemic treatment including corticosteroids. Overall, epinephrine administration was not required in any patient of this study. The dominance of mild allergic symptoms in terms of an OAS and the small number of systemic immediate-type reactions results from the instability of Bet v 1 homologue allergens against gastrointestinal digestion.³⁶ Pepsin completely abrogates the IgE-binding capacity of recombinant Mal d 1, Api g 1, and Cor a 1.04 within 1-second. Equally, trypsin destroys the IgE-binding of rMal d 1 and rApi g 1 after incubation for 15 minutes, whereas the major hazelnut allergen rCor a 1.04 shows a higher stability toward trypsinolysis. Moreover, it could previously be demonstrated that digested Mal d 1 and Cor a 1.04 still activate Bet v 1-specific T cells, whereas digested Api g 1 does not.³⁶ With regard to sequence similarity, this is lower between the major celery allergen

Api g 1 and Bet v 1 compared to the major hazelnut allergen Cor a 1.04 (61% and 79%, respectively).³⁷ However, we demonstrated *in vivo* that celery served as the most frequent elicitor of late eczematous reactions, whereas apple exclusively induced mild oropharyngeal symptoms. Likewise, carrot and hazelnut induced predominantly immediate-type reactions (91% and 80%, respectively). Hence, our data confirm previous results from Bohle et al²² who found that the degree of sequence similarity between Bet v 1-related food allergens and the major birch pollen allergen cannot be used as a marker to predict their potential to induce T cell-mediated symptoms.

Despite the fact that in all birch pollen-sensitized patients included in the present study birch pollen-related food allergy was suspected, and the overall response rate in DBPCFC was only 36%. However, those foods known to induce clear immediate-type reactions were excluded from oral food challenges. This approach probably contributed to the relatively low proportion of responders in this study.

From our data, we can also conclude that raw as well as cooked celery is able to cause a deterioration of the skin lesions. Consistently, it has already been demonstrated that thermal processing at 80°C for 15 minutes and at 100°C for 5 minutes abolishes IgE-binding and mediator release capacity of Bet v 1 homologue allergens Mal d 1, Dau c 1, and Api g 1 through changing their protein structure, whereas cooking does not influence their ability to stimulate Bet v 1-specific T cells and cause a deterioration of dermatitis.²² Thus, T-cell cross-reactivity between Bet v 1 and related food allergens occurs independently of IgE cross-reactivity *in vitro* and *in vivo*.

Data from Bohle et al²² further indicate that heating irreversibly destroys the conformational structure of Mal d 1 and Dau c 1, while Api g 1 shows a partial restoration of the protein structure. In our study, apple, carrot, and celery were cooked at 100°C for 5–15 minutes prior to the oral food challenge. This might explain previously published data^{38,39} and our finding on cooked celery that still exhibited a distinct allergenic potential with induction of immediate-type reactions in 5 of 7 patients. Surprisingly, 1 DBPCFC with heat-treated apple was accompanied by a mild immediate-type reaction of an OAS. It might be speculated that the apple used for the test meal was not heated sufficiently to abolish the IgE-binding capacity of Mal d 1. Moreover, subjective psychological factors cannot be excluded in this case as symptoms of an OAS cannot be verified by objective measurements. However, with respect to the mild severity of subjective symptoms in this birch pollen allergic patient sensitization to heat-resistant apple allergen Mal d 3 from the allergen family of lipid transfer proteins (LTP) is rather unlikely. In fact, LTP sensitization is predominantly found in nonbirch pollen-sensitized patients from Mediterranean areas exhibiting more severe clinical reactions.⁴⁰ Nevertheless, in future studies in this field, assessment of slgE sensitization to rBet v 1 and the corresponding homologue molecular food allergens from the PR-10 family (ie, rMal d 1, rDau c 1, rApi g 1 and rCor a 1) should be performed to improve the diagnostic accuracy. It might be supposed that molecular allergy diagnosis would also allow to identify and to characterize distinct sensitization patterns in birch pollen-sensitized subjects facilitating the diagnosis of birch pollen-related food allergy.

In contrast to previous studies,^{20,21} we found that the responders were significantly higher IgE-sensitized against birch pollen and apple compared to the nonresponders, while total IgE and sIgE levels against carrot, celery, and hazelnut did not differ between both groups. The latter 2 results are in line with the data from Guhl et al³⁵ who showed that sIgE testing with food extracts from celery and hazelnut is not suitable to discriminate between food-allergic and tolerant patients sensitized against Bet v 1. As expected, responders in the present study were significantly older than nonresponders and showed a significantly higher prevalence of RCA. This outcome reflects the natural course of food allergy and respiratory allergy.⁴¹ However, age, birch pollen sIgE levels, and the presence of a RCA could not be used to differentiate between responders with isolated immediate-type reactions and those with late eczematous reactions. In accordance with previous research,^{20,21} no differences could be observed between responders and nonresponders in terms of the disease severity, total IgE, or food-specific IgE against carrot, celery, or hazelnut.

The main limitation of the study that should be considered in the interpretation of our results is its retrospective and monocentric design. Therefore, our results cannot be transferred to other patient populations, especially not to those outside birch-endemic regions. Further prospective multicenter studies are needed to reproduce these results in larger cohorts from different geographical areas.

In conclusion, our data confirm that in a subpopulation of patients with AD and IgE sensitization to birch pollen corresponding foods should be considered as potential trigger for an aggravation of dermatitis. Even if the responders with immediate and/or late-type reaction tended to have higher sIgE levels against birch pollen compared to the nonresponders, the diagnostic value of the sIgE determination as a single parameter is insufficient for the diagnosis of a late-type reaction. Thus, DBPCFC cannot be replaced in patients with AD and suspected late eczematous reaction after ingestion of birch pollen-related foods.

CONFLICTS OF INTEREST

The authors declare that they have no conflicts of interest.

AUTHOR CONTRIBUTIONS

Anja Wassmann-Otto, Annice Heratizadeh, and Thomas Werfel conceived and designed the retrospective analyses. Anja Wassmann-Otto, Annice Heratizadeh, Katja Wichmann, and Thomas Werfel discussed the data. Annice Heratizadeh, Katja Wichmann, and Thomas Werfel revised the article critically. Anja Wassmann-Otto analyzed and interpreted the data and wrote the manuscript.

ORCID

A. Wassmann-Otto  <http://orcid.org/0000-0001-9923-6053>

REFERENCES

- Schäfer T. The impact of allergy on atopic eczema from data from epidemiological studies. *Curr Opin Allergy Clin Immunol*. 2008;8:418-422.
- Heratizadeh A, Wichmann K, Werfel T. Food allergy and atopic dermatitis: how are they connected? *Curr Allergy Asthma Rep*. 2011;11:284-291.
- Werfel T, Allam JP, Biedermann T, et al. Cellular and molecular immunologic mechanisms in patients with atopic dermatitis. *J Allergy Clin Immunol*. 2016;138:336-349.
- Ho MH, Wong WH, Chang C. Clinical spectrum of food allergies: a comprehensive review. *Clin Rev Allergy Immunol*. 2014;46:225-240.
- Vieths S, Scheurer S, Ballmer-Weber B. Current understanding of cross-reactivity of food allergens and pollen. *Ann N Y Acad Sci*. 2002;964:47-68.
- Breiteneder H, Ferreira F, Hoffmann-Sommergruber K, et al. Four recombinant isoforms of Cor a I, the major allergen of hazel pollen, show different IgE-binding properties. *Eur J Biochem*. 1993;212:355-362.
- Breiteneder H, Hoffmann-Sommergruber K, O'Riordain G, et al. Molecular characterization of Api g 1, the major allergen of celery (*Apium graveolens*), and its immunological and structural relationships to a group of 17-kDa tree pollen allergens. *Eur J Biochem*. 1995;233:484-489.
- Vanek-Krebitz M, Hoffmann-Sommergruber K, Da Camara L, et al. Cloning and sequencing of Mal d 1, the major allergen from apple (*Malus domestica*), and its immunological relationship to Bet v 1, the major birch pollen allergen. *Biochem Biophys Res Commun*. 1995;214:538-551.
- Scheurer S, Metzner K, Hausteiner D, Vieths S. Molecular cloning, expression and characterization of Pru a 1, the major cherry allergen. *Mol Immunol*. 1997;34:619-629.
- Hoffmann-Sommergruber K, O'Riordain G, Ahorn H, et al. Molecular characterization of Dau c 1, the Bet v 1 homologous protein from carrot and its cross-reactivity with Bet v 1 and Api g 1. *Clin Exp Allergy*. 1999;29:840-847.
- Karamloo F, Scheurer S, Wangorsch A, May S, Hausteiner D, Vieths S. Pyr c 1, the major allergen from pear (*Pyrus communis*), is a new member of the Bet v 1 allergen family. *J Chromatogr B Biomed Sci Appl*. 2001;756:281-293.
- Kleine-Tebbe J, Vogel L, Crowell DN, Hausteiner U, Vieths S. Severe oral allergy syndrome and anaphylactic reactions caused by a Bet v 1-related PR-10 protein in soybean, SAM22. *J Allergy Clin Immunol*. 2002;110:797-804.
- Mittag D, Akkerdaas J, Ballmer-Weber BK, et al. Ara h 8, a Bet v 1-homologous allergen from peanut, is a major allergen in patients with combined birch pollen and peanut allergy. *J Allergy Clin Immunol*. 2004;114:1410-1417.
- Mittag D, Vieths S, Vogel L, et al. Birch pollen-related food allergy to legumes: identification and characterization of the Bet v 1 homologue in mungbean (*Vigna radiata*), Vig r 1. *Clin Exp Allergy*. 2005;35:1049-1055.
- Etesamifar M, Wüthrich B. IgE-vermittelte Nahrungsmittelallergien bei 383 Patienten unter Berücksichtigung des oralen Allergie-Syndroms. *Allergologie*. 1998;21:451-457.
- Ballmer-Weber BK, Vieths S, Lüttkopf D, Heuschmann P, Wüthrich B. Celery allergy confirmed by double-blind, placebo-controlled food challenge: a clinical study in 32 subjects with a history of adverse reactions to celery root. *J Allergy Clin Immunol*. 2000;106:373-378.
- Ballmer-Weber BK, Wüthrich B, Wangorsch A, Fötisch K, Altmann F, Vieths S. Carrot allergy: double-blinded, placebo-controlled food challenge and identification of allergens. *J Allergy Clin Immunol*. 2001;108:301-307.

18. Mittag D, Vieths S, Vogel L, et al. Soybean allergy in patients allergic to birch pollen: clinical investigation and molecular characterization of allergens. *J Allergy Clin Immunol.* 2004;113:148-154.
19. Ballmer-Weber BK, Holzhauser T, Scibilia J, et al. Clinical characteristics of soybean allergy in Europe: a double-blind, placebo-controlled food challenge study. *J Allergy Clin Immunol.* 2007;119:1489-1496.
20. Reekers R, Busche M, Wittmann M, Kapp A, Werfel T. Birch pollen-related foods trigger atopic dermatitis in patients with specific cutaneous T-cell responses to birch pollen antigens. *J Allergy Clin Immunol.* 1999;104:466-472.
21. Breuer K, Wulf A, Constien A, Tetat D, Kapp A, Werfel T. Birch pollen-related food as a provocation factor of allergic symptoms in children with atopic eczema/dermatitis syndrome. *Allergy.* 2004;59:988-994.
22. Bohle B, Zwölfer B, Heratizadeh A, et al. Cooking birch pollen-related food: divergent consequences for IgE- and T cell-mediated reactivity in vitro and in vivo. *J Allergy Clin Immunol.* 2006;118:242-249.
23. Werfel T, Asero R, Ballmer-Weber BK, et al. Position paper of the EA ACI: food allergy due to immunological cross-reactions with common inhalant allergens. *Allergy.* 2015;70:1079-1090.
24. Muraro A, Roberts G (Eds.). *EAACI-Guidelines - Food Allergy and Anaphylaxis*. EAACI (Zurich); 2014.
25. Hanifin JM, Rajka G. Diagnostic features of atopic dermatitis. *Acta Derm Venereol (Suppl).* 1980;92:44-47.
26. Werfel T, Ballmer-Weber B, Eigenmann PA, et al. Eczematous reactions to food in atopic eczema: position paper of the EAACI and GA2LEN. *Allergy.* 2007;62:723-728.
27. Stalder JF, Taib A, Atherton DJ, et al. Severity scoring of atopic dermatitis: the SCORAD index. Consensus report of the European task force on atopic dermatitis. *Dermatology.* 1993;186:23-31.
28. Sampson HA, Gerth van Wijk R, Bindslev-Jensen C, et al. Standardizing double-blind, placebo-controlled oral food challenges: American Academy of Allergy, Asthma & Immunology-European Academy of Allergology and Clinical Immunology PRACTALL consensus report. *J Allergy Clin Immunol.* 2012;130:1260-1274.
29. Niggemann B, Beyer K, Erdmann S, et al. Standardisation of oral provocation tests on suspicion of food allergy. *Allergo J.* 2011;20:149-160.
30. Thomas L. *Labor und Diagnose. Indikation und Bewertung von Laborbefunden für die medizinische Diagnostik* (7. Aufl). Frankfurt, Germany: TH-Books-Verl.-Ges.; 2008.
31. Skamstrup Hansen K, Vestergaard H, Stahl Skov P, et al. Double-blind, placebo-controlled food challenge with apple. *Allergy.* 2001;56:109-117.
32. Worm M, Hompes S, Fiedler E, Illner A, Zuberbier T, Vieths S. Impact of native, heat-processed and encapsulated hazelnuts on the allergic response in hazelnut-allergic patients. *Clin Exp Allergy.* 2009;39:159-166.
33. Ebo DG, Verweij MM, Sabato V, Hagendorens MM, Bridts CH, De Clerck LS. Hazelnut allergy: a multi-faced condition with demographic and geographic characteristics. *Acta Clin Belg.* 2012;67:317-321.
34. Masthoff LJ, van Hoffen E, de Reus A, et al. Hazelnut allergy differs between children and adults in frequency of severity, aetiology and relevance of diagnostic parameters. *Clin Exp Allergy.* 2014;44:1539-1545.
35. Guhl EE, Hofstetter G, Lengger N, et al. IgE, IgG4 and IgA specific to Bet v 1-related food allergens do not predict oral allergy syndrome. *Allergy.* 2015;70:59-66.
36. Schimek EM, Zwölfer B, Briza P, et al. Gastrointestinal digestion of Bet v 1-homologous food allergens destroys their media-tor-releasing, but not T cell-activating, capacity. *J Allergy Clin Immunol.* 2005;116:1327-1333.
37. Jahn-Schmid B, Radakovics A, Lüttkopf D, et al. Bet v 1142-156 is the dominant T-cell epitope of the major birch pollen allergen and important for cross-reactivity with Bet v 1-related food allergens. *J Allergy Clin Immunol.* 2005;116:213-219.
38. Jankiewicz A, Baltes W, Boegl W, et al. Influence of food processing on the immunochemical stability of celery allergens. *J Sci Food Agric.* 1997;75:359-370.
39. Ballmer-Weber BK, Hoffmann A, Wüthrich B, et al. Influence of food processing on the allergenicity of celery: DBPCFC with celery spice and cooked celery in patients with celery allergy. *Allergy.* 2002;57:228-235.
40. Fernández-Rivas M, Bolhaar S, González-Mancebo E, et al. Apple allergy across Europe: how allergen sensitization profiles determine the clinical expression of allergies to plant foods. *J Allergy Clin Immunol.* 2006;118:481-488.
41. Savage S, Johns CB. Food allergy: epidemiology and natural history. *Immunol Allergy Clin North Am.* 2015;35:45-59.

How to cite this article: Wassmann-Otto A, Heratizadeh A, Wichmann K, Werfel T. Birch pollen-related foods can cause late eczematous reactions in patients with atopic dermatitis. *Allergy.* 2018;73:2046-2054. <https://doi.org/10.1111/all.13454>