

Therapeutic immunomodulators from nematode parasites

William Harnett^{1,*} and Margaret M. Harnett²

There has been an alarming increase in the incidence of autoimmune and allergic diseases in Western countries in the past few decades. However, in countries endemic for parasitic helminth infections, such diseases remain relatively rare. Hence, it has been hypothesised that helminths may protect against the development of autoimmunity and allergy. This article reviews the evidence supporting this idea with respect to helminths of the phylum Nematoda (nematodes), considering data from human studies and animal models of inflammatory disease. The nature and mode of action of nematode-derived molecules with immunomodulatory properties are considered, and their therapeutic efficacy in models of autoimmunity and allergy described. The recent and future use of nematodes and their products in treating human disease are also discussed.

The phylum *Nematoda* constitutes one of the largest and most abundant groups of animals. Although most species of these worms are free-living, this phylum also contains parasites of virtually every multicellular organism on the planet (Ref. 1). This, allied to the fact that in many cases parasitism has severe and even fatal consequences for the host, dictates that nematodes are of substantial medical, veterinary and economic importance. For this reason, parasitic nematodes have been the subject of much recent study.

Nematodes are generally small, but the existence of larger parasitic species (e.g. *Ascaris lumbricoides*, which may grow to over 40 cm in length) has been known of for thousands of years (Ref. 1). Typically, the worms are slender,

bilaterally symmetrical organisms that are encased in a noncellular cuticle. The cuticle is shed four times during the course of a nematode's life and the five stages are often referred to as L1–L5, with L5 being the adult stage (Ref. 1).

Adult parasitic nematodes can inhabit two locations within humans: the gastrointestinal tract or the tissues. The most common gastrointestinal nematodes of humans are *A. lumbricoides*, *Trichuris trichiura*, *Necator americanus* and *Ancylostoma duodenale* (the latter two species are 'hookworms') (Table 1). An infected individual may harbour more than one type of these parasites, and collectively, these four species of worm may infect more than one third of the world's population. The major tissue-dwelling

¹Strathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, Glasgow, UK

²Division of Immunology, Infection and Inflammation, Glasgow Biomedical Research Centre, University of Glasgow, 120 University Place, Glasgow, G12 8TA, UK

*Corresponding author: William Harnett, Strathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, 27 Taylor Street, Glasgow, G4 0NR, UK. Tel.: +44 141 548 3725; Fax: +44 141 552 2562; E-mail: w.harnett@strath.ac.uk

Table 1. Estimated prevalence of the major medically important human nematode parasites

Nematode	Number of people infected
<i>Ascaris lumbricoides</i>	1.5 billion
<i>Trichuris trichiura</i>	800 million
Hookworms ^a	800 million
Lymphatic filarial nematodes ^b	120 million
<i>Onchocerca volvulus</i>	35 million

^a*Ancylostoma duodenale* and *Necator americanus*;
^b*Wuchereria bancrofti* and *Brugia malayi*

nematodes of humans are the filarial nematodes, the adult worms of which may live in the lymphatic system (e.g. *Wuchereria bancrofti*, *Brugia malayi*) or the subcutaneous tissues (*Onchocerca volvulus*) (Ref. 2). Infection with gastrointestinal nematodes may arise through ingestion of eggs [*A. lumbricoides* (Ref. 3), *T. trichiura* (Ref. 4)] or penetration of the skin by L3 larvae (hookworms) (Ref. 5). Infection with filarial nematodes occurs when L3 larvae are transferred to humans by blood-feeding insects that act as vectors (Ref. 2). The majority of people infected with nematodes tend to be asymptomatic, but some species can cause very severe problems such as elephantiasis (e.g. *W. bancrofti*, *B. malayi*) and blindness (*O. volvulus*) (Ref. 2).

Parasitic nematodes can protect humans from allergy and autoimmunity

One surprising feature of nematodes is their longevity. It is reported, for example, that *W. bancrofti* can live in excess of a decade (Ref. 6) and, although a number of factors may well be involved, it is assumed that a major factor contributing to long-term infection is suppression of the host immune system. In relation to this, analysis of the immune response during human infection generally indicates that people having patent infection – that is infection with larval stages (microfilariae) in their bloodstreams – demonstrate reduced lymphocyte proliferative responses and decreased production of the cytokine interferon- γ (IFN- γ), but increased production of the cytokines interleukin- (IL) 4

and IL-10 and the antibody type IgG4 (Ref. 7). As IL-4 is a cytokine that is classically associated with helper T-cells of the type 2 category (Ref. 8) and IL-10 is an anti-inflammatory cytokine (Ref. 9), such an immune response may be characterised as being of a type-2 T-helper cell (Th2), anti-inflammatory phenotype. There is probably a benefit for humans in the generation of such an immunological phenotype because, as alluded to earlier, although filarial nematode infection can result in severe pathology, such as elephantiasis, the majority of infected individuals exhibit little evidence of an inflammatory response or overt tissue destruction or disruption (Ref. 10). Similar conclusions are now being made with respect to gastrointestinal nematodes (Ref. 11) and indeed, it is now accepted that a Th2-like, somewhat anti-inflammatory immunological phenotype is a feature of nematode infection. An interesting characteristic of this phenotype is that it has been reported in some cases to extend to subsequent responses to non-nematode antigens – for example to antigens in vaccines targeting bacterial infections that rely on Th1 responses for efficacy (Ref. 12). One immediate thought, therefore, is whether such a change in immune-responsiveness could lead to changes in susceptibility to diseases associated with aberrant Th1-associated inflammatory responses, such as the autoimmune diseases. In relation to this, there is currently great interest in the idea that the recent alarming increase in autoimmune and allergic diseases in the industrialised world, but not developing countries, reflects increased hygiene, causing a reduction in infection with pathogens such as parasitic worms, an idea known as the ‘hygiene hypothesis’ (Ref. 13). Consistent with this, it has been suspected for many years that allergic and autoimmune diseases may be less prevalent in areas of nematode transmission (see Ref. 14). Interestingly, results from human and animal studies find evidence for infection associated not only with reduced autoimmunity, but also with decreased allergy, an initially surprising observation given the Th2-associated status of this type of disease (Ref. 15).

Evidence from human studies

Although there are cases in which no association, or indeed increased susceptibility, has been observed, the majority of studies support the idea that nematode infection can protect against allergy, as measured by factors such as skin

reactivity to allergens or wheezing, and more consistently, by comparing symptoms before and after anthelmintic treatment (for reviews, see Refs 12, 14, 16, 17, 18) (Table 2). Autoimmune diseases have not been as well studied as allergic disease, but again there is evidence supporting an inverse relationship between nematode infection and conditions such as inflammatory bowel disease, type-1 diabetes, rheumatoid arthritis and multiple sclerosis (for reviews, see Refs 14, 28, 29, 30). Furthermore, an inverse relationship between infection with *Strongyloides stercoralis* and autoimmune liver disease has recently been reported (Ref. 31). Thus, overall, the idea that helminths can protect against human allergic and autoimmune inflammatory disease is compelling.

Evidence from animal studies

At the outset, it has to be appreciated that using animal models to investigate whether nematodes are likely to protect humans against allergy and autoimmunity has the limitation of differences in the pattern of exposure. Models invariably employ one or two exposures to a

significant number of nematodes, whereas humans will generally acquire their nematodes one or a few at a time, over a period of many years. There are a number of recent mouse model studies that show nematode infection can protect against allergic disease, in particular lung-associated disease (Table 3). This type of approach has also provided evidence that nematodes can protect against autoimmune diseases, such as colitis and diabetes. Hence, the use of mouse models in this area has produced a series of convincing findings supporting the hypothesis that nematodes may protect humans from allergy and autoimmunity.

Mechanism of action

The role of Th2 polarisation and regulatory T cells

As mentioned above, nematode infections are chronic and have been classically characterised by the induction of a Th2 immune response (Ref. 41). However, more recently they have also been associated with regulatory T-cell (Treg) responses (Ref. 42). It is convenient to simply view these two types of immune response in terms of the cytokines generally associated with

Table 2. Nematode infection protects humans from allergic disease

Nematode species	Observation	Ref.
Various gastrointestinal nematodes	Reduced skin reactivity to house dust mite allergen with infection	19
Various gastrointestinal nematodes	Increased skin reactivity to house dust mite allergen and serum IgE against environmental antigens following anthelmintic treatment; decreased reactivity with increased worm infection	20
<i>Ascaris lumbricoides</i> and <i>Necator americanus</i>	Inverse correlation between incidence of asthma and infection	21
<i>Ascaris lumbricoides</i> and hookworms	Inverse association between skin-test reactivity to various allergens and infection	22
Hookworm	Inverse association between recent wheeze and infection	23
<i>Ascaris lumbricoides</i> and hookworms	Reduced skin-test reactivity to various allergens with greater worm burden	24
<i>Ascaris lumbricoides</i>	Inverse association between recent wheeze and infection	25
<i>Ascaris lumbricoides</i> and <i>Trichuris trichiura</i>	Increased skin reactivity to house dust mites with anthelmintic treatment	26
<i>Ascaris lumbricoides</i> and hookworms	Inverse correlation between infection and skin sensitisation to house dust mites	27

Table 3. Nematode infection can protect mice from allergic and autoimmune diseases

Nematode species	Observation	Ref.
Allergic disease		
<i>Strongyloides stercoralis</i>	Suppression of allergic response to ovalbumin in the lungs	32
<i>Heligomosomoides polygyrus</i>	Inhibition of allergic responses to peanut extract	33
<i>Nippostrongylus brasiliensis</i>	Inhibition of ovalbumin-induced airway eosinophilia and eotaxin levels in the lungs	34
<i>Heligomosomoides polygyrus</i>	Inhibition of airway hyper-responsiveness to ovalbumin and house dust mite allergen Der p 1	35
<i>Heligomosomoides polygyrus</i>	Suppression of ovalbumin-induced airway eosinophilia and bronchial hyper-reactivity	36
Autoimmune disease		
<i>Heligomosomoides polygyrus</i> and <i>Trichuris muris</i>	Inhibition of colitis development in IL-10 ^{-/-} mice	37
<i>Trichinella spiralis</i>	Reduced colitis induced by DNBS	38
<i>Heligomosomoides polygyrus</i>	Protection from colitis in piroxicam-treated IL-10 ^{-/-} mice	39
<i>Heligomosomoides polygyrus</i> and <i>Trichinella spiralis</i>	Protection against diabetes in the NOD mouse	40

Abbreviations: DNBS, dinitrobenzene sulfonic acid sodium salt; NOD, non-obese diabetic.

them, for example IL-4, IL-5 and IL-13 for Th2 responses, and IL-10 and transforming growth factor- β (TGF- β) for (certain) regulatory responses (Fig. 1). Although the host may have evolved the latter to deal with any persistent significant antigen challenge, it is clear that nematodes are particularly adept at polarising immune responses towards Th2, in particular by their effects on dendritic cells, the sentinels of the immune system that sample and process foreign molecules for presentation to T cells. Presentation of peptides derived from antigens is accompanied by signals generated by dendritic cell surface costimulatory molecules and secreted cytokines interacting with receptors on T cells, and this combination impacts on T-cell polarisation (Ref. 43). Thus, nematode antigens must contain information that facilitates dendritic cell-driven Th2 polarisation. Interestingly, compared with antigens that induce Th1 responses, nematode-derived molecules have generally been shown to have little or only a partial effect on dendritic cell activation as measured by upregulation of

markers such as costimulatory molecules (Refs 44, 45). The induction of Th2 responses may reflect particular characteristics of nematode molecules (see Ref. 46). However, the advantage of this to the worms is uncertain, because there is evidence from rodent models of gastrointestinal nematode infection that Th2 immune responses are protective (Ref. 47). This is also supported by some human epidemiological studies (Ref. 11). Moreover, Th2 responses are associated with inflammatory pathology during gastrointestinal nematode infection (Ref. 48). Nevertheless, it is possible to perceive that the induction of an immunological environment polarised towards Th2 may provide a situation in which the autoimmune diseases generally considered to be Th1 dependent (Ref. 49) would find it difficult to flourish. Consistent with this, a reduction in inflammation is observed in IL-10^{-/-} mice with colitis upon infection with *Heligomosomoides polygyrus*, which reflects an inhibition of mucosal production of IFN- γ and IL-12 p40, and the increased production of IL-13 (Ref. 39).




Figure 1. Potential mechanisms of nematode-induced immunomodulation and suppression of inflammatory disease. Infection with nematodes can induce T-cell hypo-responsiveness by directly or indirectly (through antigen-presenting cells such as dendritic cells) suppressing T-cell activation. For example, infection can result in the induction of T regulatory (Treg) cells that are refractory to antigen and can prevent maximal T-effector responses of Th1-, Th2- or Th17-based inflammation. Moreover, worm products such as ES-62 can induce B1 cells to produce IL-10, which acts to suppress dendritic cell maturation and hence results in T-cell anergy, preventing Th1-Th2- or Th17-mediated inflammation. Alternatively, the worms can modulate dendritic cell maturation to polarise T-cell responses towards a Th2 phenotype and hence counteract Th1-based inflammation. By contrast, it has been proposed that nematodes can antagonise Th2-driven inflammation by producing polyspecific IgE or IgG4 to block allergen-induced mast-cell degranulation. Moreover, certain excretory-secretory (ES) products have been shown to directly induce mast-cell responsiveness by subverting FcεRI signalling (see Fig. 2). Abbreviations: CTLA, cytotoxic T-lymphocyte antigen; MHC, major histocompatibility complex; TCR, T-cell receptor.

Considering regulatory responses, these may be generated in response to any chronic inflammatory insult in an attempt to prevent it getting out of control, but with respect to nematodes, recent evidence intriguingly suggests that they may contribute to worm persistence during infection

via inhibition of protective immunity (Ref. 50). Of particular relevance to this review however, is the fact that regulatory responses do not only inhibit inflammatory responses to nematodes, but also to heterologous antigens. This means that responses to other infections (Ref. 12) or even

vaccines (Ref. 12) may be affected, and in addition, susceptibility to diseases associated with aberrant inflammatory responses altered. The Th2 dominance associated with nematode infection provides an attractive argument to explain the ability of the helminths to inhibit autoimmune diseases, but it obviously does not hold true with respect to allergic diseases, which are themselves Th2 dependent (Ref. 15). However, both Th1 and Th2 responses may be dampened down by anti-inflammatory regulatory responses (Ref. 42) and hence the ability of nematodes to protect against allergy may be explained in this way. Consistent with the latter, the ability of *H. polygyrus* to inhibit allergic airway hyper-responsiveness in the mouse has been shown by the use of adoptive transfer experiments to be dependent on CD4⁺ CD25⁺ Tregs (Ref. 35). In terms of cytokines, C57BL/6 strain mice showed increased IL-10 production in response to infection, whereas BALB/c mice demonstrated augmented TGF- β . The latter result may explain why the use of BALB/c-knockout mice showed no role for the anti-inflammatory molecule IL-10 in generating the protective response (Ref. 35) – a result not observed with C57BL/6 animals (Ref. 36). These results are also consistent with the ability of T-cell lines expressing IL-10 or TGF- β to protect against allergy in the ovalbumin-sensitisation model (Ref. 51).

It has already been mentioned that IL-10^{-/-} mice with colitis show a reduction in inflammation when infected with *H. polygyrus* (Ref. 39). There could be a regulatory component operating in this system, because interestingly, the protective effect can be transferred by mesenteric lymph node T cells from infected animals, and these cells were shown to have increased *Foxp3* mRNA expression – a key marker of Treg function. It was recently demonstrated that *H. polygyrus* infection is associated with the generation of a regulatory CD8⁺ T-cell population in the lamina propria (Ref. 52). Interestingly, *H. polygyrus* also causes increased expression of TLR4 on mucosal T cells, and interaction of TLR4 with its ligand LPS resulted, not in release of the expected IL-12 or TNF- α , but rather of the cytokine associated with regulatory responses TGF- β (Ref. 53).

Role of nonspecific IgE production in allergy prevention

The idea of regulatory responses preventing the development of allergic responses during

nematode infection is clearly supported by experimental evidence; however, other possible explanations have been tested. Nematodes have a tendency to induce considerable nonspecific IgE secretion during infection (which may prevent specific mast cell- and/or basophil-mediated attack) and it has been suggested that by saturating Fc ϵ RI sites, these antibodies may block mast-cell degranulation that would be mediated by allergens (reviewed in Ref. 54). Certainly, an association between high levels of circulating nonspecific IgE and protection against allergy has been shown in a number of studies (see Ref. 54). Moreover, increased cutaneous allergic responses following anthelmintic treatment are associated with a corresponding decrease in nonspecific IgE (Ref. 20). However, it has been found that *Nippostrongylus brasiliensis* infection does not reduce mast-cell degranulation in response to ovalbumin in the skin of immunised mice, in spite of the infection generating a massive increase in nonspecific IgE production (Ref. 34). It has also been reported, again using *N. brasiliensis*, that a small amount of IgE specific for the parasite antigen Nb-Ag1, is still able to facilitate mast-cell degranulation in the presence of much greater levels of nonspecific IgE (Ref. 55). Finally, studies on humans infected with a variety of filarial nematodes indicate that the relative levels of polyclonal IgE and filaria-specific IgE do not tend to reach that required for inhibition of filaria-induced mast-cell or basophil degranulation (Ref. 56). The same outcome was observed when examining dust-mite-specific IgE. Thus overall, it appears that saturation of Fc ϵ RI by polyspecific IgE antibodies during nematode infection cannot alone explain protection from allergy (Fig. 1).

Interestingly, although nematode infection increases polyspecific IgE production and as mentioned above, does not appear to inhibit the IgE response to allergens, there have been reports where the opposite is the case. Thus for example, *S. stercoralis* infection of mice results in reduced anti-ovalbumin IgE levels in the lungs (Ref. 32). Likewise, *H. polygyrus* infection results in decreased circulating IgE levels in mice in response to peanut allergen (Ref. 33). *H. polygyrus* infection has also been shown to increase ovalbumin-specific IgE as well as total IgE in a mouse model of atopic asthma (Ref. 36).

Nematode infections can also be associated with the production of high levels of IgG4, an antibody that competes with IgE for allergen binding but does not promote mast-cell or basophil degranulation (Ref. 57) (Fig. 1). However, studies investigating whether high levels of IgG4 can protect against allergic reactions in humans have produced conflicting results (Ref. 51).

IL-10, Tregs and mast-cell function

As alluded to earlier (e.g. Ref. 35) a feature of some nematode infections is the production of IL-10 and/or Tregs, and some recent papers suggest that both of these could have an impact on mast-cell function. Thus, it was shown that IL-10 inhibited the release of TNF- α , IL-8 and histamine by human mast cells activated via Fc ϵ RI (Ref. 58). Furthermore, the use of IL-10 transgenic mice demonstrated that overexpression of the cytokine reduces passive cutaneous anaphylaxis responses (Ref. 59). Also, staying with mice, studies on mast cells have revealed that Fc ϵ RI expression and IgE-mediated release of leukotriene C4 can both be inhibited by Tregs (Ref. 60). Thus, although these observations have not been related to nematode infections as yet, it is possible that nematode induction of IL-10 and/or Tregs could interfere with mast-cell function and hence allergic responses.

Molecules involved in immunomodulation in animal models

Excretory-secretory products of *Nippostrongylus brasiliensis*

An excretory-secretory (ES) extract, containing as-yet-undefined components of adult stage *N. brasiliensis*, inhibits the development of asthma (ovalbumin-hypersensitivity model) in the lungs of mice in a similar manner to infection with the parasite itself (Ref. 61). Administration of adult ES during the sensitisation period entirely blocked eosinophilia and mucus production in the lungs and also significantly inhibited airway hyper-reactivity development. Ovalbumin-specific IgG1 and IgE levels in the bloodstream and the amount of both IL-4 and IL-5 in the lungs were all reduced by exposure to the ES. No role was found for Toll-like receptor (TLR)2, TLR4, IFN- γ or IL-10. Moreover, such ES-induced parasite-specific Th2 responses develop as normal. The active agent appears

unlikely to be a protein, because it is resistant to both heat and proteinase K. Similarly, intratracheal installation of ES derived from the *N. brasiliensis* L3 stage reduces neutrophil influx in response to LPS in the lungs of Wistar rats (Ref. 62).

Ascaris suum PAS1

It was reported in 2002 that an *A. suum* extract could inhibit lung inflammation and hyper-responsiveness and suppress IL-4, IL-5 and eotaxin in response to hen's egg white (Ref. 63). More recently, it was discovered that *A. suum* contains a protein, APAS3, which, upon immunisation and challenge is able to induce Th2-dependent, eosinophilic airway hyperreactivity in BALB/c mice (Ref. 64). Interestingly, another *A. suum* product, PAS1, inhibits this response (Ref. 64). Inhibition is associated with reduced Th2 cytokines (IL-4, IL-5) and chemokines [eotaxin, RANTES (regulated upon activation, normal T-cell expressed and secreted)], and considerably elevated IL-10 levels in bronchoalveolar lavage fluid. Measurement of serum levels also showed that production of IgG1 and IgE was blocked. PAS1 is most active when administered prophylactically at both immunisation and challenge stages; administration at immunisation only reveals a partial effect, but PAS1 is completely ineffective when given only at challenge. PAS1 also induces IL-10 and TGF- β production in macrophages and this is associated with loss of proinflammatory cytokine production (Ref. 65).

Dirofilaria immitis-derived antigen (DiAg)

A purified 15 kDa protein from the canine filarial nematode, *Dirofilaria immitis* known as DiAg (*Dirofilaria immitis*-derived antigen), causes polyclonal proliferation of B cells and nonspecific IgE production (Ref. 66). Interestingly DiAg prevents the spontaneous generation of IgG anti-insulin antibodies seen in the NOD mouse, and this is associated with a lack of development of Th1-dependent autoimmune diabetes. Consistent with this, DiAg increases levels of IL-4 and IL-10 when injected into BALB/c mice. DiAg can also inhibit passive cutaneous anaphylaxis (PCA) reactions in rats (Ref. 66). This was found not to be due to an effect on the number or viability of mast cells or to differences in the amount of histamine released, but interestingly, given the relative dismissal of this mode of action in the human situation, appears to be due to nonspecific saturation of Fc ϵ RI.

ES-62

ES-62 is a phosphorylcholine (PC)-containing glycoprotein secreted by filarial nematodes, which has been shown to have a wide range of immunomodulatory properties that are largely anti-inflammatory (Ref. 67). Consequently, the molecule has been tested in mouse models of autoimmune and allergic disease and has been found to be active against collagen-induced arthritis (CIA) (Ref. 68) and type-I hypersensitivity in the skin and lungs (Ref. 69).

With respect to CIA, ES-62 was effective as a prophylactic treatment and also therapeutically, and in both cases, *ex vivo* analyses revealed that the effects were associated with inhibition of collagen-specific proinflammatory (TNF- α , IL-6) and Th1 (IFN- γ) cytokines and (IgG2a) antibody production (Ref. 68). In the

prophylactic experiment, increased production of collagen-specific IL-10 was also observed. More recent work on the CIA model, involving a comparison of ES-62, PC-free ES-62 and PC conjugated with an irrelevant protein (ovalbumin), has revealed that the antiarthritis effect of ES-62 is dependent on its PC moiety, although interestingly, not on a reduction in collagen-specific IgG2a production (Ref. 70). ES-62 was also found to suppress LPS-induced proinflammatory cytokine production by cells obtained from human synovial fluid and membranes (Ref. 68), and human cells are also susceptible to the anti-inflammatory action of PC conjugated to ovalbumin.

Turning to allergy, ES-62 was found to be active in two mouse models – ovalbumin-induced airway hypersensitivity and immediate-type




Figure 2. Filarial nematode-derived ES-62 induces hypo-responsiveness of mast cells by disrupting Fc ϵ RI signalling. Antigen (Ag)-driven ligation of IgE bound to Fc ϵ RI induces mast cell degranulation and generation of pro-inflammatory mediators and cytokines by triggering a signalling cascade involving protein kinase C- α (PKC- α) complexing with phospholipase D (PLD) to stimulate PLD-coupled, sphingosine kinase (SK)-mediated, calcium mobilisation. ES-62 directly blocks such sequestration and trafficking of PKC- α for perinuclear degradation by a caveolae/lipid raft, proteasomal-independent mechanism. Such sequestration of PKC- α uncouples Fc ϵ RI-mediated mast cell responses. Abbreviations: ES, excretory-secretory product; PA, phosphatidic acid; S-1-P, sphingosine-1-phosphate; TLR, toll-like receptor.

hypersensitivity to oxazolone in the skin (Ref. 69). In the former, ES-62 was found to reduce peribronchial inflammation and mucosal hyperplasia, inhibit eosinophilia and prevent release of IL-4, the signature cytokine required for airway inflammation development. In the latter model, a reduction in ear swelling indicated that ES-62 targeted inflammation. Subsequent in vitro analysis revealed that ES-62 directly prevented mast cell degranulation and release of mediators of allergy induced via ligation of FcεRI. The mechanism of action involved inhibition of key signal transduction events involved in mast cell activation, including phospholipase-D-coupled, sphingosine-kinase-mediated calcium mobilisation and NF-κB activation. It was found that ES-62 mediates these effects by forming a complex with TLR4, which results in the sequestration and perinuclear degradation of protein kinase C (PKC)-α, a molecule found to be critical for mast-cell activation (Ref. 65), by a caveolae or lipid raft proteosomal-independent mechanism (Fig. 2).

The finding that ES-62 can inhibit components of a Th2 response, such as IL-4 production (Ref. 69), indicates that the original hypothesis where ES-62 was solely a Th2-polarising agent (Ref. 44) was too simplistic. It is possible in fact that a target of ES-62 may perhaps be a cytokine, such as IL-17, which appears to act as a master regulator of both Th1- and Th2-type inflammation (Ref. 71) and indeed plays a critical role in CIA (Ref. 72), the first model of inflammation in which ES-62 was found to be protective.

Currently, work is underway to develop molecules that mimic the activity of ES-62 for use as drugs for treating allergic and autoimmune diseases. Unlike the parent molecule, these will be too small to be immunogenic and given its importance in promoting ES-62 activity, will be based around PC, but possibly with slight structural modifications to avoid interaction with natural anti-PC antibodies.

Clinical implications/applications

The weight of evidence in support of parasitic nematodes preventing allergy and autoimmunity not surprisingly led scientists to consider their therapeutic potential. Within the past few years therefore, parasitic nematodes have been used in clinical trials aimed at ameliorating inflammatory diseases. It has been possible to carry out these

trials because of the known or predicted lack of pathogenicity of certain species.

An example of a 'safe' species is *Trichuris suis*, which is similar to the human whipworm *T. trichiura*, but can persist for only a limited period in the human gastrointestinal tract following infection. These factors have dictated that it has been selected for testing in human trials as a treatment for inflammatory bowel disease. Encouragingly, in a series of trials, the majority of people suffering from inflammatory bowel disease have shown improvement in symptoms following ingestion of embryonated ova of this worm (reviewed in Refs 14, 30).

N. americanus has also been employed in trials, although because this human helminth has pathogenic potential, low numbers must be used to treat people. When infecting patients suffering from Crohn's disease with 50 larvae, those with moderately active (but not inactive or mild) disease showed some improvement (Ref. 73). A trial determining the effect of hookworms on asthma has also recently started (Ref. 17).

The ultimate aim in this area of research is to use defined nematode molecules (or derivatives thereof) in treating human disease. As proof that this can work, one need look no further than the recombinant nematode anticoagulant protein c2 (rNAPc2), a serine protease inhibitor derived from the original hookworm (*Ancylostoma caninum*) protein (Ref. 74). The anticoagulant activity of this protein has been found to inhibit thrombin generation in patients undergoing elective percutaneous coronary intervention (Ref. 75), to reduce the likelihood of deep venous thrombosis occurring during total knee arthroplasty (Ref. 76) and to reduce new thrombin generation and ischaemia in patients with non-ST-segment elevation acute coronary syndrome (Ref. 77). The anticoagulant activity of the protein is due to inhibition of the catalytic complex of subendothelial tissue factor/activated factor VII by a unique mechanism (see Ref. 77). It is to be hoped that nematode-derived molecules and/or their derivatives will soon also be used in the treatment of human allergic and autoimmune diseases.

Acknowledgements and funding

The authors would like to thank the Wellcome Trust, the Medical Research Council and the Biotechnology and Biological Sciences Research Council for funding their work cited in this review.

References

- 1 Schmidt, G.S. and Roberts, L.S. (1989) Foundations of Parasitology (4th edn), pp. 379-405, McGraw-Hill
- 2 Schmidt, G.S. and Roberts, L.S. (1989) Foundations of Parasitology (4th edn), pp. 488-503, McGraw-Hill
- 3 Schmidt, G.S. and Roberts, L.S. (1989) Foundations of Parasitology (4th edn), pp. 451-463, McGraw-Hill
- 4 Schmidt, G.S. and Roberts, L.S. (1989) Foundations of Parasitology (4th edn), pp. 416-427, McGraw-Hill
- 5 Schmidt, G.S. and Roberts, L.S. (1989) Foundations of Parasitology (4th edn), pp. 434-450, McGraw-Hill
- 6 Subramanian, S. et al. (2004) The dynamics of *Wuchereria bancrofti* infection: a model-based analysis of longitudinal data from Pondicherry, India. *Parasitology* 128, 467-482
- 7 Nutman, T.B. and Kumaraswami, V. (2001) Regulation of the immune response in lymphatic filariasis: perspectives on acute and chronic infection with *Wuchereria bancrofti* in South India. *Parasit Immunol* 28, 389-399
- 8 Nakamura, T. et al. (1997) Polarization of IL-4 and IFN- γ -producing CD4⁺ T cells following activation of naïve CD4⁺ T cells. *J Immunol* 158, 1085-1094
- 9 Moore, K.W. et al. (2001) Interleukin 10 and the interleukin 10 receptor. *Ann Rev Immunol* 19, 683-765
- 10 Rajan, T.V. (2005) Natural course of lymphatic filariasis: insights from epidemiology, experimental human infections and clinical observations. *Am J Trop Med Hyg* 285, 995-998
- 11 Bradley, J.E. and Jackson, J.A. (2004) Immunity, immunoregulation and the ecology of trichuriasis and ascariasis. *Parasit Immunol* 26, 429-441
- 12 van Riet, E., Hartgers, F.C. and Yazdanbakhsh, M. (2007) Chronic helminth infections induce immunomodulation: consequences and mechanisms. *Immunobiol* 212, 475-490
- 13 Wills-Karp, M., Santeliz, J. and Karp, C.L. (2001) The germless theory of allergic disease: revisiting the hygiene hypothesis. *Nat Rev Immunol* 1, 69-75
- 14 Elliott, D.E., Summers, R.W. and Weinstock, J.V. (2007) Helminths as governors of immune-mediated inflammation. *Int J Parasitol* 37, 457-464
- 15 Busse, W.W. and Lemanske, R.F. (2001) Asthma. *N Engl J Med* 344, 350-362
- 16 Cooper, P.J. (2004) Intestinal worms and human allergy. *Parasit Immunol* 26, 455-469
- 17 Falcone, F.H. and Pritchard, D.I. (2005) Parasite role reversal: worms on trial. *TRENDS Parasitol* 21, 157-160
- 18 Maizels, R.M. (2005) infections and allergy – helminths, hygiene and host immune regulation. *Curr Opin Immunol* 17, 656-661
- 19 Hagel, I. et al. (1993) Allergic reactivity of children of different socioeconomic levels in tropical populations. *Int Arch Allergy Immunol* 101, 209-214
- 20 Lynch, N.R. et al. (1993) Effect of anthelmintic treatment on the allergic reactivity of children in a tropical slum. *J Allergy Clin Immunol* 92, 404-411
- 21 Selassie, P.G. et al. (2000) Total and specific IgE (house dust mite and intestinal helminths) in asthmatics and controls from Gondar, Ethiopia. *Clin Exp Allergy* 30, 356-358
- 22 Nyan, O.A. et al. (2001) Atopy, intestinal helminth infection and total serum IgE in rural and urban adult Gambian communities. *Clin Exp Allergy* 31, 1672-1678
- 23 Scrivener, S. et al. (2001) Independent effects of intestinal parasite infection and domestic allergen exposure on the risk of wheeze in Ethiopia: a nested case-control study. *Lancet* 358, 1493-1499
- 24 Cooper, P.J. et al. (2003) Reduced risk of atopy among school age children infected with geohelminth parasites in a rural area of the tropics. *J Allergy Clin Immunol* 111, 995-1000
- 25 Dagoye, D. et al. (2003) Wheezing, allergy and parasite infection in children in urban and rural Ethiopia. *Am J Respir Crit Care Med* 167, 1369-1373
- 26 van den Biggelaar, A.H.J. et al. (2004) Long-term treatment of intestinal helminths increases mite skin-test reactivity in Gabonese schoolchildren. *J Infect Dis* 189, 892-900
- 27 Flohr, C. et al. (2006) Poor sanitation and helminth infection protect against skin sensitization in Vietnamese children: a cross-sectional study. *J Allergy Clin Immunol* 118, 1305-1311
- 28 Hunter, M.M. and McKay, D.M. (2004) Helminths as therapeutic agents for inflammatory bowel disease. *Aliment Pharmacol Ther* 19, 167-177
- 29 Zacccone, P. et al. (2006) Parasitic worms and inflammatory diseases. *Parasit Immunol* 28, 515-525
- 30 Reddy, A. and Fried, B. (2007) The use of *Trichuris suis* and other helminth therapies to treat Crohn's disease. *Parasitol Res* 100, 921-927
- 31 Aoyama, H. et al. (2007) An inverse relationship between autoimmune liver diseases and

- Strongyloides stercoralis* infection. Am J Trop Med Hyg 76, 972-976
- 32 Wang, C.C. et al. (2001) Infection of mice with the helminth *Strongyloides stercoralis* suppresses pulmonary allergic responses to ovalbumin. Clin Expl Allergy 31, 495-503
 - 33 Bashir, M.E. et al. (2002) An enteric helminth infection protects against an allergic response to dietary antigen. J Immunol 169, 3284-3292
 - 34 Wohlleben, G. et al. (2004) Helminth infection modulates the development of allergen-induced airway inflammation. Int Immunol 16, 585-596
 - 35 Wilson, M.S. et al. (2005) Suppression of allergic airway inflammation by helminth-induced regulatory T cells. J Exp Med 202, 1199-1212
 - 36 Kitagaki, K. et al. (2006) Intestinal helminths protect in a murine model of asthma. J Immunol 177, 1628-1635
 - 37 Elliott, D.E. et al. (2000) Does the failure to acquire helminth parasites predispose to Crohn's disease? FASEB J 14, 1848-1855
 - 38 Khan, W.I. et al. (2002) intestinal nematode infection ameliorates experimental colitis in mice. Infect Immun 70, 5931-5937
 - 39 Elliott, D.E. et al. (2004) *Heligomosomoides polygyrus* inhibits established colitis in IL-10-deficient mice. Eur J Immunol 34, 2690-2698
 - 40 Saunders, K.A. et al. (2007) Inhibition of autoimmune Type 1 diabetes by gastrointestinal helminth infection. Infect Immun 75, 397-407
 - 41 Finkelman, F. et al. (1997) Cytokine regulation of host defence against parasitic gastrointestinal nematodes: lessons from studies with rodent models. Ann Rev Immunol 15, 503-533
 - 42 Diaz, A. and Allen, J.E. (2007) Mapping immune response profiles: the emerging scenario from helminth immunology. Eur J Immunol 37, 3319-3326
 - 43 Banchereau, J. et al. (2000) Immunobiology of dendritic cells. Ann Rev Immunol 18, 767-811
 - 44 Whelan, M. et al. (2000) A filarial nematode secreted product signals dendritic cells to acquire a phenotype that drives development of Th2 cells. J Immunol 164, 6453-6460
 - 45 Balic, A. et al. (2004) Selective maturation of dendritic cells by *Nippostrongylus brasiliensis*-secreted proteins drives Th2 immune responses. Eur J Immunol 34, 3047-3059
 - 46 Harnett, W. and Harnett, M.M. (2006) Molecular basis of worm-induced immunomodulation. Parasit Immunol 28, 535-543
 - 47 Gause, W.C., Urban, J.F.Jr. and Stadecker, M.J. (2003) The immune response to parasitic helminths: insights from murine models. Trends Immunol 24, 269-277
 - 48 Scales, H.E. et al. (2004) Effect of inducible costimulator blockade on the pathological and protective immune responses induced by the gastrointestinal helminth *Trichinella spiralis*. Eur J Immunol 34, 2854-2862
 - 49 Liblau, R.S., Singer, S.M. and McDevitt, H.O. (1995) TH1 and TH2 CD4+ T cells in the pathogenesis of organ-specific autoimmune diseases. Imm Today 16, 34-38
 - 50 Taylor, M.D. et al. (2007) CTLA-4 and CD4+ CD25+ regulatory T cells inhibit protective immunity to filarial parasites *in vivo*. J Immunol 179, 4626-4634
 - 51 Carvalho, E.M., Bastos, L.S. and Araujo, M.I. (2006) Worms and Allergy. Parasit Immunol 28, 525-534
 - 52 Metwali, A. et al. (2006) Induction of CD8+ regulatory T cells in the intestine by *Heligomosomoides polygyrus* infection. Am J Physiol Gastrointest Liver Physiol 291, G253-G259
 - 53 Ince, M.N. et al. (2006) *Heligomosomoides polygyrus* induces TLR4 on murine mucosal T cells that produce TGFbeta after lipopolysaccharide stimulation. J Immunol 176, 726-729
 - 54 Erb, K.J. (2007) Helminths, allergic disorders and IgE-mediated immune responses: where do we stand? Eur J Immunol 37, 1170-1173
 - 55 Pochanke, V. et al. (2007) Identification and characterisation of a novel antigen from the nematode *Nippostrongylus brasiliensis* recognised by specific IgE. Eur J Immunol 37, 1275-1284
 - 56 Mitre, E., Norwood, S. and Nutman, T.B. (2005) Saturation of immunoglobulin E (IgE) binding sites by polyclonal IgE does not explain the protective effect of helminth infections against atopy. Infect Immun 73, 4106-4111
 - 57 Hussain, R., Poindexter, R.W. and Ottesen, E.A. (1992) Control of allergic reactivity in human filariasis: predominant localization of blocking antibody to the IgG4 subclass. J Immunol 148, 2731-2737.
 - 58 Royer, B. et al. (2001) Inhibition of IgE-induced activation of human mast cells by IL-10. Clin Exp Allerg 31, 694-704
 - 59 Kennedy Norton, S. et al. (2008) IL-10 suppresses mast cell IgE receptor expression and signalling *in vitro* and *in vivo*. J Immunol 180, 2848-2854

- 60 Kashyap, M. et al. (2008) CD4 T cell-mast cell interactions alter IgE receptor expression and signalling. *J Immunol* 180, 2039-2043
- 61 Trujillo-Vargas, C.M. et al. (2007) Helminth derived products inhibit the development of allergic responses in mice. *Am J Respir Crit Care Med* 175, 336-344
- 62 Keir, P.A. et al. (2004) Inhibition of neutrophil recruitment by ES of *nippostrongylus brasiliensis*. *Parasit Immunol* 26, 137-140
- 63 Lima, C. et al. (2002) Eosinophilic inflammation and airway hyper-responsiveness are profoundly inhibited by a helminth (*Ascaris suum*) extract in a murine model of asthma. *Clin Exp Allergy* 32, 1659-1666
- 64 Itami, D.M. et al. (2005) Modulation of murine experimental asthma by *Ascaris suum* components. *Clin Exp Allergy* 38, 873-879
- 65 Oshiro, T.M., Macedo, M.S. and Macedo-Soares, M.F. (2005) Anti-inflammatory activity of PAS-1, a protein component of *Ascaris suum*. *Inflamm Res* 54, 17-21
- 66 Imai, S. and Fujita, K. (2004) Molecules of parasites as immunomodulatory drugs. *Curr Top Med Chem* 4, 539-560
- 67 Harnett, W. and Harnett, M.M. (2006) Filarial nematode secreted product ES-62 is an anti-inflammatory agent: therapeutic potential of small molecule derivatives and ES-62 peptide mimetics. *Clin Exp Phar Phys* 33, 511-518
- 68 McInnes, I.B. et al. (2003) A novel therapeutic approach targeting articular inflammation using the filarial nematode-derived phosphorylcholine-containing glycoprotein ES-62. *J Immunol* 171, 2127-2133
- 69 Melendez, A.J. et al. (2007) Inhibition of FcεRI-mediated mast cell responses by ES-62, a product of parasitic filarial nematodes. *Nat Med* 13, 1375-1381
- 70 Harnett, M.M. et al. (2007) The phosphorylcholine moiety of the filarial nematode immunomodulator ES-62 is responsible for its anti-inflammatory action in arthritis. *Ann Rheum Dis* 67, 518-523
- 71 Kollis, J.K. and Linden, A. (2004) Interleukin 17 family members and inflammation. *Immunity* 21, 467-478
- 72 Nakae, S., Nambu, A., Sudo, K. and Iwakura, Y. (2003) Suppression of immune induction of collagen-induced arthritis in IL-17-deficient mice. *J Immunol* 171, 6173-6177
- 73 Croese, J. et al. (2006) A proof of concept study establishing *Necator americanus* in Crohn's patients and reservoir donors. *Gut* 55, 136-137
- 74 Stassens, P. et al. (1996) Anticoagulant repertoire of the hookworm *Ancylostoma caninum*. *Proc Natl Acad Sci U S A* 93, 2149-2154
- 75 Moons, A.H. et al. (2003) Recombinant nematode anticoagulant protein c2, an inhibitor of the tissue factor/factor VIIa complex, in patients undergoing elective coronary angioplasty. *J Am Coll Cardiol* 41, 2147-2153
- 76 Lee, A. et al. (2001) Dose-response study of recombinant factor VIIa/tissue factor inhibitor recombinant nematode anti-coagulant protein c2 in prevention of postoperative venous thromboembolism in patients undergoing total knee replacement. *Circulation* 104, 74-78
- 77 Giugliano, R.P. et al. (2007) Recombinant nematode anticoagulant protein c2 in patients with non-ST-segment elevation acute coronary syndrome. *J Am Coll Cardiol* 49, 2398-2407

Further reading, resources and contacts

Publications

Schmidt, G.S. and Roberts, L.S. (2004) *Foundations of Parasitology* (7th edn), McGraw-Hill
This is a useful reference book for all aspects of nematode parasitology.

Websites

The nematode pages of the Cambridge University Schistosomiasis Research Group website provide background information on the various species:

http://www.path.cam.ac.uk/~schisto/General_Parasitology/Hm.helminths.html

The Wikipedia site on helminthic therapy offers a good summary and reference to some other related links:

http://en.wikipedia.org/wiki/Helminthic_therapy

Features associated with this article

Figures

Figure 1. Potential mechanisms of nematode-induced immunomodulation and suppression of inflammatory disease.

Figure 2. Filarial nematode-derived ES-62 induces hyporesponsiveness of mast cells by disrupting FcεRI signalling.

Tables

Table 1. Estimated prevalence of the major medically important human nematode parasites.

Table 2. Nematode infection protects humans from allergic disease.

Table 3. Nematode infection can protect mice from allergic and autoimmune diseases.

Citation details for this article

William Harnett and Margaret M. Harnett (2008) Therapeutic immunomodulators from nematode parasites. Expert Rev. Mol. Med. Vol. 10, e18, June 2008, doi:10.1017/S1462399408000720