



NLR4 inflammasomopathies

Neil Romberg^a, Tiphany P. Vogel^b, and Scott W. Canna^c

Purpose of review

The purpose of the review is to highlight developments in autoinflammatory diseases associated with gain-of-function mutations in the gene encoding NLR-family CARD-containing protein 4 (NLRC4), the NLRC4-inflammasomopathies.

Recent findings

Three years since the identification of the first autoinflammation with infantile enterocolitis (AIFEC) patients, there is an improved understanding of how the NLRC4 inflammasome and interleukin 18 (IL-18) contribute to gut inflammation in myeloid and also intestinal epithelial cells. This information has opened new therapeutic avenues to treat AIFEC patients with targeted agents like recombinant IL-18 binding protein and anti-interferon- γ antibodies. Additional phenotypes traditionally associated with NLRP3 mutations like familial cold autoinflammatory syndrome and neonatal onset multisystem inflammatory disease (NOMID), have now also been associated with gain-of-function *NLRC4* mutations. Finally, *NLRC4* somatic mosaicism has now been identified in a NOMID and an AIFEC patient, a finding emphasizing nontraditional modes of inheritance in autoinflammatory diseases.

Summary

The NLRC4 inflammasomopathies constitute a growing autoinflammatory disease category that spans a broad clinical spectrum from cold urticaria to NOMID and the often fatal disease AIFEC. Rapid case identification with biomarkers like elevated serum IL-18 concentrations and early intervention with targeted immunomodulatory therapies are key strategies to improving outcomes for AIFEC patients.

Keywords

autoinflammation with infantile enterocolitis, FCAS4, IL-18, NLR-family CARD domain-containing protein 4, neonatal onset multisystem inflammatory disease

INTRODUCTION

An inflammasome is a pro-inflammatory cytoplasmic structure formed upon detection of exogenous pathogen-associated or danger-associated molecular patterns (PAMPs/DAMPs). The inflammasomopathies describe a group of mechanistically related diseases, each, the result of inappropriate inflammasome activation. Inflammasomopathies can be protean disorders; for instance, activating mutations in the gene encoding NLR family pyrin domain containing 3 (NLRP3) may present with cold urticaria, or progressive hearing loss or neonatal onset multisystem inflammatory disease (NOMID) [1].

In 2014, two groups independently identified gain-of-function mutations in the gene encoding NLR-family caspase activation and recruitment domain (CARD)-containing protein 4 (NLRC4) in four patients displaying recurrent, life-threatening episodes of autoinflammation and infantile enterocolitis (AIFEC) [2,3]. Since that time, the identification of additional AIFEC patients and their

successful treatment with targeted therapies have revealed important insights into AIFEC pathophysiology. Also, new reports have described additional clinical phenotypes associated with novel germline or somatic *NLRC4* mutations. Herein, we summarize the growing body of literature describing *NLRC4* mutation-associated autoinflammatory diseases, the NLRC4 inflammasomopathies.

^aDivision of Immunology and Allergy, The Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, ^bDivision of Immunology, Allergy and Rheumatology, Department of Pediatrics, Baylor College of Medicine and Center for Human Immunobiology, Texas Children's Hospital, Houston, Texas and ^cRK Mellon Institute for Pediatric Research/Pediatric Rheumatology, Children's Hospital of Pittsburgh of UPMC, Pittsburgh, Pennsylvania, USA

Correspondence to Scott W. Canna, RK Mellon Institute for Pediatric Research/Pediatric Rheumatology, Children's Hospital of Pittsburgh of UPMC, 8124 Rangos Research Bldg, 4401 Penn Avenue, Pittsburgh, PA 15224, USA. E-mail: scott.canna@chp.edu

Curr Opin Allergy Clin Immunol 2017, 17:398–404

DOI:10.1097/ACI.0000000000000396

KEY POINTS

- The NLRC4 inflammasomopathies are caused by gain-of-function mutations in the gene encoding the innate immune protein NLRC4. Mutant NLRC4 promotes spontaneous formation of the NLRC4 inflammasome, production of IL1 family cytokines (IL-1 β and IL-18) and inflammatory cell death.
- Three NLRC4 inflammasomopathy phenotypes have been described: AIFEC, NOMID and FCAS4. AIFEC is associated with very early onset enterocolitis, macrophage activation syndrome and neonatal death.
- Rapid case identification using disease biomarkers like elevated serum IL-18 concentrations, characteristic skin biopsy findings and ultimately diagnostic gene sequencing, combined with timely application of immune modulatory therapy are key strategies to improving AIFEC patient outcomes.

NLRC4 INFLAMMASOME BIOLOGY

There are several canonical inflammasomes, and each is organized similarly: cytosolic PAMP/DAMP detectors are linked via an adaptor protein, apoptosis-associated speck-like protein containing a CARD (ASC), to the cleaved, active form of pro-caspase-1 [4]. Upon activation, the inflammasome rapidly forms a large wheel-shaped structure [5], exhausting cellular ASC stores. Inflammasome formation initiates pyroptosis, a form of inflammatory cell death [6] and also proteolytically activates pro-interleukin 1 family cytokines (IL-1 β and IL-18) into their cleaved, active forms [7–9].

Inflammasome identity and specificity are determined by eponymous detector proteins, which include: absent in melanoma 2 (AIM2) [10], NLRP3 [11] and NLRC4 [12]. Like the NLRP3 inflammasome, which responds to numerous cytosolic DAMPs/PAMPs [13–15], the human NLRC4 inflammasome recognizes at least two bacterial ligands, flagellin and the type three secretion system (T3SS) [16]. NLRC4 is distinct from NLRP3 and AIM2 because it does not directly interact with its ligands. Instead, NLRC4 is activated via contact with the sensor protein NLR family of apoptosis inhibitory protein (NAIP), and it is NAIP that physically binds either flagellin or a T3SS [17]. This arrangement suggests NLRC4 might be better categorized as a scaffolding protein rather than a PAMP detector, although NLRC4 might also be considered an adaptor. Unlike NLRP3 and AIM2, NLRC4 contains a CARD allowing it to directly contact pro-caspase-1 without ASC [18]. Notably, absent ASC, the NLRC4 inflammasome is functionally altered favoring pyroptosis over cytokine production [3,16].

NLRC4 inflammasome biology has primarily been studied in myeloid cells including circulating monocytes and neutrophils but as NLRC4 detects components of lung and gut-trophic pathogens, its behavior in mucosal tissues is also of vital interest. Recently, a specialized host defense role of the NLRC4 inflammasome was identified in mouse intestinal epithelial cells (IECs). Upon detection of *Salmonella* spp. within IEC cytoplasm, the NLRC4 inflammasome rapidly forms producing IL-18 and diarrhea-causing eicosanoids [19^{***}]. Instead of pyroptosis, *Salmonella* spp. containing IECs undergo IL-18-independent, caspase-dependent, nonlytic cell death with subsequent expulsion into the colonic lumen [20]. Although this adaptation produces secretory diarrhea, vascular leak and shock, it likely prevents catastrophic, invasive bacterial infections.

NLRC4 inflammasome initiation is exquisitely sensitive; a single ligand-bound NAIP molecule is sufficient to propagate NLRC4 oligomerization [21], yet as systemic inflammation impacts host survival, the process is highly regulated. One level of regulation occurs intrinsically through the autoinhibitory structure of the NLRC4 molecule [22]. NLRC4 consists of a CARD, a ligand binding and NAIP-interacting leucine-rich repeat (LRR) and a regulatory nucleotide-binding oligomerization domain (NOD) (Fig. 1) [23]. Within the NOD, helical domain 1 (HD1), winged helix domain (WHD) and helical domain 2 (HD2) form a specialized adenosine diphosphate (ADP)-binding pocket that stabilizes NLRC4 in its inactive conformation [12,22,24]. Upon LRR detection of ligand-bound NAIP, the NOD undergoes a conformational change that promotes ADP for adenosine triphosphate (ATP) exchange, NLRC4 oligomerization and inflammasome assembly [22]. A second regulatory layer controls cytokine production and is inflammasome-extrinsic, as production of inactive pro-IL-1 family cytokines requires signals from cell surface PAMP/DAMP receptors including tumor necrosis factor (TNF) receptor 1 and toll-like receptor 4 (TLR4) [25]. Physiologically inert, these pro-cytokines accumulate intracellularly until an inflammasome cleaves them into immunologically active compounds.

There are 11 IL-1 cytokine family members; only two, IL-1 β and IL-18, are synthesized as inactive precursors [25]. IL-1 β mediates the classic signs and symptoms of the febrile response. The IL-1 receptor binds to either IL-1 β or its endogenously produced competitive inhibitor, IL-1 receptor agonist (IL-1RA) [26,27]. Canonically, IL-18 amplifies lymphocyte functions, increasing cytotoxicity through IFN γ production. Circulating IL-18 is




FIGURE 1. Disease-associated NLRC4 variants cluster near the ADP-binding site. (a) The functional domains of NLRC4 are depicted and color-coded: CARD domain black; NBD = purple; HD-1 = pink; WHD = yellow; HD2 = green; LRR = gray. Variants associated with predominant skin phenotypes are marked by the symbol *, the NOMID-associated variant is marked by ‡, and AIFEC mutations by #. (b) A color-coded NLRC4 crystal structure without a CARD is displayed as per Ref. [22]. ADP (red) is depicted within its binding pocket (inset, right). Disease-associated residues (white) are labeled.

sequestered by an endogenous-binding protein (IL-18BP), preventing receptor binding [25]. The full effects of IL-18 are incompletely understood, but recent work suggests IL-18 equilibrium controls intestinal barrier function in mice and that excessive free IL-18 potentiates colitis [28^{••}].

AUTOINFLAMMATION WITH INFANTILE ENTEROCOLITIS

In 2014, two investigative groups simultaneously and independently reported gain-of-function *NLRC4* mutations in four patients from two unrelated families [2,3]. The disease, called by one group as NLRC4 macrophage activation syndrome (NLRC4-MAS) and by the other as syndrome of enterocolitis and autoinflammation associated with mutation of *NLRC4* (SCAN4), was later designated autoinflammation with infantile enterocolitis (AIFEC) by Online Mendelian Inheritance in Man (OMIM, #616050, *606831) [29]. As the syndromes originally described by each team were remarkably similar and shared a common cause, we will use the umbrella term AIFEC, in this review, to avoid confusion and build consensus.

AIFEC is a chronic inflammatory disease punctuated by episodes of extreme acuity. All seven

reported AIFEC patients have required hospitalizations in intensive care units during inflammatory episodes. Two infantile cases have been fatal. AIFEC flares have been appropriately compared with MAS; both entities share signature IL-1 β symptomology (fever, tachycardia) and interferon gamma (IFN γ)-related histopathology (hemophagocytosis). Peripheral blood transcriptional profiles generated from two AIFEC patients in flare suggest MAS-like myeloid cell activation and cytotoxic T-cell dysfunction [2,30^{••}]. In its most extreme form, an AIFEC flare can be confused with primary hemophagocytic lymphohistiocytosis (HLH), with hypertriglyceridemia, coagulopathy, multilineage cytopenias, elevated soluble IL-2 receptor and poor in-vitro cytotoxicity [2,3,30^{••},31[•],32[•]]. Importantly, cytotoxic function normalizes in AIFEC patients after flares resolve, indicating intact granule-dependent cytotoxicity machinery. Another feature and useful biomarker distinguishing AIFEC from primary HLH is extremely elevated serum IL-18 concentrations (more than 10⁴ pg/ml). IL-18 elevations of this degree are described in only a small cadre of inflammatory diseases including AIFEC, X-linked inhibitor of apoptosis (XIAP) deficiency and systemic juvenile idiopathic arthritis/adult-onset Stills disease-related MAS [33,34].

Table 1. Patients reported with *NLRC4* mutation-associated diseases

Variant	Number of patients	Origin of mutation	Age of onset	Phenotype	Disease-related mortality	References
p.S171F	One	Somatic mosaicism	In utero	AIFEC	Deceased at 2 months	[31 [■]]
p.T177A	One	Somatic mosaicism	Birth	NOMID	Alive at 19 years	[37]
p.T337S	One	De novo	6 Months	AIFEC	Alive at 7 years	[2]
p.T337N	One	De novo	3 Weeks	AIFEC	Alive at 8 months	[32 [■]]
p.V341A	One	De novo	6 Weeks	AIFEC	Alive at 1 year	[30 [■]]
p.V341A	Three in one family	De novo, then inherited	1–2 Weeks	AIFEC	Deceased at 23 days (pro-band) or alive at 7 and 46 years	[3]
p.H443P	13 in one family	Inherited	2–3 Months	FCAS4	Alive into adulthood	[38]
p.S445P	13 in one family	Inherited	infancy to childhood	FCAS-like	Alive into adulthood	[39 [■]]

AIFEC, autoinflammation with infantile enterocolitis; FCAS, familial cold autoinflammatory syndrome; NOMID, neonatal onset multisystem inflammatory disease.

The trigger for AIFEC flares is unknown. At least two older AIFEC patients experienced flares with physical and emotional stressors [3]. Another possible trigger could be activation of the mutant *NLRC4* inflammasome with flagellin or T3SS expressing bacteria. *In vitro*, AIFEC macrophages infected with T3SS-expressing *Salmonella* spp. or *Pseudomonas* spp. form numerous altered assemblies of inflammasome components that favor pyroptosis over cytokine production [3]. Nonetheless, such bacteria have not yet been recovered from flaring AIFEC patients.

One area where MAS and AIFEC do not overlap is gastrointestinal disease. Diarrhea is an uncommon MAS symptom whereas severe, secretory, neonatal diarrhea is the most common AIFEC presentation. In one case, there is evidence AIFEC enterocolitis began in utero [31[■]]. Intestinal biopsies and autopsy specimens from all reported AIFEC patients consistently show a mixed inflammatory infiltrate, villous flattening with tissue edema, epithelial erosions and tissue autolysis [2,3,30[■],35]. Activated macrophages have been visualized in some [31[■]] but not all AIFEC intestinal tissue samples [2,3,30[■]], raising the possibility that mutant *NLRC4* in IECs, not myeloid cells, is the primary driver of gut disease. Because of feeding intolerance, many AIFEC patients temporarily require parenteral nutrition, including the oldest known AIFEC patient, now 46 years old. His course was first described by Shwachman and coworkers in a 1973 case series of 16 infants with undifferentiated, protracted diarrhea supported by this nutritive method [36]. For this and other surviving AIFEC patients, symptomatic enterocolitis spontaneously normalized after the first year of life even though serum IL-18 concentrations remain persistently elevated into adulthood.

In addition to enterocolitis and MAS-like episodes, AIFEC patients can develop numerous

organ-specific symptoms secondary to lymphohistiocytic inflammation. For instance, ephemeral maculopapular and urticarial rashes associated with lymphohistiocytic infiltrates can occur (unpublished data). Astute clinicians may elect to biopsy and stain skin lesions in suspected AIFEC cases for rapid diagnostic information, as this histologic finding is unusual. Similarly, central nervous system dysfunction (obtundation, seizures) and hepatic dysfunction correlate with activated macrophages in related tissues [3,31[■]].

AIFEC is caused by heterozygous *NLRC4* gain-of-function mutations, which exhibit complete penetrance. There are now four published AIFEC-associated *NLRC4* mutations: V341A, T337S, T337N and S171F (Table 1 and Fig. 1). The V341A variant is described in four individuals from two unrelated pedigrees, whereas the remaining variants occurred in single cases. Interestingly, the AIFEC patient harboring the S171F variant was mosaic for this allele. Although the variant accounted for only a quarter of *NLRC4* peripheral blood transcripts, the patient's in-utero disease onset and fatal outcome suggest it to be highly pathogenic. Amino acid positions 337 and 341 are both located in the autoinhibitory HD-1 subdomain of NOD. *NLRC4* crystal structure analysis suggests that hydrophobic residues at position 341 are important for closing the 'lid' on the ADP-binding pocket to prevent ADP–ATP exchange [3,22]. Similarly, position 337 may stabilize the pocket's tertiary structure through interactions with residues 170 and 173 or 173 only [2,22]. Presumably, substitution of phenylalanine for serine at position 171 alters these interactions [31[■]].

Although infantile diarrhea and MAS-like episodes are the signature features of AIFEC, it is a chronic inflammatory disease. Patients surviving infancy are of short stature and exhibit anemia of chronic disease [3]. Between flares, untreated AIFEC

patients display moderately elevated acute phase reactants and highly elevated serum IL-18 concentrations [2,3]. Amyloidosis, a long-term complication of many untreated autoinflammatory disorders, has not yet been described in AIFEC patients but could be a concern. Longitudinal observation of surviving AIFEC pediatric cases and identification of more adult AIFEC cases may be informative.

EXTENDED NLRC4 PHENOTYPES

Shortly after publication of the original AIFEC cases, 13 members of a Japanese pedigree were reported with a syndrome of neonatal-onset fever, cold-induced urticarial rash and arthralgias [38]. A heterozygous NLRC4 H443P variant segregated with disease. Symptoms were sufficiently mild that most affected members did not require treatment. Serum IL-18 concentrations were not assessed, but patient cells exposed to cold *in vitro* spontaneously produced IL-1 β . Given the clear phenotypic similarities to the NLRP3-associated familial cold autoinflammatory syndrome (FCAS1), OMIM designated this disease as FCAS4 (#616115; *606831) [29]. In 2017, a second large Dutch kindred was reported with prominent skin manifestations [39[¶]]. A heterozygous S445P NLRC4 variant segregated with disease. Like FCAS4 patients, these patients developed inflammatory symptoms early in life that did not significantly alter long-term survival; the eldest affected member was age 88 years. Uniquely, in these kindred, most patients developed conjunctivitis or uveitis, and many had a nodular and urticarial rash. Although affected individuals lacked MAS or infantile enterocolitis, two developed intestinal inflammation in adulthood. Functional studies of the S445P variant were not conducted, but many individuals displayed highly elevated serum IL-18 concentrations. Unlike FCAS1 patients whose biopsied skin lesions are characteristically neutrophilic [40], skin infiltrates in Dutch patients' biopsies were lymphohistiocytic as seen in AIFEC [39[¶]].

In addition to the traditional patterns of disease inheritance, two patients have been described with NLRC4 somatic mosaicism. One, discussed above, presented with prenatal AIFEC [31[¶]], the other presented with cardinal NOMID features (fever, rash, inflammatory bone lesions, sensory neural hearing loss and structural brain defects) but he was NLPR3 mutation-negative and displayed chronically elevated serum IL-18 [37]. Functional analysis of patient pluripotent stem cell-derived monocytes revealed two distinct populations, one with aberrant and the other with normal IL-1 β secretion. The aberrant IL-1 β producing cells harbored a novel

T177A NLRC4 variant. Notably, this mutation was initially missed on whole exome sequencing analysis.

It is unclear how gain-of-function mutations in the same gene can produce such disparate clinical phenotypes, but a genotype/phenotype relationship may exist. The FCAS phenotype appears to correlate with WHD subdomain variants (H443P and S445P), whereas AIFEC and NOMID associate with HD-1 and NBD subdomain variants (S171F, T177A, T337S, T337N and V341A) (Fig. 1). Recent functional work suggests pathologic NLRC4 variants in WHD may differentially promote caspase 8-mediated cell death whereas variants in HD-1 do not [41].

TREATMENT OF NLRC4 INFLAMMASOMOPATHIES

Although all NLRC4 inflammasomopathies are categorically autoinflammatory disorders, they manifest across a broad severity spectrum. Without a NLRC4-specific therapy, treating physicians have chosen to target downstream inflammatory mediators based upon their patients' clinical needs. For instance, many patients with FCAS4 were well controlled with only nonsteroidal anti-inflammatory drugs or nothing at all [39[¶]]. Other non-AIFEC patients were treated with recombinant IL-1RA (anakinra) and although skin manifestations were completely responsive, other disease features were not. Anakinra was also highly effective in treating NOMID-like symptoms because of NLRC4 mutation [37]. Used prophylactically, anakinra reduced the severity and frequency of MAS episodes more effectively than low-dose steroids and colchicine in one AIFEC patient with mild gastrointestinal disease [2], but was not efficacious treating another AIFEC patient in flare [30[¶]].

The most therapeutically challenging AIFEC manifestations are MAS and enterocolitis. Sadly, NLRC4 mutations were found posthumously in several patients with overwhelming neonatal AIFEC, and their aggressive presentation and rapid progression clouded assessment of treatment efficacy [3,31[¶]]. MAS episodes were managed successfully with corticosteroids, cyclosporine and IVIg in at least one adult patient prior to his AIFEC diagnosis [3]. Notably, this same patient has experienced prolonged periods of treatment-free quiescence between life-threatening MAS flares.

Several agents specifically targeting deranged AIFEC immunologic pathways are under development. For example, we observed dramatic efficacy using recombinant IL-18BP in a critically ill neonatal AIFEC patient whose disease was refractory to combined corticosteroids, cyclosporine, IL-1

inhibition, TNF-inhibition and integrin-inhibition [30^{''}]. The infant's clinical improvement corresponded with a precipitous drop in free but not total IL-18 emphasizing the importance of endogenous IL-18BP in humans. This case prompted broader evaluation of IL-18BP in an ongoing clinical trial (NCT03113760). Blockade of IFN γ was also efficacious in one AIFEC flare [32^{*}].

A growing literature links the intestinal ecosystem with NLRC4 activation and IL-18 production [19^{''},20,28^{''},42]. As such, early gut colonization may promote excessive IL-1 family cytokine production in AIFEC patients. Likewise, spontaneous resolution of AIFEC enterocolitis may coincide with maturation of the gut, mucosal immunity and/or the adoption of a less inflammatory microbiota [42]. Without more data, we cannot recommend either intestinal decontamination or fecal transplantation in AIFEC patients. Similarly, as NLRC4 is expressed in both myeloid and intestinal epithelia cells, we would not anticipate hematopoietic stem cell transplantation by itself or intestinal transplantation by itself, would be curative.

CONCLUSION

The NLRC4 inflammasomopathies constitute a growing category of autoinflammatory diseases that span a broad clinical spectrum from cold urticaria to NOMID and the often-fatal disease AIFEC. Since 2014, 34 patients have been reported with *NLRC4* gain-of-function mutation-associated diseases. Of these, most are from two unrelated families with *NLRC4* variants that do not appear to confer a significant survival disadvantage.

There are seven published case studies of AIFEC patients; two were fatal cases and all the patients experienced significant morbidity due, in part, to diagnostic delay. Even now that it has been established as a distinct clinical entity, AIFEC continues to pose a diagnostic challenge because most cases are sporadic. Accordingly, early recognition of tell-tale AIFEC symptoms and a rapid diagnosis, using disease biomarkers like serum IL-18 concentrations, characteristic skin biopsy findings and ultimately gene sequencing, will be paramount to improving disease outcomes. Likewise, the timely implementation of anti-inflammatory therapies, either already approved (anakinra, steroids, IVIg, cyclosporine) or available through ongoing clinical trials (recombinant IL-18BP; NCT03113760 or anti-IFN γ monoclonal antibodies; NCT02069899), will continue to be a key determinate of survival.

Acknowledgements

None.

Financial support and sponsorship

The authors would like to thank their funding sources including the Jeffrey Modell Foundation (N.R.) and the RK Mellon Foundation (S.W.C.).

Conflicts of interest

S.W.C. is a paid consultant for AB2Bio, Ltd and T.P.V. attended their clinical trial investigator meeting.

REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Hoffman HM, Mueller JL, Broide DH, *et al.* Mutation of a new gene encoding a putative pyrin-like protein causes familial cold autoinflammatory syndrome and Muckle-Wells syndrome. *Nat Genet* 2001; 29:301–305.
 2. Canna SW, de Jesus AA, Gouni S, *et al.* An activating NLRC4 inflammasome mutation causes autoinflammation with recurrent macrophage activation syndrome. *Nat Genet* 2014; 46:1140–1146.
 3. Romberg N, Al Moussawi K, Nelson-Williams C, *et al.* Mutation of NLRC4 causes a syndrome of enterocolitis and autoinflammation. *Nat Genet* 2014; 46:1135–1139.
 4. Broz P, Dixit VM. Inflammasomes: mechanism of assembly, regulation and signalling. *Nat Rev Immunol* 2016; 16:407–420.
 5. Cheng J, Waite AL, Tkaczyk ER, *et al.* Kinetic properties of ASC protein aggregation in epithelial cells. *J Cell Physiol* 2010; 222:738–747.
 6. Brennan MA, Cookson BT. Salmonella induces macrophage death by caspase-1-dependent necrosis. *Mol Microbiol* 2000; 38:31–40.
 7. Ghayur T, Banerjee S, Hugunin M, *et al.* Caspase-1 processes IFN- γ -inducing factor and regulates LPS-induced IFN- γ production. *Nature* 1997; 386:619–623.
 8. Li P, Allen H, Banerjee S, *et al.* Mice deficient in IL-1 β -converting enzyme are defective in production of mature IL-1 β and resistant to endotoxic shock. *Cell* 1995; 80:401–411.
 9. Kuida K, Lippke JA, Ku G, *et al.* Altered cytokine export and apoptosis in mice deficient in interleukin-1 β converting enzyme. *Science* 1995; 267:2000–2003.
 10. Fernandes-Alnemri T, Yu J-W, Datta P, *et al.* AIM2 activates the inflammasome and cell death in response to cytoplasmic DNA. *Nature* 2009; 458:509–513.
 11. Agostini L, Martinon F, Burns K, *et al.* NALP3 forms an IL-1 β -processing inflammasome with increased activity in Muckle-Wells autoinflammatory disorder. *Immunity* 2004; 20:319–325.
 12. Poyet JL, Srinivasula SM, Tnani M, *et al.* Identification of Ipaf, a human caspase-1-activating protein related to Apaf-1. *J Biol Chem* 2001; 276:28309–28313.
 13. Mariathasan S, Weiss DS, Newton K, *et al.* Cryopyrin activates the inflammasome in response to toxins and ATP. *Nature* 2006; 440:228–232.
 14. Martinon F, Pétrilli V, Mayor A, *et al.* Gout-associated uric acid crystals activate the NALP3 inflammasome. *Nature* 2006; 440:237–241.
 15. Cassel SL, Eisenbarth SC, Iyer SS, *et al.* The Nalp3 inflammasome is essential for the development of silicosis. *Proc Natl Acad Sci USA* 2008; 105:9035–9040.
 16. Sutterwala FS, Mijares LA, Li L, *et al.* Immune recognition of *Pseudomonas aeruginosa* mediated by the IPAF/NLRC4 inflammasome. *J Exp Med* 2007; 204:3235–3245.
 17. Kofoed EM, Vance RE. Innate immune recognition of bacterial ligands by NALPs determines inflammasome specificity. *Nature* 2011; 477:592–595.
 18. Mariathasan S, Newton K, Monack DM, *et al.* Differential activation of the inflammasome by caspase-1 adaptors ASC and Ipaf. *Nature* 2004; 430:213–218.
 19. Rauch I, Deets KA, Ji DX, *et al.* NAIP-NLRC4 inflammasomes coordinate intestinal epithelial cell expulsion with eicosanoid and IL-18 release via activation of caspase-1 and -8. *Immunity* 2017; 46:649–659.
- This report emphasizes a unique role for the NLRC4 inflammasome in intestinal epithelial cells, both for preventing invasive bacterial infections and for driving intestinal and systemic immunopathology.
20. Sellin ME, Müller AA, Felmy B, *et al.* Epithelium-intrinsic NAIP/NLRC4 inflammasome drives infected enterocyte expulsion to restrict *Salmonella* replication in the intestinal mucosa. *Cell Host Microbe* 2014; 16:237–248.
 21. Hu Z, Zhou Q, Zhang C, *et al.* Structural and biochemical basis for induced self-propagation of NLRC4. *Science* 2015; 350:399–404.
 22. Hu Z, Yan C, Liu P, *et al.* Crystal structure of NLRC4 reveals its autoinhibition mechanism. *Science* 2013; 341:172–175.

23. Damiano JS, Stehlik C, Pio F, *et al.* CLAN, a novel human CED-4-like gene. *Genomics* 2001; 75:77–83.
24. Danot O, Marquenet E, Vidal-Ingiglieri D, Richet E. Wheel of life, wheel of death: a mechanistic insight into signaling by STAND proteins. *Struct Lond Engl* 2009; 17:172–182.
25. Dinarello CA. Immunological and inflammatory functions of the interleukin-1 family. *Annu Rev Immunol* 2009; 27:519–550.
26. Carter DB, Deibel MR, Dunn CJ, *et al.* Purification, cloning, expression and biological characterization of an interleukin-1 receptor antagonist protein. *Nature* 1990; 344:633–638.
27. Sims JE, Acres RB, Grubin CE, *et al.* Cloning the interleukin 1 receptor from human T cells. *Proc Natl Acad Sci USA* 1989; 86:8946–8950.
28. Nowarski R, Jackson R, Gagliani N, *et al.* Epithelial IL-18 equilibrium controls ■ barrier function in colitis. *Cell* 2015; 163:1444–1456.
- This report establishes the role of free IL-18 and IL-18BP deficiency in murine intestinal inflammation.
29. Online Mendelian Inheritance in Man, OMIM®. Baltimore, MD: Johns Hopkins University [MIM number: 616050: 05/16/2017. World Wide Web URL: <https://omim.org/>.
30. Canna SW, Girard C, Malle L, *et al.* Life-threatening NLRC4-associated ■ hyperinflammation successfully treated with IL-18 inhibition. *J Allergy Clin Immunol* 2017; 139:1698–1701.
- This report is the first description of recombinant IL-18BP as a treatment for AIFEC, highlighting the utility of free IL-18 and differences between intestinal and systemic IL-18 regulation.
31. Liang J, Alfano DN, Squires JE, *et al.* Novel NLRC4 mutation causes a ■ syndrome of perinatal autoinflammation with hemophagocytic lymphohistiocytosis, hepatosplenomegaly, fetal thrombotic vasculopathy, and congenital anemia and ascites. *Pediatr Dev Pathol Off J Soc Pediatr Pathol Paediatr Pathol Soc* 2017. [Epub ahead of print]
- This report describes a fatal AIFEC case, an infant with somatic mosaicism for a novel NLRC4 variant. The authors report systemic inflammation likely commenced in utero.
32. Bracaglia C, Gatto A, Pardeo M, *et al.* Anti interferon-gamma (IFNγ) mono- clonal antibody treatment in a patient carrying an NLRC4 mutation and severe hemophagocytic lymphohistiocytosis. *Pediatr Rheumatol* 2015; 13:O68.
- This abstract describes AIFEC in an infant with a novel NLRC4 variant. It also is the first description of treating AIFEC with anti-IFNγ monoclonal antibodies.
33. Maeno N, Takei S, Nomura Y, *et al.* Highly elevated serum levels of interleukin-18 in systemic juvenile idiopathic arthritis but not in other juvenile idiopathic arthritis subtypes or in Kawasaki disease: comment on the article by Kawashima *et al.* *Arthritis Rheum* 2002; 46:2539–2541.
34. Wada T, Kanegane H, Ohta K, *et al.* Sustained elevation of serum interleukin-18 and its association with hemophagocytic lymphohistiocytosis in XIAP deficiency. *Cytokine* 2014; 65:74–78.
35. Shwachman H, Lloyd-Still JD, Khaw K-T, Antonowicz I. Protracted diarrhea of infancy treated by intravenous alimentation: II. Studies of small intestinal biopsy results. *Am J Dis Child* 1973; 125:365–368.
36. Lloyd-Still JD, Shwachman H, Filler RM. Protracted diarrhea of infancy treated by intravenous alimentation: I. Clinical studies of 16 infants. *Am J Dis Child* 1973; 125:358–364.
37. Kawasaki Y, Oda H, Ito J, *et al.* Identification of a high-frequency somatic NLRC4 mutation as a cause of autoinflammation by pluripotent cell-based phenotype dissection. *Arthritis Rheumatol* 2017; 69:447–459.
38. Kitamura A, Sasaki Y, Abe T, *et al.* An inherited mutation in *NLRC4* causes autoinflammation in human and mice. *J Exp Med* 2014; 211: 2385–2396.
39. Volker-Touw CML, de Koning HD, Giltay JC, *et al.* Erythematous nodes, ■ urticarial rash and arthralgias in a large pedigree with NLRC4-related auto-inflammatory disease, expansion of the phenotype. *Br J Dermatol* 2017; 176:244–248.
- This report describes a large, and phenotypically diverse, Dutch kindred including novel features like urticarial and nodular rashes, conjunctivitis, arthritis and IBD. Skin biopsies displayed a lymphohistiocytic infiltrate distinct from FCAS1 and similar to AIFEC.
40. Kolivras A, Theunis A, Ferster A, *et al.* Cryopyrin-associated periodic syn- drome: an autoinflammatory disease manifested as neutrophilic urticarial dermatosis with additional perieccrine involvement. *J Cutan Pathol* 2011; 38:202–208.
41. Raghawan AK, Sripada A, Gopinath G, *et al.* A disease-associated mutant of NLRC4 shows enhanced interaction with *SUG1* leading to constitutive FADD dependent caspase-8 activation and cell death. *J Biol Chem* 2017; 292:1218–1230.
42. Chudnovskiy A, Mortha A, Kana V, *et al.* Host-protozoan interactions protect from mucosal infections through activation of the inflammasome. *Cell* 2016; 167:444–456; e14.