

Changing prevalence of wheeze, rhinitis and allergic sensitisation in late childhood: findings from 2 Isle of Wight birth cohorts 12 years apart

V. K. Patil^{1,2}, R. J. Kurukulaaratchy¹, C. Venter^{1,3}, J. Grundy¹, G. Roberts^{1,4,5}, T. Dean^{1,3} and S. H. Arshad^{1,4,5}

¹The David Hide Asthma and Allergy Research Centre, St Mary's Hospital, Newport Isle of Wight, ²MSc Allergy Student, Faculty of Medicine, University of Southampton, UK, ³School of Health Sciences and Social Work, University of Portsmouth, Portsmouth, ⁴NIHR Respiratory Biomedical Research Unit, University Hospital Southampton NHS Foundation Trust, Southampton, UK, and ⁵Clinical and Experimental Medicine, Faculty of Medicine, University of Southampton, UK.

Clinical & Experimental Allergy

Summary

Background While the prevalence of asthma in children is decreasing or remaining the same, time trends in the prevalence of rhinitis in children are not known. Understanding sensitisation trends may help inform about trends in asthma and rhinitis prevalence.

Objective To assess time trends of wheeze, rhinitis and aero-allergen sensitisation prevalence at 10 years of age, we compared two birth cohorts established 12 years apart. To gain insight into differences in disease prevalence, we assessed association of family history, early life exposures and sensitisation with wheeze and rhinitis in each cohort.

Methods The IoW (Isle of Wight) and FAIR (Food Allergy and Intolerance Research) unselected birth cohorts were established in 1989 and 2001 respectively in IoW. Identical ISAAC questionnaire and skin prick test data were collected and compared at 10 years of age.

Results Over the 12-year period from 2001 to 2012, prevalence of lifetime wheeze, current wheeze and those ever treated for asthma decreased by 15.9% (45.5 vs. 29.6, $P < 0.001$), 3.9% (18.9 vs. 15, $P = 0.020$) and 8.2% (31.7 vs. 23.5, $P = 0.001$), respectively. Conversely, current rhinitis and lifetime rhinitis prevalence increased by 5.5% (22.6 vs. 28.1, $P = 0.004$) and 13% (18.6 vs. 31.7, $P < 0.001$), respectively. Atopic status remained stable; however, house dust mite (HDM) sensitisation decreased by 5.6% (19.2 vs. 13.6, $P = 0.004$) and grass sensitisation increased by 3.5% (12.9 vs. 16.4, $P = 0.054$). Male sex, parental history of asthma and HDM sensitisation were significantly associated with lifetime wheeze in both cohorts, while maternal smoking during pregnancy was a significant risk factor only in the earlier IoW cohort. Parental history of rhinitis and grass sensitisation was significantly associated with lifetime rhinitis in both cohorts, while HDM sensitisation was significant only for the IoW cohort.

Conclusion Contrasting changes were noted with falling wheeze and HDM sensitisation but rising rhinitis and grass sensitisation prevalence. Changing prevalence of aero-allergen sensitisations may explain the different time trends observed in these cohorts.

Keywords allergic rhinitis, asthma prevalence, hayfever prevalence, sensitisation prevalence, time trends in UK

Submitted 4 June 2014; revised 26 January 2015; accepted 17 February 2015

Correspondence:

Professor S Hasan Arshad, MP 810, South Block, Southampton General Hospital, Tremona Road, Southampton, United Kingdom.

E-mail: sha@soton.ac.uk

Cite this as: V. K. Patil, R. J.

Kurukulaaratchy, C. Venter, J. Grundy, G. Roberts, T. Dean and S. H. Arshad, *Clinical & Experimental Allergy*, 2015 (45) 1430–1438.

Introduction

Asthma is the most common chronic disease affecting children [1], and rhinitis remains a significant health problem affecting quality of life [2]. There is global variation in asthma prevalence, with the point prevalence being higher in more affluent regions compared

to less affluent regions [3, 4]. Globally asthma and rhinitis prevalence increased in the latter part of the 20th century [5]. However, global time trend studies in the last decade showed that while childhood asthma prevalence continued to increase in less affluent regions with a low point prevalence, it was reaching a plateau in more affluent regions with higher point

prevalence [6]. The United Kingdom (UK) has one of the highest asthma prevalence globally [4]. In UK, asthma and rhinitis prevalence increased from the 1970s to mid-1990s [5, 7]. However, time trends in asthma prevalence during the first decade of the 21st century are unclear, with some studies showing increase and others showing plateauing or decrease in asthma prevalence in schoolchildren [7–11].

It has been challenging to identify mechanisms driving changes in prevalence of allergic airway disease [12]. Asthma is a complex disease where interaction of genetics and environment play an important role [13]. The temporal changes in asthma prevalence are too quick to be explained by naturally occurring genetic alterations in populations [6]. Allergic sensitisation to aero-allergens plays an important role; house dust mite (HDM) sensitisation is strongly associated with asthma [14–16]. Recent European studies of time trends in asthma prevalence have found that similar to asthma prevalence, aero-allergen sensitisation has remained stable from 1991 to 2001–02 in schoolchildren [17, 18]. Similarly, allergic sensitisation is a strong risk factor for rhinitis [19], and it is known that grass pollen sensitisation is associated with rhinitis in the UK [20].

Meaningful analyses of time trends in changing prevalence of allergic disease require studies employing the same methodology and in the same geographical location over time. Arshad *et al.* have shown that HDM sensitisation is an important risk factor for asthma with an odds ratio of 8.07 (4.60–14.14) and sensitisation to grass was found to be an important risk factor for rhinitis with an odds ratio of 5.02 (2.21–11.41). We have previously reported the prevalence of current wheeze in the Isle of Wight (IoW) birth cohort at 10 years as 18.9% in 1999–2000 with associations to allergic sensitisation and sex [16]. We hypothesise that (i) the prevalence of wheeze and rhinitis will increase over a 12-year period and (ii) the increase in wheeze and rhinitis will be associated with a change in the same direction for house dust mite and grass pollen sensitisation, respectively. Trends in prevalence of wheeze and rhinitis in UK have been reported till 2005; this study looks at the continued trends till 2012 and also looks at time trends of aero-allergen sensitisations which have not been reported before in UK population. To investigate this, we compared current wheeze, lifetime wheeze, ever treated for asthma, current rhinitis symptoms, lifetime rhinitis and prevalence of allergic sensitisation to common aero-allergens at 10 years of age in two unselected population-based birth cohorts established 12 years apart in the same county, the Isle of Wight, United Kingdom. Results in this study are based on data of follow-ups at 10 years of age of the IoW birth cohort [21] (1999–

2000) and Food Allergy and Intolerance Research (FAIR) birth cohort [22] (2011–12).

Methods

Study population/birth cohorts

This is a repeat cross-sectional study performed in two separate cohorts of children aged 10 years, resident in the same locality and separated by 12 years. The Isle of Wight birth cohort (IoW cohort) is a population-based birth cohort established in 1989/90 for prospective study of asthma and allergic diseases. Children born between January 1989 and February 1990 on the Isle of Wight were recruited at birth ($N = 1536$). Children have been reviewed at 1, 2, 4, 10 and 18 years [23]. The ten-year follow-up was carried out between 1999 and 2000. Food Allergy and Intolerance Research (FAIR) is another population-based birth cohort ($N = 969$) established on the Isle of Wight in 2001/02 for prospective study of allergic diseases [22]. Children were seen at 1, 2, 3 [24] and 10 year. The 10-year follow-up was completed between 2011 and 2012. Table 1 gives a general description of both cohorts at their 10-year follow-ups. Ethics approval was obtained at each follow-ups for both cohorts by local research ethics committees (FAIR: 10/H0504/11, IoW: No. 18/98), and informed consent was obtained from parents and assent from participants.

Questionnaires

ISAAC (International Study of Asthma and Allergies in Childhood) and study-specific questionnaires were used in both studies. Prevalence rates are based on parental responses to ISAAC questionnaires [25] used at 10-year follow-ups in both cohorts. Lifetime wheeze symptoms were based on response to 'Has your child ever had wheezing or whistling in the chest at any time in the

Table 1. Descriptions of IoW and FAIR cohorts and 10-year follow-up rates

	IoW cohort	FAIR cohort
Birth year	1989–1990	2001–2002
Recruited at birth	1456	969
Male: Female	53%: 47%	51%: 49%
Period of 10-year FU	1999–2000	2011–2012
Questionnaire completed at 10 year	1373 (94%)	827 (85%)
SPT performed at 10 year	1036 (71%)	588 (61%)

FU, follow-up; SPT, skin prick test; IoW, Isle of Wight; FAIR, Food Allergy and Intolerance Research.

past?’ and current wheeze was based on ‘Has your child had wheezing or whistling in the chest in the last 12 months?’ and ever treated for asthma was based on ‘Has your child been treated for asthma?’ Current rhinitis symptoms and lifetime rhinitis were derived from the responses to ‘In the past 12 months, has your child had a problem with sneezing, or a runny or blocked nose when he/she did not have a cold or the flu?’ and ‘Has your child ever had hayfever?’ respectively. Risk factors such as family history of asthma or rhinitis, prenatal smoking and pet exposures were prospectively collected.

Sensitisation

Allergic sensitisation was defined by positive skin prick test (SPT) indicated by a wheal size of 3 mm or more than the negative control (saline). Sensitisation to aero-allergens that were tested in both studies was compared; HDM (*Dermatophagoides pteronyssinus*), cat (*Felis domesticus*) and grass (mixed grasses) pollen. SPT was performed using standardised allergen reagents and methodology (ALK-Abellø, Hørsholm, Denmark) [16, 22] and by the same research team.

Statistical analysis

The prevalence of wheeze, rhinitis and allergen sensitisation is given in proportions with 95% confidence intervals (95% CI). The significance of difference in prevalence between the two cohorts was tested using chi-square tests. Logistic regression analyses (univariate and multivariate) were used to test the association of

sex, family history, prenatal exposures and aero-allergen sensitisation with prevalence of wheeze and rhinitis. Backward stepwise model was used for multivariate regression analysis. Analyses were carried out using SPSS version 19 (IBM, Chicago, IL, USA). CIA (Confidence Interval Analysis software) was used for 95% CI for proportions and the difference in prevalence. *P* value of < 0.05 was considered significant.

Results

Study population

Of the 1456 children available for follow-up in the IoW cohort, 1373 (94%) provided questionnaire-based information and 1036 (71%) underwent SPT at 10 years of age in 1999–2000. In the FAIR follow-up at 10 years of age in 2011–12, 827 (85%) of 969 available children responded to the questionnaire-based information and 588 (61%) underwent SPT. Characteristics of the cohorts are displayed in Table 1. The subsets who underwent SPT in both cohorts were representative of the respective cohorts (Table S1).

Trend in prevalence of wheeze, rhinitis and sensitisation

Prevalence of lifetime wheeze, current wheeze and ever treated asthma all decreased significantly from 1999–2000 to 2011–12. Conversely, current rhinitis symptoms and lifetime rhinitis both increased significantly between these two time points (Table 2).

Changes in prevalence for individual aero-allergen sensitisation varied according to allergen (Table 3). HDM

Table 2. Time trends in wheeze, asthma and rhinitis prevalence

Characteristic	IoW 1999–2000 % (95% CI) <i>n/N</i>	FAIR 2011–12 % (95% CI) <i>n/N</i>	Difference % (95% CI)	<i>P</i> value
Lifetime Wheeze	45.5% (42.9 to 48.2) 625/1373	29.6% (26.6 to 32.8) 245/828	–15.9 (–19.9 to –11.8)	< 0.001
Current Wheeze	18.9% (16.9 to 21.0) 259/1373	15% (12.7 to 17.6) 124/827	–3.9 (–7 to –0.6)	0.020
Ever treated for asthma	31.7% (29.2 to 34.3) 402/1267	23.5% (20.1 to 27.4) 120/510	–8.2 (–12.5 to –3.6)	0.001
Current rhinitis	22.6% (20.5 to 24.9) 308/1362	28.1% (25.1 to 31.2) 232/826	5.5 (1.7 to 9.3)	0.004
Lifetime rhinitis	18.6% (16.7 to 20.8) 256/1373	31.7% (26.9 to 36.8) 107/338	13.0 (7.8 to 18.5)	< 0.001

n, proportion with feature; *N*, total number with responses to question.

sensitisation decreased significantly, while grass pollen sensitisation increased (not significant) and cat sensitisation remained unchanged during the same period.

There was no difference in time trends of wheeze and rhinitis prevalence between IoW and FAIR cohorts when stratified by sex (Table S2). The only difference noticed was in the increase in grass sensitisation from IoW to FAIR cohort which was higher in girls compared to boys. Interactions of sex by cohorts were not significant for both lifetime wheeze and rhinitis (Table S3).

Family history and prenatal risk factors

Overall, parental history of asthma was significantly higher in the FAIR cohort compared to the IoW cohort by 5.2% ($P = 0.012$), which was significant for maternal (4.6%, $P = 0.009$) but not for paternal (2.5%, $P = 0.118$) history. However, sibling history of asthma significantly decreased by 5.1% ($P = 0.017$). Parental history of rhinitis significantly increased by 14.1% ($P < 0.001$), which was significant for both maternal and paternal history of rhinitis. Sibling history of rhinitis showed non-significant increase of 1.7% ($P = 0.349$). Maternal smoking during pregnancy showed a drop by 2.8% comparing the two time points, but this was not significant, and there was no difference in prenatal pet exposure in the two cohorts (Table 4).

Association of sex, family history and prenatal factors and aero-allergen sensitisation to wheeze and rhinitis

Lifetime prevalence of wheeze. Male sex, parental history, sibling history, prenatal smoke exposure, HDM, grass and cat sensitisation were associated with wheeze

in both cohorts in univariate analysis. Using backward stepwise multivariate logistic regression, male sex ($P = 0.038$), parental history ($P < 0.001$), sibling history ($P = 0.008$), maternal smoking during pregnancy ($P = 0.018$) and HDM sensitisation ($P < 0.001$) remained independently significant in the IoW cohort, while sex ($P = 0.007$), parental history ($P < 0.001$) and HDM sensitisation ($P = 0.004$) remained independently significant in the FAIR cohort (Table 5). Therefore, sex, parental history and HDM sensitisation were the common factors significantly associated with lifetime wheeze in both cohorts.

Lifetime prevalence of rhinitis. Parental history and sensitisations to HDM, cat and grass were associated with lifetime rhinitis in both cohorts. Sibling history and prenatal maternal smoking in IoW and prenatal pet exposure in the FAIR cohort were also associated with lifetime rhinitis using a univariate regression analysis. Using a multivariate regression with backward stepwise analysis, parental history ($P < 0.001$), HDM ($P < 0.001$) and grass ($P < 0.001$) sensitisation remained significant in the IoW cohort, whereas only parental history of rhinitis ($P < 0.001$) and grass sensitisation ($P < 0.001$) remained significant in the FAIR cohort (Table 6). Thus, parental history of rhinitis and sensitisation to grass were common factors significantly associated with lifetime rhinitis in both cohorts.

Risk factors such as HDM sensitisation and parental history of asthma for lifetime wheeze and grass sensitisation and parental history of rhinitis for lifetime rhinitis were tested for interaction with cohorts. Only cohort and parental history of asthma were significant ($P = 0.045$) and all other interaction terms were not significant as summarised in Table S4.

Table 3. Time trends in prevalence of aero-allergen sensitisation

Aero-allergens	IoW 1999–2000 % (95% CI) n/N	FAIR 2011–2012 % (95% CI) n/N	Difference % (95% CI)	P value
HDM	19.2 (16.9 to 21.7) 199/1035	13.6 (11.1 to 16.6) 80/587	–5.6 (–9.2 to –1.8)	0.004
Cat	7.9 (6.4 to 9.7) 82/1035	7.7 (5.8 to 10.1) 45/586	–0.2 (–2.9 to 2.6)	0.861
Grass	12.9 (11.0 to 15.1) 134/1036	16.4 (13.6 to 19.6) 96/585	3.5 (–0.1 to 7.2)	0.054
Any aero-allergen	25.7 (23.1 to 28.4) 266/1035	24.1 (20.8 to 27.7) 141/587	–1.6 (–5.9 to 2.8)	0.512

n, number of positive SPT to the allergen; N, total number tested for the allergen.

Table 4. Prevalence of family history and prenatal exposures in both cohorts (irrespective of asthma or rhinitis in children)

Factor	IoW 1999–2000 % (95% CI) n/N	FAIR 2011–12 % (95% CI) n/N	Difference % (95% CI)	P value
Mother Hx of asthma	17.7 (15.7 to 19.8) 233/1317	22.3 (19.6 to 25.3) 184/825	+4.6 (1.2 to 8.2)	0.009
Father Hx of asthma	13.9 (12.1 to 15.9) 183/1317	16.4 (14.0 to 19.1) 132/806	+2.5 (–0.6 to 5.7)	0.118
Parental Hx of asthma	30.4 (28.0 to 33.0) 401/1317	35.7 (32.5 to 39.0) 289/810	+5.3 (1.1 to 9.4)	0.012
Sibling Hx of asthma	31.4 (29.0 to 34.0) 414/1318	26.4 (23.3 to 29.7) 190/721	–5.0 (–9.1 to –0.9)	0.017
Mother Hx of rhinitis	24.3 (22.0 to 26.7) 320/1318	33.9 (30.8 to 37.2) 280/825	+9.6 (5.7 to 13.6)	< 0.001
Father Hx of rhinitis	16.4 (14.5 to 18.5) 216/1316	25.1 (22.2 to 28.2) 202/804	+8.7 (5.2 to 12.4)	< 0.001
Parental Hx of rhinitis	37.1 (34.5 to 39.7) 489/1318	51.2 (47.7 to 54.6) 415/811	+14.1 (9.7 to 18.4)	< 0.001
Sibling Hx of rhinitis	19.3 (17.3 to 21.6) 255/1318	21.1 (18.3 to 24.2) 152/721	+1.8 (–1.9 to 5.5)	0.349
Mother smoked during pregnancy	25.2 (23.1 to 27.5) 384/1521	22.4 (19.9 to 25.2) 210/937	–2.8 (–6.2 to 0.7)	0.111
Any pet during pregnancy	50.4 (47.9 to 52.9) 765/1517	50.4 (47.2 to 53.5) 488/969	0 (–4.1 to 4.0)	0.974

Hx; history, n; number with factor, N total number

Discussion

This study comparing upper and lower allergic airway disease symptom prevalence at 10 years of age in two population-based birth cohorts 12 years apart in the same location has shown falling prevalence of current and lifetime wheeze plus ever being treated for asthma, associated with HDM sensitisation which decreased during the same period. By contrast in the same period from 1999–2000 to 2011–12, there was an increase in current and lifetime rhinitis prevalence, which was associated with grass sensitisation which increased during the same period. Male sex, parental history of asthma and HDM sensitisation for wheeze and parental history of rhinitis and grass sensitisation remained significant risk factors in both cohorts for asthma and rhinitis, respectively. Maternal smoking in pregnancy was significantly associated with lifetime wheeze only in the earlier IOW cohort.

Time trend analysis can be challenging due to a number of factors including lack of uniformity in meth-

odology and definitions of outcomes. We used identical validated questions (ISAAC) and the same methodology [25] which makes prevalence in our 2 cohorts comparable. Moreover, both cohorts in this study are population based, from the same geographical location and assessed by the same research team, thus restricting the plausible biases and the influence of environmental differences when comparing the prevalence from wider geographical regions. Both cohorts have shown high follow-up rates, and the risk of significant bias is relatively low. Other important aero-allergens such as tree pollen and common moulds are missing in this study which can be one of the limitations of this paper, but the three aero-allergens studied usually explain majority of the atopic status [26]. Another potential limitation is the generalisability of the prevalence rates estimated on the IoW. Although it remains a theoretical possibility, the population on the IoW is neither socially, not geographically, isolated. The prevalence rates for allergic diseases we have reported previously

Table 5. Logistic regression analyses for association of sex, aero-allergen sensitisations, heredity and prenatal exposures with lifetime wheeze

		IoW Cohort (1999–2000)					FAIR Cohort (2011–2012)					
		Univariate		Multivariate					Univariate		Multivariate	
	% <i>n</i> / <i>N</i>	OR (95% CI)	<i>P</i>	OR (95% CI)	<i>P</i>	%	OR (95% CI)	<i>P</i>	OR (95% CI)	<i>P</i>		
<i>Lifetime wheeze</i>												
Male sex	50.2 350/697	1.47 (1.19–1.82)	< 0.001	1.32 (1.02–1.72)	0.038	32.9 140/425	1.39 (1.03–1.88)	0.030	1.81 (1.18–2.45)	0.007		
HDM Sensitisation	67.8 135/199	2.76 (1.99–3.83)	< 0.001	2.28 (1.58–3.31)	< 0.001	56.3 45/80	3.84 (2.36–6.23)	< 0.001	2.61 (1.36–5.03)	0.004		
Grass Sensitisation	63.4 85/134	2.06 (1.42–3.00)	< 0.001	–	NS	41.7 40/95	1.95 (1.24–3.06)	0.004	–	NS		
Cat Sensitisation	68.3 56/82	2.50 (1.54–4.05)	< 0.001	1.93 (1.08–3.44)	0.026	50.0 22/44	2.631 (1.42–4.89)	0.002	–	NS		
Parental History	59.1 237/401	2.21 (1.74–2.81)	< 0.001	2.00 (1.51–2.65)	< 0.001	45.8 132/288	3.31 (2.42–4.54)	< 0.001	2.82 (1.83–4.36)	< 0.001		
Sibling history	54.1 224/414	1.66 (1.32–2.10)	< 0.001	1.47 (1.11–1.96)	0.008	41.1 78/190	2.08 (1.47–2.95)	< 0.001	–	NS		
Pet during pregnancy	45.5 317/697	1.01 (0.81–1.25)	0.944	–	NS	28.1 121/430	0.87 (0.64–1.17)	0.342	–	NS		
Maternal smoking*	52.7 168/319	1.46 2(1.14–1.88)	0.003	1.47 (1.11–1.96)	0.018	35.7 60/168	1.44 (1.00–2.06)	0.049	–	NS		

Multivariate; backward stepwise model. % – proportion of the characteristic with wheeze (n/N), OR, odds ratio; 95% CI, 95% confidence interval; HDM, house dust mite.

*Mother smoking during pregnancy.

are very similar to the developed world in general and United Kingdom in particular [16]. Therefore, the time trends seen in our cohorts should be broadly generalisable. Also results from other islands have been comparable to the reference region on the mainland [27].

Anderson *et al.* looked at ISAAC study results from UK centres from 1995 to 2002 in 12- to 14-year-old UK children and reported an increase in 'lifetime asthma' (from 20.6% to 25.9%), but decrease in current wheeze (from 33.9% to 27.5%) [9]. In the Aberdeen schools asthma survey of 9 to 12 year-old children, Malik *et al.* assessed the time trends at three time points (1999, 2004 and 2009). Current wheeze (in the past 3 years) steadily decreased from 1999 to 2009 (27.9% in 1999, 25.2% in 2004, 22.2% in 2009), while 'lifetime asthma' prevalence showed temporal increase (from 24.3% to 28.4%) between 1999 and 2004 and then decreased from 2004 to 2009 (22.1%) [28]. Our results in similar age children (10 years old) confirm the continued decrease in prevalence of wheeze in the United Kingdom till 2012. The decrease in current and lifetime wheeze prevalence in our study is further substantiated by a concurrent fall in the prevalence of ever treated for asthma. In contrast to wheeze prevalence, we observed a significant rise in the prevalence of current rhinitis symptoms and lifetime rhinitis. Similar increasing trends for rhinitis have been reported in the United Kingdom looking at the primary care data sets from 2001 to 2005 [29]; however, the Aberdeen schools asthma survey also showed a temporal

increase in prevalence of 'lifetime rhinitis' from 1999 to 2004 (15.4% to 26.5%) but not much change between 2004 and 2009 (25.7%) [11].

As reported by Zollner *et al.* in Germany and Braun-Fahrlander *et al.* in Switzerland, we also found stable overall aero-allergen sensitisation on the Isle of Wight. However, both studies reported mixed inhalant screen and differences in individual aero-allergen sensitisation were not investigated [17, 18]. We report a significant decrease in HDM sensitisation by 5.6% and a non-significant increase in grass sensitisation by 3.5%. In the Isle of Wight prevention study (different cohort from the 2 cohorts compared in this paper), reduction in the level of HDM was associated with reduced level of allergic sensitisation to HDM [30]. Based on this finding, it might be that falling HDM exposure levels may have contributed to the falling HDM sensitisation observed in our study. However, without HDM exposure information at two time points, this remains purely speculative but an important area for future research.

Allergic conditions are complex diseases where multiple environmental exposures, hereditary factors and gene–environment interactions play important roles [13]. One of the criticisms for this paper would be not looking at all the plausible factors such as decreasing birth rate, changes in family sizes, changes in living conditions and respiratory viral infections [12, 31]. Parental history is a known risk factor for developing allergic airway diseases [32, 33], and prenatal smoke exposure is associated with

Table 6. Logistic regression analysis for association of sex, aero-allergen sensitisations, heredity and prenatal exposures with lifetime rhinitis

	IoW Cohort (1999–2000)				FAIR Cohort (2011–2012)			
	Univariate		Multivariate		Univariate		Multivariate	
	%	OR (95% CI)	P		OR (95% CI)	P	OR (95% CI)	P
<i>Lifetime rhinitis</i>								
Male Sex	18.8 131/697	1.020 (0.78–1.34)	0.885	NS	1.38 (0.87–2.19)	0.178	–	NS
HDM Sensitisation	44.2 88/199	4.55 (3.25–6.39)	< 0.001	< 0.001	4.83 (2.74–8.51)	< 0.001	–	NS
Grass Sensitisation	64.9 87/134	11.51 (7.70–17.20)	< 0.001	< 0.001	18.79 (10.17–34.73)	< 0.001	15.89 (7.711–32.76)	< 0.001
Cat Sensitisation	56.1 46/82	6.06 (3.80–9.67)	< 0.001	NS	8.93 (4.03–19.76)	< 0.001	–	NS
Parental History	26.8 131/489	2.32 (1.75–3.07)	< 0.001	< 0.001	4.72 (2.72–8.20)	< 0.001	4.21 (2.01–8.80)	< 0.001
Sibling history	25.5 65/255	1.69 (1.22–2.34)	0.002	NS	1.26 (0.71–2.23)	0.428	–	NS
Pet during pregnancy	17.5 122/697	0.86 (0.66–1.13)	0.290	NS	0.59 (0.37–0.94)	0.028	–	NS
Maternal smoking*	14.1 45/319	1.52 (1.07–2.15)	0.020	NS	0.81 (0.45–1.49)	0.502	–	NS

Multivariate; backward stepwise model. % – proportion of the characteristic with rhinitis, OR, odds ratio; 95% CI, 95% confidence interval; HDM, house dust mite.

*Mother smoking during pregnancy.

the development of asthma [33]. Prenatal smoke exposure declined which could be due to a better awareness of harmful effects of smoking particularly during pregnancy, and the association analysis suggested declining effect of prenatal smoke exposure on wheeze and rhinitis between these two cohorts. Parental history remained significantly associated with wheeze and rhinitis in both cohorts. We have shown that the prevalence of wheeze decreased in the current generation (in the cohort studied) and the sibling history of asthma also decreased, whereas the parental history of asthma increased between the two cohorts, suggesting different time trends in prevalence in two generations. For mothers of IoW cohort, date of birth ranges from 1941 to 1973 (mean 1962) and for mothers of FAIR from 1956 to 1986 (mean 1973). It is plausible that the increase in the parent generation reflects increase in asthma prevalence observed in the latter half of 20th century.

HDM sensitisation is associated with lifetime wheeze and sensitisation to grass with lifetime rhinitis in both cohorts. Li *et al.* show that asthma prevalence continues to increase predominantly in HDM-sensitised urban Chinese children [34]. An association does not necessarily imply causality; however, one interpretation of our results is of a possible role of decreasing HDM and increasing grass sensitisations, driving the contrasting time trends of wheeze and rhinitis prevalence, respectively. Recently, a synergistic effect of family history and second-hand smoking has been shown for adult-onset asthma [35]. There is a possibility of synergic effect of decreasing HDM sensitisation, decreasing prenatal smoke exposure and family history of asthma on decreasing wheeze prevalence, and this provides an area for further research. We did observe a decrease in the maternal smoking during pregnancy from IoW to FAIR cohort, but this change was not significant, so the interpretations have to be made with caution.

In conclusion, this is the first study reporting contrasting time trends for childhood allergic airway diseases along with aero-allergen sensitisations in the same geographical location; falling wheeze and rising rhinitis associated with falling time trends of HDM sensitisation with rise in grass pollen sensitisation respectively. Changing associations with aero-allergen sensitisation and smoke exposure partly contribute to changing prevalence of wheeze and rhinitis.

Acknowledgements

We would like to thank all the participants and the parents for their continued support for both birth cohorts. Authors thank all the research nurses and study coordinators involved in both cohorts.

Statement of Contribution: VKP contributed in conceiving the idea for the paper, contributed to study

design and conduct, conducted data analysis and wrote the first draft of the manuscript. RJK contributed to study conduct, design and preparation of manuscript. CV contributed to study conduct and in manuscript preparation. JG contributed to study conduct; GR contributed to study design and manuscript preparation; TD contributed to the study design and manuscript preparation; SHA contributed in conceiving the idea for the paper, contributed to study design and manuscript preparation.

Funding: The FAIR Follow-up was funded by NIHR Postdoctoral Fellowship Award Ref. NIHR/DF/01/055. The IoW 10-year follow-up was funded by National Asthma Campaign, UK (Grant No 364, PI: S. Hasan Arshad).

Conflict of interest

The authors declare no conflict of interest.

References

- WHO. Asthma. 2012. Available from: <http://www.who.int/mediacentre/factsheets/fs307/en/index.html>. (Last accessed 12 October 2012)
- Canonica GW, Bousquet J, Mullol J, Scadding GK, Virchow JC. A survey of the burden of allergic rhinitis in Europe. *Allergy* 2007; 62(Suppl. 85):17–25.
- Lai CK, Beasley R, Crane J *et al.* Global variation in the prevalence and severity of asthma symptoms: phase three of the International Study of Asthma and Allergies in Childhood (ISAAC). *Thorax* 2009; 64:476–83.
- ISAAC Steering Committee. Worldwide variation in prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and atopic eczema: ISAAC. The International Study of Asthma and Allergies in Childhood (ISAAC) Steering Committee. *Lancet* 1998; 351:1225–32.
- Burr ML, Butland BK, King S, Vaughan-Williams E. Changes in asthma prevalence: two surveys 15 years apart. *Arch Dis Child* 1989; 64:1452–6.
- Asher MI, Montefort S, Björkstén B *et al.* Worldwide time trends in the prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and eczema in childhood: ISAAC Phases One and Three repeat multicountry cross-sectional surveys. *Lancet* 2006; 368:733–43.
- Gupta R, Sheikh A, Strachan DP, Anderson HR. Time trends in allergic disorders in the UK. *Thorax* 2007; 62:91–6.
- Burr ML, Wat D, Evans C, Dunstan FD, Doull IJ. British thoracic society research C. Asthma prevalence in 1973, 1988 and 2003. *Thorax* 2006; 61:296–9.
- Anderson HR, Ruggles R, Strachan DP *et al.* Trends in prevalence of symptoms of asthma, hay fever, and eczema in 12–14 year olds in the British Isles, 1995–2002: questionnaire survey. *BMJ* 2004; 328:1052–3.
- Shamssain M. Trends in the prevalence and severity of asthma, rhinitis and atopic eczema in 6- to 7- and 13- to 14-yr-old children from the north-east of England. *Pediatr Allergy Immunol* 2007; 18:149–53.
- Malik G, Tagiyeva N, Aucott L, McNeill G, Turner SW. Changing trends in asthma in 9–12 year olds between 1964 and 2009. *Arch Dis Child* 2011; 96:227–31.
- Butland BK, Strachan DP, Lewis S, Bynner J, Butler N, Britton J. Investigation into the increase in hay fever and eczema at age 16 observed between the 1958 and 1970 British birth cohorts. *BMJ* 1997; 315:717–21.
- Su MW, Tung KY, Liang PH, Tsai CH, Kuo NW, Lee YL. Gene-gene and gene-environmental interactions of childhood asthma: a multifactor dimension reduction approach. *PLoS ONE* 2012; 7:e30694.
- Lau S, Illi S, Sommerfeld C *et al.* Early exposure to house-dust mite and cat allergens and development of childhood asthma: a cohort study. *Lancet* 2000; 356:1392–7.
- Arshad SH, Tariq SM, Matthews S, Hakim E. Sensitization to common allergens and its association with allergic disorders at age 4 years: a whole population birth cohort study. *Pediatrics* 2001; 108:art. no. -e33.
- Kurukulaaratchy RJ, Fenn M, Twisleton R, Matthews S, Arshad SH. The prevalence of asthma and wheezing illnesses amongst 10-year-old schoolchildren. *Respir Med* 2002; 96:163–9.
- Zollner IK, Weiland SK, Piechowski I *et al.* No increase in the prevalence of asthma, allergies, and atopic sensitization among children in Germany: 1992–2001. *Thorax* 2005; 60:545–8.
- Braun-Fahrlander C, Gassner M, Grize L *et al.* No further increase in asthma, hay fever and atopic sensitisation in adolescents living in Switzerland. *Eur Respir J* 2004; 23:407–13.
- Kurukulaaratchy RJ, Karmaus W, Arshad SH. Sex and atopy influences on the natural history of rhinitis. *Curr Opin Allergy Clin Immunol* 2012; 12:7–12.
- Arshad SH, Karmaus W, Matthews S *et al.* Association of allergy-related symptoms with sensitisation to common allergens in an adult European population. *J Investig Allergol Clin Immunol* 2001; 11:94–102.
- Arshad SH, Hide DW. Effect of environmental factors on the development of allergic disorders in infancy. *J Allergy Clin Immunol* 1992; 90:235–41.
- Dean T, Venter C, Pereira B *et al.* Patterns of sensitization to food and aero-allergens in the first 3 years of life. *J Allergy Clin Immunol* 2007; 120:1166–71.
- Raza A, Kurukulaaratchy R, Grundy JD, Clayton CB *et al.* What does adolescent undiagnosed wheeze represent? Findings from the Isle of Wight Cohort. *Eur Respir J* 2012; 40:580–8.
- Venter C, Pereira B, Voigt K *et al.* Prevalence and cumulative incidence of food hypersensitivity in the first 3 years of life. *Allergy* 2008; 63:354–9.
- Lukrafka JL, Fuchs SC, Moreira LB, Picon RV, Fischer GB, Fuchs FD. Performance of the ISAAC questionnaire to establish the prevalence of asthma in adolescents: a population-based study. *J Asthma* 2010; 47:166–9.
- Arshad SH, Tariq SM, Matthews S, Hakim E. Sensitization to common allergens and its association with allergic disorders at age 4 years: a whole population birth cohort study. *Pediatrics* 2001; 108:E33.

- 27 Jeffs D, Grainger R, Powell P. Is childhood allergy more common amongst an island population? *J R Soc Promot Health* 2000; 120:236–41.
- 28 Panickar JR, Dodd SR, Smyth RL, Courriel JM. Trends in deaths from respiratory illness in children in England and Wales from 1968 to 2000. *Thorax* 2005; 60:1035–8.
- 29 Ghouri N, Hippisley-Cox J, Newton J, Sheikh A. Trends in the epidemiology and prescribing of medication for allergic rhinitis in England. *J R Soc Med* 2008; 101:466–72.
- 30 Scott M, Roberts G, Kurukulaaratchy RJ, Matthews S, Nove A, Arshad SH. Multifaceted allergen avoidance during infancy reduces asthma during childhood with the effect persisting until age 18 years. *Thorax* 2012; 67:1046–51.
- 31 Urquhart DS, Anderson AK, McKenzie SA. Fewer colds, less asthma? A hypothesis to explain the fall in childhood asthma in the UK. *J Epidemiol Community Health* 2008; 62:921–5.
- 32 Arshad SH, Karmaus W, Raza A *et al*. The effect of parental allergy on childhood allergic diseases depends on the sex of the child. *J Allergy Clin Immunol* 2012; 130:427–34 e6.
- 33 Kurukulaaratchy RJ, Matthews S, Waterhouse L, Arshad SH. Factors influencing symptom expression in children with bronchial hyperresponsiveness at 10 years of age. *J Allergy Clin Immunol* 2003; 112:311–6.
- 34 Li J, Wang H, Chen Y, Zheng J, Wong GWK, Zhong N. House dust mite sensitization is the main risk factor for the increase in prevalence of wheeze in 13- to 14-year-old schoolchildren in Guangzhou city, China. *Clin Exp Allergy* 2013; 43:1171–9.
- 35 Lajunen TK, Jaakkola JJ, Jaakkola MS. The synergistic effect of heredity and exposure to second-hand smoke on adult-onset asthma. *Am J Respir Crit Care Med* 2013; 188:776–82.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Table S1. Description of the subset with SPT and the whole cohorts

Table S2. Prevalence of wheeze, rhinitis and sensitization by sex

Table S3. Sex by cohort interaction effects on lifetime wheeze and rhinitis

Table S4. Interactions of risk factors and cohorts for lifetime wheeze and rhinitis.