

Management of ocular allergy

Andrea Leonardi¹ | Diana Silva² | Daniel Perez Formigo^{3,4} | Banu Bozkurt⁵ |
Vibha Sharma⁶ | Pia Allegri⁷ | Carmen Rondon⁸ | Virginia Calder⁹ | Dermot Ryan¹⁰ |
Marek L. Kowalski¹¹ | Luis Delgado² | Serge Doan¹² | Jean L. Fauquert¹³

¹Ophthalmology Unit, Department of Neuroscience, University of Padua, Padua, Italy

²Basic and Clinical Immunology, Department of Pathology, Faculty of Medicine, Serviço de Imunoalergologia, Centro Hospitalar São João, University of Porto, EPE – Porto, Porto, Portugal

³Department of Ophthalmology, Hospital Universitario de Torrejón, Madrid, Spain

⁴Faculty of Medicine, University of Francisco de Vitoria (UFV), Pozuelo de Alarcón, Madrid, Spain

⁵Department of Ophthalmology, Faculty of Medicine, Selçuk University, Konya, Turkey

⁶Department of Paediatric Allergy and Immunology, Royal Manchester Children's Hospital, Manchester University NHS Foundation Trust, Manchester, UK

⁷Allergic Conjunctivitis Unit, Ocular Inflammatory Diseases Referral Center, Rapallo Hospital Ophthalmology Department, Genova, Italy

⁸Allergy Unit, Regional University Hospital of Malaga, IBIMA, UMA, Malaga, Spain

⁹Department of Ocular Biology & Therapeutics, UCL Institute of Ophthalmology, London, UK

¹⁰Allergy and Respiratory Research Group, Medical School, Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, Edinburgh, UK

¹¹Department of Immunology, Rheumatology, and Allergy, Medical University of Lodz, Lodz, Poland

¹²Service d'Ophthalmologie, Hôpital Bichat and Fondation A. de Rothschild, Paris, France

¹³CHU Estaing, Unité d'Allergologie de l'Enfant, Clermont-Ferrand Cedex1, France

Correspondence

Andrea Leonardi, Ophthalmology Unit,
Department of Neuroscience, University of
Padua, Padua, Italy.
Email: andrea.leonardi@unipd.it

Funding information

This work was done under the approval of
EAACI with a TF budget 2015-18.

Abstract

The treatment and management of ocular allergy (OA) remain a major concern for different specialties, including allergists, ophthalmologists, primary care physicians, rhinologists, pediatricians, dermatologists, clinical immunologists, and pharmacists. We performed a systematic review of all relevant publications in MEDLINE, Scopus, and Web Science including systematic reviews and meta-analysis. Publications were considered relevant if they addressed treatments, or management strategies of OA. A further wider systematic literature search was performed if no evidence or good quality evidence was found. There are effective drugs for the treatment of OA; however, there is a lack of an optimal treatment for the perennial and severe forms. Topical antihistamines, mast cell stabilizers, or double-action drugs are the first choice of treatment. All of them are effective in reducing signs and symptoms of OA. The safety and optimal dosing regimen of the most effective topical anti-inflammatory drugs, corticosteroids, are still a major concern. Topical calcineurin inhibitors may be used in steroid-dependent/resistant cases of severe allergic keratoconjunctivitis. Allergen-specific immunotherapy may be considered in cases of failure of first-line treatments or to modify the natural course of OA disease. Based on the current wealth of publications and on the collective experience, recommendations on management of OA have been proposed.

KEYWORDS

management, ocular allergy, allergic conjunctivitis, systematic review, treatment

1 | INTRODUCTION

Ocular allergy (OA) represents a collection of ocular hypersensitivity disorders affecting the eyelid, conjunctiva, and cornea. OA includes seasonal and perennial allergic conjunctivitis (SAC and PAC), vernal and atopic keratoconjunctivitis (VKC and AKC), and contact blepharoconjunctivitis (CBC).¹ These clinical subtypes may be diagnosed and managed by ophthalmologists, allergists, pediatricians, and rhinologists, with or without experience in managing allergies, considering clinical history and signs and symptoms, aided by in vivo and in vitro tests.¹⁻³ Although several studies suggest a high comorbidity of conjunctivitis and rhinitis, conjunctival symptoms are often perceived by clinicians as a minor problem and suboptimally treated. A recent survey revealed that daily treatment of OA has little concordance with current recommendations.⁴ Topical ocular decongestants and corticosteroids were used in the majority of cases. This was independent of the specific diagnosis of OA subtype and severity. Topical antihistamines and mast cell stabilizers, which are the first-line therapy in most published recommendations, were used less frequently.⁴ The incorrect management of OA may increase the risk of local and systemic treatment-related side effects.⁴ This systematic review (SR) intends to provide a comprehensive overview of the currently available treatments for OA and of ocular comorbidities and/or complication, and suggest recommendations for their management using best available evidence in published literature.

2 | METHODS

2.1 | General search strategy

The literature search was based on the systematic literature search in MEDLINE, Scopus, and Web Science. First, we performed a SR of all SR and meta-analyses that addressed OA treatments, according to the search query. For each subtype of OA or treatment, if good quality SR or systematic analysis of randomized controlled trials (RCTs) were found, no further systematic search was performed. If no evidence or poor quality of evidence was found, a further systematic search was performed. The included literature was selected with respect to their hierarchy in the "evidence pyramid."

2.2 | Eligibility criteria

We included SR of observational and interventional studies regarding treatment of OA (SAC, PAC, VKC, AKC, and CBC). The following treatments were included the following: antihistamines, mast cell stabilizers, dual-acting agents (topical mast cell stabilizers and antihistamines), non-steroidal anti-inflammatory drugs (prostaglandin

and leukotriene inhibitors), steroids, calcineurin inhibitors, allergen-specific immunotherapy, and biologics. A SR was defined as a review of the literature with a predetermined and transparent search strategy, where the search strategy and inclusion and exclusion criteria were explicitly described and included guidelines or position papers containing information regarding quality of evidence. Our systematic literature search included nonpharmacological interventions including surgery, psychological, lid hygiene, and lubricants.

Specific search strategy, selection of the study, and assessment of the quality of the evidence are reported as Appendix S1 and S1.

3 | RESULTS

The flowchart of the selection strategy is shown in Figure 1.




FIGURE 1 Flowchart of the selection strategy of the systematic reviews potentially relevant for the purpose of the position paper. Of the 432 publications, 28 were selected and included (see Tables 2-6)

3.1 | Overview of the available pharmacological classes of anti-allergic drugs

There is a wide range of treatment options for OA, some of which are off-label. Currently available topical drugs for OA can be classified into different pharmacological classes based on their mechanism of action (Table 1): antihistamines, mast cell stabilizers, dual-acting agents (topical mast cell stabilizers and antihistamines), alpha-adrenergic agonists (vasoconstrictors), non-steroidal anti-inflammatory drugs (prostaglandin inhibitors), corticosteroids, and calcineurin inhibitors. Immunomodulatory treatments for OA include allergen-specific immunotherapy and biologicals.

3.1.1 | Topical antihistamines, mast cell stabilizers, and dual-acting agents

Three SR addressed the use of topical antihistamines, topical mast cell stabilizers, or topical dual-acting agents for the

treatment of SAC and PAC (Table 2): One included 23 RCTs,⁵ the second 30 RCTs⁶ in a head-to-head study, and the third 41 RCTs.⁷ All the three reviews concluded that these drugs were effective in reducing ocular symptoms vs placebo.⁷ Direct comparisons of different antihistamines and mast cell stabilizers showed insufficient evidence to recommend one drug over another⁶ even though the peer-reviewed literature suggested that olopatadine may be clinically superior to the other anti-allergic molecules,⁸ and alcaftadine may be superior to olopatadine in reducing ocular itch.⁵ A fourth SR⁹ comparing olopatadine with other topical antihistamines (epinastine, ketotifen, and alcaftadine) showed a significant benefit from the use of alcaftadine in reducing symptoms scores when compared to the others drugs. Alcaftadine is only currently approved and available in the United States. Since the publication of the last SR, a further clinical trial has been published demonstrating efficacy of epinastine in controlling symptoms of birch pollen allergic patients.¹⁰

TABLE 1 Topical ocular allergy approved medications

Class	Drug	Dosing	Indication	Considerations
Antihistamines (second generation)	Levocabastine Emedastine	4× daily	- Relief of itching - Relief of signs and symptoms	- Short duration of action - Frequently not enough to treat alone the entire disease
Mast cell stabilizers	Cromolyn Nedocromil Lodoxamide NAAGA	4× daily	Relief of signs and symptoms	- Long-term usage - Slow onset of action - Prophylactic dosing - Frequently not enough to treat alone the entire disease
Dual-acting agents (Antihistamine/ mast cell stabilizers)	Alcaftadine Azelastine Bepotastine Epinastine Ketotifen Olopatadine	2× daily	- Relief of itching - Relief of signs and symptoms	- Bitter taste (azelastine) - No reported serious side effects - Frequently not enough to treat alone the entire disease
Vasoconstrictor/ vasoconstrictor-anti-histamine combinations	Naphazoline/ Pheniramine	2-4× daily	- Rapid onset of action - Episodic itching and redness	- Short duration of action - Tachyphylaxis - Mydriasis - Ocular irritation - Hypersensitivity - Systemic hypertension - Potential for inappropriate patient use
Corticosteroids (listed in ascending potency order)	Hydrocortisone Loteprednol Fluorometholone Desonide Rimexolone Prednisolone Dexamethasone Betamethasone	As required	- Treatment of allergic inflammation - Use in moderate-to-severe forms	- Risk of long-term side effects - No mast cell stabilization - Potential for inappropriate patient use - Requires close monitoring
Calcineurin inhibitors	Cyclosporine A Tacrolimus	2-4× daily	Treatment of severe VKC and AKC not responding anti-allergic drugs	- Off label in OA (tacrolimus approved for VKC only in Japan) - CsA 0.1% received marketing authorization by EMA in July 2018 for severe VKC - Magistral/officials preparations are different from center to center - Quality control and availability of magistral preparations are poor

TABLE 2 Topical antihistamines, mast cell stabilizers, dual-acting agents

Author, Y	Studies design	Participants	Intervention comparisons	Outcomes	Main results	Author's conclusion	Amstar score
Topical antihistamines							
Kam et al 2016 ⁵	23 RCT	Allergic conjunctivitis (n = 3388 eyes)	Olopatadine 0.05%-0.20% vs placebo; ketotifen; alcaftadine	Symptoms scores (itching, hyperemia)	Olopatadine vs placebo Itch SMD -1.33 [-1.43; -1.23]/Itch score -2.62 [-3.25; -1.99] Hyperemia -0.92 [-1.19; -0.65]/-1.92 [-2.67; -1.17] Olopatadine vs Epinastine no significant differences Olopatadine vs Ketotifen no significant differences Olopatadine vs Alcaftadine SMD 0.30 [0.28; 0.50]	Topical olopatadine is safe and effective, but alcaftadine appears to be superior	High
Castillo et al, 2015 ⁶	30 RCT 4 Meta-analysis	Seasonal and perennial allergic conjunctivitis (n = 4344)	Topical antihistamines Mast cell stabilizers (alone and in combination)	Symptoms scores (itching, irritation, watering eyes or photophobia)	Mast cell stabilizers vs placebo (8 RCT), not pooled, nedocromil sodium or sodium cromoglycate is more effective than placebo in improving ocular symptoms Azelastine vs placebo (9 RCT), not pooled, individuals studies improved symptoms Levocabastine vs placebo (5 RCT), not pooled, individual studies with improvement symptoms Olopatadine vs ketotifen (4 RCT) MSD -0.32 [-0.59; -0.06] for itching; MSD -0.06 [-0.35; 0.22] for tearing Nedocromil vs levocabastine (2 RCT), not pooled Azelastine vs levocabastine (2 RCT), not pooled Olopatadine vs control (2 RCT), not pooled, evidence from two small trials may improve symptoms Nedocromil vs azelastine (1 RCT), no difference Olopatadine vs nedocromil (1 RCT), olopatadine more effective Levocabastine vs antazoline and tetryzoline, no difference Ketotifen vs placebo (1 RCT) more effective Ketotifen vs levocabastine (1 RCT) Ketotifen is superior Levocabastine and pemirolast vs levocabastine, combination is more effective Levocabastine vs mequitazine (1 RCT) equally effective Bepotastine vs placebo (1 RCT), effective Bepotastine vs olopatadine (1 RCT), insufficient	All topical antihistamines and mast cell stabilizers reduce symptoms and signs of seasonal allergic conjunctivitis when compared with placebo in the short term are safe and well tolerated. Olopatadine may be more effective than ketotifen	High
Mahvan TD, 2012 ⁹	2 RCT	Allergic conjunctivitis (n = 228)	Alcaftadine 0.05%-0.25% vs placebo	Ocular itching Ocular redness	Improved ocular itching improvement, no improvement in ocular redness, not pooled	Alcaftadine is safe and effective	Critically low
Rosenwasser et al 2005 ⁸	9 CT	Allergic conjunctivitis (n = 714)	Olopatadine 0.1% and 0.5% vs placebo; ketorolac, nedocromil, ketotifen, azelastine and epinastine	Not pooled	No significant side effects; no ocular dryness or irritation with topical use. Reduces redness, itching and swelling. Comparison with ketorolac 0.5%, nedocromil, ketotifen, azelastine and epinastine- not pooled	Olopatadine is clinically superior to the other anti-allergic molecules	Critically low

(Continues)

TABLE 2 (Continued)

Author, Y	Studies design	Participants	Intervention comparisons	Outcomes	Main results	Author's conclusion	Amstar score
Topical mast cell stabilizers							
Owen et al, 2004 ⁷	40 RCT	Seasonal allergic conjunctivitis (n = 790)	Topical mast cell stabilizers; topical antihistamines and topical mast cell stabilizers with antihistamines	Ocular symptom score (itching, burning, lacrimation and soreness)	<i>Sodium cromoglycate vs placebo</i> (8 RCT) OR = 1.7 [4; 78] to benefit from treatment <i>Nedocromil sodium vs placebo</i> (5 RCT) OR = 1.8 [1.3; 2.6] <i>Lodoxamide tromethamine vs placebo</i> (1 RCT) <i>Topical antihistamines vs placebo</i> (9 RCT; 6 levocabastine; 1 azelastine; 1 emedastine; 1 antazoline) not pooled Topical mast cell stabilizers vs topical antihistamines (8 RCT) Levocabastine vs mast cell stabilizers OR 1.3 [0.8; 2.2]	Confirm the benefit of topical mast cell stabilizers and antihistamines over placebo for the treatment of allergic conjunctivitis. There is insufficient evidence to recommend the use of one type of medication over another.	Moderate
Mantelli et al, 2007 ¹²	27 RCT 10 meta-analysis	Vernal Keratoconjunctivitis (n = 1092 patients, 2184 eyes) Mean age 13.3 (4.5 y)	Mast cell stabilizers vs placebo (n = 10) Mast cell stabilizers vs another (n = 8) Mast cell stabilizers vs corticosteroids (n = 2) Mast cell stabilizers vs antihistamines (n = 1) NSAID vs placebo (n = 2) NSAID vs CCT (n = 1) Antimitotic drug (n = 1)	Itching, tearing, photophobia, hyperemia, tarsal papillae, limbal disease and corneal involvement	Itching (mitomycin; sodium cromoglycate, cyclosporine 2%, ketorolac) SMD -1.43 [-1.76; -1.10] Tearing (mitomycin, sodium cromoglycate, cyclosporine 2%) SMD -0.84 [-1.20; -0.49] Photophobia (mitomycin, sodium cromoglycate, cyclosporine 2%) -0.27 [-0.82; 0.39] Total signs (cyclosporine and sodium cromoglycate) SMD -0.94 [-1.34; -0.54] Total symptoms (cyclosporine 2% and sodium cromoglycate) SMD -0.73 [-1.14; -0.32] Tarsal papillae (mitomycin, sodium cromoglycate, cyclosporine) SMD -0.32 [-0.64; -0.00] Corneal involvement (mitomycin, sodium cromoglycate, cyclosporine) SMD -1.15 [-1.50; -0.80] Limbal disease (mitomycin, sodium cromoglycate, cyclosporine) SMD -1.47 [-1.50; -0.83] Hyperemia (mitomycin, sodium cromoglycate, cyclosporine, mipragoside 0.5) SMD -1.07 [-1.38; -0.76]	The currently available topical drugs are effective in treating acute phases of VKC. There is a lack of evidence to support the recommendation of one specific type of medication for treating this disorder	Low

CCT, corticosteroids; NSAID, non-steroidal anti-inflammatory drug; RCT, randomized clinical trials; SMD, standardized mean difference.

Nedocromil sodium or sodium cromoglycate, olopatadine, ketotifen, azelastine, emedastine, levocabastine (or levocabastine), mequitazine, bepotastine besilate, combination of antazoline and tetrazoline, combination of levocabastine and pemirolast potassium.

TABLE 3 Non-steroidal anti-inflammatory drugs (topical prostaglandin and oral leukotriene inhibitors)

Author, Y	Studies design	Participants	Intervention comparisons	Outcomes	Main results	Author's conclusion	Amstar score
Non-steroidal anti-inflammatory drug							
Swamy et al, 2007 ¹⁵	8 RCT	Allergic conjunctivitis (n = 712)	Ophtalmic NSAID ((ketorolac, diclofenac, flurbiprofen) piroxicam and hydrocortisone) vs placebo	Symptoms, ocular itching, and conjunctival injection Side effects	Itching SMD -0.54 [-0.84; -0.24] Lacrimation SMD -0.21 [-0.41; -0.01] Conjunctival injection SMD -0.52 [-0.97; -0.05] Ocular discomfort with treatment SMD 3.97 [2.67; 5.89]	NSAID are more effective than placebo in reducing conjunctival itching and improving a cardinal sign	Low
Wilson et al, 2015 ¹⁶	8 clinical trials 2 reviews	Seasonal allergic conjunctivitis (number of participants not specified)	Ophtalmic non-steroidal anti-inflammatory drugs (Ketorolac 0.5% and diclofenac 0.1%)	Symptoms score change	Decrease short-term treatment in comparison with placebo (7-14 d) in 3 studies; 2 cross-over without benefit ketorolac 0.5%. Diclofenac 0.1% better than ketorolac 0.5% in 1 trial for symptoms. Data not pooled.	Effective in decreasing short-term symptoms	Critically low
Leukotriene antagonist (montelukast)							
Gane et al, 2013 ¹⁷	18 RCT 6 meta-analysis	Ocular eye disease 12 SAC, 5 PAC (n = 9017 adult) (n = 175 children 2-14 y)	Leukotriene receptor antagonists (montelukast)	Ocular symptom scores	LTRA vs placebo (6 RCT) SMD -0.10 [-0.14; -0.07] LTRA vs oral antihistamine (3 RCT) 0.08 [0.02; 0.14] in favor of antihistamines LTRA and oral antihistamine vs placebo (2 RCT) -0.30 [-0.38; -0.21]	In seasonal AC, LTRAs are more efficacious than placebo but less efficacious than oral antihistamines in adult patients.	Moderate

AC, allergic conjunctivitis; CCT, corticosteroids; LTRA, leukotriene receptor antagonists; NSAID, non-steroidal anti-inflammatory drug; PAC, perennial allergic conjunctivitis; RCT, randomized clinical trials; SAC, seasonal allergic conjunctivitis; SMD, standardized mean difference.

TABLE 4 Systemic antihistamines

Author, Y	Studies design	Participants	Intervention comparisons	Outcomes	Main results	Author's conclusion	Amstar score
Compalati et al, 2013 ²⁰	4 RCT	Allergic rhinoconjunctivitis (n = 1135)	Rupatadine vs ebastine; placebo (n = 473) Rupatadine vs loratadine; placebo (n = 283) Rupatadine vs desloratadine; placebo (n = 379)	Itchy and watery eyes	Itch eyes SMD: -0.29, 95% CI -0.45 to -0.14 Watery eyes reduction SMD: -0.25, 95% CI -0.45 to -0.06	Improvement ocular symptom of rupatadine vs placebo.	Low

RCT, randomized clinical trials; SMD, standardized mean difference.

Overall, topical antihistamines and mast cell stabilizers appear to be safe and well tolerated.^{6,11} The most frequently reported side effects from the use of these agents were burning and stinging sensation, blurred vision and unacceptable aftertaste.^{5,6} Data on their long-term efficacy and safety are still lacking. To minimize possible toxic effects of preservative compounds on the ocular surface, single-dose preservative-free eye drops should be used whenever possible.

A SR of 20 RCT evaluated the efficacy of topical antihistamines and mast cell stabilizers for the treatment of VKC showing an improvement of ocular symptoms score with use of all these drugs.¹² The pooled data were unable to recommend use of one agent over the other. A RCT published after this SR showed benefit with improved inflammatory biomarkers and total symptom score using preservative-free N-acetyl-aspartyl-glutamic acid (NAAGA) compared to levocabastine.¹³

The treatment of AKC with antihistamines and mast cell stabilizers has only been reported in a few case reports and case series.¹⁴

Recommendations

- All topical drugs are effective in reducing signs and symptoms ⊕⊕⊕⊕↑↑
- Topical antihistamines and dual-acting drugs may have lead to a quicker onset symptom relief when compared to mast cell stabilizers ⊕○○○↑?
- Dual-acting agents with combined mast cell stabilizer and antihistaminic function provide better symptom control ⊕○○○↑?
- Mast cell stabilizers such as chromones require multiple daily doses and have a delayed onset of action, hence are less preferable ⊕○○○↑?
- SAC and PAC can be managed using the same drugs ⊕⊕⊕⊕↑↑
- The duration of treatment is longer in PAC compared to SAC ⊕○○○↑?
- Topical antihistamines and mast cell stabilizers can be used in VKC ⊕⊕○○↑ and AKC ⊕○○○↑?
- All these drugs can be used in combination ⊕○○○↑?

3.1.2 | Topical alpha-adrenergic agonists (vasoconstrictors)

Topical decongestants are frequently used as first-line treatment due to their availability over the counter.⁴ They merely alleviate

hyperemia, having little to no relief from itch and a short duration of action.⁷ In a recent randomized controlled trial comparing several treatment options, the use of naphazoline/antazoline was associated with lower tolerability profile of all treatment. They may cause side effects such as rebound redness, chronic follicular conjunctivitis, and tachyphylaxis. In older formulations, ocular decongestants are paired with topical first-generation antihistamines, such as pheniramine and antazoline, to relieve both itching and redness.

Recommendations

- Vasoconstrictors alleviate only hyperemia ⊕○○○↑?
- They should be used with caution and for a short period of 5-7 days because of side effects and tachyphylaxis (TF expert opinion)↑↑

3.1.3 | Non-steroidal anti-inflammatory drugs NSAID (topical prostaglandin and oral leukotriene inhibitors)

One SR¹⁵ based on 8 RCTs concluded that topical NSAIDs are more effective than placebo in reducing ocular itching and redness. Use of varied outcome parameters did not permit a comparison^{15,16} (Table 3). NSAIDs are rarely used due to their local side effects, such as burning/stinging after application.

The oral leukotriene inhibitor montelukast has shown to be useful in the treatment of ocular symptoms in SAC and PAC, but less effective than oral antihistamines.¹⁷

Recommendations from the TF group

- NSAIDs are effective for their short-term use but do not target specific inflammatory mechanisms ⊕○○○↓?
- In adult SAC patients, leukotriene inhibitors are less efficacious than oral antihistamines ⊕○○○↑?

3.1.4 | Systemic antihistamines

Oral antihistamines are frequently used in case of allergic comorbidities such as rhinitis and are used in almost one-third of the patients with ocular symptoms.⁴ Drugs such as loratadine, desloratadine,¹⁸ and fexofenadine¹⁹ are highly effective for the treatment of allergic

TABLE 5 Corticosteroids and calcineurin inhibitors

Author, Y	Studies design	Participants	Intervention comparisons	Outcomes	Main results	Author's conclusion	Amstar score
Topical corticosteroids							
O'Gallagher et al, 2017 ⁸⁴	1 RCT	Blepharoconjunctivitis (n = 137 children)	Loteprednol etabonate	Improvement of symptoms Elimination clinical signs Adverse effects	No sufficient data on the improvement in symptoms or signs; no difference on adverse effects	No high-quality evidence regarding safety and efficacy of topical treatments for BKC	High
Sheppard et al 2016 ²⁴	40 clinical trials SAC (4) PAC (1) VKC (1) BKC (3)	SAC (n = 856) PAC (n = 159) VKC (n = 37) BKC (n = 355)	Loteprednol etabonate	Incidence of IOP	0.2% suspension there was no increase in IOP for SAC or PAC; 0.5% suspension there was no increase in VKC or SAC (both in comparison with vehicle)	Favorable IOP safety profile for loteprednol etabonate with both short-term and long-term use	Critically low
Wu et al, 2015 ²³	8 RCT SAC (4 RCT), GPC (3 RCT) VKC (1 RCT)	(n = 1445)	Loteprednol etabonate vs placebo; topical olopatadine; topical fluorometholone acetate	Ocular symptoms (ocular itching) Sign (bulbar conjunctival injection or papillae) Increased intraocular pressure (IOP)	Compared to placebo: Sign severity SMD -0.85 [-1.35; -0.35] (all) SAC -0.45 [-0.62; -0.28]; only one trial GPC and VKC Weighted mean difference -0.66 [-0.97; -0.35] (all) SAC -0.43 [-0.46; -0.31]; only one trial GPC and VKC IOP OR 3.03 [1.04; 8.80] Compared to topical olopatadine: Sign severity SMD -3.78 [-10.61; 3.04] WMD -0.98 [-2.00; 0.05] (all interventions)	Topical loteprednol etabonate is effective in treating allergic conjunctivitis	High
Nasal corticosteroids							
Weiner et al, 1998 ²⁶	16 RCT	Allergic rhinoconjunctivitis (n = 2267)	Nasal corticosteroids vs oral antihistamines (terfenadine; astemizole; loratadine; cetirizine)	Ocular symptoms (included 11 studies)	Nasal steroid vs oral antihistamines SMD -0.043 [-0.157; 0.072]	There was no significant difference between intranasal corticosteroids and oral antihistamines on ocular symptoms	Low
Hong et al, 2011 ²⁷	32 studies RCT	SAC and PAC (n = 6573)	Nasal corticosteroids	Total ocular symptom score (TOSS)	Data not pooled Improvement of TOSS in 9 out of 10 studies Total eye symptoms 5 out of 13 studies	Intranasal corticosteroids have a positive impact on the TOSS of AR	Critically low

(Continues)

TABLE 5 (Continued)

Author, Y	Studies design	Participants	Intervention comparisons	Outcomes	Main results	Author's conclusion	Amstar score
Naclerio et al, 2008 ²⁸	3 SR 5 RCT	Allergic rhinoconjunctivitis TAA (n = 375) FP (n = 1645) MF (n = 1198) FF (n = 940) BUD (n = 280)	Nasal corticosteroids Triamcinolone acetonide (n = 375) Propionate Fluticasone (7 RCT data association) Mometasone furoate (2 RS 1 RCT) Furoate fluticasone (2 RCT) Budesonide (1 RCT)	Total ocular symptom score	Data not pooled, an improvement of total eye symptom score was seen vs placebo was seen in all clinical trials and meta-analysis	Intranasal corticosteroids are effective and well tolerated in the treatment of ocular symptoms associated with allergic rhinitis.	Critically low
Yáñez et al, 2002 ²⁹	4 RCT	Allergic rhinitis Beclomethasone (n = 166) BUD (n = 31) Fluticasone (n = 193) Flunisolide (n = 38) Azelastine (n = 240) Levocabastine (n = 408)	Beclomethasone or flunisolide vs azelastine (n = 2 RCT) Budesonide or fluticasone vs levocabastine (n = 2 RCT)	Eye symptoms	Nasal steroid vs nasal antihistamines SMD -0.07 [-0.27; 0.12, P = 0.4]	There was no difference between interventions in the relief of ocular symptoms	Critically low

Topical cyclosporine

Wan et al, 2013 ³²	7 RCT	Allergic conjunctivitis (n = 306 eyes of 153 patients) 3 studies recruited steroid-dependent allergic conjunctivitis	Topical cyclosporine in concentration from 0.05% to 2%	Composite sign score (average of at least one of the following signs: hyperemia, swelling, papillae and giant papillae on the tarsal conjunctiva, hyperemia and edema of the bulbar conjunctiva, or corneal involvement) Composite symptom score (average of: redness, tearing, burning, discomfort, foreign body sensation, discharge, and photophobia) Medication use	Compared to placebo: Composite sign score: SMD -1.21 95% CI, [-1.80; -0.62] Composite symptom score: SMD -0.84 95% CI [-1.51; -0.16] Reduction on steroid eye drop (3 RCT) SMD -61.2 [-101.6; -20.7]	Topical cyclosporine could be an effective and safe treatment method for allergic conjunctivitis	Low
-------------------------------	-------	--	--	---	---	--	-----

(Continues)

TABLE 5 (Continued)

Author, Y	Studies design	Participants	Intervention comparisons	Outcomes	Main results	Author's conclusion	Amstar score
González-López et al, 2012 ³³	3 RCT	Atopic keratoconjunctivitis (n = 58)	Cyclosporine 0.05% or cyclosporine 2% in maize oil vs preservative-free artificial tears or placebo	Symptoms improvement (reported by the participant) itching, tearing, discomfort, mucous discharge, photophobia or pain Topical steroid use Clinical signs Adverse effects	Not pooled Symptoms composite score significantly improved for all associated, but not for specific symptoms in one study (Clinical signs improved in the composite score in one study) Reduction of topical steroid use in one study	Topical CsA may provide clinical and symptomatic relief in AKC and may help to reduce topical steroid use in patients with steroid-dependent or steroid-resistant AKC. No serious adverse events were reported	High
Mantelli et al, 2007 ¹²	27 RCT 10 meta-analysis	Vernal Keratoconjunctivitis (n = 1092 patients, 2184 eyes) mean age 13.3 (4.5 y)	Mast cell stabilizers vs placebo (n = 10) Mast cell stabilizers vs another (n = 8) Mast cell stabilizers vs corticosteroids (n = 2) Mast cell stabilizers vs antihistamines (n = 1) NSAID vs placebo (n = 2) NSAID vs CCT (n = 1) Antimitotic drug (n = 1)	Itching, tearing, photophobia, hyperemia, tarsal papillae, limbal disease, and corneal involvement	Itching (mitomycin, sodium cromoglicate, cyclosporine 2%, ketorolac) SMD -1.43 [-1.76; -1.10] Tearing (mitomycin, sodium cromoglicate, cyclosporine 2%) SMD -0.84 [-1.20; -0.49] Photophobia (mitomycin, sodium cromoglicate, cyclosporine 2%) -0.27 [-0.82; 0.39] Total signs (cyclosporine and sodium cromoglicate) SMD -0.94 [-1.34; -0.54] Total symptoms (cyclosporine 2% and sodium cromoglicate) SMD -0.73 [-1.14; -0.32] Tarsal papillae (mitomycin, sodium cromoglicate, cyclosporine) SMD -0.32 [-0.64; -0.00] Corneal Involvement (mitomycin, sodium cromoglicate, cyclosporine) SMD -1.15 [-1.50; -0.80] Limbal disease (mitomycin, sodium cromoglicate, cyclosporine) SMD -1.17 [-1.50; -0.83] Hyperemia (mitomycin, sodium cromoglicate, cyclosporine, miprofoside 0.5) SMD -1.07 [-1.38; -0.76]	The currently available topical drugs are effective in treating acute phases of VKC. However, there is a lack of evidence to support the recommendation of one specific type of medication for treating this disorder	Low

(Continues)

TABLE 5 (Continued)

Author, Y	Studies design	Participants	Intervention comparisons	Outcomes	Main results	Author's conclusion	Amstar score
Tacrolimus							
Zhai et al 2010 ³⁶	2 RCT 4 Case series	VKC (n = 87) AKC (n = 35) PAC (n = 20)	0.1% tacrolimus ophthalmic suspension 0.03% tacrolimus Tacrolimus ointment vs clobetasol AKC	Total objective score Symptom score Ulcer improvement	Not pooled Improvement of total score with tacrolimus vs placebo (-5.6 ± 5.1 for tacrolimus ophthalmic suspension vs -0.1 ± 4.5 for placebo) Similar effect to topical corticosteroids for AKC	Needed more studies	Critically low

CCT, corticosteroids; IOP, intraocular pressure; NSAID, non-steroidal anti-inflammatory drug; RCT, randomized clinical trials; TOSS, total ocular symptom score.

rhinoconjunctivitis (ARC) (Table 4). Most of the SRs have addressed total symptoms scores,^{18,19} without evaluating impact on specific ocular symptoms. Itching and watery eye symptoms significantly improved after rupatadine treatment compared to placebo.²⁰

While the drowsiness so commonly noted with the older first-generation systemic antihistamines has improved in the newer second-generation antihistamines, some of the new molecules still inhibit muscarinic receptors, leading to mucosal dryness.^{21,22} Moreover, patients with dry eye have reduced barrier function at the mucosal interface against environmental allergens and pollutants and possibly a lower threshold for allergen response. Some oral antihistamines may exacerbate OA by lowering the barrier defense offered by a healthy tear film.

Recommendations

- Systemic antihistamines should be used in case of comorbidities that require it use ⊕○○○↑?
- Some systemic antihistamines may induce drying effects, particularly relevant at the ocular surface barrier ⊕○○○↑?

3.1.5 | Corticosteroids

Should not be the first choice of therapy for OA. In clinical practice, they are the most effective anti-inflammatory agents in active OA. Because of potential adverse effects (increased intraocular pressure, with a potential evolution toward glaucoma, cataract formation, bacterial, viral and fungal superinfections), their use must be monitored by an ophthalmologist (especially in prolonged treatments). A SR (Table 5) on the use of a loteprednol eye drops for treating SAC (4 RCTs) and VKC (1 RCT) reached a high level of confidence using AMSTAR2 score,²³ supporting the efficacy of this treatment. Loteprednol 0.5% and 0.2% were considered effective in treating signs and symptoms of SAC, but should be used with caution due to the higher incidence of intraocular pressure (IOP) elevation (pooled odds ratio = 3.03) compared with placebo and olopatadine.²³ A second review demonstrated significantly lower rates of IOP elevation (≥ 10 mm Hg) when compared to topical prednisolone 1% or dexamethasone 0.1%, suggesting a favorable IOP safety profile for loteprednol with both short-term and long-term use. However, this review received a critically low AMSTAR score.²⁴

A wide variety of corticosteroid eye drops of different potencies are available across the world (Table 1).¹ There are no studies directly comparing formulation, strength or regimen of any specific corticosteroid over another for the treatment of OA. There are two main regimens used in OA: (a) pulsed therapy of 3-4 drops per day for 3-5 days and (b) prolonged treatment of 1-3 weeks, tapered slowly over several days. Pulsed therapy is the favored treatment of acute exacerbations of VKC and AKC, especially when the cornea is involved. The potency and treatment duration of the topical corticosteroid should be chosen clinically based on the severity of ocular inflammation and corneal involvement.

The beneficial effect of *intranasal corticosteroids* (INCs) on ocular symptoms has been demonstrated in several studies suggesting that

TABLE 6 Allergen-specific immunotherapy

Author, Y	Studies design	Participants	Intervention comparisons	Outcomes	Main results	Author's conclusion	Amstar score
Immunotherapy							
Dhami et al, 2017 ⁴⁵	160, 134 DBRCT 61 SCIT; 71 SLIT	Allergic rhinoconjunctivitis or allergic rhinitis 6379 patients SCIT 13 636 patients SLIT	Weed, tree and grass pollens, molds, cat and dog dander, and house dust mites	Difference symptom score Difference medication score Difference QoL	Symptom score [SMD -0.53 (95% CI -0.63 to -0.42); SCIT -0.65 (-0.86 to -0.36) vs SLIT -0.48 (95% CI -0.61 to -0.36); Seasonal allergens -0.37 (95% CI -0.45 to -0.28)] Medication score SMD of -0.38 (95% CI -0.49, -0.26) [SCIT SMD -0.52 (95% CI -0.75, -0.29); SLIT -0.31 (95% CI -0.44, -0.18)]	AIT is effective in achieving clinically important short-term improvements in symptom, medication, and combined symptom and medication scores	High
Di Bona et al ⁴⁷ 2015	13 RCT	Seasonal allergic rhinoconjunctivitis (n = 4659)	Grass pollen SLIT vs placebo Phleum p5 or 5 grass extracts	Difference in symptom score Difference medication score Number adverse events	Symptom score (SMD, -0.28; 95% CI, -0.37 to -0.19; <i>P</i> < 0.001) Medication score (SMD, -0.24; 95% CI, -0.31 to -0.17; <i>P</i> < 0.001) OR adverse events 2.91	Small benefit of grass SLIT tablets in the treatment of SARC	Moderate
Nelson et al, 2015 ⁴⁸	37 RCT 14 SLIT tablet; 14 SLIT drop; 9 SCIT	Allergic rhinitis treated with IT (n = 4016) and placebo (n = 3743)	SCIT vs SLIT grass pollen IT vs placebo SLIT tablets vs SLIT drops	Difference in symptom score (rhinoconjunctivitis score preferred) Difference medication score	Symptom score [SCIT vs placebo SMDs (95% CI): -0.32 (-0.45 to -0.18); SLIT vs placebo SMDs (95% CI): -0.32 (-0.41 to -0.23)] Medication score [SCIT vs placebo SMDs (95% CI): -0.33 (-0.52 to -0.13); SLIT vs Placebo SMDs (95% CI): -0.44 (-0.83 to -0.06)] No difference between SCIT or SLIT	Comparable reduction in allergic conjunctivitis symptoms with SLIT and SCIT	Moderate
Devillier et al, 2014 ⁴⁹	38 DBRCT 28 med 10 SLIT	Seasonal allergic rhinitis (n = 21 223)	Antihistamines (desloratadine; bilastine; loratadine; fexofenadine; cetirizine); Nasal CCT; montelukast and grass pollen SLIT tablet,	Symptoms score Combined symptom-medication score Relative clinical impact Reported post-treatment or season-long nasal Total symptom scores. 100 × (score Placebo - score Active)/score Placebo)	Five grass pollen SLIT tablets [-0.30 (-0.36, -0.23)]; -29.6% (-23% to -37%) Timothy SLIT tablets -19.2% (-6% to -29%) Nasal corticosteroids [-0.55 (-0.63, -0.47)] -23.5% (-7% to -54%) azelastine-fluticasone [SMD -1.00 (-1.10, -0.90)] -17.1% (-15% to -20%) H1-antihistamines [SMD -0.39 (-0.43, -0.35)] -15.0% (-3% to -26%) Montelukast [-0.23 (-0.30, -0.16)] -6.5% (-3% to -10%)	Grass pollen SLIT tablets had a greater mean relative clinical impact than second-generation antihistamines and montelukast and similar to the mean relative clinical impact of nasal corticosteroids	Moderate
Lin et al, 2013 ⁵⁰	RCT 13 SLIT	Conjunctivitis (n = 1074)	SLIT Grass mix, dust mite, parietaria, timothy grass, olive, tree mix	Symptom score	No pooled All but two studies showed a greater improvement in SLIT	Moderate strength in support of SLIT for treating allergic conjunctivitis	High

(Continues)

TABLE 6 (Continued)

Author, Y	Studies design	Participants	Intervention comparisons	Outcomes	Main results	Author's conclusion	Amstar score
Calderon et al, 2011 ⁵¹	RCT 42	Allergic conjunctivitis seasonal and perennial (n = 3399)	SLIT Grass, mites, weeds vs placebo	TOSS Individual symptom score Medication score Conjunctival immediate allergen sensitivity	TOSS (n = 36 studies) SMD -0.41 [-0.53; -0.28], not significant for perennial allergens (n = 6) and significant for children and adults Itch SMD -0.31 [-0.42; -0.20] Watery eyes SMD -0.23 [-0.34; -0.11] Red eyes SMD -0.33 [-0.45; -0.22] No significant differences in medication score	SLIT is effective in reducing total and individual symptom score in subjects with ARC or conjunctivitis.	High
Kim et al, 2013 ⁵²	RCT 3 SCIT 5 SLIT	Conjunctivitis (n = 513)	AIT pollen and dust mite	Percent difference in pre-to-post change for conjunctivitis symptoms (<15% weak; 15-40 moderate; >40% strong)	Without specific values of percent of increase for conjunctivitis. SCIT showed an improvement vs placebo in all studies and 4 out of 5 studies showed an improvement with SLIT.	The strength of evidence is low for SCIT and moderate for SLIT in conjunctivitis symptoms improvement	Moderate
Röder et al, 2008 ⁵⁵	RCT 6 SCIT 4 LNIT 7 OIT 11 SLIT	Allergic rhinoconjunctivitis (n = 1619)	19 trials with seasonal allergen (grass pollen) 9 trials (house dust mite)	Symptoms scores Medication scores	Did not present specific data; not pooled	Insufficient evidence that immunotherapy in any administration has a positive effect on symptoms and/or medication	Low

DBRCT, double-blind randomized controlled trial; IT, allergen immunotherapy; LNIT, nasal immunotherapy; OIT, oral immunotherapy; RCT, randomized clinical trials; SCIT, subcutaneous immunotherapy; SLIT, sublingual immunotherapy; SMD, standardized mean difference.

their reduction is mediated via the ocular-nasal reflex inhibition. The variability of the effect depends on the affinity of the drug to its glucocorticoid receptor.²⁵ Four SR evaluated the use of INCs for the treatment of ocular symptoms associated with allergic rhinitis^{26,27} (Table 5) showing that INCs are well tolerated and effective in reducing the total ocular symptom score (TOSS), even though the outcome measures were not designed to focus specifically on ocular symptoms. It is noted that oral/topical antihistamines are not superior to INCs in reducing TOSS.^{28,29} However, despite large patient cohorts, all SR had a low or critically low confidence rating of results according to AMSTAR2. A recent metanalysis of 3 RCTs noted benefit of a topical nasal combination, fluticasone propionate, and azelastine on TOSS in patients with seasonal ARC.³⁰

There are no studies specifically comparing INCs against each other for the treatment of ocular symptoms. Although data are scarce, there is no evidence that INCs used for prolonged periods of several months increase the risk of cataract formation, intraocular hypertension, and glaucoma, since they have little or no systemic absorption (fluticasone and mometasone).³⁰

The use of corticosteroids as dermatologic applications in OA is reserved for AKC and CBC.¹ Lowest appropriate potency corticosteroids, such as hydrocortisone or budesonide on the eyelid skin, are recommended for the treatment of severe acute eyelid eczema.

Supra-tarsal injections of dexamethasone sodium phosphate, triamcinolone acetonide, or hydrocortisone sodium succinate have been proposed to treat recalcitrant AKC and VKC cases,³¹ but should only be used by specialists with caution in severe patients unresponsive to other treatments.

Systemic corticosteroids may be used as short course in selected severe hyperacute exacerbations involving either eyelid skin or cornea especially in VKC and AKC.

Recommendations

- Topical corticosteroids eye drops should be used with caution under ophthalmologist's monitoring and preferably for shorter duration due to the high risk of local and potential blinding side effects ⊕○○○↑↑
- For the treatment of SAC and PAC, topical corticosteroids are rarely needed ⊕○○○↓↓
- Corticosteroid eye drops can be used preferably as short, pulsed therapy in acute exacerbations of OA, especially in VKC and AKC or when the cornea is involved under ophthalmologist supervision (TF expert opinion) ↑↑
- INCs are effective and well tolerated in the treatment of ocular symptoms associated with ARC ⊕○○○↑↑
- INCs should not be used if only ocular signs and symptoms are present (TF expert opinion) ↓↓
- Topical skin corticosteroid applications should be used in the acute phase of eyelid eczema, with a preference for low potency corticosteroids (TF expert opinion) ↑↑

3.1.6 | Calcineurin inhibitors

Topical calcineurin inhibitors are the most frequently used treatments as steroid-sparing agents in steroid-dependent cases of

VKC and AKC. Two SR evaluated the use of topical cyclosporine (CsA) in VKC and AKC^{32,33} (Table 5). The first one showed that topical CsA is effective in alleviating the signs and symptoms of VKC and AKC, reducing the dependency on topical steroid eye drops while maintaining similar safety profile as of placebo. The second SR highlighted the relative scarcity of RCTs assessing the efficacy of topical CsA in AKC and suggested that CsA provides clinical and symptomatic improvement and may help in reducing topical steroid use in patients with steroid-dependent or non-steroid-responsive AKC.³³

Compounded formulations of CsA are prepared in many countries by hospital and retail pharmacies with differing excipients, processes, and quality. Drug concentrations range from 0.05% to 2% and posology from 1 to 6 instillations daily.

Cyclosporine 0.1% cationic emulsion (CE) is commercially available for the treatment of severe dry eye disease.³⁴ The same formulation has obtained in 2018 the marketing authorization by EMA for the treatment of severe VKC. Severe VKC patients treated with this formulation achieved significant improvements in signs, symptoms, and QoL compared with patients who received vehicle alone.³⁵

Tacrolimus 0.03%-0.1% eye drops or ointments have been proposed for the treatment of severe, refractory cases of AKC and VKC. A commercial eye drop preparation is available only in Asia with the indication of severe AKC and VKC. One review, with a critically low quality of evidence score, highlighted the benefits of tacrolimus over placebo in 2 RCTs and 4 case series³⁶ (Table 5). A RCT comparing the effect of tacrolimus 0.1% vs CsA 2%³⁷ showed that both drugs were effective in treating VKC without significant differences between the two. In a second RCT, CsA-resistant VKC patients,³⁸ treated with tacrolimus 0.1%, showed a significant improvement in clinical scores over CsA 1%. A recent trial comparing the effect of 0.1% topical tacrolimus alone or in combination with topical corticosteroids in refractory allergic ocular diseases also showed a potential steroid-sparing effect.³⁹ In addition, tacrolimus skin ointments 0.03% or 0.1% have been shown to be beneficial in the treatment of lid eczema in AKC patients.⁴⁰⁻⁴² Tolerability of topical calcineurin inhibitors is a concern as burning sensation is frequently reported. Infections with molluscum contagiosum, papilloma virus, and herpes are infrequent but are recognized risks.

A systemic immunosuppressive treatment may be prescribed in most refractory cases of AKC threatening vision. Cyclosporine is the most frequently used drug.⁴³ Tacrolimus and mycophenolate mofetil are alternative options.

Recommendations

- CsA eye drops are not recommended for SAC and PAC (TF expert opinion) ↑↑
- CsA eye drops may be used as a steroid-sparing agent in steroid-dependent cases of VKC or AKC ⊕⊕○○↑↑
- Tacrolimus off-label eye drops/ointment should be reserved for use in severe VKC and AKC cases refractory to CsA ⊕○○○↑↑

3.1.7 | Allergen-specific immunotherapy

Since in most patients OA is associated with AR, criteria for allergen immunotherapy (AIT) should follow the recommendations given by the EAACI guidelines.⁴⁴ AIT should be considered only when IgE-mediated allergy is evidenced and when all of following criteria are met the following: moderate-to-severe symptoms strongly suggestive of ARC, which interfere with usual daily activities or sleep despite regular and appropriate pharmacotherapy and/or avoidance strategies and evidencing of IgE sensitization (positive SPT and/or serum-specific IgE) to one or more clinically relevant allergens.^{44,45} In addition, conjunctival allergen provocation test may be helpful in detection of the most relevant allergen before initiating and as a follow-up tool in assessing response of AIT.² AIT should also be considered in less severe ARC to take advantage of the long-term benefit on AR and potential prevention of asthma.⁴⁶ Seven of the 8 selected SR with high and moderate AMSTAR2 scores (Table 6) recommended the use of SLIT and SCIT for moderate improvement of ocular symptoms in the treatment of ARC.^{45,47-52} In two recent RCT regarding house dust mite immunotherapy, an improvement was seen in ocular symptoms score.^{53,54} Only one SR (Table 6) (28 RCT including 1619 children and adolescents with ARC) showed low evidence of the efficacy of SLIT and SCIT on ocular symptoms.⁵⁵

Meta-analysis showed evidence for AIT, with some heterogeneity, in both adults and children, with both SLIT and SCIT,⁵² drop and tablet formulations, in perennial and seasonal allergies, in pre/co-seasonal therapy, and with continuous therapy and in various formulations. Concerns were focused on standardization of allergen extracts and formulation of SLIT preparations.

In cases of isolated allergic conjunctivitis, AIT may be considered. TOSS was evaluated as the primary outcome parameter in 36 studies (1725 SLIT and 1674 placebo)⁵¹; TOSS was significantly reduced when compared with placebo, as well as individual ocular symptoms scores (redness, itchy and watery eyes). No significant reduction in ocular eye drops use was observed, whereas the threshold dose for conjunctival immediate allergen sensitivity was increased. Two other SRs focusing on ocular symptoms⁵⁰ concluded that the evidence was of moderate strength in support of SLIT and low for SCIT for treating allergic conjunctivitis. No publication was found assessing impact of AIT in VKC and AKC.

Recommendation

- AIT may be considered in cases of failure of first-line treatments or to modify the natural course of ocular allergic disease ⊕⊕○○↑?
- AIT can only be considered only when IgE-mediated hypersensitivity is evidenced ⊕⊕⊕⊕ ↑↑
- Before AIT is recommended, control of symptoms of allergic conjunctivitis and other systemic symptoms to assess suitability should be taken into account. ⊕⊕○○↑?
- AIT is effective for the treatment of allergic conjunctivitis due to grass pollen (⊕⊕⊕⊕ ↑↑) and house dust mite (⊕⊕⊕⊕↑?)

- SLIT is effective in reducing total and individual ocular symptom score in subjects with allergic conjunctivitis ⊕⊕⊕⊕ ↑?
- There are no studies on AIT in VKC and AKC patients. In these forms, AIT requires case-to-case assessment by experts (TF expert opinion)↑?

3.1.8 | Biologicals

Omalizumab, a systemic anti-IgE antibody approved for severe asthma, has been used in refractory VKC and AKC and reported in a few case reports/series.⁵⁶ Control of the disease was partial or complete in most patients, but poor response was noted in some with very severe presentation.⁵⁷

Dupilumab is a promising intervention in the management of atopic dermatitis and asthma; however, dupilumab-associated ocular inflammation leading to cicatricial ectropion has been reported suggesting that this drug may not be ideal for the treatment of AKC with eyelid eczema.⁵⁸

3.2 | Nonpharmacological management

Patients and caregivers should receive educative support regarding the anticipated duration and prognosis of the OA, and possible complications from suboptimal control.¹ The first line of management is the identification of offending allergens and avoidance measures. Particularly during exacerbations in VKC, to minimize the exposure to nonspecific triggering factors, such as sun, wind, and salty water, patients should use measures such as sunglasses, hats with visors, and swimming goggles. Frequent hand, face, lid hygiene, and eye washing should also be suggested. Cold compresses may provide decongestant effect. Tear substitutes aid in stabilization of the tear film providing a better mucosal barrier against allergens, acting as an eyewash and diluting the concentration of mediators in the tear film in contact with the ocular surface. Products with herbal extracts such as chamomile-containing eye drops should be avoided as they may cross-react with allergens (for example, *Artemisia vulgaris*).⁵⁹

Psychological support may be necessary in severe cases of VKC and AKC. The psychodynamic research on OA is currently poor. For patients with AKC and VKC, a collaborative approach between the family doctor, the medical specialist, the psychologist, and occupational therapists should be considered.⁶⁰ There are reports of impact on QoL in different types of OA. There is a dearth of reported interventions of mitigation of psychological impact of the disease.³

3.3 | Management in specific populations

3.3.1 | Pregnancy

Few reports are available in literature concerning the management of OA in pregnant or lactating women. Careful evaluation of allergic status and need of drug administration is warranted. Allergen

TABLE 7 Practical treatment of ocular allergy

A. How to treat IgE-mediated diseases SAC and PAC	Avoidance of clinically relevant allergens is the first step in the prevention of ocular allergy symptoms Topical antihistamines, mast cell stabilizers, or double-action drugs are the first choice of treatment, and all effective in reducing signs and symptoms Dual-acting agents with combined mast cell stabilizer and antihistaminic function increase the possibility of symptom improvement and have a faster relief of symptoms compared to mast cell stabilizers Avoid topical corticosteroids, as they are rarely needed Intranasal corticosteroids are effective and well tolerated in the treatment of ocular symptoms associated with ARC, but should not be used if only ocular signs and symptoms are present Topical vasoconstrictors alleviate only hyperemia and should be used with caution for a short period of 5-7 d because of side effects and tachyphylaxis Systemic antihistamines should be used in acute forms or when ocular symptoms are associated with other allergic comorbidities Leukotriene inhibitors are reported to be less efficacious than oral antihistamines in adult SAC patients Consider SIT when specific sensitization is the main cause of ocular allergy, as it is effective for the treatment of ARC to seasonal allergens and perennial allergens SLIT has been shown to be effective in reducing total and individual ocular symptom score in subjects with conjunctivitis
B. How to treat persistent/chronic forms (IgE- and non-IgE-mediated) VKC and AKC	Avoidance of specific and nonspecific triggers is the first step in the prevention of ocular allergy symptoms Use cold compresses, good eyelid hygiene, and lubricants Topical antihistamines, mast cell stabilizers, or double-action drugs are the first treatment choice and may be used in combination. They should be used frequently during the day and during the whole season Systemic anti-allergic drugs should be used when ocular symptoms are associated with other allergic comorbidities Topical corticosteroids should be used as short, pulsed therapy, in acute exacerbations or when the cornea is involved, under ophthalmologist's monitoring Topical calcineurin inhibitors, preferentially cyclosporine A (0.1% on-label treatment in the EU), may be used as a steroid-sparing agent in steroid-dependent patients followed in specialized centers; tacrolimus 0.1% eye drops should be reserved for severe VKC and AKC cases refractory to CsA (off-label treatment in the EU) A systemic immunosuppressive treatment should be prescribed in most refractory cases of AKC with visual threat. Cyclosporine is the most frequently used drug. Tacrolimus and mycophenolate mofetil are alternative options
C. How to treat non-IgE-mediated diseases CBC	Avoidance of irritants and/or sensitizing antigens Eyelid hygiene Emollients and skin moisturizing agents Oral antihistamines can be used to alleviate eyelid itching and inflammation Topical corticosteroids ointments or dermatological creams should be used in the acute phases of eyelid eczema, with a preference for low potency corticosteroids, such as hydrocortisone, desonide, and triamcinolone acetonide Topical calcineurin inhibitors skin ointments 0.03% or 0.1% have been shown to be useful in the treatment of lid eczema in AKC patients. Tolerability is a concern as burning sensation is frequently reported and secondary infections, although infrequent, have been recognized

AKC, atopic keratoconjunctivitis; CBC, contact blepharoconjunctivitis; PAC, perennial allergic conjunctivitis; SAC, seasonal allergic conjunctivitis; VKC, vernal keratoconjunctivitis.

avoidance and environmental measures are the first step, before mast cells stabilizers eye drops can be used. Topical antihistamines or double-acting drugs can be safely tried. As yet there is no evidence of severe adverse events with their use, although US-FDA has assigned many of them to the C category (use with caution if benefits outweigh risks). Short courses of topical corticosteroids if required are cautiously permitted. Vasoconstrictors and decongestants are generally avoided during pregnancy. The use of systemic medications should be minimized if possible. Pregnant (especially in the first trimester) and lactating women can receive

second-generation oral antihistamine treatment (no teratogenic effects have been described), and low concentrations of these drugs are secreted in breast milk.^{61,62}

Immunotherapy may be continued but not initiated in pregnancy.⁶³

3.3.2 | Children

Topical eye drops used for adults are also approved in children over the age of 3 years, giving the advice to the subject to close the punctum with a finger to avoid systemic absorption.




FIGURE 2 Treatment of different forms and different severities of ocular allergies based on recommendations given by the TF. AH, antihistamines; AKC, atopic keratoconjunctivitis; CS, steroids; DA, dual actions; MCS, mast cell stabilizers; PAC, perennial allergic conjunctivitis; SAC, seasonal allergic conjunctivitis; T, topical; TCS, topical corticosteroids; VKC, vernal keratoconjunctivitis

First-generation antihistamines are not indicated because of the sedative effect; second-generation antihistamines display a good long-term safety profile in the pediatric population. The use of these drugs is not licensed in children under the age of 6 months.⁶²

3.3.3 | Elderly people

Ocular allergy may persist into older age and can occasionally make its initial appearance in the elderly.^{64,65} The complex process of the aging immune system affects both the innate and the adaptive immunity, also on the ocular surface.⁶⁶ However, structural changes of the eyelid, eyelid margin, lacrimal system, conjunctiva, and the cornea more frequently induce a variety of ocular surface dysfunctions that can be broadly included under the umbrella of the *dry eye disease* (DED). Descriptions of OA in elderly are scarce. Therefore, OA treatment options in elderly patients may be limited by comorbidities and drug interactions.⁶⁴ Systemic and local corticosteroids in these patients should be limited to short-time administration because of their known side effects (diabetes, hypertension, osteoporosis, cataract, and glaucoma). Topical and systemic decongestants, and systemic antihistamines may cause dryness and should be avoided. In elderly patients, cumulative use of antihistamines and other anticholinergic drugs needs to be taken into consideration.⁶⁷

3.4 | Treatments of ocular comorbidities and complications

Ocular allergy and DED are distinct clinical entities but some overlapping features suggest a complex interaction of mechanisms involving the immune, endocrine, and nervous systems.⁶⁸ For example, mucosal hyperresponsiveness to nonspecific environmental stimuli has been described in both OA and DED. However, OA is mostly a disease of youth while DED is more common at an older age when signs and symptoms of allergy generally disappear. Artificial tears, routinely used for DED patients, may improve symptoms in all the clinical varieties of OA. Both OA and DED may show a favorable response to topical anti-inflammatory agents such as steroids and CsA.⁶⁹

Corneal epithelial erosions, shield ulcers, and plaques, frequently observed in AKC and VKC, occur as a result of mediators released from inflammatory cells and partially by the mechanical trauma from upper tarsal conjunctival giant papillae (GP).⁷⁰ Delayed epithelial healing may lead to secondary infections, corneal opacities, and amblyopia.

Giant papillae resection and cryotherapy are usually not necessary. GP excision with intraoperative application of 0.02% mitomycin-C may be helpful in preventing recurrent corneal complications in severe AKC and VKC cases.

The treatment of corneal ulcer can be based on the Cameron clinical grading of shield ulcers.^{71,72} Contact lenses and/or amniotic

membrane grafts may be a useful treatment option in the management of refractory vernal ulcers.⁷³

Allergic patients in childhood may develop *keratoconus*,⁷⁴ a progressive, noninflammatory disorder of the cornea characterized by thinning and steepening in the central or paracentral cornea causing irregular astigmatism and subsequent decrease in visual acuity. Corneal cross-linking, consisting in the topical application of a 0.1% riboflavin 5-phosphate solution to the de-epithelized corneal surface followed by exposure to UVA radiation, seems to be a safe and effective surgical option to arrest disease progression,⁷⁵ which may be very aggressive in children.⁷⁶

Visual rehabilitation in early and moderate stages consists of spectacles, contact lenses, and intracorneal ring implantation.⁷⁷

Although the clinical outcome of corneal transplantation in keratoconus with and without VKC is comparable, postoperative complications are more common in VKC.^{78,79} Atopy is a risk factor for complications after corneal grafting.

Prolonged treatment with topical steroids should be avoided since *glaucoma* can occur in all age-groups.^{24,80} Withdrawal of steroids and addition of antiglaucoma medications is effective in controlling IOP in the majority of patients.⁸⁰ Glaucoma surgery is rarely necessary.⁸¹

Limbal stem cell deficiency (LSCD) is a rare complication of longstanding VKC and AKC, contributing to severe visual impairment.⁸² It is characterized by conjunctival epithelial ingrowth on the cornea, neovascularization, ocular surface inflammation, and/or recurrent corneal epithelial defects. Fibrovascular pannus resection with amniotic membrane transplantation⁸³ or allolimbal transplantation with systemic immunosuppression has been reported in severe patients.⁸³

3.5 | Final TF recommendations

Table 7 and Figure 2.

4 | CONCLUSIONS AND UNMET NEEDS

There are effective drugs for the treatment of OA; however, there is a lack of an optimal treatment for the perennial and severe forms, especially for AKC and VKC. The safety and optimal dosing regimen of the most effective topical anti-inflammatory drugs, corticosteroids, are still a major concern but no specific randomized clinical trials have ever been performed because of the lack of marketing interest. There are no guidelines or consensus from scientific societies on how, when and duration of use of topical formulations of immunomodulators such as cyclosporine and tacrolimus.

Pharmacological and immunological research has identified new possibilities to modifying the allergic immune response. Hopefully, this progress will be applied to the eye and eventually lead to complete control of moderate-to-severe forms of OA. However, just looking in the in www.clinicaltrials.gov for "ongoing clinical trials in allergic conjunctivitis," it seems that the OA indication has not attracted investors and the pharmaceutical industry.

CONFLICTS OF INTEREST

The authors declare that they have no conflicts of interest.

AUTHOR CONTRIBUTION

AL, DS, JLF, LD, VC, and SD conceived and designed the study. DS, AL, LD, JLF, and SD analyzed and interpreted the data. AL, DS, SD, JLF, BB, DPF, PA, LD, VS, and CR wrote the article. DR, MK, and VC critically revised the article. AL, SD, JLF, LD, SD, DR, and MK made final approval of the article. DS, DPF, AL, LD, SD, and JLF collected the data. AL, DS, LD, SD, and JLF made provision of materials, patients, or resources. DS statistically expertized. AL obtained funding. DS, DPF, AL, SD, JLF, BB, PA, FM, and CR involved in the literature search. AL, DS, JLF, VC, and LD gave administrative, technical, or logistic support.

ORCID

Andrea Leonardi  <https://orcid.org/0000-0002-7246-8580>

Carmen Rondon  <https://orcid.org/0000-0003-0976-3402>

Dermot Ryan  <https://orcid.org/0000-0002-4115-7376>

Marek L. Kowalsky  <https://orcid.org/0000-0002-8442-2774>

REFERENCES

- Leonardi A, Bogacka E, Fauquert JI, et al. Ocular allergy: recognizing and diagnosing hypersensitivity disorders of the ocular surface. *Allergy*. 2012;67(11):1327-1337.
- Fauquert J-I, Jedrzejczak-Czechowicz M, Rondon C, et al. Conjunctival allergen provocation test: guidelines for daily practice. *Allergy*. 2017;72(1):43-54.
- Leonardi A, Doan S, Fauquert JI, et al. Diagnostic tools in ocular allergy. *Allergy*. 2017;72(10):1485-1498.
- Leonardi A, Piliego F, Castegnaro A, et al. Allergic conjunctivitis: a cross-sectional study. *Clin Exp Allergy*. 2015;45(6):1118-1125.
- Kam KW, Chen LJ, Wat N, Young AL. Topical olopatadine in the treatment of allergic conjunctivitis: a systematic review and meta-analysis. *Ocul Immunol Inflamm*. 2016;25:663-677.
- Castillo M, Scott NW, Mustafa MZ, Mustafa MS, Azuara-Blanco A. Topical antihistamines and mast cell stabilisers for treating seasonal and perennial allergic conjunctivitis. *Cochrane Database Syst Rev*. 2015;6:CD009566.
- Owen CG, Shah A, Henshaw K, Smeeth L, Sheikh A. Topical treatments for seasonal allergic conjunctivitis: systematic review and meta-analysis of efficacy and effectiveness. *Br J Gen Pract*. 2004;54(503):451-456.
- Rosenwasser LJ, O'Brien T, Weyne J. Mast cell stabilization and anti-histamine effects of olopatadine ophthalmic solution: a review of pre-clinical and clinical research. *Curr Med Res Opin*. 2005;21(9):1377-1387.
- Mahvan TD, Buckley WA, Hornecker JR. Alcaftadine for the prevention of itching associated with allergic conjunctivitis. *Ann Pharmacother*. 2012;46(7-8):1025-1032.
- Tagawa Y, Namba K, Nakazono Y, Iwata D, Ishida S. Evaluating the efficacy of epinastine ophthalmic solution using a conjunctivitis allergen challenge model in patients with birch pollen allergic conjunctivitis. *Allergol Int*. 2017;66(2):338-343.

11. Elieh Ali Komi D, Rambasek T, Bielory L. Clinical implications of mast cell involvement in allergic conjunctivitis. *Allergy*. 2018;73(3):528-539.
12. Mantelli F, Santos MS, Petitti T, et al. Systematic review and meta-analysis of randomised clinical trials on topical treatments for vernal keratoconjunctivitis. *Br J Ophthalmol*. 2007;91(12):1656-1661.
13. Leonardi A, Bremond-Gignac D, Bortolotti M, et al. Clinical and biological efficacy of preservative-free NAAGA eye-drops versus levocabastine eye-drops in vernal keratoconjunctivitis patients. *Br J Ophthalmol*. 2007;91(12):1662-1666.
14. Jay JL. Clinical features and diagnosis of adult atopic keratoconjunctivitis and the effect of treatment with sodium cromoglycate. *Br J Ophthalmol*. 1981;65(5):335-340.
15. Swamy BN, Chilov M, McClellan K, Petsoglou C. Topical non-steroidal anti-inflammatory drugs in allergic conjunctivitis: meta-analysis of randomized trial data. *Ophthalmic Epidemiol*. 2007;14(5):311-319.
16. Wilson DJ, Schutte SM, Abel SR. Comparing the efficacy of ophthalmic NSAIDs in common indications: a literature review to support cost-effective prescribing. *Ann Pharmacother*. 2015;49(6):727-734.
17. Gane J, Buckley R. Leukotriene receptor antagonists in allergic eye disease: a systematic review and meta-analysis. *J Allergy Clin Immunol Pract*. 2013;1(1):65-74.
18. Canonica GW, Tarantini F, Compalati E, Penagos M. Efficacy of desloratadine in the treatment of allergic rhinitis: a meta-analysis of randomized, double-blind, controlled trials. *Allergy*. 2007;62(4):359-366.
19. Compalati E, Baena-Cagnani R, Penagos M, et al. Systematic review on the efficacy of fexofenadine in seasonal allergic rhinitis: a meta-analysis of randomized, double-blind, placebo-controlled clinical trials. *Int Arch Allergy Immunol*. 2011;156(1):1-15.
20. Compalati E, Canonica GW. Efficacy and safety of rupatadine for allergic rhinoconjunctivitis: a systematic review of randomized, double-blind, placebo-controlled studies with meta-analysis. *Curr Med Res Opin*. 2013;29(11):1539-1551.
21. Ousler GW 3rd, Workman DA, Torkildsen GL. An open-label, investigator-masked, crossover study of the ocular drying effects of two antihistamines, topical epinastine and systemic loratadine, in adult volunteers with seasonal allergic conjunctivitis. *Clin Ther*. 2007;29(4):611-616.
22. Ousler GW, Wilcox KA, Gupta G, Abelson MB, Fink K. An evaluation of the ocular drying effects of 2 systemic antihistamines: loratadine and cetirizine hydrochloride. *Ann Allergy Asthma Immunol*. 2004;93(5):460-464.
23. Wu LQ, Chen X, Lou H, Cheng JW, Wei RL. Loteprednol etabonate in the treatment of allergic conjunctivitis: a meta-analysis. *Curr Med Res Opin*. 2015;31(8):1509-1518.
24. Sheppard JD, Comstock TL, Cavet ME. Impact of the topical ophthalmic corticosteroid loteprednol etabonate on intraocular pressure. *Adv Ther*. 2016;33(4):532-552.
25. Baroody FM, Shenaq D, DeTineo M, Wang J, Naclerio RM. Fluticasone furoate nasal spray reduces the nasal-ocular reflex: a mechanism for the efficacy of topical steroids in controlling allergic eye symptoms. *J Allergy Clin Immunol*. 2009;123(6):1342-1348.
26. Weiner JM, Abramson MJ, Puy RM. Intranasal corticosteroids versus oral H1 receptor antagonists in allergic rhinitis: systematic review of randomised controlled trials. *BMJ*. 1998;317(7173):1624-1629.
27. Hong J, Bielory B, Rosenberg JL, Bielory L. Efficacy of intranasal corticosteroids for the ocular symptoms of allergic rhinitis: a systematic review. *Allergy Asthma Proc*. 2011;32(1):22-35.
28. Naclerio R. Intranasal corticosteroids reduce ocular symptoms associated with allergic rhinitis. *Otolaryngol Head Neck Surg*. 2008;138(2):129-139.
29. Yanez A, Rodrigo GJ. Intranasal corticosteroids versus topical H1 receptor antagonists for the treatment of allergic rhinitis: a systematic review with meta-analysis. *Ann Allergy Asthma Immunol*. 2002;89(5):479-484.
30. Carr W, Bernstein J, Lieberman P, et al. A novel intranasal therapy of azelastine with fluticasone for the treatment of allergic rhinitis. *J Allergy Clin Immunol*. 2012;129(5):1282-1289.
31. Singh S, Pal V, Dhull CS. Supratarsal injection of corticosteroids in the treatment of refractory vernal keratoconjunctivitis. *Indian J Ophthalmol*. 2001;49(4):241-245.
32. Wan KH, Chen LJ, Rong SS, Pang CP, Young AL. Topical cyclosporine in the treatment of allergic conjunctivitis: a meta-analysis. *Ophthalmology*. 2013;120(11):2197-2203.
33. Gonzalez-Lopez JJ, Lopez-Alcalde J, Morcillo Laiz R, Fernandez Buenaga R, Rebolleda Fernandez G. Topical cyclosporine for atopic keratoconjunctivitis. *Cochrane Database Syst Rev*. 2012;9:CD009078.
34. Leonardi A, Van Setten G, Amrane M, et al. Efficacy and safety of 0.1% cyclosporine A cationic emulsion in the treatment of severe dry eye disease: a multicenter randomized trial. *Eur J Ophthalmol*. 2016;26(4):287-296.
35. Leonardi A, Doan S, Amrane M, et al. Controlled trial of cyclosporine A cationic emulsion in pediatric vernal keratoconjunctivitis: the VEKTIS study. *Ophthalmology*. 2018. <https://doi.org/10.1016/j.optha.2018.12.027>. [Epub ahead of print]
36. Zhai J, Gu J, Yuan J, Chen J. Tacrolimus in the treatment of ocular diseases. *BioDrugs*. 2011;25(2):89-103.
37. Labcharoenwongs P, Jirapongsananuruk O, Visitsunthorn N, Kosrirukvongs P, Saengin P, Vichyanond P. A double-masked comparison of 0.1% tacrolimus ointment and 2% cyclosporine eye drops in the treatment of vernal keratoconjunctivitis in children. *Asian Pac J Allergy Immunol*. 2012;30(3):177-184.
38. Pucci N, Caputo R, di Grande L, et al. Tacrolimus vs. cyclosporine eyedrops in severe cyclosporine-resistant vernal keratoconjunctivitis: a randomized, comparative, double-blind, crossover study. *Pediatr Allergy Immunol*. 2015;26(3):256-261.
39. Miyazaki D, Fukushima A, Ohashi Y, et al. Steroid-sparing effect of 0.1% tacrolimus eye drop for treatment of shield ulcer and corneal epitheliopathy in refractory allergic ocular diseases. *Ophthalmology*. 2017;124(3):287-294.
40. Nivenius E, van der Ploeg I, Jung K, Chrysanthou E, van Hage M, Montan PG. Tacrolimus ointment vs steroid ointment for eyelid dermatitis in patients with atopic keratoconjunctivitis. *Eye (Lond)*. 2007;21(7):968-975.
41. Zribi H, Descamps V, Hoang-Xuan T, Crickx B, Doan S. Dramatic improvement of atopic keratoconjunctivitis after topical treatment with tacrolimus ointment restricted to the eyelids. *J Eur Acad Dermatol Venereol*. 2009;23(4):489-490.
42. Al-Amri AM. Long-term follow-up of tacrolimus ointment for treatment of atopic keratoconjunctivitis. *Am J Ophthalmol*. 2014;157(2):280-286.
43. Hoang-Xuan T, Prisant O, Hannouche D, Robin H. Systemic cyclosporine A in severe atopic keratoconjunctivitis. *Ophthalmology*. 1997;104(8):1300-1305.
44. Roberts G, Pfaar O, Akdis Ca, et al. Guidelines on allergen immunotherapy: allergic rhinoconjunctivitis. *Allergy*. 2018;73(4):765-798.
45. Dhami S, Nurmatov U, Arasi S, et al. Allergen immunotherapy for allergic rhinoconjunctivitis: a systematic review and meta-analysis. *Allergy*. 2017;72(11):1597-1631.
46. Halken S, Agertoft L, Seidenberg J, et al. Five-grass pollen 300IR SLIT tablets: efficacy and safety in children and adolescents. *Pediatr Allergy Immunol*. 2010;21(6):970-976.
47. Di Bona D, Plaia A, Leto-Barone MS, La Piana S, Di Lorenzo G. Efficacy of grass pollen allergen sublingual immunotherapy tablets for seasonal allergic rhinoconjunctivitis: a systematic review and meta-analysis. *JAMA Intern Med*. 2015;175(8):1301-1309.

48. Nelson H, Cartier S, Allen-Ramey F, Lawton S, Calderon MA. Network meta-analysis shows commercialized subcutaneous and sublingual grass products have comparable efficacy. *J Allergy Clin Immunol Pract*. 2015;3(2):256-266.
49. Devillier P, Dreyfus JF, Demoly P, Calderon MA. A meta-analysis of sublingual allergen immunotherapy and pharmacotherapy in pollen-induced seasonal allergic rhinoconjunctivitis. *BMC Med*. 2014;12:71.
50. Lin SY, Erekesima N, Kim JM, et al. Sublingual immunotherapy for the treatment of allergic rhinoconjunctivitis and asthma: a systematic review. *JAMA*. 2013;309(12):1278-1288.
51. Calderon MA, Penagos M, Sheikh A, Canonica GW, Durham SR. Sublingual immunotherapy for allergic conjunctivitis: cochrane systematic review and meta-analysis. *Clin Exp Allergy*. 2011;41(9):1263-1272.
52. Kim Jm, Lin Sy, Suarez-Cuervo C, et al. Allergen-specific immunotherapy for pediatric asthma and rhinoconjunctivitis: a systematic review. *Pediatrics*. 2013;131(6):1155-1167.
53. Okamoto Y, Fujieda S, Okano M, Yoshida Y, Kakudo S, Masuyama K. House dust mite sublingual tablet is effective and safe in patients with allergic rhinitis. *Allergy*. 2017;72(3):435-443.
54. Ziegelmayer P, Nolte H, Nelson HS, et al. Long-term effects of a house dust mite sublingual immunotherapy tablet in an environmental exposure chamber trial. *Ann Allergy Asthma Immunol*. 2016;117(6):690-696.
55. Roder E, Berger MY, de Groot H, van Wijk RG. Immunotherapy in children and adolescents with allergic rhinoconjunctivitis: a systematic review. *Pediatr Allergy Immunol*. 2008;19(3):197-207.
56. Williams PB, Sheppard JD Jr. Omalizumab: a future innovation for treatment of severe ocular allergy? *Expert Opin Biol Ther*. 2005;5(12):1603-1609.
57. Doan S, Amat F, Gabison E, Saf S, Cochereau I, Just J. Omalizumab in severe refractory vernal keratoconjunctivitis in children: case series and review of the literature. *Ophthalmol Ther*. 2017;6(1):195-206.
58. Barnes AC, Blandford AD, Perry JD. Cicatricial ectropion in a patient treated with dupilumab. *Am J Ophthalmol Case Rep*. 2017;7:120-122.
59. Subiza J, Subiza JL, Alonso M, et al. Allergic conjunctivitis to chamomile tea. *Ann Allergy*. 1990;65(2):127-132.
60. Burrioni AG, Maio M. Ocular allergies: a psychodynamic approach. *Curr Opin Allergy Clin Immunol*. 2008;8(5):461-465.
61. Etwel F, Djokanovic N, Moretti ME, Boskovic R, Martinovic J, Koren G. The fetal safety of cetirizine: an observational cohort study and meta-analysis. *J Obstet Gynaecol*. 2014;34(5):392-399.
62. Gilbert C, Mazzotta P, Loebstein R, Koren G. Fetal safety of drugs used in the treatment of allergic rhinitis: a critical review. *Drug Saf*. 2005;28(8):707-719.
63. Pitsios C, Demoly P, Bilò Mb, et al. Clinical contraindications to allergen immunotherapy: an EAACI position paper. *Allergy*. 2015;70(8):897-909.
64. Cardona V, Guilarde M, Luengo O, Labrador-Horrillo M, Sala-Cunill A, Garriga T. Allergic diseases in the elderly. *Clin Trans Allergy*. 2011;1(1):11.
65. Ventura MT, Scichilone N, Paganelli R, et al. Allergic diseases in the elderly: biological characteristics and main immunological and non-immunological mechanisms. *Clin Mol Allergy*. 2017;15:2.
66. Mashaghi A, Hong J, Chauhan SK, Dana R. Ageing and ocular surface immunity. *Br J Ophthalmol*. 2017;101(1):1-5.
67. Gray SL, Anderson ML, Dublin S, et al. Cumulative use of strong anticholinergics and incident dementia: a prospective cohort study. *JAMA Intern Med*. 2015;175(3):401-407.
68. Bron AJ, de Paiva CS, Chauhan SK, et al. TFOS DEWS II pathophysiology report. *Ocul Surf*. 2017;15(3):438-510.
69. Lambiase A, Leonardi A, Sacchetti M, Deligianni V, Sposato S, Bonini S. Topical cyclosporine prevents seasonal recurrences of vernal keratoconjunctivitis in a randomized, double-masked, controlled 2-year study. *J Allergy Clin Immunol*. 2011;128(4):896-897.
70. Reddy JC, Basu S, Saboo US, Murthy SI, Vaddavalli PK, Sangwan VS. Management, clinical outcomes, and complications of shield ulcers in vernal keratoconjunctivitis. *Am J Ophthalmol*. 2013;155(3):550-559.
71. Cameron JA. Shield ulcers and plaques of the cornea in vernal keratoconjunctivitis. *Ophthalmology*. 1995;102(6):985-993.
72. Guo P, Kheirkhah A, Zhou WW, Qin L, Shen XL. Surgical resection and amniotic membrane transplantation for treatment of refractory giant papillae in vernal keratoconjunctivitis. *Cornea*. 2013;32(6):816-820.
73. Quah SA, Hemmerdinger C, Nicholson S, Kaye SB. Treatment of refractory vernal ulcers with large-diameter bandage contact lenses. *Eye Contact Lens*. 2006;32(5):245-247.
74. Merdler I, Hassidim A, Sorkin N, Shapira S, Gronovich Y, Korach Z. Keratoconus and allergic diseases among Israeli adolescents between 2005 and 2013. *Cornea*. 2015;34(5):525-529.
75. Li W, Wang B. Efficacy and safety of transepithelial corneal collagen crosslinking surgery versus standard corneal collagen crosslinking surgery for keratoconus: a meta-analysis of randomized controlled trials. *BMC Ophthalmol*. 2017;17(1):262.
76. Kankariya VP, Kymionis GD, Diakonis VF, Yoo SH. Management of pediatric keratoconus - evolving role of corneal collagen cross-linking: an update. *Indian J Ophthalmol*. 2013;61(8):435-440.
77. El-Khoury S, Abdelmassih Y, Hamade A, et al. Pediatric keratoconus in a tertiary referral center: incidence, presentation, risk factors, and treatment. *J Refract Surg*. 2016;32(8):534-541.
78. Egrilmez S, Sahin S, Yagci A. The effect of vernal keratoconjunctivitis on clinical outcomes of penetrating keratoplasty for keratoconus. *Can J Ophthalmol*. 2004;39(7):772-777.
79. Mahmood MA, Wagoner MD. Penetrating keratoplasty in eyes with keratoconus and vernal keratoconjunctivitis. *Cornea*. 2000;19(4):468-470.
80. Kaur S, Dhiman I, Kaushik S, Raj S, Pandav SS. Outcome of ocular steroid hypertensive response in children. *J Glaucoma*. 2016;25(4):343-347.
81. Ang M, Ho CL, Tan D, Chan C. Severe vernal keratoconjunctivitis requiring trabeculectomy with mitomycin C for corticosteroid-induced glaucoma. *Clin Exp Ophthalmol*. 2012;40(4):e149-55.
82. Saboo US, Basu S, Tiwari S, Mohamed A, Vemuganti GK, Sangwan VS. Clinical and cytologic evidence of limbal stem cell deficiency in eyes with long-standing vernal keratoconjunctivitis. *Asia Pac J Ophthalmol (Phila)*. 2013;2(2):88-93.
83. Sangwan VS, Jain V, Vemuganti GK, Murthy SI. Vernal keratoconjunctivitis with limbal stem cell deficiency. *Cornea*. 2011;30(5):491-496.
84. O'Gallagher M, Bunce C, Hingorani M, Larkin F, Tuft S, Dahlmann-Noor A. Topical treatments for blepharokeratoconjunctivitis in children. *Cochrane Database Syst Rev*. 2017;2:CD011965.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Leonardi A, Silva D, Perez Formigo D, et al. Management of ocular allergy. *Allergy*. 2019;74:1611-1630. <https://doi.org/10.1111/all.13786>

Table 1S. Classification of level of evidence and strength of recommendation accordingly GRADE guidelines[1-3].

Level of evidence	Symbol	Strength of recommendation	Symbol
High	⊕⊕⊕⊕	Strong for an intervention	↑↑
Moderate	⊕⊕⊕○	Weak for an intervention	↑?
Low	⊕⊕○○	Weak against an intervention	↓?
Very Low	⊕○○○	Strong against an intervention	↓↓

[1] J. Andrews, G. Guyatt, A.D. Oxman, P. Alderson, P. Dahm, Y. Falck-Ytter, M. Nasser, J. Meerpohl, P.N. Post, R. Kunz, J. Brozek, G. Vist, D. Rind, E.A. Akl, H.J. Schunemann, GRADE guidelines: 14. Going from evidence to recommendations: the significance and presentation of recommendations, J Clin Epidemiol 66(7) (2013) 719-25.

[2] J.C. Andrews, H.J. Schunemann, A.D. Oxman, K. Pottie, J.J. Meerpohl, P.A. Coello, D. Rind, V.M. Montori, J.P. Brito, S. Norris, M. Elbarbary, P. Post, M. Nasser, V. Shukla, R. Jaeschke, J. Brozek, B. Djulbegovic, G. Guyatt, GRADE guidelines: 15. Going from evidence to recommendation-determinants of a recommendation's direction and strength, J Clin Epidemiol 66(7) (2013) 726-35.

[3] B.J. Schünemann H, Guyatt G, Oxman A, GRADE handbook for grading quality of evidence and strength of recommendations., in: T.G.W. Group (Ed.) 2013.

Management of ocular allergy

Supplementary information

Specific Search Strategy

We searched the following databases: Medline, SCOPUS and WebScience.

A previous validated search query was used regarding PubMed's SR search strategy, using the PubMed clinical queries. The search strategies combined a term of a specific OA disease coupled with treatment options. If no papers were retrieved, the specific ocular condition coupled with the treatment searching strategy was used without filters. Reference lists of pertinent studies found during the systematic review and in three journals relevant in the field of allergy (Allergy, Journal of Allergy and Clinical Immunology, Pediatric Allergy and Immunology) were hand searched (from 2012 to 2018) to identify any relevant studies.

Selection of Studies

After excluding duplicates, all abstracts were screened by two independent researchers. Where differences in judgment occurred, the full texts were accessed. Additionally, references of included articles were screened for potential inclusion. Experts were consulted to include further guidelines and position papers reporting management of OA.

Assessment of the Quality of the Evidence

Studies were evaluated using the GRADE approach (1). The key principle of this approach was to provide transparency as well as clear and explicit criteria for assessing the quality of evidence and for grading the strength of recommendations based on risks vs benefits.

The retrieved SR and meta-analyses were further assessed for study methodological quality by the AMSTAR tool (A measurement Tool to Assess the Methodological quality of Systematic Reviews)(2-4). Reviews were grouped by treatment options and OA disorders. Studies were rated using AMSTAR2 checklist and classified as high, moderate, low or critically low based on the overall confidence in the results of the review(4). The strength of a recommendation was performed using the Evidence-To-Decision Framework (5)(supplementary Table 1). The quality of supporting evidence was assessed independently for each recommendation, taking into consideration benefits versus adverse effects, reduction of symptoms, practicability, feasibility and costs.

1. Brozek JL, Akl EA, Alonso-Coello P, Lang D, Jaeschke R, Williams JW, et al. Grading quality of evidence and strength of recommendations in clinical practice guidelines. Part 1 of 3. An overview of the GRADE approach and grading quality of evidence about interventions. *Allergy*. 2009;64(5):669-77.
2. Shea BJ, Grimshaw JM, Wells GA, Boers M, Andersson N, Hamel C, et al. Development of AMSTAR: a measurement tool to assess the methodological quality of systematic reviews. *BMC Med Res Methodol*. 2007;7:10.
3. Shea BJ, Hamel C, Wells GA, Bouter LM, Kristjansson E, Grimshaw J, et al. AMSTAR is a reliable and valid measurement tool to assess the methodological quality of systematic reviews. *J Clin Epidemiol*. 2009;62(10):1013-20.
4. Shea BJ, Reeves BC, Wells G, Thuku M, Hamel C, Moran J, et al. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ*. 2017;358:j4008.
5. Alonso-Coello P, Schunemann HJ, Moher J, Brignardello-Petersen R, Akl EA, Davoli M, et al. GRADE Evidence to Decision (EtD) frameworks: a systematic and transparent

50	approach to making well informed healthcare choices. 1: Introduction. BMJ.
51	2016;353:i2016.
52	