

DR. INGER FEMKE ASTRA BOCCA-TJEERTES (Orcid ID : 0000-0001-9184-2227)

DR. ANNICK A.J.M. VAN DE VEN (Orcid ID : 0000-0001-7032-9571)

Article type : News and Views: Legends of Allergy and Immunology

Medical Algorithm: peri-operative management of mastocytosis patients

Manuscript Acceptance Date: 25-Apr-2021

Bocca-Tjeertes IFA¹MD PhD, van de Ven AAJM² MD PhD, Koppelman GH¹ MD PhD, Sprickelman AB¹ MD PhD, Oude Elberink JNG² MD PhD.

1. Division of Pediatric Pulmonology and Allergology, Beatrix Children's Hospital, University Medical Center Groningen, The Netherlands
2. Division of Allergology and Clinical Immunology, University Medical Center Groningen, The Netherlands

Mastocytosis is a clonal disorder characterized by the proliferation and accumulation of mast cells (MCs) in various tissue types, preferentially skin and bone marrow (BM). Mastocytosis consists of cutaneous and systemic forms in both pediatric and adult patients. Both the excess and increased propensity of MCs to release mediators leads to a higher frequency and severity of immediate hypersensitivity reactions.¹⁻⁴

In the peri-operative setting, systemic mastocytosis is associated with a higher risk of severe anaphylaxis.¹⁻

⁷Here, drugs such as nonsteroidal anti-inflammatory drugs, opioids and a broad range of anesthetics, temperature change, friction/pressure to the skin, as well as stress, can cause mainly non-IgE mediated MC (mediator) release.¹⁻⁷

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/ALL.14891](https://doi.org/10.1111/ALL.14891)

This article is protected by copyright. All rights reserved

Prospective controlled studies on peri-operative hypersensitivity reactions in mastocytosis are lacking.⁸ Nevertheless, expert-based advice has been given on prohibited drugs.¹ Moreover, antimediation therapy such as anti-H1, anti-H2, and corticosteroids is commonly prescribed peri-operatively.¹⁻⁷ Albeit common practice, evidence supporting the value of premedication with antihistamines and corticosteroids is lacking, nor is there evidence of the benefit of specific drug regimens. However, there is also no evidence to the contrary.

The potentially serious adverse outcomes in the peri-operative setting cause uncertainty in both doctor and patient. Position papers have discussed the available evidence in detail, resulting in recommendations for mastocytosis patients undergoing surgery under general anesthesia.^{3,4} We provide an algorithm which serves as a tool for the practical management of both cutaneous and systemic, pediatric and adult mastocytosis patients in need of procedures requiring (local or general) anesthesia in line with these recommendations (Figure 1).^{3,4}

Preferably, surgery in patients with systemic mastocytosis requiring general anesthesia should take place in a mastocytosis reference center. The medical team should be aware of and able to counteract severe allergic reactions before, during, and after surgery. To limit uncertainties, an individual plan should be made for each patient considering previous events and including decisions on premedication, type of anesthesia, and intra- and postoperative analgesia. In patients that have no record of anesthesia or analgesia, or when in doubt, medication with low capacity to elicit MC mediator release is preferred.

Given uncertainties in premedication strategies, it may be useful to limit premedication to a risk group with previous anaphylaxis and/or extensive skin involvement in which excessive MC mediator release is most expected, undergoing general anesthesia.

Premedication can consist of antihistamines and corticosteroids. If premedication is chosen to be administered, antihistamines can be given orally or intravenously shortly (<30 minutes) before starting anesthesia. Corticosteroids may be added, especially if H1 and H2 antihistamines are not available. The choice of corticosteroids depends on the clinical setting; elective procedures allow for oral administration 12 and 2 hours before surgery, while emergency procedures depend on intravenous use. The choice of corticosteroids is preferably also made upon the desired duration of effect.

Additionally, prevention of nonspecific triggers is as much needed as any other precautions. Consider benzodiazepines for stress reduction and its probable direct beneficial regulatory effect on MCs.^{8,9}

Cutaneous MC proliferation in mastocytosis patients requires extra safety measures (Figure 1). Most pediatric and many adult patients have cutaneous lesions, of which some with blistering. Friction and pressure to the skin can cause excessive MC mediator release. Therefore, applying plasters or adhesives on lesions should be avoided and the use of adhesive remover is recommended upon (slow) removal.

During surgery, anaphylaxis may consist of hypotension, often without skin involvement, or respiratory events. In case of hypotension, consider anaphylaxis as a cause.

Following surgery, MC mediator release may still occur in the next 24 hours.⁷ Therefore, patients should restart their regular anti-mediators as soon as possible and monitored carefully.

Whereas this algorithm is based on clinical guidelines, published evidence and the authors' experience, future prospective controlled studies on peri-operative hypersensitivity reactions are needed to further improve peri-operative management of mastocytosis patients.

Figure1:

Algorithm for peri-operative management of mastocytosis patients

Patients with a suspicion for underlying mastocytosis may undergo diagnostic testing needed for diagnosing mastocytosis (e.g. bone marrow biopsy), requiring general anesthesia. This especially applies to pediatric patients. In this case, systemic mastocytosis is suspected, but not yet officially confirmed. It is however useful to create awareness of this probable underlying condition and prevent massive mediator release. This algorithm does not apply for adult patients with elevated tryptase and confirmed Hereditary Alpha Tryptasemia.

* No severe reactions are seen using local anesthetics, however be aware of mast cell mediator release by friction, pressure, and (removing) plasters.

** In case of previous anaphylaxis: always use premedication (both antihistamines and corticosteroids).

*** Analgesics:

Recommended: Sufentanil, Remifentanil, Oxytocin, Alfentanil, Acetaminophen, Fentanyl[°],

Unclear: Morphine^{°°}, NSAIDs^{°°°}

Discouraged: Codeine, Nefonam.

[°] One case of severe anaphylaxis to fentanyl in our center (pediatric patient with systemic mastocytosis).

^{°°} Titrate slowly if morphine is used.

^{°°°} Do not use NSAIDs if they were not used previously .¹

**** Only in systemic mastocytosis. The use of both antihistamines and a benzodiazepine is recommended. When antihistamines are not available, use corticosteroids. In cutaneous mastocytosis, no premedication is required.

*****For children: use dosage according to body weight.

References

1. Hermans MA, Arends NJ, et al. Management around invasive procedures in mastocytosis, an update. *Ann Allergy Asthma Immunol* 119 (2017) 304e309
2. Escribano L, Akin C, et al. Mastocytosis: Current concepts in diagnosis and treatment. *Ann Hematol* 2002; 81:677–90
3. Carter MC, Metcalfe DD, et al. Adverse reactions to drugs and biologics in patients with clonal mast cell disorders: A Work Group Report of the Mast Cells Disorder Committee, American Academy of Allergy, Asthma & Immunology. *J Allergy Clin Immunol* 2019 Mar;143(3):880-893.
4. Bonadonna P, Pagani M, et al. Drug hypersensitivity in clonal mast cell disorders: ENDA/EAACI position paper. *Allergy*. 2015 Jul;70(7):755-63.
5. Mertes PM, Malinovsky JM, et al. Reducing the risk of anaphylaxis during anesthesia: 2011 updated guidelines for clinical practice. *J Investig Allergol Clin Immunol*. 2011;21(6):442-53.
6. Brockow K, Bonadonna P. Drug allergy in mast cell disease. *Curr Opin Allergy Clin Immunol*. 2012 Aug;12(4):354-60.
7. Dewachter P, Castells MC, et al. Perioperative Management of Patients with Mastocytosis. *Anesthesiology* March 2014, Vol. 120, 753–759.
8. Matito A, Morgado JM, et al. Management of Anesthesia in Adult and Pediatric Mastocytosis: A Study of the Spanish Network on Mastocytosis (REMA) Based on 726 Anesthetic Procedures. *Int Arch Allergy Immunol*. 2015;167(1):47-56.
9. Yousefi OS, Wilhelm T, et al. The 1,4-benzodiazepine Ro5-4864 (4-chlorodiazepam) suppresses multiple pro-inflammatory mast cell effector functions. *Cell Communication and Signaling*. Volume 11, Article number: 13 (2013).

Acknowledgements

- Dr. Bocca-Tjeertes has nothing to disclose.
- Dr. van de Ven has nothing to disclose.

- Prof dr. Koppelman reports grants from Lung Foundation of the Netherlands, GSK, TEVA the Netherlands, UbboEmmius Foundation, TETRI foundation, Vertex, European Union (H2020 program), outside the submitted work; and he has participated in advisory boards to GSK and PURE-IMS (Money to institution, outside the submitted work).
- Dr. Sprikkelman reports grants from Aimmune, outside the submitted work.
- Dr. Oude Elberink has nothing to disclose.

Dr Bocca-Tjeertes is the corresponding author. E-mail: i.tjeertes@umcg.nl