

Allergy and mental health among pregnant women in the Japan Environment and Children's Study

Kiwako Yamamoto-Hanada, MD, PhD^a,
Kazue Ishitsuka, MD, MPH^a, Kyongsun Pak, BPharm^b,
Mayako Saito, MD^a, Tadayuki Ayabe, MD, PhD^a,
Hidetoshi Mezawa, MD, PhD^a, Mizuho Konishi, PhD^a,
Limin Yang, MD, PhD^a, Kenji Matsumoto, MD, PhD^a,
Hirohisa Saito, MD, PhD^a, and Yukihiro Ohya, MD, PhD^a,
for the Japan Environment and Children's Study (JECS)
Group



Clinical Implications

- Allergy increased the risk of depression and lower quality of life among pregnant women across Japan. We should provide additional support to pregnant women with allergy, to help them maintain optimal mental health.

TO THE EDITOR:

The Japan Environment and Children's Study (JECS) is a nationwide, multicenter, prospective birth cohort study conducted by the Ministry of Environment of Japan.¹ The association of mental health with allergy in pregnant women has not been well documented. The clinical question of this study was that allergic immunological features might affect mental status among pregnant women. We aimed to investigate the association of maternal mental health, including depression, anxiety, and quality of life, with allergic features during pregnancy among JECS participants.

This is a cross-sectional study of a birth cohort. Participants in the JECS were recruited between January 2011 and March 2014. Maternal information and blood samples were obtained from women during pregnancy, using self-administered questionnaires. Lifetime prevalence of allergic disease (asthma, allergic rhinitis, eczema, food allergy, conjunctivitis, drug allergy, and contact dermatitis) was assessed on the basis of a self-reported doctor's diagnosis, obtained from the questionnaire. Serum total and allergen-specific IgE titers assayed by ImmunoCAP (Thermo Fisher Scientific, Inc, Uppsala, Sweden) were analyzed at a contract clinical laboratory. Specific titers were detected for the following allergens: *Der p 1* (*Dermatophagoides pteronyssinus*), animal allergen mixes (cat dander, dog dander, guinea pig epithelium, rat, and mouse), Japanese cedar, birch, moth, and egg white. Positive IgE sensitization to a specific allergen was defined as allergen-specific IgE value of greater than or equal to 0.35 UA/mL. We defined high sensitization if the total IgE level was 170 IU/mL or above.² As for psychological assessment, Kessler's K-6 Non-Specific Psychological Distress Scale (K-6)³ was used to evaluate maternal anxiety and depression. The K6 scale consists of 6 questions. All variables were summed to yield a total score that range from 0 to 24. Scores of 13 or higher indicate severe depression,⁴ and scores of 5 or higher indicate any

depression.⁵ Health-related quality of life was measured using the Medical Outcomes Survey Short Form-8 questionnaire (SF-8).⁶ Two summary scores of a physical component summary (PCS) and a mental component summary (MCS) are indicated as SD scores (the national standard score is 50). Higher scores mean a higher quality health status.

Figure E1 in this article's Online Repository at www.jaci-inpractice.org shows the flowchart of this study. We examined the backgrounds of participants (n = 97,058) with no missing variables for the K-6 and the SF-8 and appropriate gestational age. Among these, we performed statistical analysis for the data with no missing variables (n = 86,085). Logistic regression and linear regression were used to examine the associations of K-6 and SF-8 scores (MSC and PSC scores), respectively, with allergic features, using the same potential confounders. We considered the following factors as possible confounders in the regression analyses: maternal age, place of residence, marital status, having another child, history of abnormal pregnancy, current smoking status, employment status, and maternal education level. This study was an exploratory study to estimate the magnitude of the association between maternal allergy status and mental health. Therefore, we had not defined any statistically significant level of *P* values at the beginning of the study. In the resulting tables, we show the estimates of the coefficients adjusted or nonadjusted by the confounders and the 95% CIs. We also show the *P* values of the estimates in the tables, for reference. The JECS protocol was reviewed and approved by the Ministry of Environment's Institutional Review Board on Epidemiological Studies and by the ethics committees of all participating institutions. Written informed consent was obtained from all participants.

K-6 scores of 13 or more and 5 or more were seen in 3.5% and 31.9% of participants, respectively (see Table E1); IgE sensitization was found in 73.9%. The prevalence of asthma, allergic rhinitis, eczema, food allergy, and drug allergy was 10.9%, 36.1%, 15.8%, 4.8%, and 2.6%, respectively. Among the confounding factors, marital status and smoking status may have strongly affected the K-6 and SF-8 scores (PCS) (data not shown). Table I presents the associations of K-6 with allergic features. *Der p 1* (adjusted odds ratio [aOR], 1.05; 95% CI, 1.02-1.08), animal allergen mixes (aOR, 1.05; 95% CI, 1.01-1.09), and moth (aOR, 1.05; 95% CI, 1.01-1.08) sensitizations increased the risk of any depression (K-6 scores ≥ 5). Allergic diseases increased the risk of any depression even more. The aOR for contact dermatitis (aOR, 1.55; 95% CI, 1.40-1.72) was the highest among all allergic diseases, including asthma (aOR, 1.28; 95% CI, 1.22-1.34), rhinitis (aOR, 1.20; 95% CI, 1.17-1.24), eczema (aOR, 1.17; 95% CI, 1.12-1.22), and food allergy (aOR, 1.33; 95% CI, 1.24-1.42). Severe depression (K-6 score ≥ 13) was statistically associated with allergic diseases (aOR, 1.25; 95% CI, 1.15-1.35), including asthma (aOR, 1.49; 95% CI, 1.34-1.66), rhinitis (aOR, 1.11; 95% CI, 1.02-1.20), eczema (aOR, 1.22; 95% CI, 1.10-1.35), food allergy (aOR, 1.40; 95% CI, 1.20-1.64), and contact dermatitis (aOR, 1.53; 95% CI, 1.19-1.96). For PSC and MCS scores of the SF-8, allergic diseases were negatively associated with PCS and MCS (see Table II). With regard to sensitization, only *Der p 1* sensitization showed a

TABLE I. Univariate and multivariate logistic regression results for allergic features associated with depression (K-6) in pregnant women across Japan (n = 78,705)

Outcome	Allergy features	Univariate				Multivariate			
		OR	Lower 95% CI	Upper 95% CI	P value	aOR	Lower 95% CI	Upper 95% CI	P value
K-6 (≥ 13)	None (reference)								
Severe depression									
	log_IgE, nonspecific (continuous)	1.03	1.01	1.06	.0172	1.01	0.98	1.04	.4935
	IgE (nonspecific) ≥ 170 UA/mL	1.16	1.06	1.27	.0015	1.08	0.98	1.18	.1124
	IgE_sensitization to any specific allergen	0.96	0.88	1.05	.3217	0.94	0.86	1.03	.1855
	IgE_Der p 1	1.04	0.96	1.12	.3514	1.01	0.93	1.09	.8083
	IgE_Japanese cedar	0.93	0.86	1.01	.0892	0.90	0.83	0.98	.0134
	IgE_egg white	1.57	1.14	2.16	.0059	1.53	1.11	2.11	.0010
	IgE_animal allergen mixes	1.08	0.98	1.19	.1052	1.03	0.93	1.13	.6135
	IgE_moth	1.11	1.02	1.21	.0180	1.08	0.99	1.17	.1023
	Any allergic disease	1.20	1.11	1.30	<.0001	1.25	1.15	1.35	<.0001
	Asthma	1.57	1.41	1.75	<.0001	1.49	1.34	1.66	<.0001
	Allergic rhinitis	1.05	0.97	1.14	.2025	1.11	1.02	1.20	.0132
	Eczema	1.22	1.11	1.35	<.0001	1.22	1.10	1.35	.0002
	Allergic conjunctivitis	1.18	1.05	1.34	.0073	1.26	1.12	1.43	.0002
	Food allergy	1.44	1.23	1.69	<.0001	1.40	1.20	1.64	<.0001
	Drug allergy	1.38	1.11	1.71	.0032	1.54	1.24	1.91	<.0001
	Contact dermatitis	1.35	1.06	1.73	.0163	1.53	1.19	1.96	.0008
K-6 (≥ 5)	None (reference)								
Any depression									
	log_IgE, nonspecific (continuous)	1.02	1.01	1.03	.0001	1.01	1.00	1.02	.0511
	IgE (nonspecific) ≥ 170 UA/mL	1.05	1.01	1.09	.0112	1.05	0.98	1.05	.4299
	IgE_sensitization to any specific allergen	1.02	0.98	1.05	.2901	1.01	0.97	1.05	.6236
	IgE_Der p 1	1.06	1.03	1.10	<.0001	1.05	1.02	1.08	.0014
	IgE_Japanese cedar	1.00	0.97	1.04	.7775	0.98	0.95	1.02	.2798
	IgE_egg white	1.04	0.89	1.21	.6205	1.02	0.88	1.19	.7962
	IgE_animal allergen mixes	1.08	1.04	1.12	.0001	1.05	1.01	1.09	.0212
	IgE_moth	1.06	1.02	1.09	.0012	1.05	1.01	1.08	.0093
	Any allergic diseases	1.23	1.19	1.27	<.0001	1.25	1.21	1.29	<.0001
	Asthma	1.31	1.25	1.37	<.0001	1.28	1.22	1.34	<.0001
	Allergic rhinitis	1.18	1.14	1.21	<.0001	1.20	1.17	1.24	<.0001
	Eczema	1.18	1.14	1.23	<.0001	1.17	1.12	1.22	<.0001
	Allergic conjunctivitis	1.28	1.22	1.35	<.0001	1.32	1.26	1.39	<.0001
	Food allergy	1.35	1.26	1.44	<.0001	1.33	1.24	1.42	<.0001
	Drug allergy	1.28	1.17	1.40	<.0001	1.35	1.23	1.48	<.0001
	Contact dermatitis	1.45	1.31	1.61	<.0001	1.55	1.40	1.72	<.0001

Adjustors for multivariate analysis: maternal age, place of residence, marital status, having another child, history of abnormal pregnancy, current smoking status, employment status, and maternal education level. Statistically significant results shown in bold font.

significant negative association with PCS (regression coefficient of -0.145). Drug allergy and contact dermatitis were more significantly associated with MCS than were asthma and eczema (regression coefficients of -0.799 , -0.933 , -0.513 , and -0.375 , respectively).

To our knowledge, this is the first report to evaluate the associations of mental health with allergic features among Japanese pregnant women from a nationwide population-based birth cohort study. Compared with allergic sensitization, allergic diseases seemed to be related to a higher risk of depressive symptoms and deterioration in quality of life from the multiple regression analysis. A population-based study of adults in China demonstrated that IgE sensitization was not associated with depression but that depression was associated with asthma and eczema⁷; our study findings coincide with these results. The

reason why more than 70% pregnant women had IgE sensitization to any allergens in our study is considered to be that allergy to Japanese cedar and/or house dust mites is very common among Japanese populations.⁸

A case series study of 695 patients with asthma, aged 20 to 44 years, from 29 medical centers in Japan reported a weak correlation between the severity of asthma and PCS of the SF-8 ($r = -0.144$; $P = .003$).⁹ The data from an international survey by the World Health Organization found that diagnosed depression and 30-day anxiety were associated with adult asthma in 65% and 40% of countries, respectively.¹⁰ Taken together, the association between asthma and depression or anxiety appears to exist at a global level. A systematic review and meta-analysis reported that IL-4, IL-6, and IL-10, produced by TH2 cells, were significantly higher among patients

TABLE II. Univariate and multivariate linear regression results for allergic features associated with quality of life (SF-8) in pregnant women across Japan (n = 77,639)

Outcome	Allergy features	Univariate				Multivariate			
		Coefficient	Lower 95% CI	Upper 95% CI	P value	Coefficient	Lower 95% CI	Upper 95% CI	P value
SF-8 (PCS)	None (reference)								
	log_IgE.nonspecific (continuous)	0.039	0.003	0.075	.0345	0.010	−0.026	0.046	.5812
	IgE (nonspecific) ≥170 UA/mL	0.163	0.041	0.285	.0087	0.163	−0.062	0.181	.3371
	IgE_sensitization to any specific allergen	−0.043	−0.16	0.074	.4721	−0.034	−0.151	0.083	.5653
	IgE_Der p 1	−0.099	−0.202	0.004	.0607	−0.145	−0.247	−0.042	.0056
	IgE_Japanese cedar	0.012	−0.092	0.115	.8232	0.024	−0.084	0.132	.6664
	IgE_egg	0.174	−0.341	0.689	.5076	0.184	−0.327	0.695	.4807
	IgE_animal allergen mixes	0.031	−0.096	0.158	.6333	−0.029	−0.157	0.098	.6514
	IgE_moth	−0.051	−0.165	0.064	.3875	−0.089	−0.203	0.025	.1261
	Any allergic disease	−0.904	−1.007	−0.801	<.0001	−0.840	−0.942	−0.738	<.0001
	Asthma	−0.934	−1.100	−0.769	<.0001	−1.006	−1.170	−0.841	<.0001
	Allergic rhinitis	−0.827	−0.934	−0.721	<.0001	−0.756	−0.863	−0.650	<.0001
	Eczema	−0.606	−0.747	−0.465	<.0001	−0.587	−0.727	−0.447	<.0001
	Allergic conjunctivitis	−1.105	−1.275	−0.934	<.0001	−1.025	−1.195	−0.855	<.0001
	Food allergy	−1.19	−1.429	−0.952	<.0001	−1.187	−1.424	−0.949	<.0001
	Drug allergy	−1.705	−2.025	−1.386	<.0001	−1.501	−1.819	−1.184	<.0001
	Contact dermatitis	−1.501	−1.868	−1.133	<.0001	−1.353	−1.719	−0.988	<.0001
SF-8 (MCS)	None (reference)								
	log_IgE. nonspecific (continuous)	−0.046	−0.082	−0.010	.0115	−0.022	−0.058	0.014	.2290
	IgE (nonspecific) ≥170 UA/mL	−0.113	−0.233	0.008	.0664	−0.113	−0.159	0.081	.5252
	IgE_sensitization to any specific allergen	−0.021	−0.137	0.095	.7242	0.007	−0.109	0.123	.9085
	IgE_Der p 1	−0.125	−0.227	−0.023	.0160	−0.091	−0.192	0.011	.0799
	IgE_Japanese cedar	−0.038	−0.140	0.064	.4690	0.004	−0.103	0.111	.9380
	IgE_egg	−0.32	−0.828	0.189	.2177	−0.253	−0.758	0.252	.3265
	IgE_animal allergen mixes	−0.113	−0.239	0.012	.0772	−0.027	−0.153	0.099	.6764
	IgE_moth	−0.066	−0.179	0.047	.2521	−0.029	−0.142	0.084	.6161
	Any allergic disease	−0.514	−0.615	−0.412	<.0001	−0.56	−0.661	−0.459	<.0001
	Asthma	−0.574	−0.738	−0.410	<.0001	−0.513	−0.676	−0.350	<.0001
	Allergic rhinitis	−0.424	−0.529	−0.318	<.0001	−0.489	−0.594	−0.383	<.0001
	Eczema	−0.385	−0.524	−0.246	<.0001	−0.375	−0.514	−0.237	<.0001
	Allergic conjunctivitis	−0.599	−0.768	−0.430	<.0001	−0.676	−0.844	−0.508	<.0001
	Food allergy	−0.576	−0.811	−0.340	<.0001	−0.527	−0.761	−0.292	<.0001
	Drug allergy	−0.676	−0.992	−0.360	<.0001	−0.799	−1.113	−0.484	<.0001
	Contact dermatitis	−0.783	−1.146	−0.420	<.0001	−0.933	−1.294	−0.572	<.0001

Adjustors for multivariate analysis: maternal age, place of residence, marital status, having another child, history of abnormal pregnancy, current smoking status, employment status, and maternal education level. Statistically significant results shown in bold font.

with bipolar disorder compared with healthy controls.¹¹ Immunological changes owing to allergy may serve as a mediating pathway to neuroprogression, defined as the pathological reorganization of the central nervous system along the course of a severe mental disorder, in depression, and in other psychiatric diseases.¹²

Matud¹³ reported sex differences with respect to stress events and coping styles. They found that female participants had more chronic stress and more minor daily stressors than male participants. Our study suggests that allergy in pregnant women is associated with mental disorders, such as depression, and contribute to lower quality of life. Our results highlight the importance of addressing allergy—mental health comorbidity as a public health and clinical management priority, to improve overall health conditions. We should provide additional support

to pregnant women with allergy, to help them maintain good mental health.

Acknowledgement

We thank the Edanz Group (www.edanzediting.com/ac) for editing a draft of this manuscript.

^aMedical Support Center for the Japan Environment and Children's Study, National Center for Child Health and Development, Tokyo, Japan

^bDepartment of Clinical Medicine (Biostatistics), Kitasato University School of Pharmacy, Tokyo, Japan

This study was funded and supported by the Ministry of Environment, Japan. The findings and conclusions of this article are solely the responsibility of the authors and do not represent the official views of the above government agency.

Conflicts of interest: This study was funded and supported by the Ministry of Environment, Japan. The authors declare that they have no competing interests related to the contents of this article.

Received for publication July 28, 2017; revised December 1, 2017; accepted for publication December 11, 2017.

Available online February 1, 2018.

Corresponding author: Yukihiro Ohya, MD, PhD, Division of Allergy, Department of Medical Subspecialties, Medical Support Center for Japan Environment and Children's Study (JECS), National Center for Child Health and Development, 2-10-1, Okura, Setagayaku, Tokyo 157-8535, Japan. E-mail: ohya-y@ncchd.go.jp. 2213-2198

© 2017 American Academy of Allergy, Asthma & Immunology

<https://doi.org/10.1016/j.jaip.2017.12.006>

REFERENCES

1. Kawamoto T, Nitta H, Murata K, Toda E, Tsukamoto N, Hasegawa M, et al. Rationale and study design of the Japan Environment and Children's Study (JECS). *BMC Public Health* 2014;14:1.
2. Shimazu S, Enomoto T. Serum IgE levels in healthy children. *Allergol Immunol* 1995;2:920-5.
3. Kessler RC, Andrews G, Colpe LJ, Hiripi E, Mroczek DK, Normand S-L, et al. Short screening scales to monitor population prevalences and trends in non-specific psychological distress. *Psychol Med* 2002;32:959-76.
4. Kessler RC, Barker PR, Colpe LJ, Epstein JF, Gfroerer JC, Hiripi E, et al. Screening for serious mental illness in the general population. *Arch Gen Psychiatr* 2003;60:184-9.
5. Sakurai K, Nishi A, Kondo K, Yanagida K, Kawakami N. Screening performance of K6/K10 and other screening instruments for mood and anxiety disorders in Japan. *Psychiatry Clin Neurosci* 2011;65:434-41.
6. Ware JE, Kosinski M, Dewey JE, Gandek B. How to score and interpret single-item health status measures: a manual for users of the SF-8 health survey. Lincoln, RI: QualityMetric Inc; 2001:4-19.
7. Yang YW, Tseng KC, Chen YH, Yang JY. Associations among eczema, asthma, serum immunoglobulin E and depression in adults: a population-based study. *Allergy* 2010;65:801-2.
8. Yamamoto-Hanada K, Yang L, Ishitsuka K, Ayabe T, Mezawa H, Konishi M, et al. Allergic profiles of mothers and fathers in the Japan Environment and Children's Study (JECS): a nationwide birth cohort study. *World Allergy Organ J* 2017;10:24.
9. Kimura T, Yokoyama A, Kohno N, Nakamura H, Eboshida A. Perceived stress, severity of asthma, and quality of life in young adults with asthma. *Allergol Int* 2009;58:71-9.
10. Wong KO, Hunter Rowe B, Douwes J, Senthilselvan A. Asthma and wheezing are associated with depression and anxiety in adults: an analysis from 54 countries. *Pulm Med* 2013;2013:929028.
11. Modabbernia A, Taslimi S, Brietzke E, Ashrafi M. Cytokine alterations in bipolar disorder: a meta-analysis of 30 studies. *Biol Psychiatry* 2013;74:15-25.
12. Berk M, Williams LJ, Jacka FN, O'Neil A, Pasco JA, Moylan S, et al. So depression is an inflammatory disease, but where does the inflammation come from? *BMC Med* 2013;11:200.
13. Matud MP. Gender differences in stress and coping styles. *Pers Individual Diff* 2004;37:1401-15.

APPENDIX

Members of the Japan Environment and Children's Study (JECS), as of 2017 (principal investigator, Toshihiro Kawamoto): Reiko Kishi (Hokkaido Regional Center for JECS, Hokkaido University, Sapporo, Japan); Nobuo Yaegashi (Miyagi Regional Center for JECS, Tohoku University, Sendai, Japan); Koichi Hashimoto (Fukushima Regional Center for JECS, Fukushima Medical University, Fukushima, Japan); Chisato Mori (Chiba Regional Center for JECS, Chiba University, Chiba, Japan); Shuichi Ito (Kanagawa Regional Center for JECS, Yokohama City University, Yokohama, Japan); Zentaro Yamagata (Koshin Regional Center for JECS, University of Yamagata, Chuo, Japan); Hidekuni Inadera (Toyama Regional Center

for JECS, University of Toyama, Toyama, Japan); Michihiro Kamijima (Aichi Regional Center for JECS, Nagoya City University, Nagoya, Japan); Toshio Heike (Kyoto Regional Center for JECS, Kyoto University, Kyoto, Japan); Hiroyasu Iso (Osaka Regional Center for JECS, Osaka University, Suita, Japan); Masayuki Shima (Hyogo Regional Center for JECS, Hyogo College of Medicine, Nishinomiya, Japan); Yasuaki Kawai (Tottori Regional Center for JECS, Tottori University, Yonago, Japan); Narufumi Suganuma (Kochi Regional Center for JECS, Kochi University, Nankoku, Japan); Koichi Kusuhara (Fukuoka Regional Center for JECS, Kyushu University, Fukuoka, Japan); and Takahiko Katoh (South Kyushu/Okinawa Regional Center for JECS, Kumamoto University, Kumamoto, Japan).




FIGURE E1. Flowchart of participant selection in this study.

TABLE E1. Summary of participant baseline characteristics
(n = 97,058)

Variable	n (%)	Proportion missing (%)
Age (y), mean $\pm$ SD	30.8 $\pm$ 5	0
Has another child	56,194 (57.9)	0
Current marriage	92,476 (95.7)	0.4
History of abnormal pregnancy	6,050 (6.2)	0
Current smoking	4,617 (4.8)	0.7
Employment status	56,343 (61.6)	5.8
Education completed		2.1
Junior high school	4,520 (4.7)	
High school	69,782 (71.9)	
University	20,702 (21.3)	
Place of recruitment		0
Hokkaido	7,664 (7.9)	
Fukushima/Miyagi	21,236 (21.9)	
Chiba/Kanagawa	11,721 (12.1)	
Koshin	6,949 (7.2)	
Toyama	5,271 (5.4)	
Aichi	5,406 (5.6)	
Kyoto/Osaka/Hyogo	16,141 (16.6)	
Tottori	2,929 (3)	
Kochi	6,765 (7)	
Fukuoka/Kumamoto/Miyazaki	12,121 (12.5)	
Okinawa	855 (0.9)	
K-6, min, median, max	0, 3, 24	0.1
K-6, severe depression (≥ 13)	3,375 (3.5)	0.1
K-6, any depression (≥ 5)	30,951 (31.9)	0.1
SF-8 (PCS), min, median, max	5.6, 45.9, 67.39	1.6
SF-8 (MCS), min, median, max	8.02, 46.96, 72.61	1.6
IgE (nonspecific), ≥ 170	20,260 (23.3)	10.4
log_IgE (nonspecific), percentile 25, median, percentile 75	3, 4.1, 5.1	10.4
IgE_sensitization to any specific allergen, ≥ 0.35	63, 627 (73.9)	11.3
IgE_Der p 1	40,460 (47)	11.3
IgE_Japanese cedar	47,912 (55.7)	11.3
IgE_Hen egg	884 (1)	11.3
IgE_Animal allergen mixes	17,680 (20.5)	11.3
IgE_Moth	24,077 (28)	11.3
Any allergic disease	49,547 (50)	0
Asthma	10,600 (10)	0
Allergic rhinitis	35,064 (40)	0
Eczema	15,309 (20)	0
Allergic conjunctivitis	9,672 (10)	0
Food allergy	4,685 (4.8)	0
Drug allergy	2,531 (2.6)	0
Contact dermatitis	1,874 (1.9)	0

Reproduced with permission of copyright owner. Further reproduction prohibited without permission.