

Accepted Manuscript

Skin microbiome prior to development of atopic dermatitis: early colonization with commensal staphylococci at 2 months is associated with a lower risk of atopic dermatitis at 1 year

Elizabeth A. Kennedy, BS, Jennifer Connolly, MSc, Jonathan O'B Hourihane, DM, Padraic G. Fallon, PhD, WH Irwin McLean, DSc FRS, Deirdre Murray, PhD, Jay-Hyun Jo, PhD, Julia A. Segre, PhD, Heidi H. Kong, MD, MHSc, Alan D. Irvine, MD DSc

PII: S0091-6749(16)30893-4

DOI: [10.1016/j.jaci.2016.07.029](https://doi.org/10.1016/j.jaci.2016.07.029)

Reference: YMAI 12319

To appear in: *Journal of Allergy and Clinical Immunology*


Received Date: 6 May 2016

Revised Date: 20 July 2016

Accepted Date: 27 July 2016

Please cite this article as: Kennedy EA, Connolly J, O'B Hourihane J, Fallon PG, McLean WI, Murray D, Jo J-H, Segre JA, Kong HH, Irvine AD, Skin microbiome prior to development of atopic dermatitis: early colonization with commensal staphylococci at 2 months is associated with a lower risk of atopic dermatitis at 1 year, *Journal of Allergy and Clinical Immunology* (2016), doi: 10.1016/j.jaci.2016.07.029.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



**Skin microbiome prior to development of atopic dermatitis:
early colonization with commensal staphylococci at 2 months is
associated with a lower risk of atopic dermatitis at 1 year**

Elizabeth A. Kennedy BS¹, Jennifer Connolly MSc², Jonathan O'B Hourihane DM²,
Padraic G. Fallon PhD^{3,4}, WH Irwin McLean DSc FRS⁵, Deirdre Murray PhD², Jay-
Hyun Jo, PhD¹, Julia A. Segre, PhD⁶, Heidi H. Kong MD, MHSc^{1*}, Alan D. Irvine MD
DSc^{3,4,7*}

1. Dermatology Branch, CCR, NCI, NIH
2. Paediatrics and Child Health, University College, Cork and the Irish Centre for
Fetal and Neonatal Translational (INFANT) Research, Cork University Maternity
Hospital, Ireland
3. Clinical Medicine, Trinity College Dublin
4. National Children's Research Centre, Our Lady's Children's Hospital Crumlin,
Dublin
5. Dermatology and Genetic Medicine, University of Dundee
6. Translational and Functional Genomics Branch, NHGRI, NIH
7. Paediatric Dermatology, Our Lady's Children's Hospital Crumlin, Dublin

*These authors contributed equally to this work.

Correspondence to:

Alan D. Irvine MD
National Children's Research Centre
Our Lady's Children's Hospital Crumlin, Dublin 12
IRELAND
irvinea@tcd.ie
Phone 3531 428 2532
Fax 3531-428-2651

or:

Heidi H. Kong MD, MHSc
Dermatology Branch/CCR/NCI/NIH
10 Center Dr, Bldg 10/MSB 1908
Bethesda, MD 20892-1908
USA
konghe@mail.nih.gov
Fax 1-301-496-5370
Phone 1-301-496-2481

Funding Sources

ADI is funded by the National Children's Research Centre, Dublin, Ireland. The National Children's Research Centre (NCRC) is the primary funding source for the on-going Cork BASELINE birth cohort. Additional support came from the UK Food Standards Agency (FSA). Work also supported by Intramural Research Program of the NIH National

45 Cancer Institute (EAK, J-HJ, HHK) and National Human Genome Research Institute
46 (JAS).

Abstract**Background**

Disease flares of established atopic dermatitis (AD) are generally associated with a low-diversity skin microbiota and *Staphylococcus aureus* dominance. The temporal transition of the skin microbiome between early infancy and the dysbiosis of established AD is unknown.

Methods

We randomly selected 50 children from the Cork BASELINE longitudinal birth cohort for microbiome sampling at three times in the first six months of life, at four skin sites relevant to AD: the antecubital and popliteal fossae, nasal tip, and cheek. We identified ten infants who developed AD and compared them with ten randomly selected control infants with no AD. We performed bacterial 16S ribosomal RNA sequencing and analysis directly from clinical samples.

Results

Bacterial community structures and diversity shifted over time, suggesting that age strongly affects the skin microbiome in infants. Unlike established AD, these infantile AD patients did not have noticeably dysbiotic communities prior to or with disease and were not colonized by *S. aureus*. In comparing patients and controls, infants who had affected skin at month 12 had statistically significant differences in bacterial communities on the antecubital fossa at month 2 compared to infants who were unaffected at month 12. In particular, commensal staphylococci were significantly less abundant in infants

affected at month 12, suggesting that this genus may protect against the later development of AD.

Conclusions

This study suggests that 12-month-old infants with AD were not colonized with *Staphylococcus aureus* before developing AD. Additional studies are needed to confirm if colonization with commensal staphylococci modulates skin immunity and attenuates development of AD.

Clinical Implications or Key Messages

S. aureus colonization was absent in infants with AD. Colonization by commensal staphylococcal species may protect against development of eczema.

Capsule Summary

Staphylococcus aureus colonization does not predate development of AD in infants; colonization by commensal staphylococci may protect against later AD development. Bacterial communities shifted with age during the first year of life.

Key words*Staphylococcus aureus*

Atopic dermatitis

Skin

microbiome

Longitudinal birth cohort

16S sequencing

Abbreviations

AD (Atopic Dermatitis)

BASELINE (Babies After Scope: Evaluating the Longitudinal Impact Using

Neurological and Nutritional Endpoints)

CUMH (Cork University Maternity Hospital)

INTRODUCTION

Atopic dermatitis (AD) is a common inflammatory skin condition that begins early in life. AD patients with established disease experience frequent colonization and increased infections with *Staphylococcus aureus* as well as potentially life-threatening eczema herpeticum with herpes simplex virus. The hygiene hypothesis relates the development of atopic disorders (AD, allergic rhinitis, and asthma) to reduced microbial exposure at a young age¹. Epidemiological studies examining the incidence of asthma have linked exposure to farming environments to lower rates of allergic disorders²⁻⁴. However, the potential role of microbe exposure in early childhood to the development of AD and the subsequent atopic march towards the development of allergic rhinitis and asthma remains to be elucidated.

There is significant interest in the potential effects of microbes on the development of skin immunity as well as disease⁵⁻⁹. Recent work in mice has shown that cutaneous exposure to commensal bacteria early in life can induce tolerance to these microbes⁶. Given these epidemiologic associations between environmental exposure and development of atopic diseases, we investigated the skin microbiome in a birth cohort. We analyzed bacterial 16S rRNA gene sequences from swabs collected from four skin sites in infants in a birth cohort (Babies After Scope: Evaluating the Longitudinal Impact Using Neurological and Nutritional Endpoints/BASELINE) at three different time points to determine if the differences in the skin microbiome were associated with AD development.

METHODS

Study subjects

The Cork Babies After Scope: Evaluating the Longitudinal Impact Using Neurological and Nutritional Endpoints (BASELINE) birth cohort study is the pediatric follow-on from the Cork Centre for the Screening for Pregnancy Endpoints (SCOPE) study^{10, 11}. The Cork BASELINE birth cohort study recruited within a white Irish population in Cork, Ireland from August 2009 to October 2011. These women were subject to the inclusion criteria of the SCOPE study: low-risk primigravidae mothers with singleton pregnancies, who delivered at or near term. Maternal consent was obtained at 20 weeks gestation, and verified at delivery. Ethical approval for the Cork BASELINE birth cohort Study was granted by the Clinical Research Ethics Committee of the Cork Teaching Hospitals, ref ECM 5 (9) 01/07/2008. The BASELINE study is registered with the United States National Institutes of Health Clinical Trials Registry (<http://www.clinicaltrials.gov>), ID: NCT01498965.

Clinical diagnosis of Atopic Dermatitis

All infants were assessed at birth, and at 2, 6, 12, and 24 months of age. Assessment included parental questionnaires and physical exam. Screening questions specific for atopic dermatitis were included in the questionnaires administered at 2, 6, and 12 months. AD was diagnosed (at 6, 12, and 24 months) by experienced healthcare personnel using the UK Working Party diagnostic criteria¹²⁻¹⁴. When AD was present, the SCORAD (SCORing Atopic Dermatitis) clinical tool was used to assess severity^{15, 16}. The

Nottingham Eczema Severity Score (NESS) was also used to assess AD severity at 12 months¹⁷. Demographic data and clinical details are shown in Table 1 and Table E1.

Filaggrin (FLG) genotyping

Cord blood samples were collected at birth and stored for analysis. *FLG* genotyping was carried out on all study subjects with testing for the four most common Irish/European mutations as previously described¹⁸. None of the subjects in this study were found to have *FLG* mutations.

Sampling for microbiome analysis

We randomly selected 50 infants from the birth cohort and obtained skin swabs at day 2, month 2, and month 6. Skin samples and negative controls were collected using pre-moistened swabs as previously described¹⁹. After all infants had been assessed at 1 year, ten infants with clinical AD at months 2, 6 and/or 12 were selected for analysis as ‘AD patients’. Healthy controls were ten infants without AD at any study time points selected at random. Sample sites were selected based on the presentation of AD at different ages. Cheeks (Ch) are a site of AD predilection in infants, and the nasal tip (Nt) is typically spared. Antecubital fossae (Af) and popliteal fossae (Pf) are typical sites of AD predilection in children and adolescents.

Sample processing/sequencing

16S rRNA V1-V3 sequencing was performed on swab samples as previously described¹⁹. Swabs were incubated in Yeast Cell Lysis Solution (Epicenter MasterPure kit, MPY80200) and Ready-Lyse Solution (Epicenter R1802M) for 1 hour at 37°C. Two 5-mm stainless steel beads (Qiagen) were added and processed in a TissueLyser (Qiagen) for 2 min at 30 Hz. The solution was treated with MPC Protein Precipitation Reagent (Epicenter MasturePure kit MPY80200) to remove cellular debris. Subsequent steps were performed using the PureLink Genomic DNA kit (Invitrogen). Barcoded primers flanking V1 (27F, 59-AGAGTTTGATCCTGGCTCAG-39) and V3 (534R, 59-ATTACCGCGGCTGCTGG-39) were used for PCR. PCR products were purified using the Agencourt AMPure XP Kit (A63880) and quantitated using the Quant-iT dsDNA high-sensitivity assay kit (Invitrogen, Q33120); equivalent amounts of these PCR products were pooled, purified with a Qiagen MinElute column (Qiagen, 28004) into 30 µL TE, and sequenced at the NIH Intramural Sequencing Center on a 454 GS FLX (Roche) platform. Reagents and collection procedure controls were tested and demonstrated no significant background contamination.

Data analysis

Sequences were preprocessed using mothur version 1.35.1²⁰. Briefly, 454 flowgram data were trimmed and denoised and chimera-checking was completed using the mothur implementation of UCHIME²¹. Sequences were classified using the Ribosomal Database Project naïve Bayesian classifier²². **Sequences classified as Chloroplast or Mitochondria were discarded.** Site-specific definition of operational taxonomic units (OTUs, or groups of sequences that share a specific level of similarity) and downstream

analyses was performed in mothur. Within the samples from each time point or site, pairwise distances were calculated and OTUs defined at 97% nucleotide similarity. Within-sample (Shannon diversity) and between-sample (theta index) measurement were performed based on these OTU definitions with subsampling to 1000 sequences per sample²³. Rarefaction curves level off by this value, suggesting adequate sequencing coverage; any samples with fewer than 1000 sequences after preprocessing were removed from analysis (Fig E1). Differentially abundant OTUs were detected using the metastats command in mothur.

The sequences classified to the *Staphylococcus* genus by the RDP naïve Bayesian classifier were then placed on a phylogenetic reference tree using “-keep-at-most 1000 max-pitches 1000”. Taxonomy was assigned using the guppy program in pplacer²⁴ with a likelihood cutoff set to 0.65 as previously described¹⁹.

Statistics

All data analysis was performed in R; results are presented as mean $\pm$ SEM unless otherwise indicated. The 95% confidence intervals were estimated. Post-hoc tests (i.e. pairwise comparisons in AMOVA) were adjusted using a Bonferroni correction. For detection of differentially abundant OTUs, metastats results are filtered for OTUs with a mean abundance of 0.05% or greater and p-values calculated using an FDR adjustment.

RESULTS

Site-specific bacterial colonization patterns

Since different skin microenvironments and anatomic regions harbor distinct microbial communities in adults and older children^{25, 26}, we initially compared the major bacterial taxa present on the four sites on infants. Relative abundances of these bacterial taxa showed differences between the two facial sites (Ch and Nt) and the extremity sites (Af and Pf) (Fig E2). Calculation of differentially abundant species between the site types confirmed that *Staphylococcus* spp. were relatively more abundant on extremity sites at all time points, and *Gemella* spp. were relatively more abundant on facial sites. Other taxa were only differentially represented at some time points, with facial sites in *Propionibacterium* spp. at day 2 and *Streptococcus* spp. at later time points (Table E2).

We validated these findings with biodiversity calculations, examining samples from each time point. We analyzed how similar the bacterial community structures were between the samples using the theta similarity index, which accounts for both the presence and proportion of bacterial species²³. A theta index value of 1 indicates that the two bacterial communities have identical structures; a value of 0 indicates maximally dissimilarity. In principal coordinates analyses (PCoA) based on these theta values, samples that are more similar to each other cluster more closely together. At each time point, the facial site samples clustered together (AMOVA p-value > 0.05) but distinctly separate from the extremity site samples (p-value < 0.006). The extremity sites clustered together at day 2 and month 2, but had different centroids at month 6 (p-value=<0.006) (Fig 1 and Fig E3).

Changes in bacterial colonization over time

Skin microbiomes differ between children and adults; however, studies with longitudinal skin sampling in infants are infrequent^{27, 28}. Alterations in the skin bacterial abundances at the different sampling time points were apparent in our cohort (Fig E4). For each skin site, the bacterial community structures showed striking shifts based on sampling timepoint (Fig 2AB, Fig E5). At both extremity sites, the samples clustered separately between day 2 and month 6 (AMOVA p-value = 0.024 for Af, 0.003 at Pf). For both facial sites, day two and month 6 samples clustered significantly as well ($p < 0.003$ for each), and between month 2 and month 6 ($p < 0.003$ at Ch, $p = 0.06$ at Nt).

To examine interpersonal variation, we calculated the mean similarity between samples at a single site and time point. For both facial sites, bacterial communities between subjects were most similar at month 6, converging to a more common bacterial population across subjects. Extremity sites did not present this same pattern; instead the most similar bacterial community structures were observed at month 2 (Fig 2C).

We analyzed the bacterial diversity of all samples, using the Shannon diversity index (a higher value signifies more taxonomic groups and/or a more even distribution of these groups). At each time point, diversity was similar between Af, Ch, and Nt (Wilcoxon Rank-Sum test p-value > 0.05 ; Fig 3, Table E3). Pf had a substantially altered pattern, significantly different from the other sites at all time points, except Af at day 2. At the facial sites, bacterial diversity increased significantly over the time studied (Wilcoxon Rank-Sum test p-value < 0.001 for each site between Day 2 and Month 6; Table E3).

Combined with the increasingly similar bacterial community structures on the face, this suggests that over time the microbial population converges and stabilizes at facial sites. Samples from the antecubital fossa also significantly increased in diversity between the time points ($p=0.033$).

Colonization of antecubital fossa with commensal staphylococci at month 2 is associated with decreased incidence of AD at 1 year

To identify any bacterial differences associated with AD in this cohort, we compared infants who developed AD at any time within the first year of life versus controls for each site and sampling time. At all time points, the bacterial community structures of infants who developed AD at any time within the first year of life did not cluster separately from control infants, and no significant differences in Shannon diversity were identified between the groups. Since the patients had clinical disease presenting at different time points (Table E1), we also compared samples based on whether the subjects presented with disease at each time point. Overall, there was almost no distinction between affected and unaffected samples in within- or between-sample diversity, either before or at the time point when the patients were affected (Tables E3, E4).

Interestingly, the month 2 Af samples demonstrated statistically significant clustering, grouped by those infants that went on to be affected at month 12 in this study (AMOVA p -value = 0.003). OTU-based analysis suggested that a single OTU was differentially abundant between the groups; this OTU was classified as *Staphylococcus* (Fig 4A). When considering all sequences classified to the *Staphylococcus* genus, the relative abundances

were significantly different between the two groups, with subjects that went on to be affected colonized by significantly less staphylococci (mean 0.065, 95% CI: 0.035-0.094) as compared to those that went on to be unaffected (mean 0.495, 95% CI 0.458-0.531). ($p < 0.003$ for Wilcoxon rank-sum test) (Fig 4B).

Given the specific association between *S. aureus* and AD flares, we classified the *Staphylococcus* sequences to the species level; in these samples, the most prevalent species were *S. epidermidis* and *S. cohnii* (Fig 4C). In contrast to older patients with AD²⁹, essentially no *S. aureus* sequences were present in the samples in our cohort, even at the sites and times that patients were affected (Fig E6-7). There were no statistically significant differences within individual *Staphylococcus* spp. levels in the month 2 Af samples between the later-affected and later-unaffected samples.

Birth Method and Feeding Method Have Little Effect on Skin Microbiota

Differences have previously been reported between the skin microbiota of infants born by C-section versus vaginal birth²⁸. We investigated whether birth method was associated with differences in skin microbiota in our cohort. There was no clustering of samples at any site or time point based on birth method, except Af samples at day 2 (Fig E8AB). No statistically significant differences in skin colonization based on birth method were observed at the earliest timepoint in this cohort, the second day of life. Shannon diversity was similar between the two birth methods as well (Fig E8C). Feeding method has been associated with differences in the intestinal microbiome composition of infants^{30, 31}.

However, feeding method – breast, formula, or combination – and gender did not affect skin bacterial colonization patterns in this cohort (Tables E3, 4)

Discussion

While infections with *S. aureus* and herpes simplex virus can complicate the course of established AD, the role of microbes in the etiology, genesis and pathogenesis of AD remains unclear. Recent murine studies have shown that cutaneous microbes can influence the development of skin immunity and disease^{5, 6, 9}. Determining if cutaneous microbes play a role in the initiation of AD could provide an opportunity to reduce the development of atopic disorders. To investigate the skin microbiome in infants prior to the development of AD, we used 16S rRNA gene sequencing of skin samples from a birth cohort and determined that shifts occur in the skin microbiome over the first six months of life, with site-specific bacteria communities changing in composition and diversity over time. We also identified a difference in staphylococcal colonization at a site of AD predilection that predates the presentation of disease, with patients who went on to be affected at a later date colonized by fewer *Staphylococcus* spp. Birth method and feeding method did not appear to affect skin bacterial communities at the sites and time points studied in this cohort, but other studies are needed to confirm these findings.

Prior studies of human skin have shown that skin microbial communities are site-specific^{32, 33}. While there is heterogeneity of bacterial communities across the skin surface, specific skin sites in different individuals often share common patterns of bacterial composition. This biogeography of the skin microbiome has been observed in older children and adults^{26, 34}. In previous infant skin microbiome studies, site-specific

differences were not evident in the first 3 months of life because infants were studied at a single timepoint immediately after delivery or were sampled at a single timepoint and analyzed in age cohorts of 1-3, 4-6, and 6-12 months^{27, 28}. The present study differed by sampling the same cohort of infants over a six-month interval (day 2, month 2, and month 6) and observed site-specific differences as early as the second day of life, a timepoint not previously investigated. The bacterial diversity of one skin site, the popliteal fossa, shifted at time points differentially from the three other sites studied. Since this specific skin site has not been examined in a cohort this young, the results may be related to a unique aspect of infant skin physiology and/or exposure, or specific to this cohort. Interestingly, the body site differences in bacterial communities also reflect observed site differences in immune cell density and composition from human skin³⁵⁻³⁸. Investigating site-specific differences in host-microbial interactions may enhance our understanding of the predilection of certain skin regions for dermatologic diseases.

In addition to the biogeography of the skin microbiome, skin bacterial communities can shift significantly during different periods of the life cycle, such as puberty²⁶. The physiology of infant skin changes over the first year of life with alterations in stratum corneum hydration, skin pH, and sebum production³⁹. In this study, the shifts in skin bacterial communities in the first months of life were the inverse of skin microbiome alterations that have been observed later in childhood. The increasing Shannon diversity observed in the first year of life in this infant cohort supports previous work that showed increased evenness, or similar numbers in each taxa, in bacterial communities from three skin sites in a cross-sectional study²⁷. During puberty, significant shifts in skin bacterial communities likely reflect the changes in skin

physiology and systemic hormones²⁶. The changes in the skin microbiome observed in these infants potentially reflect the influences of waning maternal hormones as well as the continued development of infant skin. For example, lipophilic *Propionibacterium* are relatively abundant on the facial sites at neonatal day 2, but decrease substantially at later time points. This corresponds to the high sebaceous activity triggered by maternal hormones in the first days of life, which wane significantly in the weeks after birth⁴⁰. These findings lead to additional questions, including whether neonatal skin disorders, e.g. cephalic pustulosis (aka neonatal acne), attributed to maternal hormones potentially may also be affected by alterations in skin bacteria.

A previous study in mice reported that developing antigen-specific tolerance to commensals depends on early colonization, suggesting that there is a ‘critical window’ for inducing regulatory T cells that prevent a later inflammatory response to these bacteria⁶. Scharschmidt et al. showed that application of a commensal species of *Staphylococcus* on neonatal skin induced these immunomodulatory effects. The relatively low abundance of pathogenic staphylococcal species on the antecubital fossa of two-month-old infants who later had AD at twelve months of age is intriguing in the context of this prior work in mice. Whether cutaneous exposure to commensal staphylococci during early infancy may have a similar effect remains unknown and further investigation is needed to understand if this may influence the development of AD. The absence of *S. aureus* at AD lesions in this cohort was somewhat surprising, given that this species is associated with AD^{29, 41-43}. A culture-based analysis of infant skin demonstrated *S. aureus* colonization in approximately 21% of AD lesions among infants in their first year of life⁴⁴. The differences may be related to inherent differences in the study populations

and/or the severity of sampled skin lesions between the study groups.

Differences in birth method have been studied in relation to the incidence of atopy and to the neonatal skin microbiome^{28, 45}. An earlier study showed skin microbiome differences based on birth method in neonates sampled a few minutes after delivery. The small sample size and rare number of Caesarean deliveries in the current study potentially contribute to the lack of statistically significant differences between the skin microbiota of infants born vaginally or by Caesarean at the earliest time point in this study, day 2 of life. While this study analyzed different sites over a longer time frame than the previous work, a larger study would be needed to address this question. Birth method may determine skin colonization very early in life; however, environmental exposures and skin physiology may predominate in shaping bacterial communities after this initial delivery. The skin barrier and *FLG* mutations are additional aspects of skin physiology that have been studied in relation to atopy. While approximately 10% of subjects in BASELINE publications and the Irish population have *FLG* mutations, the current cohort had fewer *FLG* mutations than expected due to sampling effects. Since a large proportion of patients with atopic dermatitis do not carry *FLG*-null alleles, the results in the current cohort avoid the potential effects of *FLG* mutations and remain relevant to AD. With interest in the potential immunological effects of neonatal exposures to skin microbes^{6, 46}, characterizing the early skin microbiome in neonates with and without *FLG* mutations and the timeframe for possible development of immunotolerance would be of significant clinical importance.

There are increasing efforts to understand the potential relationship between the skin microbial landscape and the development of skin immunity and human disease.

Early studies of the skin microbiome will identify possible associations between specific microbes and human health and disease but need extensive further research will be required to unravel the pathophysiology and key mechanisms involved. Longitudinal sampling of the same individuals as internal controls, and the initiation of sampling soon after birth were features of this study that improve the ability to identify distinct microbial patterns that could provide insight into the skin microbial milieu prior to the development of skin disease. As a result, we were able to define the site-specificity and the longitudinal shifts of the skin microbiome in the first six months of life, as well as the difference in relative abundances of commensal staphylococci prior to the development of AD. Additional investigations are needed to test whether site-specific differences in skin microbes influence the development of atopic dermatitis.

422 **Acknowledgements**

423 We thank Sean Conlan and Weng-Ian Ng for their underlying efforts; NISC Comparative
424 Sequencing Program for sequencing; and Mark C. Udey for discussions. This work
425 utilized the computational resources of the NIH HPC Biowulf cluster
426 (<http://hpc.nih.gov>).

427

Table 1. Demographic data for study subjects

		Healthy controls	AD patients
Female:Male		6:4	5:5
C-section:Vaginal		1:9	3:7
BF:FF:C*		2:3:5	1:2:7
Rural:Urban		5:5	4:6
Pet:No pet		5:5	3:7
Emollient use (Month 2) – Y:N		2:8	6:4
Bathing frequency (Month 2) – ≤weekly : > weekly		5:5	6:4
Antibiotic use (Month 2) – Y:N		1:9	1:9
Antibiotic use (6 months) – Y:N		5:5	3:7
TEWL	Day 2	9.668 ± 0.776	9.749 ± 0.618

	Month 2	10.402 ± 1.619	11.124 ± 2.135
	Month 6	11.412 ± 2.149	10.08 ± 1.342

431 *BF – breast-fed exclusively; FF – formula-fed exclusively; C – combination feeding

432

433

Figure legends

Fig 1: Site-specificity of bacterial community composition

All samples at day two clustered by principal coordinates analysis based on theta similarity coefficients. At day two, Af and Pf clustered together (AMOVA p-value=1), as did Ch and Nt (p=1), but each clustered distinctly from the other site pair ($p < 0.006$). P-values adjusted with Bonferonni correction (n=6)

Fig 2: Skin microbial communities on infants undergo site-specific shifts in composition with age.

- (a) Antecubital fossa samples clustered by principal coordinates analysis based on theta similarity coefficients. Using AMOVA, samples clustered distinctly between day 2 and month 6 ($p=0.024$), but not significantly between month 2 and month 6 ($p=0.12$) or between day 2 and month 2 ($p=0.21$)*
- (b) Cheek samples clustered by principal coordinates analysis based on theta similarity coefficients. Using AMOVA, samples clustered significantly between day 2 and month 6 ($p < 0.003$), between month 2 and month 6 ($p < 0.003$), but not between day 2 and month 2 ($p=0.102$)*
- (c) Mean theta similarity coefficients of comparisons within samples of each time point. Higher theta values signify greater similarity (Wilcoxon Rank-Sum *: $p < 0.05$, **: $p < 0.01$, *** $p < 0.001$).

Post-hoc p-values adjusted with Bonferonni correction (n=3)

Fig 3: Changes in bacterial biodiversity with age

Mean Shannon diversity at each time point; Shannon diversity is calculated based on richness and evenness of taxa within the community. (Wilcoxon Rank-Sum *: $p < 0.05$, **: $p < 0.01$, *** $p < 0.001$).

Fig 4: Antecubital fossa microbial community differences predate AD presentation.

- (a) Month 2 antecubital fossa samples by principal coordinates analysis of theta similarity coefficient. Samples clustered by those that went on to be affected at month 12 and those that were unaffected at month 12 (AMOVA p-value = 0.003).
- (b) Relative abundance of major taxa; subjects that went on to be affected at month 12 had significantly lower proportions of *Staphylococcus* than those that went on to be unaffected (Wilcoxon Rank-Sum p-value = 0.008)
- (c) Relative abundance of staphylococcal species. *S. aureus* was essentially absent in these communities.

REFERENCES

1. Strachan DP. Hay fever, hygiene, and household size. *Bmj* 1989; 299:1259-60.
2. Braun-Fahrlander C, Gassner M, Grize L, Neu U, Sennhauser FH, Varonier HS, et al. Prevalence of hay fever and allergic sensitization in farmer's children and their peers living in the same rural community. SCARPOL team. Swiss Study on Childhood Allergy and Respiratory Symptoms with Respect to Air Pollution. *Clin Exp Allergy* 1999; 29:28-34.
3. Riedler J, Braun-Fahrlander C, Eder W, Schreuer M, Waser M, Maisch S, et al. Exposure to farming in early life and development of asthma and allergy: a cross-sectional survey. *Lancet* 2001; 358:1129-33.
4. Riedler J, Eder W, Oberfeld G, Schreuer M. Austrian children living on a farm have less hay fever, asthma and allergic sensitization. *Clin Exp Allergy* 2000; 30:194-200.
5. Naik S, Bouladoux N, Wilhelm C, Molloy MJ, Salcedo R, Kastenmuller W, et al. Compartmentalized control of skin immunity by resident commensals. *Science* 2012; 337:1115-9.
6. Scharschmidt TC, Vasquez KS, Truong HA, Gearty SV, Pauli ML, Nosbaum A, et al. A Wave of Regulatory T Cells into Neonatal Skin Mediates Tolerance to Commensal Microbes. *Immunity* 2015; 43:1011-21.
7. Jiang X, Clark RA, Liu L, Wagers AJ, Fuhlbrigge RC, Kupper TS. Skin infection generates non-migratory memory CD8⁺ T(RM) cells providing global skin immunity. *Nature* 2012; 483:227-31.

8. Sanford JA, Gallo RL. Functions of the skin microbiota in health and disease. *Semin Immunol* 2013; 25:370-7.
9. Kobayashi T, Glatz M, Horiuchi K, Kawasaki H, Akiyama H, Kaplan DH, et al. Dysbiosis and *Staphylococcus aureus* Colonization Drives Inflammation in Atopic Dermatitis. *Immunity* 2015; 42:756-66.
10. AB. MSN. Welcome to the BASELINE study. 2008. Available at: <http://www.baselinestudy.net/>. Accessed June 2016].
11. SCOPE Consortium and MedSciNet AB. 2004. Available at: <http://www.scopestudy.net/>. Accessed June 2016.].
12. Williams HC, Burney PG, Hay RJ, Archer CB, Shipley MJ, Hunter JJ, et al. The U.K. Working Party's Diagnostic Criteria for Atopic Dermatitis. I. Derivation of a minimum set of discriminators for atopic dermatitis. *Br J Dermatol* 1994; 131:383-96.
13. Williams HC, Burney PG, Pembroke AC, Hay RJ. The U.K. Working Party's Diagnostic Criteria for Atopic Dermatitis. III. Independent hospital validation. *Br J Dermatol* 1994; 131:406-16.
14. Williams HC, Burney PG, Strachan D, Hay RJ. The U.K. Working Party's Diagnostic Criteria for Atopic Dermatitis. II. Observer variation of clinical diagnosis and signs of atopic dermatitis. *Br J Dermatol* 1994; 131:397-405.
15. Severity scoring of atopic dermatitis: the SCORAD index. Consensus Report of the European Task Force on Atopic Dermatitis. *Dermatology* 1993; 186:23-31.

- 524 16. Kunz B, Oranje AP, Labreze L, Stalder JF, Ring J, Taieb A. Clinical validation
525 and guidelines for the SCORAD index: consensus report of the European Task
526 Force on Atopic Dermatitis. *Dermatology* 1997; 195:10-9.
- 527 17. Emerson RM, Charman CR, Williams HC. The Nottingham Eczema Severity
528 Score: preliminary refinement of the Rajka and Langeland grading. *Br J*
529 *Dermatol* 2000; 142:288-97.
- 530 18. Sandilands A, Terron-Kwiatkowski A, Hull PR, O'Regan GM, Clayton TH,
531 Watson RM, et al. Comprehensive analysis of the gene encoding filaggrin
532 uncovers prevalent and rare mutations in ichthyosis vulgaris and atopic
533 eczema. *Nat Genet* 2007; 39:650-4.
- 534 19. Oh J, Freeman AF, Program NCS, Park M, Sokolic R, Candotti F, et al. The
535 altered landscape of the human skin microbiome in patients with primary
536 immunodeficiencies. *Genome Res* 2013; 23:2103-14.
- 537 20. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, et al.
538 Introducing mothur: open-source, platform-independent, community-
539 supported software for describing and comparing microbial communities.
540 *Appl Environ Microbiol* 2009; 75:7537-41.
- 541 21. Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R. UCHIME improves
542 sensitivity and speed of chimera detection. *Bioinformatics* 2011; 27:2194-
543 200.
- 544 22. Wang Q, Garrity GM, Tiedje JM, Cole JR. Naive Bayesian classifier for rapid
545 assignment of rRNA sequences into the new bacterial taxonomy. *Appl*
546 *Environ Microbiol* 2007; 73:5261-7.


- 547 23. Yue JC, Clayton MK. A Similarity Measure Based on Species Proportions.
548 Communications in Statistics - Theory and Methods 2005; 34:2123-31.
- 549 24. Matsen FA, Kodner RB, Armbrust EV. pplacer: linear time maximum-
550 likelihood and Bayesian phylogenetic placement of sequences onto a fixed
551 reference tree. BMC Bioinformatics 2010; 11:538.
- 552 25. Grice EA, Kong HH, Conlan S, Deming CB, Davis J, Young AC, et al.
553 Topographical and temporal diversity of the human skin microbiome.
554 Science 2009; 324:1190-2.
- 555 26. Oh J, Conlan S, Polley EC, Segre JA, Kong HH. Shifts in human skin and nares
556 microbiota of healthy children and adults. Genome Med 2012; 4:77.
- 557 27. Capone KA, Dowd SE, Stamatas GN, Nikolovski J. Diversity of the human skin
558 microbiome early in life. J Invest Dermatol 2011; 131:2026-32.
- 559 28. Dominguez-Bello MG, Costello EK, Contreras M, Magris M, Hidalgo G, Fierer
560 N, et al. Delivery mode shapes the acquisition and structure of the initial
561 microbiota across multiple body habitats in newborns. Proc Natl Acad Sci U S
562 A 2010; 107:11971-5.
- 563 29. Kong HH, Oh J, Deming C, Conlan S, Grice EA, Beatson MA, et al. Temporal
564 shifts in the skin microbiome associated with disease flares and treatment in
565 children with atopic dermatitis. Genome Res 2012; 22:850-9.
- 566 30. Madan JC, Hoen AG, Lundgren SN, Farzan SF, Cottingham KL, Morrison HG, et
567 al. Association of Cesarean Delivery and Formula Supplementation With the
568 Intestinal Microbiome of 6-Week-Old Infants. JAMA Pediatr 2016:1-8.



- 569 31. Jakobsson HE, Abrahamsson TR, Jenmalm MC, Harris K, Quince C, Jernberg C,
 570 et al. Decreased gut microbiota diversity, delayed Bacteroidetes colonisation
 571 and reduced Th1 responses in infants delivered by caesarean section. Gut
 572 2014; 63:559-66.
- 573 32. Costello EK, Lauber CL, Hamady M, Fierer N, Gordon JL, Knight R. Bacterial
 574 community variation in human body habitats across space and time. Science
 575 2009; 326:1694-7.
- 576 33. Grice EA, Kong HH, Conlan S, Deming CB, Davis J, Young AC, et al.
 577 Topographical and temporal diversity of the human skin microbiome.
 578 Science 2009; 324:1190-2.
- 579 34. Oh J, Byrd AL, Deming C, Conlan S, Program NCS, Kong HH, et al.
 580 Biogeography and individuality shape function in the human skin
 581 metagenome. Nature 2014; 514:59-64.
- 582 35. Omine Y, Hinata N, Yamamoto M, Kasahara M, Matsunaga S, Murakami G, et
 583 al. Regional differences in the density of Langerhans cells, CD8-positive T
 584 lymphocytes and CD68-positive macrophages: a preliminary study using
 585 elderly donated cadavers. Anat Cell Biol 2015; 48:177-87.
- 586 36. Weber A, Knop J, Maurer M. Pattern analysis of human cutaneous mast cell
 587 populations by total body surface mapping. Br J Dermatol 2003; 148:224-8.
- 588 37. Grimaldeston MA, Simpson A, Finlay-Jones JJ, Hart PH. The effect of
 589 ultraviolet radiation exposure on the prevalence of mast cells in human skin.
 590 Br J Dermatol 2003; 148:300-6.

38. Thomas JA, Biggerstaff M, Sloane JP, Easton DF. Immunological and histochemical analysis of regional variations of epidermal Langerhans cells in normal human skin. *Histochem J* 1984; 16:507-19.
39. Fluhr JW, Darlenski R, Taieb A, Hachem JP, Baudouin C, Msika P, et al. Functional skin adaptation in infancy - almost complete but not fully competent. *Exp Dermatol* 2010; 19:483-92.
40. Agache P, Blanc D, Barrand C, Laurent R. Sebum levels during the first year of life. *Br J Dermatol* 1980; 103:643-9.
41. Leyden JJ, Marples RR, Kligman AM. Staphylococcus aureus in the lesions of atopic dermatitis. *Br J Dermatol* 1974; 90:525-30.
42. Leung DY, Guttman-Yassky E. Deciphering the complexities of atopic dermatitis: shifting paradigms in treatment approaches. *J Allergy Clin Immunol* 2014; 134:769-79.
43. Bieber T. Atopic dermatitis. *N Engl J Med* 2008; 358:1483-94.
44. Skov L, Halkjaer LB, Agner T, Frimodt-Moller N, Jarlov JO, Bisgaard H. Neonatal colonization with Staphylococcus aureus is not associated with development of atopic dermatitis. *Br J Dermatol* 2009; 160:1286-91.
45. Bager P, Wohlfahrt J, Westergaard T. Caesarean delivery and risk of atopy and allergic disease: meta-analyses. *Clin Exp Allergy* 2008; 38:634-42.
46. Nagao K, Segre JA. "Bringing Up Baby" to Tolerate Germs. *Immunity* 2015; 43:842-4.



ACCEPTED MANUSCRIPT

Day Two





Mean biodiversity (Shannon)








Table E1: Metadata for each subject

Subject ID	Subject type	Presence of eczema				Mode of delivery	Method of feeding	Pet Exposure during pregnancy and 1st two months of life	2 month Urban/Rural	2 month - frequency of washing	2 month - emollient use	2 month - antibiotic use	6 month - antibiotic use
		Month 2 (affected site)	Month 6 (affected site)	Month 12 (affected site)	Month 24 (severity)								
AD1	AD patient	-	+ (Ch, Pc)	+ (skinfold at front of both ankles)	SCORAD 19 NESS mild	elective c section	Formula Fed only	No	Rural	weekly	No	No	No
AD2	AD patient	-	-	+ (back: under both shoulders)	-	normal vaginal delivery	Breast Fed (received occasional formula)	Yes (cat)	Rural	2-3 times/week	No	No	No
AD3	AD patient	-	-	+ (front of left knee)	NESS mild	normal vaginal delivery	Breast Fed (received occasional formula)	No	Urban	weekly	No	No	No
AD4	AD patient	-	+ (Ch, Pc)	+ (cheeks, forearms, Rt thigh)	NESS moderate	normal vaginal delivery	Formula Fed only	No	Rural	weekly	Yes	No	Yes (2 times)
AD5	AD patient	-	+ (Pc)	-	-	normal vaginal delivery	Breast fed for 10 days only	Yes (fish)	Urban	2-3 times/week	Yes	No	No
AD6	AD patient	-	+ (Ch)	-	-	elective c section	Breast fed for 41 days only	No	Urban	weekly	Yes	No	No
AD7*	AD patient	+ (?)	-	-	-	normal vaginal delivery	Combination feeding	No	Rural	weekly	Yes	No	Yes (1 time)
AD8	AD patient	-	+ (Ac, Pc)	-	-	elective c section	Breast fed for 13 days only	No	Urban	2-3 times/week	No	No	No
AD9*	AD patient	+ (?)	+ (Ch, Ac)	-	SCORAD 36 NESS moderate	normal vaginal delivery	Breast fed for 42 days	Yes (dog)	Urban	daily	Yes	No	No
AD10*	AD patient	+ (?)	+ (Ch)	-	NESS mild	normal vaginal delivery	Breast Fed Only	No	Urban	weekly	Yes	Yes (1 time)	Yes (same as 2 months)
C1	control				-	normal vaginal delivery	Breast Fed Only	Yes (dog)	Rural (farm)	every 10days	No	No	Yes (1 time)

C2	control				-	normal vaginal delivery	Formula Fed only	Yes (fish)	Urban	daily	No	Yes (IV)	Yes (same as 2 months)
C3	control				-	vacum delivery	Combination feeding	No	Urban	2-3 times/week	Yes	No	No
C4	control				-	normal vaginal delivery	Combination feeding	No	Rural	2-3 times/week	No	No	Yes (1 time)
C5	control				-	normal vaginal delivery	Combination feeding	No	Rural	weekly	No	No	Yes (1 time)
C6	control				-	normal vaginal delivery	Combination feeding	No	Rural	2-3 times/week	No	No	No
C7	control				no data	emergency c section	Formula Fed only	Yes (dog)	Rural	weekly	Yes	No	Yes (1 time)
C8	control				-	normal vaginal delivery	Breast Fed (received occasional formula)	Yes (dog)	Urban	3-5 times/week	No	No	No
C9	control				-	normal vaginal delivery	Formula Fed only	Yes (dog, cat, fish)	Urban	weekly	No	No	No
C10	control				-	vacum delivery	Breast Fed Only	No	Urban	weekly	No	No	No

Table E2: Metastats by site type

Time point	OTU	Variance -			Mean -	Variance -	SE -	p-value	p-value (FDR- corrected)	OTU size	OTU classification	Higher in
		Mean - Face	Face	SE - Face	Extremity	Extremity	Extremity					
Day 2	Otu0002	0.079975	0.009553	0.015454	0.2406	0.070343	0.041935	0.000999	0.00724275	65073	Staphylococcus(100)	Extremity
Day 2	Otu0003	0.247075	0.067132	0.040967	0.018725	0.003489	0.009339	0.000999	0.00724275	48068	Gemella(100)	Face
Day 2	Otu0006	0.0019	1.70E-05	0.000653	0.065875	0.026238	0.025611	0.000999	0.00724275	21437	Streptococcus(100)	Extremity
Day 2	Otu0009	0.040075	0.003209	0.008956	0.0063	0.000311	0.002786	0.000999	0.00724275	9229	Rothia(100)	Face
Day 2	Otu0005	0.097675	0.049571	0.035203	0.004125	0.000123	0.001751	0.002997	0.01655486	29524	Streptococcus(100)	Face
Day 2	Otu0013	0.00375	6.40E-05	0.001265	0.02245	0.001789	0.006689	0.003996	0.01655486	5615	Corynebacterium(100)	Extremity
Day 2	Otu0023	0.00055	2.00E-06	0.00024	0.009675	0.000807	0.004491	0.003996	0.01655486	1911	Enterococcus(100)	Extremity
Day 2	Otu0001	0.2323	0.054002	0.036743	0.109125	0.019272	0.02195	0.00999	0.03621375	69564	Propionibacterium(100)	Face
Day 2	Otu0029	0.000825	5.00E-06	0.000368	0.007675	0.000346	0.002941	0.013986	0.045066	1609	Corynebacterium(100)	Extremity
Month 2	Otu0001	0.056395	0.006899	0.013474	0.4912	0.11718	0.054125	0.000999	0.0033966	114867	Staphylococcus(100)	Extremity
Month 2	Otu0002	0.276474	0.036481	0.030984	0.05235	0.009377	0.015311	0.000999	0.0033966	49332	Streptococcus(100)	Face
Month 2	Otu0006	0.052895	0.002803	0.008589	0.00955	0.000283	0.002658	0.000999	0.0033966	9053	Gemella(100)	Face
Month 2	Otu0008	0.057289	0.006236	0.01281	0.002875	4.60E-05	0.001071	0.000999	0.0033966	7726	Veillonella(100)	Face
Month 2	Otu0010	0.033737	0.00107	0.005307	0.009825	0.000363	0.003011	0.000999	0.0033966	7201	Veillonella(100)	Face
Month 2	Otu0015	0.027974	0.002144	0.007511	0.002725	6.00E-05	0.001221	0.000999	0.0033966	5623	Prevotella(100)	Face
Month 2	Otu0018	0.035184	0.005951	0.012514	0.00085	3.00E-06	0.000283	0.000999	0.0033966	4486	Porphyromonas(100)	Face
Month 2	Otu0025	0.000632	1.00E-06	0.000157	0.011575	0.000931	0.004824	0.000999	0.0033966	1858	Paracoccus(56)	Extremity
Month 2	Otu0029	0.000632	1.00E-06	0.000143	0.0092	0.000824	0.004539	0.000999	0.0033966	1315	Kocuria(64)	Extremity
Month 2	Otu0030	0.011789	0.000339	0.002988	0.00075	4.00E-06	0.00032	0.000999	0.0033966	1307	Actinomyces(100)	Face
Month 2	Otu0007	0.058237	0.019089	0.022413	0.00175	3.30E-05	0.00091	0.001998	0.003996	7949	Prevotella(99)	Face
Month 2	Otu0009	0.032737	0.004386	0.010743	0.002525	6.50E-05	0.001274	0.001998	0.003996	7221	Simonsiella(100)	Face
Month 2	Otu0016	0.027816	0.000981	0.005081	0.006275	0.000384	0.0031	0.001998	0.003996	4977	Rothia(100)	Face
Month 2	Otu0022	5.30E-05	0	5.30E-05	0.011975	0.002619	0.008092	0.001998	0.003996	2110	Bacteroides(100)	Extremity
Month 2	Otu0027	0.011158	0.000613	0.004016	6.00E-04	3.00E-06	0.00027	0.001998	0.003996	1635	Prevotella(100)	Face
Month 2	Otu0031	0.000895	3.00E-06	0.000289	0.00765	0.000156	0.001974	0.001998	0.003996	1303	Chryseobacterium(100)	Extremity
Month 2	Otu0032	0.000605	5.00E-06	0.000359	0.006375	0.000199	0.00223	0.001998	0.003996	1088	Rhizobium(84)	Extremity
Month 2	Otu0005	0.071789	0.030372	0.028271	0.00285	8.00E-05	0.001415	0.004995	0.00893842	10639	Corynebacterium(100)	Face
Month 2	Otu0021	0.000342	1.00E-06	0.000193	0.01825	0.003418	0.009244	0.004995	0.00893842	2248	Porphyromonas(100)	Extremity
Month 2	Otu0003	0.010026	0.000632	0.004079	0.076875	0.032224	0.028383	0.005994	0.00970457	17518	Anaerococcus(100)	Extremity
Month 2	Otu0012	0.000447	4.00E-06	0.000322	0.021275	0.008443	0.014529	0.005994	0.00970457	6278	Bacteroides(100)	Extremity
Month 2	Otu0033	0.0015	9.00E-06	0.00049	0.005025	7.80E-05	0.001398	0.00999	0.01543909	1083	Corynebacterium(100)	Extremity
Month 2	Otu0023	0.015368	0.002454	0.008036	0.001575	2.20E-05	0.000744	0.017982	0.02658209	1958	Corynebacterium(100)	Face
Month 2	Otu0019	0.018395	0.000617	0.00403	0.006525	0.000374	0.003058	0.018981	0.02688975	3571	Actinomyces(100)	Face
Month 2	Otu0020	0	0	0	0.00745	0.001592	0.006309	0.01998	0.0271728	2581	Clostridium_sensu_stricto(100)	Extremity
Month 2	Otu0013	0.000816	5.00E-06	0.000365	0.01925	0.009254	0.01521	0.022977	0.03004685	5928	Propionibacterium(100)	Extremity

Month 6	Otu0001	0.007625	8.00E-05	0.00141	0.336825	0.134304	0.057945	0.000999	0.00164835	73260 Staphylococcus(100)	Extremity
Month 6	Otu0002	0.254225	0.023266	0.024118	0.071225	0.006845	0.013082	0.000999	0.00164835	59238 Streptococcus(100)	Face
Month 6	Otu0004	0.09245	0.010219	0.015983	0.007225	0.000166	0.002038	0.000999	0.00164835	16867 Prevotella(97)	Face
Month 6	Otu0006	0.070025	0.002215	0.007441	0.0072	0.000113	0.001683	0.000999	0.00164835	12845 Veillonella(100)	Face
Month 6	Otu0007	0.058475	0.001472	0.006066	0.01065	0.000199	0.00223	0.000999	0.00164835	10903 Gemella(100)	Face
Month 6	Otu0009	0.0489	0.001125	0.005304	0.0108	0.000311	0.002791	0.000999	0.00164835	9899 Rothia(100)	Face
Month 6	Otu0010	0.047225	0.002046	0.007152	0.008575	0.000563	0.003752	0.000999	0.00164835	9638 Prevotella(100)	Face
Month 6	Otu0011	0.042525	0.001943	0.00697	0.01175	0.000403	0.003175	0.000999	0.00164835	9591 Actinomyces(100)	Face
Month 6	Otu0013	0.000775	5.00E-06	0.000369	0.0422	0.012416	0.017618	0.000999	0.00164835	7696 Paracoccus(90)	Extremity
Month 6	Otu0015	0.038325	0.001231	0.005546	0.003475	1.70E-05	0.000656	0.000999	0.00164835	6530 Porphyromonas(99)	Face
Month 6	Otu0019	0.02775	0.004834	0.010993	0.001675	5.60E-05	0.001183	0.000999	0.00164835	4811 Simonsiella(100)	Face
Month 6	Otu0020	0.00185	1.00E-05	0.000508	0.02005	0.002534	0.007959	0.000999	0.00164835	4354 Chryseobacterium(100)	Extremity
Month 6	Otu0022	0.024525	0.002298	0.007579	0.00145	1.70E-05	0.000656	0.000999	0.00164835	3309 Soonwooa(100)	Face
Month 6	Otu0023	0.01385	0.000922	0.004802	0.00125	1.40E-05	0.00059	0.000999	0.00164835	2661 Corynebacterium(100)	Face
Month 6	Otu0024	0.00015	0	6.70E-05	0.011875	0.000854	0.00462	0.000999	0.00164835	2608 Bacteroides(100)	Extremity
Month 6	Otu0025	0.000925	2.00E-06	0.000194	0.011475	0.000514	0.003585	0.000999	0.00164835	2301 Kocuria(67)	Extremity
Month 6	Otu0026	0.010325	7.10E-05	0.001328	0.00155	5.00E-06	0.000358	0.000999	0.00164835	2064 Streptococcus(100)	Face
Month 6	Otu0028	4.00E-04	1.00E-06	0.000118	0.009225	0.000599	0.003868	0.000999	0.00164835	1939 Rhizobium(66)	Extremity
Month 6	Otu0030	0.008025	0.000114	0.001692	0.000975	3.00E-06	0.000254	0.000999	0.00164835	1670 Prevotella(100)	Face
Month 6	Otu0032	2.50E-05	0	2.50E-05	0.00565	0.000189	0.002172	0.000999	0.00164835	1215 Bacteroides(100)	Extremity
Month 6	Otu0014	0.000475	1.00E-06	0.000164	0.02845	0.013563	0.018414	0.001998	0.002997	6760 Anaerococcus(100)	Extremity
Month 6	Otu0027	0.00025	1.00E-06	0.000155	0.011075	0.00142	0.005958	0.001998	0.002997	2021 Blautia(100)	Extremity
Month 6	Otu0034	0.00035	1.00E-06	0.000154	0.005825	0.000367	0.003029	0.002997	0.00430004	1213 Brevundimonas(55)	Extremity
Month 6	Otu0003	0.081225	0.002872	0.008474	0.0362	0.008324	0.014426	0.006993	0.00923076	20922 Veillonella(100)	Face
Month 6	Otu0029	0.000125	0	7.30E-05	0.00895	0.001267	0.005628	0.006993	0.00923076	1726 Exiguobacterium(100)	Extremity
Month 6	Otu0018	8.00E-04	3.00E-06	0.000282	0.02695	0.012451	0.017643	0.007992	0.009768	5116 Xanthomonas(96)	Extremity
Month 6	Otu0031	0.000475	2.00E-06	0.000209	0.00605	0.000474	0.003443	0.007992	0.009768	1437 Corynebacterium(100)	Extremity
Month 6	Otu0008	1.00E-04	0	6.00E-05	0.03165	0.020618	0.022704	0.031968	0.03767657	10102 Corynebacterium(100)	Extremity

Table E3: Shannon comparisons based on timepoint, gender, birth method

Highlighted values: p-values <0.05*

Shannon - time		Shannon - gender		Shannon - patient vs. control		Shannon - affected vs. unaffected	
Comparison	p-value	Comparison	p-value	Comparison	p-value	Comparison	p-value
Af-Day 2::Af-Month 2	0.78412628	Af Day 2 female::Af Day 2 male	0.71030007	Af Day 2 Patient::Af Day 2 Control	0.43587218	Af Day 2	0.47894737
Af-Month 2::Af-Month 6	0.08969498	Pf Day 2 female::Pf Day 2 male	0.60267921	Pf Day 2 Patient::Pf Day 2 Control	0.43587218	Pf Day 2	0.30526316
Af-Day 2::Af-Month 6	0.03276825	Ch Day 2 female::Ch Day 2 male	0.20136937	Ch Day 2 Patient::Ch Day 2 Control	0.52884886	Ch Day 2	0.25789474
Pf-Day 2::Pf-Month 2	0.00048256	Nt Day 2 female::Nt Day 2 male	0.20136937	Nt Day 2 Patient::Nt Day 2 Control	0.43587218	Nt Day 2	0.30526316
Pf-Month 2::Pf-Month 6	0.3682766	Af Month 2 female::Af Month 2 male	0.88198381	Af Month 2 Patient::Af Month 2 Control	0.91179718	Af Month 2	0.11754386
Pf-Day 2::Pf-Month 6	0.00365448	Pf Month 2 female::Pf Month 2 male	0.41189569	Pf Month 2 Patient::Pf Month 2 Control	0.97051246	Pf Month 2	0.76491228
Ch-Day 2::Ch-Month 2	0.10838318	Ch Month 2 female::Ch Month 2 male	0.23698524	Ch Month 2 Patient::Ch Month 2 Control	0.43628137	Ch Month 2	0.42647059
Ch-Month 2::Ch-Month 6	0.02081299	Nt Month 2 female::Nt Month 2 male	0.06744463	Nt Month 2 Patient::Nt Month 2 Control	0.19031588	Nt Month 2	0.92105263
Ch-Day 2::Ch-Month 6	0.00016785	Af Month 6 female::Af Month 6 male	1	Af Month 6 Patient::Af Month 6 Control	0.14314014	Af Month 6	0.25789474
Nt-Day 2::Nt-Month 2	0.13272667	Pf Month 6 female::Pf Month 6 male	0.04645154	Pf Month 6 Patient::Pf Month 6 Control	0.48125095	Pf Month 6	Month 2 0.41578947
Nt-Month 2::Nt-Month 6	0.00485992	Ch Month 6 female::Ch Month 6 male	0.3702191	Ch Month 6 Patient::Ch Month 6 Control	0.63052891	Ch Month 6	Affected::Month 2 Unaffected 0.84210526
Nt-Day 2::Nt-Month 6	8.20E-05	Nt Month 6 female::Nt Month 6 male	0.3702191	Nt Month 6 Patient::Nt Month 6 Control	0.57874169	Nt Month 6	2 Unaffected 0.25789474
						Af Day 2	0.2749226
						Pf Day 2	0.53555212
						Ch Day 2	0.75730134
						Nt Day 2	0.4377967
						Af Month 2	1
						Pf Month 2	0.08111455
						Ch Month 2	0.12594268
						Nt Month 2	0.13480392
						Af Month 6	0.31137771
						Pf Month 6	Month 6 0.87729618
						Ch Month 6	Affected::Month 6 Unaffected 0.81679567
						Nt Month 6	6 Unaffected 0.4377967
						Af Day 2	0.24850361
						Pf Day 2	0.38472652
						Ch Day 2	0.4371517
						Nt Day 2	0.61671827
						Af Month 2	0.49411765
						Pf Month 2	0.55356037
						Ch Month 2	0.65441176
						Nt Month 2	0.81981424
						Af Month 6	0.49411765
						Pf Month 6	Month 12 0.81981424
						Ch Month 6	Affected::Month 12 Unaffected 0.89164087
						Nt Month 6	12 Unaffected 0.49411765
						Af Day 2	0.00980392
						Pf Day 2	0.92413632
						Ch Day 2	0.24603175
						Nt Day 2	0.50280112
						Af Month 2	0.92413632
						Pf Month 2	0.56629318
						Ch Month 2	0.82692308
						Nt Month 2	0.92413632
						Af Month 6	0.09453782
						Pf Month 6	Month 24 0.92413632
						Ch Month 6	Affected::Month 24 Unaffected 0.33590103
						Nt Month 6	24 Unaffected 1

***no p-value correction

Table E4: AMOVA comparisons for skin sites and timepoint

		AMOVA p-values (theta)						
		Month 2	Month 6	Month 12	Month 24			
		Aff::Month 2	Aff::Month 6	Aff::Month 12	Aff::Month 24			
		Patient::Control	Unaff	Unaff	Unaff	Male::Female	Vaginal::C-section	
Af	Day 2	0.93	0.313	0.471	0.575	0.979	0.458	0.002*
	Month 2	0.41	0.102	0.974	0.002*	0.418	0.379	0.909
	Month 6	0.172	0.039*	0.204	0.281	0.05	0.968	0.168
Pf	Day 2	0.997	0.647	0.251	0.969	0.868	0.252	0.094
	Month 2	0.671	0.424	0.266	0.466	0.093	0.726	0.358
	Month 6	0.361	0.526	0.441	0.377	0.8	0.044*	0.159
Nt	Day 2	0.711	0.811	0.301	0.888	0.432	0.687	0.36
	Month 2	0.544	0.99	0.554	0.533	0.797	0.127	0.283
	Month 6	0.581	0.553	0.648	0.556	0.291	0.511	0.905
Ch	Day 2	0.671	0.369	0.219	0.759	0.263	0.871	0.299
	Month 2	0.689	0.914	0.517	0.602	0.266	0.284	0.78
	Month 6	0.88	0.285	0.834	0.446	0.086	0.548	0.311

***no p-value correction

Fig E1

Rarefaction curves for sampling at each site to a cutoff of 1000 sequences; OTUs calculated at a cutoff of 97% nucleotide similarity. Each point represents mean $\pm$ SEM for all subjects at the site and sampling time indicated.

Fig E2

Relative abundance of major taxa; each bar represents a subject sampled at a single site and time point.

Fig E3

All samples month two and month six clustered by principal coordinates analysis based on theta similarity coefficients. At month two, Af and Pf had similar centroids (AMOVA p-value=0.18), as did Ch and Nt ($p=0.276$), and each still clustered distinctly from the other site pair ($p < 0.006$). At month six, Af and Pf had distinct centroids (AMOVA $p < 0.006$), but Ch and Nt clustered together ($p=1$); and the two site pairs clustered distinctly ($p < 0.006$).

Post-hoc p-values adjusted with Bonferonni correction ($n=6$)

Fig E4

Mean of relative abundance of major taxa; each bar represents the mean $\pm$ SEM of all subjects at a single site and time point.

Fig E5

- (a) Popliteal fossa samples clustered by principal coordinates analysis based on theta similarity coefficients. Using AMOVA, samples clustered significantly between day two and month six ($p = 0.003$), between day two and month six ($p = 0.042$), but not between month two and month 6 ($p=1$).
- (b) Nasal tip samples clustered by principal coordinates analysis based on theta similarity coefficients. Using AMOVA, samples clustered distinctly between day two and month two ($p < 0.003$), between day two and month six ($p < 0.003$), and between month two and month six ($p=0.06$).

Post-hoc p-values adjusted with Bonferonni correction ($n=3$)

Fig E6


Relative abundance of Staphylococcal species; each bar represents a subject sampled at a single site and time point.

Fig E7


Mean of relative abundance of Staphylococcal species; each bar represents the mean $\pm$ SEM of all subjects at a single site and time point.

Fig E8


- (a) Day 2 antecubital fossa samples clustered by principal coordinates analysis based on theta similarity coefficients. Samples clustered by birth method (AMOVA p-value = 0.005)
- (b) Day 2 cheek samples by principal coordinates analysis of theta similarity coefficient. Samples did not cluster by birth method (AMOVA p-value = 0.337).
- (c) Shannon diversity was generally similar between the birth methods, with only the popliteal fossa significantly different at day two (Wilcox rank-sum test p-value = 0.016)




Day 2
Month 2
Month 6






Month Two




Month Six





A**B**



Mean proportion

